

Received: February 27, 2022 Revised: May 2, 2022 Accepted: May 2, 2022

<https://doi.org/10.1016/j.neurom.2022.05.002>

Effects of Multisession Transcranial Direct Current Stimulation on Stress Regulation and Emotional Working Memory: A Randomized Controlled Trial in Healthy Military Personnel

Fenne M. Smits, MSc^{1,2} ; Elbert Geuze, PhD^{1,2} ; Guido J. de Kort, MSc¹; Karlijn Kouwer, MSc¹ ; Lisa Geerlings, MSc¹; Jack van Honk, PhD^{3,4,5} ; Dennis J.L.G. Schutter, PhD³ 

ABSTRACT

Objectives: Top-down stress regulation, important for military operational performance and mental health, involves emotional working memory and the dorsolateral prefrontal cortex (DLPFC). Multisession transcranial direct current stimulation (tDCS) applied over the DLPFC during working memory training has been shown to improve working memory performance. This study tested the hypothesis that combined tDCS with working memory training also improves top-down stress regulation. However, tDCS response differs between individuals. Resting-state electrophysiological brain activity was post hoc explored as a possible predictor of tDCS response. The predictive value of the ratio between slow-wave theta oscillations and fast-wave beta oscillations (theta/beta ratio) was examined, together with the previously identified tDCS response predictors age, education, and baseline working memory performance.

Materials and Methods: Healthy military service members ($n = 79$) underwent three sessions of real or sham tDCS over the right DLPFC (anode: F4, cathode: behind C2) at 2 mA for 20 minutes during emotional working memory training (N -back task). At baseline and within a week after the tDCS training sessions, stress regulation was assessed by fear-potentiated startle responses and subjective fear in a threat-of-shock paradigm with instructed emotional downregulation. Results were analyzed in generalized linear mixed-effects models.

Results: Threat-of-shock responses and emotional working memory performance showed no significant group-level effects of the real vs sham tDCS training intervention ($p > 0.07$). In contrast, when considering baseline theta/beta ratios or the other tDCS response predictors, exploratory results showed a trait-dependent beneficial effect of tDCS on emotional working memory training performance during the first session ($p < 0.01$).

Conclusions: No evidence was found for effectivity of the tDCS training intervention to improve stress regulation in healthy military personnel. The emotional working memory training results emphasize the importance of studying the effects of tDCS in relation to individual differences.

Clinical Trial Registration: This study was preregistered on September 16, 2019, at the Netherlands Trial Register (www.trialregister.nl) with ID: NL8028.

Keywords: Dorsolateral prefrontal cortex (DLPFC), emotional working memory, military, stress regulation, transcranial direct current stimulation (tDCS)

Conflict of Interest: The authors reported no conflict of interest.

Address correspondence to: Fenne M. Smits, MSc, Centraal Militair Hospitaal, Lundlaan 1, 3584 EZ, Utrecht, The Netherlands. Email: F.M.Smits-2@umcutrecht.nl

¹ Brain Research & Innovation Centre, Ministry of Defence, Utrecht, The Netherlands;

² Department of Psychiatry, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht, The Netherlands;

³ Department of Experimental Psychology, Helmholtz Institute, Utrecht University, Utrecht, The Netherlands;

⁴ Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa; and

⁵ Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa

For more information on author guidelines, an explanation of our peer review process, and conflict of interest informed consent policies, please see the journal's [Guide for Authors](#).

Source(s) of financial support: This project was supported by the Dutch Ministry of Defence. The Dutch Ministry of Defence were not involved in study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

INTRODUCTION

Military personnel risk exposure to a variety of stressors, including physical danger and witnessing severe human suffering.^{1,2} Prolonged and high levels of stress can impair operational performance³ and contribute to the development of mental health problems such as anxiety and posttraumatic stress disorder (PTSD).^{4,5} Adequate top-down regulation of stress-related reactions and emotions contributes to psychological resilience against these adverse effects of stress.^{6–10} Cognitive strategies for top-down stress regulation involve, for instance, reevaluating the value or meaning of a stressful situation to reduce its emotional impact.¹¹

However, the effectiveness of top-down stress regulation may be compromised when stress levels are too high.¹² Both acute and chronic stress levels interfere with the functioning of the prefrontal cortex (PFC), especially in the dorsolateral parts (DLPFC) that play a substantial role in stress regulation^{13,14} and cognitive functions such as working memory.¹⁵ Instead, better stress regulation has been associated with better working memory performance, specifically in the emotional domain.^{16–19} Emotional working memory plays a role by actively keeping threat-related information available and allowing for the selection and updating of this information to deploy effective stress regulation strategies.²⁰ Accordingly, the right DLPFC has been identified as a target region for noninvasive brain stimulation to improve symptoms of stress-related disorders, including PTSD.²¹ In addition, in healthy volunteers, several studies demonstrated that stress regulation improved after the application of transcranial direct current stimulation (tDCS) over the right DLPFC or the neighboring ventrolateral PFC.^{22–27}

Transcranial direct current stimulation modulates subthreshold cortical excitability and plasticity by polarizing nerve tissue using low-intensity electrical currents (1–2.5 mA) administered over the scalp. Cortical excitability is generally assumed to increase by anodal tDCS and decrease by cathodal tDCS, although the exact mechanisms underlying tDCS effects are still unclear (further reading^{28,29}). Single-session tDCS, however, yields transient neurophysiological effects that typically fade out within a few hours,³⁰ and single-session tDCS over the DLPFC does not always effectively modulate stress regulation.^{31–33}

More consistent and sustained effects on higher-order cognitive functions and on symptoms of depression and PTSD are suggested to follow from multiple sessions of tDCS over the DLPFC or the ventrolateral PFC, particularly when applied during a neuro-cognitive training or therapy that activates the tDCS-targeted brain region.^{34–41} One way to activate the right DLPFC is by a working memory task that is shown to activate several PFC regions involved in stress regulation.^{18,42} The idea that working memory performance can be improved by anodal tDCS over the DLPFC is supported by converging evidence from a recent meta-analysis.⁴³ In addition, several studies suggest that multisession tDCS during working memory training may lead to long-lasting performance gains in working memory and other cognitive functions depending on working memory,^{44–47} indicating a potential benefit for stress regulation capacity.

However, not all studies of combined tDCS with working memory training find these beneficial effects.^{48,49} These negative findings could be related to the considerable variability between individuals in tDCS effects on cognitive functions such as working memory.^{37,50} This variability has been associated with factors such as age,^{51–53} baseline cognitive performance,^{54–58} and education.⁵⁹

Variability in tDCS response may be even better explained by neural processes that interact with the effects of tDCS. Such processes are, for instance, indicated by markers from electroencephalography (EEG).^{60–62} One EEG marker that could be related to tDCS effects on emotional working memory is the power ratio between slow-wave theta band activity (4–7 Hz) and fast-wave beta band activity (13–30 Hz) in resting-state EEG, that is, the theta/beta ratio.^{63–65} The theta/beta ratio is thought to reflect the balance between subcortical-based emotional and motivational drives and cortical-based cognitive control, based on associations between the theta/beta ratio and cognitive control over emotional input,^{66–68} reward-motivated learning on cognitive tasks,^{69–71} and working memory training gains.⁷² In addition, the theta/beta ratio has been associated with effects on cognitive performance of a tDCS-related technique, transcranial random noise stimulation.^{73,74}

Following these lines of evidence, the primary objective of this study was to test whether multisession anodal tDCS of the right DLPFC during emotional working memory (WM) training could improve top-down stress regulation in healthy military personnel. The secondary objective was to explore the predictive value of the theta/beta ratio on interindividual variability in tDCS effects, in addition to the previously identified tDCS response predictors age, baseline performance, and education.

MATERIALS AND METHODS

Participants

Active-duty military personnel (aged 18–60 years, uncorrected normal hearing) were recruited between January 2020 and April 2021. This study was part of a double-blind randomized controlled trial that was preregistered at The Netherlands Trial Register (www.trialregister.nl) with ID: NL8028. The a priori-computed required sample size to detect tDCS effects on stress regulation was 62. [Supplementary Data section 1.1](#) describes how the required sample size was computed. Exclusion criteria were alcohol or drug dependence, psychoactive medication or drug use within the past two weeks, (history of) a psychiatric or neurologic disorder (except for attention-deficit/hyperactivity disorder) or serious head trauma, large or ferromagnetic metal parts in the head, implanted cardiac pacemaker or neurostimulator, pregnancy, neurostimulation in the past month, or skin damage or diseases at intended electrode sites. All participants provided written informed consent and received €65 for participation. The authors assert that all procedures contributing to this work complied with the ethical standards of the relevant national and institutional committees on human experimentation and with the Declaration of Helsinki of 1975, as revised in 2008. The medical ethical committee of the University Medical Center Utrecht approved the study. [Table 1](#) shows demographics and baseline psychological characteristics for each tDCS group.

Noninvasive Brain Stimulation

Transcranial direct current stimulation was administered at an intensity of 2.0 mA (impedance < 10 k Ω) for a duration of 20 minutes with a DC-stimulator Plus (NeuroConn GmbH, Ilmenau, Germany). Anodal tDCS was concentrated on an area in the right DLPFC that has been shown to be activated by both WM performance and top-down emotion regulation.^{13,18,75} The electrode montage to target this area was selected based on simulations of the electric field distribution in SimNIBS 3.2.3⁷⁶ ([Figure 1b](#)). A 3 × 3 cm saline-soaked sponge-covered anode was placed over EEG

Table 1. Participant Characteristics (Frequency or Mean \pm SD).

Variable name	Real tDCS	Sham tDCS
Sex		
Male	35	33
Female	2	2
Age (years)	34.0 \pm 10.7	36.1 \pm 11.1
Educational level*		
High school Diploma	2	3
Vocational degree	20	21
Associate's degree	2	0
Bachelor's degree	6	4
Master's degree	7	7
Number of deployments		
Never deployed	16	11
1 deployment	6	8
2–3 deployments	8	9
\geq 4 deployments	6	7
Rank		
Officer	8	6
Student-officer	3	3
Senior NCO	17	20
Junior NCO	9	6
Handedness†		
Right-handed	33	30
Left-handed	4	4
No preference	0	1
ASI-3 (rating 0–4)		
Physical	0.3 \pm 0.4	0.3 \pm 0.4
Cognitive	0.2 \pm 0.3	0.2 \pm 0.3
Social	0.8 \pm 0.5	0.9 \pm 0.7
ACS (rating 1–4)		
Focusing	2.7 \pm 0.4	2.6 \pm 0.5
Shifting	2.8 \pm 0.3	2.8 \pm 0.4
ERQ (rating 1–7)		
Reappraisal	4.7 \pm 0.9	4.8 \pm 0.8
Suppression	3.6 \pm 1.0	3.7 \pm 1.0
PANAS (rating 1–5)		
PA	3.7 \pm 0.5	3.7 \pm 0.4
NA	1.5 \pm 0.4	1.5 \pm 0.5
Shock intensity		
Current (mA)	8.2 \pm 3.5, range: 3–30	8.3 \pm 4.6, range: 2–39
Duration (ms)	754 \pm 452, range: 200–2000	644 \pm 424, range: 200–2000
Fear of shock (rating 0–10)	2.7 \pm 1.6	3.2 \pm 1.8
Pain of shock (rating 0–10)	1.7 \pm 0.9	2.8 \pm 1.4
Start performance <i>N</i> -back task (block 1, session 1)		
d' ($z_{\text{hits}} - z_{\text{false alarms}}$)	1.35 \pm 1.0	1.29 \pm 1.0

ACS, attentional control scale; ASI-3, 18-item Anxiety Sensitivity Inventory; ERQ, emotion regulation questionnaire; NA, negative affect; NCO, noncommissioned officer; PA, positive affect; PANAS, Positive and Negative Affect Schedule.

*Educational levels were assessed based on the Dutch educational system and for international interpretability converted to the best corresponding American degree.

†Participants were asked to identify themselves as left-handed, right-handed, or no preference.

position F4, and a 5 \times 7 cm cathode was placed dorsal of C2 (Fig. 1b). Sham tDCS involved a 16-second fade-in fade-out stimulation at the start and end of the stimulation period, interleaved by 15-millisecond pulses of 0.11 mA. Changes in emotional state

(STAI-6⁷⁹) and possible tDCS adverse effects⁸⁰ (scored from 1, “absent,” to 4, “severe”) were assessed in each session.

Emotional Working Memory Training

During tDCS, participants performed an emotional WM task based on the visuospatial/auditory *N*-back task from Schweizer et al.¹⁸ In each trial, participants indicated whether the location of an angry face in a 4 \times 4 grid on a computer screen or a one-syllable negative word (eg, “death,” “fear,” “hate”) matched with *N* trials before (Fig. 1b). Based on response accuracy, *N* increased or decreased by one in the consecutive block. The task contained ten blocks of 20 + *N* trials per block, with six target trials. To further increase emotional arousal during the task, unpredictable aversive screams (~80 dB, ~1 second) and negative fictitious performance feedback were presented during six of the blocks,⁸¹ as illustrated in Figure 1b and further described in Supplementary Data section 1.2.

At postassessment, *N*-back task performance was tested on four prespecified WM load levels (*N* = 1–4).

Primary Outcome Measure: Threat-of-Shock Paradigm

Stress-related responses were assessed by a threat-of-shock test with three conditions: “no-shock” (N), “predictable-shock” (P) and “unpredictable-shock” (U), abbreviated as the NPU-threat test.⁸² The test contained two seven-minute sequences. Per sequence, three no-shock blocks were alternated by two predictable- and two unpredictable-shock blocks. In a workup procedure, electrical shocks were tuned to an intensity rated by the participant as 4 on a 5-point Likert scale (1: “I feel no shock,” 5: “the shock feels very uncomfortable but not painful”). During each 60-second block, three cues were presented (4 seconds), interleaved by variable intertrial-intervals (ITI) (3–30 seconds). One or two shocks were delivered per threat block at a computer-randomized moment during the last second of cue presentation (predictable shock) or during the ITI (unpredictable shock). In total, six shocks were delivered per threat condition. Supplementary Data section 1.3.1. describes further test details.

Physiological threat responses were assessed by the eyeblink fear-potentiated startle reflex.⁸³ Eyeblink startles were recorded by electromyography (EMG) of two active 4-mm flat surface Ag/AgCl electrodes (BioSemi BV, Amsterdam, The Netherlands) filled with conductive gel, placed ~1 cm apart on the left orbicularis oculi muscle.⁸⁴ A startle probe of 50 milliseconds of white noise at ~100 dB was delivered through 3A insert earphones (E-A-RTONE™, 3M™) at a computer-randomized moment during the first three seconds of each cue, and during each ITI. EMG data were preprocessed (Supplementary Data section 1.3.3.), and startle amplitudes were quantified as the maximum amplitude between 20 and 120 milliseconds after probe onset (baseline-corrected and within-subject standardized).⁸⁴ Subjective fear for each condition and context (cue, ITI) was self-reported after every sequence on a visual analogue scale from 0: “I did not feel nervous or anxious at all” to 100: “I felt very nervous or anxious.”

In line with previous research of top-down stress regulation,^{8,12} psychoeducation was provided before the NPU-threat test. Stress regulation strategies were explained by instructing participants to view a situation with a “detached, objective, impartial and scientific mindset” or think of more positive aspects of the situation,¹¹ as described in more detail in Supplementary Data section 1.3.2. Participants were instructed to use these strategies to downregulate threat-related emotional responses during the NPU-threat test.




Figure 1. a. Overview of the study design. b. Overview of a tDCS training session. The tDCS montage and example of electrical field distribution are simulated in SimNIBS 3.2.3⁷⁶ with twenty brains obtained from a publicly available magnetic resonance imaging data set of neurologically healthy individuals.⁷⁷ The figure displays the average electrical field distribution across these 20 simulations. The example of an *N*-back trial sequence represents a WM load of *N* = 2. Angry face stimuli are derived from the Chicago face data base.⁷⁸ [Color figure can be viewed at www.neuromodulationjournal.org]

Questionnaires

Top-down stress regulation tendencies, anxiety sensitivity, emotional symptoms, and cognitive control of attention were assessed by self-report on the Dutch translation of the Emotion Regulation Questionnaire,^{85,86} the 18-item Anxiety Sensitivity Inventory (ASI-3),^{87,88} the Positive and Negative Affect Schedule,⁸⁹ and the Attentional Control Scale (item 12 about attention during “lectures” was adapted to “lessons”).⁹⁰

Exploratory Measures

Educational level and age were recorded during the baseline visit. Baseline WM performance was defined as a Start Performance score based on the sensitivity ($d' = Z_{\text{hits}} - Z_{\text{false alarms}}$) at the start of

the training (block 1, session 1, where all participants performed a 1-back task). Start performance scores did not significantly differ between groups (mean \pm SD: real tDCS 1.35 ± 1.0 , sham tDCS 1.29 ± 1.0 , $t_{70} = 0.26$, $p = 0.799$).

The theta/beta ratio was extracted from 4-minute resting-state EEG (alternating 1-minute eyes open, 1-minute eyes closed), recorded at the start of the baseline and postassessment visits (experimenters left the room). EEG data were recorded and amplified with a BioSemi ActiveTwo system (BioSemi BV, Amsterdam, The Netherlands) at 2048 Hz, relative to a Common Mode Sense active electrode in combination with a Driven Right Leg passive electrode, from channels Fp1, Fp2, AF3, AF4, F7, F8, F3, F4, Fz, Cz, FC1, FC2, FC5, FC6, C3, C4, CP1, CP2, CP5, CP6, P7, P8, P3, P4,

Pz, PO3, PO4, O1, O2, and Oz. Offline preprocessing was done with custom MATLAB scripts, EEGLAB v2021.0⁹¹ and ERPLAB v8.10.⁹² Continuous data were segmented in 1-second epochs. Eye blinks were identified and removed based on the EEGLAB ICA function. Epochs containing artifacts caused by movement or facial muscle contractions were automatically marked (> 30 mV difference between adjacent samples, > 100-mV difference per 200-millisecond signal, or absolute amplitude larger than ± 75 mV) and deleted after visual inspection. A fast Fourier transform was applied per epoch using Welch's method (Hanning taper, 50% overlap, 0.25-Hz resolution). The power spectral density (mV^2/Hz) was averaged over epochs and log-transformed. Following previous research, the frontocentral theta/beta ratio was calculated as the average theta power (4–7 Hz) divided by the average beta power (13–30 Hz) from channels Fz and Cz (data collapsed across eyes-open and eyes-closed conditions).^{66,69} Theta/beta ratios tended to be higher in the real tDCS group than in the sham tDCS group (mean \pm SD: real tDCS 7.7 ± 3.1 , sham tDCS 6.5 ± 2.4 , $t_{70} = 1.82$, $p = 0.074$).

Procedure

Participants were recruited, and study visits were carried out at several military bases in The Netherlands. After provision of study information, screening, and obtaining of informed consent, eligible individuals were randomly allocated to real or sham tDCS (1:1) by selecting the next-available stimulator-activating code from a list. This list contained 20 codes for real tDCS and 20 codes for sham tDCS, which were randomized with the MATLAB function "randsample." Experimenters and participants were blind for code-to-condition correspondence. Baseline and postassessment visits took place one to six days before and after the tDCS training intervention and included a resting-state EEG recording and the NPU-threat test. Self-report questionnaires were completed online. Three tDCS training sessions took place one to six days apart during working hours (between 6 AM and 9 PM, depending on working shift). Because it is not yet clear whether online tDCS or offline tDCS is most effective to modulate cognitive performance,^{93–95} tDCS (20 minutes) was turned on approximately 10 minutes before the emotional *N*-back training was started such that tDCS continued during the first half of *N*-back task performance and was turned off during the second half of the task, shown in Figure 1b. The emotional *N*-back test version was carried out during the postassessment visit. Participants were debriefed about their tDCS condition (real or sham) and the fictitious nature of the *N*-back performance feedback after data collection was completed.

Data Reduction and Statistical Analysis

Data were analyzed in generalized linear mixed-effects models (GLMMs) based on a gamma distribution in R⁹⁶ using the "lme4" package.⁹⁷ Within-subject outliers in dependent variables (> 3 standard deviations from the mean) were excluded. Effects are reported as significant when $p < 0.05$ (two-tailed). Significant interaction effects were followed up by post hoc comparisons of the estimated marginal means using the "emmeans" package⁹⁸ and were reported with Cohen's *d* effect sizes

Emotional Working Memory

The effects of tDCS were examined for both *N*-back training performance and *N*-back postassessment performance. *N*-back

training performance was operationalized as the achieved WM load level (*N*) per block and analyzed by fixed effects for Group (real vs sham tDCS), Session (sessions 1, 2, and 3), Block (1–10), and the quadratic term Block² to model the typical nonlinear learning curve during training sessions. Interindividual variability in performance levels and in learning rates was modeled by a random intercept for Participant and a random slope for Block. *N*-back postassessment performance was operationalized as the correct-response median reaction time (RT) and the sensitivity ($d' = Z_{\text{hits}} - Z_{\text{false alarms}}$) reflecting the ability to distinguish target trials from nontarget trials,⁹⁹ analyzed by fixed effects for Group, WM load ($N = 1-4$), and the maximum WM load during training (Train Max), together with a random intercept for Participant.

NPU-Threat Test

Startle amplitudes and fear ratings were significantly higher in the threat than in the safe conditions, as shown in Figure 2a and Supplementary Data section 2.4.1. To test the effects of tDCS, threat cue responses were analyzed by fixed effects for Group, Threat condition (predictable or unpredictable shock), Time (baseline, postassessment), Sequence Number (1 or 2), and Probe Number (1–6). Variability in threat responsivity and startle habituation was modeled by a random intercept for Participant (fear ratings) and a random slope for Probe Number (startle amplitudes).

Exploratory Analyses

To explore the predictive value of the theta/beta ratio on tDCS effects during the tDCS training sessions, the theta/beta ratio was entered to the GLMM analyzing *N*-back training performance as described above. Interactions between the theta/beta ratio and effects of tDCS group over time (over blocks or sessions) would indicate a predictive value of the theta/beta ratio. In addition, the predictive values of age, education, and start performance were evaluated following the same procedure.

RESULTS

Participants tolerated tDCS well. We noticed a small skin lesion on the anode location in one participant, likely resulting from an insufficiently soaked sponge pad during administration of tDCS in session 3. The lesion healed within six days. Some participants reported mild burning, itching, and tingling sensations, rated on average between 1 ("absent") and 2 ("mild"). No significant group differences appeared in these or other tDCS adverse effects or emotional state fluctuations during tDCS sessions, as shown in Supplementary Data section 2.1.

Three of the 79 included volunteers failed to comply with the *N*-back task instructions, and four dropped out prematurely because of restrictions related to coronavirus disease-2019 ($n = 2$), lack of time ($n = 1$), or no reason provided ($n = 1$). Ten participants showed insufficient (< 30%) valid startle responses ($n = 2$) or encountered technical issues during NPU-threat test recordings ($n = 8$), resulting in a sample of $n = 62$ for analysis of the primary outcome measure. For the other outcome measures, data from 72 participants were available for statistical analyses (real tDCS: $n = 37$, sham tDCS: $n = 35$). Supplementary Data Figure S1 shows the full CONSORT (Consolidated Standards of Reporting Trials) flow diagram.




Figure 2. Startle response amplitudes and fear ratings per condition of the NPU-threat test (a) and per tDCS group (b). Error bars represent the standard deviation. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$. Full model outcomes represent the estimated effects (b) with associated standard errors. Bars with a star (*) represent statistically significant effects. n.s., not significant. [Color figure can be viewed at www.neuromodulationjournal.org]

Emotional Working Memory Performance

TDCS Training Sessions

Results showed significant interaction effects of Group \times Session and Group \times Block² (p 's < 0.016 , [Supplementary Data Table S2.2.1](#) and [Fig. 3a](#)). In separate GLMMs per session, the Group \times Block² interaction showed a trend-like effect in tDCS training sessions 1 and 3 (p 's < 0.059 , [Supplementary Data Table S2.2.2](#)). [Figure 3a](#) shows that N -back training performance tended to be higher in the real tDCS group than in the sham tDCS group in the first tDCS training session ($p = 0.081$). This trend was not observed in the subsequent sessions (p 's > 0.40 , [Supplementary Data Table S2.2.3](#)). Results did not significantly differ when data were analyzed separately for performance during online tDCS (blocks 1–5) or offline tDCS (blocks 6–10), as shown in [Supplementary Data section 2.2.1](#). Together, these results show no significant group difference but suggest that the real tDCS group tended to improve N -back performance faster compared with the sham tDCS group during the first tDCS session.

Postassessment

No significant main or interaction effects of Group were found for RT and d' (p 's > 0.29), as shown in [Figure 3b](#) and [Supplementary](#)

[Data Table S2.3.1](#). Excluding data from participants who achieved a relatively low maximum WM load level (Train max < 4 , real tDCS: $n = 9$, sham tDCS: $n = 3$) did not significantly change the results.

NPU-Threat Test

Significant Group \times Time interactions were found for startle amplitudes and fear ratings (p 's < 0.038 , [Fig. 2b](#) and [Supplementary Data Table S2.4.3](#)). Follow-up comparisons revealed that the real tDCS group showed lower baseline startle amplitudes (p 's < 0.035) and lower baseline fear ratings ($p = 0.050$) in response to the Predictable shock cues compared with the sham tDCS group, as shown in [Figure 2b](#) and [Supplementary Data Table S2.4.4](#). Postassessment results showed no significant effects of real vs sham tDCS on startle amplitudes (p 's > 0.141) or fear ratings (p 's > 0.075).

Exploratory Analyses: Predictors of tDCS Effects

Exploratory results indicated that the baseline theta/beta ratio influenced the effect of real tDCS on N -back training performance. In the full model, significant three-way interactions were observed of Theta/beta ratio with Group and Session (Group \times Session \times Theta/beta ratio: $b(\text{SE}) = -0.11(0.04)$, $p = 0.013$) and with Group and Block (Group \times Block² \times Theta/beta ratio: $b(\text{SE}) = -0.11(0.04)$, $p = 0.013$, [Supplementary Data Table S2.5.1](#)). To interpret these interactions, N -back training performance was plotted per session, separately for median-split subsamples based on baseline theta/beta ratio ([Figure 3c](#)). Follow-up group comparisons indicated that the real tDCS group only showed significantly improved performance relative to the sham tDCS group during session 1 (not during later sessions, [Supplementary Data Table S2.5.3](#)) in participants with a higher baseline theta/beta ratio. Results showed no evidence for changes in theta/beta ratios from baseline to post-assessment; no significant effect on theta/beta ratios was observed for Time, Group, or their interaction (Time: $b(\text{SE}) = 0.52(0.36)$, $p = 0.149$; Group: $b(\text{SE}) = 0.06(0.06)$, $p = 0.284$; Time \times Group: $b(\text{SE}) = -0.13(0.08)$, $p = 0.111$).

In addition, the results of the second part of these exploratory analyses followed previous findings by showing an influence of age, education, and start performance on the effect of real tDCS on N -back training performance. For all three predictors, results showed significant four-way interactions with Group, Session, and Block (p 's < 0.010 , [Supplementary Data Table S2.5.2](#)). Visual inspection of N -back training performance per median-split subsamples and follow-up group comparisons indicated that improved performance in the real vs sham tDCS group during session 1 was shown by participants with lower educational level, younger age, and higher start performance ([Supplementary Data Table S2.5.3](#)). This predictor-dependent group difference was most pronounced for age ([Supplementary Data Table S2.5.3](#)), as illustrated in [Figure 3c](#).

DISCUSSION

This study examined changes in top-down stress regulation in healthy military personnel after three sessions of anodal tDCS applied over the right DLPFC at 2 mA combined with emotional WM training. Contrary to our hypothesis, results indicated no significant effect of real vs sham tDCS combined with WM training on stress regulation; stress-related responses to a threat-of-shock paradigm with instructed emotional downregulation did not




Figure 3. a. *N*-back performance level during training per session. Error bars represent the standard error around the mean. Ribbons represent the standard deviation. b. RT and *d'* per WM load in the postassessment *N*-back test. Error bars represent the standard deviation. c. WM load level during emotional *N*-back training per session, separated for median-split subsamples based on the theta/beta ratio (above) and age (below). **p* < 0.005; ·*p* < 0.10. Full model outcomes represent the estimated effects (b) with associated standard errors. Bars with a star (*) represent statistically significant effects. n.s., not significant. [Color figure can be viewed at www.neuromodulationjournal.org]

differ between groups. Moreover, at group level, results yielded no significant effect of real vs sham tDCS on emotional WM performance during the training or at postassessment. Interestingly, however, post hoc exploratory analyses of potential predictors of the tDCS response, including the theta/beta ratio, suggested a trait-dependent effect of tDCS on performance during the first tDCS WM training session. These findings suggest that tDCS as applied here may have only a short-lasting and trait-dependent effect during the early stages of the tDCS training intervention.

tDCS Effects on Stress Regulation

No significant real vs sham tDCS effects were observed at postassessment in the intensity or habituation of threat-related responses during the NPU-threat test. Although the NPU-threat manipulation was successful, our sample showed on average relatively low startle amplitudes (< 30 mV) and fear ratings (< 30 on a 0–100 scale) in response to the threat conditions, compared with other healthy participant samples (average startle amplitudes of almost 50 mV, fear ratings between 4 and 5 on a 0–10 scale).^{100,101}

Although the stress regulation instructions may have lowered the intensity of threat responses in this study,^{8,9} the low threat responses also could be a result of overall lower anxiety sensitivity in our sample (ASI-3 total scores mean \pm SD: 8.1 ± 6.0) than in other nonmilitary Dutch healthy participant samples (mean \pm SD: 10.7 ± 8.1).⁸⁸ Hence, our participants may have required little improvement in top-down stress regulation to attenuate the already low threat responses. Considering that stress resilience is part of military training and selection, threat manipulations that elicit stronger stress responses may be necessary for studies in healthy military populations. Moreover, it should be noted that our results may not generalize to individuals with higher threat sensitivity or stress regulation problems.

tDCS Effects on Emotional Working Memory Training

When considering predictive factors of tDCS response, tDCS showed beneficial effects on emotional WM performance, but this effect was limited to the first training session. This short-lived tDCS effect contrasts with results from four previous studies showing significant tDCS-induced performance gains that accumulated during WM training and that were sustained in the days or even months after the tDCS training intervention.^{44–46,102} These studies administered anodal tDCS over the right^{44,45,102} or left^{44–46} DLPFC over multiple (four to seven) sessions in healthy volunteers, as in this study. Unlike this study, the effects of tDCS were tested in students, and different reference (cathode) locations were used (over the parietal cortex, contralateral to the anode, or surrounding the anode, ie, high-definition tDCS). However, because two of the same research groups did not replicate their results in later studies using similar samples and electrode montages,^{48,49} these differences in study characteristics do not seem to explain the difference in study results.

The short-lasting tDCS effect in this study also could reflect that participants already reached ceiling performance during the first *N*-back training session. It has been shown that *N*-back ceiling performance can be achieved within 20 minutes of the task,¹⁰³ and we observed a significant flattening of performance improvement toward the end of the first session. In contrast, the emotional dual *N*-back task was based on a WM training study by Schweizer et al.¹⁸ Maximum performance levels in that study (*N* between 4 and 5) were higher than the performance levels reached in our sample at the end of the first tDCS training session (*N* between 2 and 3, Fig. 2a). In addition, our participants further increased in *N*-back performance from the first to the second and third session (session 3: *N* between 3 and 4, Fig. 2a). These results suggest that participants did not reach ceiling performance during the first tDCS training session and therefore do not support the idea that ceiling performance explains the lack of tDCS effects on WM performance beyond the first tDCS training session.

Interestingly, our results concur with several previous studies showing similar short-lasting tDCS effects. For example, three studies in healthy students showed that multiple (two, three, or ten) sessions of anodal tDCS over the left DLPFC (1–2 mA) significantly enhanced WM performance during the first session, but not in subsequent sessions.^{104–106} Another study showed that anodal tDCS over the left DLPFC (1.5 mA) only significantly enhanced cognitive task performance when real tDCS was applied in the first of two experimental sessions in a crossover design.¹⁰⁷ The authors of this study proposed that tDCS may primarily

facilitate performance at the onset of learning new cognitive skills. In line with this idea, our results show that the early-stage tDCS effect was driven by a steeper learning curve, suggesting accelerated learning. Evidence from motor cortex stimulation research supports the idea that performing a learning task during tDCS mediates tDCS-related changes in task performance.^{108,109} Moreover, effects of noninvasive brain stimulation, including tDCS effects on WM performance, have been shown to depend on arousal and stress levels.^{110–113} In this study, central nervous system arousal could have been elevated in particular during the first session owing to the novelty of tDCS and the training task. Altogether, factors such as novel learning processes and arousal could have interacted with the neurophysiological effects of tDCS specifically during the early stages of the tDCS training intervention. More insights into these potential interactions may help to unravel how beneficial tDCS effects can be extended beyond the short-lasting performance enhancement observed here.

Influence of Individual Characteristics on tDCS Response

In addition to the confirmatory group-level analyses, post hoc analyses were performed to explore sources of individual differences in tDCS response. These exploratory analyses indicated that the early-stage tDCS effect on emotional WM performance was stronger in relation to higher theta/beta ratios. Higher theta/beta ratios have been repeatedly associated with higher reward motivation.^{69,70,114} Earlier findings showed that reward motivation increases working memory performance and related PFC activity¹¹⁵ and may promote tDCS effects on working memory,^{116,117} which might explain the predictive value of the theta/beta ratio in this study. Moreover, higher theta/beta ratios have been associated with lower cognitive control, possibly indicating that individuals with dominant subcortical-based drives relative to cortical-based cognitive control benefited more from tDCS.^{67,68} In addition, in line with previous findings, higher baseline WM performance,^{54,118} younger age,^{52,53} and lower educational level were associated with a stronger early-stage tDCS effect. Accordingly, several findings suggest that factors such as a shorter brain-to-skull distance¹¹⁹ and higher levels of neural plasticity^{120,121} may contribute to the higher effectivity of tDCS observed in the younger adults. Instead, results on the influence of education and baseline WM performance on tDCS response are mixed.^{55,57,58} Our finding of larger tDCS response in individuals with higher baseline WM performance is in line with some previous studies,^{54,59} and may indicate that higher recruitment of the targeted frontoparietal pathways facilitates tDCS effects.⁵⁴ However, such explanations remain largely speculative because this study does not identify the processes underlying the interaction between tDCS and these factors. Moreover, our results do not elucidate to what extent the four examined predictors reflect overlapping or distinct factors that influence tDCS response.

Clearly, replication of these exploratory results is required before firm conclusions can be drawn. Nevertheless, these findings highlight the importance of involving individual state- and trait-dependent factors in understanding the effects of tDCS. Interestingly, by replicating the association between the theta/beta ratio and effects of transcranial electric stimulation on cognitive performance,^{73,74} our results motivate further research to establish whether this resting-state EEG readout is a useful predictor of tDCS response.

Limitations

This study is disadvantaged by some methodological drawbacks. First, ramp-up-ramp-down stimulation was applied in the sham tDCS condition. Although this sham method is commonly used,¹²² results about blinding success have been mixed.^{123–125} Blinding success was not formally assessed in this study, and a possible effect of unsuccessful blinding can therefore not be completely ruled out. Second, the effectivity of anodal tDCS applied over the DLPFC on working memory tasks performance depends on the electric field strength and consequent excitability changes in a relatively ventrally located area of the DLPFC.^{38,43} Based on a priori electric field modeling estimations (Fig. 1b), the applied electrode montage should induce peak level electrical field strength in this region. Unfortunately, the actual electrical field strength in this region in individual participants in our study is not known and could not be estimated because of the lack of anatomical scans. Third, a tDCS training intervention with three sessions is comparable to previous tDCS intervention studies (two–ten sessions, as discussed above). However, not all studies show a significant effect of tDCS starting from the first session onward. For example, careful examination of findings in two previous studies shows that a clear effect of tDCS on WM performance started to manifest after three or more sessions.^{44,46} Together with the potential ceiling performance issues on the NPU-threat test and N-back task, this suggests that future studies may benefit from applying more tDCS training sessions and using more sensitive outcomes measures (eg, more variety or cognitive challenge in the tasks) to get better insight into the effect of multisession tDCS on WM training. Finally, this study applied tDCS both online and offline, that is, tDCS administration only covered the first half of the N-back training in each session. Online and offline tDCS were combined because it is not yet clear which timing of tDCS has better effects on cognitive performance.^{93–95} However, many previous tDCS WM training studies in healthy individuals applied tDCS online only^{44–46,104,105} or offline only.¹⁰² The generalizability of our results may therefore be limited to this specific online/offline tDCS application.

CONCLUSIONS

This study in healthy military personnel showed no evidence for the hypothesized beneficial effect on top-down stress regulation of multisession anodal tDCS over the right DLPFC during emotional WM training. Instead, the results suggest that tDCS had a short-term beneficial effect on emotional WM performance in the early stages of the training. This effect was moderated by the theta/beta ratio and other previously identified predictors of tDCS response, emphasizing the importance of state- and trait-dependent factors in effects of tDCS on cognitive performance.

Acknowledgements

The authors thank Florian Bruckmann, Carolien Bogaard, and Dayenne Teulings for their help in data collection; Anouck Schippers for her help in developing the software; and Tim Varkevisser and Nadia Leen for the graphical optimization of the face stimuli. Furthermore, they thank Fenna Leijten, Marieke Langemeyer, and their military colleagues at the Dutch Ministry of Defence who facilitated participant recruitment and experimental work on location. The authors also thank two anonymous reviewers for their

constructive feedback on the manuscript. Finally, and most of all, they thank their participants for their valuable time and effort; this study would not have been possible without them.

Authorship Statements

Fenne M. Smits, Elbert Geuze, Jack van Honk, and Dennis J.L.G. Schutter designed the study conceptually. Fenne M. Smits and Guido J. de Kort designed the practical aspects of the study. Fenne M. Smits, Karlijn Kouwer, and Lisa Geerlings conducted the study, including participant recruitment and data collection, with the supervision of Elbert Geuze and Dennis J.L.G. Schutter. Fenne M. Smits performed data analysis with the supervision of Elbert Geuze and Dennis J.L.G. Schutter. Fenne M. Smits drafted the manuscript with important intellectual input from Elbert Geuze, Jack van Honk, and Dennis J.L.G. Schutter. All authors reviewed the drafted manuscript. All authors approved the final manuscript.

How to Cite This Article

Smits F.M., Geuze E., de Kort G.J., Kouwer K., Geerlings L., van Honk J., Schutter D.J.L.G. 2023. Effects of Multisession Transcranial Direct Current Stimulation on Stress Regulation and Emotional Working Memory: A Randomized Controlled Trial in Healthy Military Personnel. *Neuromodulation* 2023; 26: 817–828.

SUPPLEMENTARY DATA

To access the supplementary material accompanying this article, visit the online version of *Neuromodulation: Technology at the Neural Interface* at www.neuromodulationjournal.org and at <https://doi.org/10.1016/j.neurom.2022.05.002>.

REFERENCES

1. Reijnen A, Rademaker AR, Vermetten E, Geuze E. Prevalence of mental health symptoms in Dutch military personnel returning from deployment to Afghanistan: a 2-year longitudinal analysis. *Eur Psychiatry*. 2015;30:341–346. <https://doi.org/10.1016/j.eurpsy.2014.05.003>.
2. Bartone PT, Adler AB, Vaitkus MA. Dimensions of psychological stress in peacekeeping operations. *Mil Med*. 1998;163:587–593. <https://doi.org/10.1093/milmed/163.9.587>.
3. Martin K, McLeod E, Périard J, Rattray B, Keegan R, Pyne DB. The impact of environmental stress on cognitive performance: a systematic review. *Hum Factors*. 2019;61:1205–1246. <https://doi.org/10.1177/0018720819839817>.
4. Sareen J, Cox BJ, Afifi TO, et al. Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care: findings from a large representative sample of military personnel. *Arch Gen Psychiatry*. 2007;64:843–852. <https://doi.org/10.1001/archpsyc.64.7.843>.
5. van der Wal SJ, Vermetten E, Elbert G. Long-term development of post-traumatic stress symptoms and associated risk factors in military service members deployed to Afghanistan: results from the PRISMO 10-year follow-up. *Eur Psychiatry*. 2020;64:e10. <https://doi.org/10.1192/j.eurpsy.2020.113>.
6. Webb TL, Miles E, Sheeran P. Dealing with feeling: a meta-analysis of the effectiveness of strategies derived from the process model of emotion regulation. *Psychol Bull*. 2012;138:775–808. <https://doi.org/10.1037/a0027600>.
7. Gross JJ. Emotion regulation: affective, cognitive, and social consequences. *Psychophysiology*. 2002;39:281–291. <https://doi.org/10.1017/S0048577201393198>.
8. Shurick AA, Hamilton JR, Harris LT, Roy AK, Gross JJ, Phelps EA. Durable effects of cognitive restructuring on conditioned fear. *Emotion*. 2012;12:1393–1397. <https://doi.org/10.1037/a0029143>.
9. Lissek S, Orme K, McDowell DJ, et al. Emotion regulation and potentiated startle across affective picture and threat-of-shock paradigms. *Biol Psychol*. 2007;76:124–133. <https://doi.org/10.1016/j.biopsycho.2007.07.002>.

10. Kalisch R, Müller MB, Tüscher O. A conceptual framework for the neurobiological study of resilience. *Behav Brain Sci.* 2015;38:e92. <https://doi.org/10.1017/S0140525X1400082X>.
11. Denny BT, Ochsner KN. Behavioral effects of longitudinal training in cognitive reappraisal. *Emotion.* 2014;14:425–433. <https://doi.org/10.1037/a0035276>.
12. Raio CM, Oredru TA, Palazzolo L, Shurick AA, Phelps EA. Cognitive emotion regulation fails the stress test. *Proc Natl Acad Sci USA.* 2013;110:15139–15144. <https://doi.org/10.1073/pnas.1305706110>.
13. Kohn N, Eickhoff SB, Scheller M, Laird AR, Fox PT, Habel U. Neural network of cognitive emotion regulation — an ALE meta-analysis and MACM analysis. *Neuroimage.* 2014;87:345–355. <https://doi.org/10.1016/j.neuroimage.2013.11.001>.
14. Berboth S, Morawetz C. Amygdala-prefrontal connectivity during emotion regulation: a meta-analysis of psychophysiological interactions. *Neuro-psychologia.* 2021;153:107767. <https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2021.107767>.
15. Arnsten AFT. Stress weakens prefrontal networks: molecular insults to higher cognition. *Nat Neurosci.* 2015;18:1376–1385. <https://doi.org/10.1038/nn.4087>.
16. Schmeichel BJ, Volokhov RN, Demaree HA. Working memory capacity and the self-regulation of emotional expression and experience. *J Pers Soc Psychol.* 2008;95:1526–1540. <https://doi.org/10.1037/a0013345>.
17. Schmeichel BJ, Demaree HA. Working memory capacity and spontaneous emotion regulation: high capacity predicts self-enhancement in response to negative feedback. *Emotion.* 2010;10:739–744. <https://doi.org/10.1037/a0019355>.
18. Schweizer S, Grahn J, Hampshire A, Mobbs D, Dalgleish T. Training the emotional brain: improving affective control through emotional working memory training. *J Neurosci.* 2013;33:5301–5311. <https://doi.org/10.1523/JNEUROSCI.2593-12.2013>.
19. Schweizer S, Hampshire A, Dalgleish T. Extending brain-training to the affective domain: increasing cognitive and affective executive control through emotional working memory training. *PLoS One.* 2011;6:e24372. <https://doi.org/10.1371/journal.pone.0024372>.
20. Pe ML, Raes F, Kuppens P. The cognitive building blocks of emotion regulation: ability to update working memory moderates the efficacy of rumination and reappraisal on emotion. *PLoS One.* 2013;8:e69071. <https://doi.org/10.1371/JOURNAL.PONE.0069071>.
21. Cirillo P, Gold AK, Nardi AE, et al. Transcranial magnetic stimulation in anxiety and trauma-related disorders: a systematic review and meta-analysis. *Brain Behav.* 2019;9:e01284. <https://doi.org/10.1002/brb3.1284>.
22. Doerig N, Seinsche RJ, Moisa M, Seifritz E, Ruff CC, Kleim B. Enhancing reappraisal of negative emotional memories with transcranial direct current stimulation. *Sci Rep.* 2021;11:14760. <https://doi.org/10.1038/s41598-021-93647-1>.
23. Feeser M, Prehn K, Kazzer P, Mungee A, Bajbouj M. Transcranial direct current stimulation enhances cognitive control during emotion regulation. *Brain Stimul.* 2014;7:105–112. <https://doi.org/10.1016/j.brs.2013.08.006>.
24. He Z, Lin Y, Xia L, Liu Z, Zhang D, Elliott R. Critical role of the right VLPFC in emotional regulation of social exclusion: a tDCS study. *Soc Cogn Affect Neurosci.* 2018;13:357–366. <https://doi.org/10.1093/scan/nsy026>.
25. Vanderhasselt M-A, Sanchez A, Josephy H, Baeken C, Brunoni AR, De Raedt R. Anodal tDCS over the right dorsolateral prefrontal cortex modulates cognitive processing of emotional information as a function of trait rumination in healthy volunteers. *Biol Psychol.* 2017;123:111–118. <https://doi.org/10.1016/j.biopsycho.2016.12.006>.
26. Herrmann MJ, Simons BSE, Horst AK, Boehme S, Straube T, Polak T. Modulation of sustained fear by transcranial direct current stimulation (tDCS) of the right inferior frontal cortex (rIFC). *Biol Psychol.* 2018;139:173–177. <https://doi.org/10.1016/j.biopsycho.2018.10.013>.
27. Herrmann MJ, Beier JS, Simons B, Polak T. Transcranial direct current stimulation (tDCS) of the right inferior frontal gyrus attenuates skin conductance responses to unpredictable threat conditions. *Front Hum Neurosci.* 2016;10:352. <https://doi.org/10.3389/fnhum.2016.00352>.
28. Chan MMY, Yau SSY, Han YMY. The neurobiology of prefrontal transcranial direct current stimulation (tDCS) in promoting brain plasticity: a systematic review and meta-analyses of human and rodent studies. *Neurosci Biobehav Rev.* 2021;125:392–416. <https://doi.org/10.1016/J.NEUBIOREV.2021.02.035>.
29. Knotkova H, Nitsche MA, Bikson M, Woods AJ, eds. *Practical Guide to Transcranial Direct Current Stimulation: Principles, Procedures and Applications.* Springer; 2019. <https://doi.org/10.1007/978-3-319-95948-1>.
30. Nitsche MA, Cohen LG, Wassermann EM, et al. Transcranial direct current stimulation: state of the art 2008. *Brain Stimul.* 2008;1:206–223. <https://doi.org/10.1016/J.BRS.2008.06.004>.
31. Marques LM, Morello LYN, Boggio PS. Ventrolateral but not dorsolateral Prefrontal Cortex tDCS effectively impact emotion reappraisal – effects on emotional experience and interbeat interval. *Sci Rep.* 2018;8, 15295. <https://doi.org/10.1038/s41598-018-33711-5>.
32. Clarke PJF, Van Bockstaele B, Marinovic W, Howell JA, Boyes ME, Notebaert L. The effects of left DLPFC tDCS on emotion regulation, biased attention, and emotional reactivity to negative content. *Cogn Affect Behav Neurosci.* 2020;20:1323–1335. <https://doi.org/10.3758/s13415-020-00840-2>.
33. Clarke PJF, Haridas SMP, Van Bockstaele B, Chen NTM, Saleminck E, Notebaert L. Frontal tDCS and emotional reactivity to negative content: examining the roles of biased interpretation and emotion regulation. *Cogn Ther Res.* 2021;45:19–30. <https://doi.org/10.1007/s10608-020-10162-9>.
34. Segrave RA, Arnold S, Hoy K, Fitzgerald PB. Concurrent cognitive control training augments the antidepressant efficacy of tDCS: a pilot study. *Brain Stimul.* 2014;7:325–331. <https://doi.org/10.1016/j.brs.2013.12.008>.
35. Berryhill ME, Martin D. Cognitive effects of transcranial direct current stimulation in healthy and clinical populations: an overview. *J ECT.* 2018;34:e25–e35. <https://doi.org/10.1097/YCT.0000000000000534>.
36. Bikson M, Rahman A. Origins of specificity during tDCS: anatomical, activity-selective, and input-bias mechanisms. *Front Hum Neurosci.* 2013;7:688. <https://doi.org/10.3389/FNHUM.2013.00688>.
37. Dedoncker J, Baeken C, de Raedt R, Vanderhasselt MA. Combined transcranial direct current stimulation and psychological interventions: state of the art and promising perspectives for clinical psychology. *Biol Psychol.* 2021;158:107991. <https://doi.org/10.1016/J.BIOPSYCHO.2020.107991>.
38. Pisoni A, Mattavelli G, Papagno C, Rosanova M, Casali AG, Romero Lauro LJ. Cognitive enhancement induced by anodal tDCS drives circuit-specific cortical plasticity. *Cereb Cortex.* 2018;28:1132–1140. <https://doi.org/10.1093/cercor/bhx021>.
39. Li LM, Violante IR, Leech R, et al. Brain state and polarity dependent modulation of brain networks by transcranial direct current stimulation. *Hum Brain Mapp.* 2019;40:904–915. <https://doi.org/10.1002/hbm.24420>.
40. Monte-Silva K, Kuo MF, Hesselthaler S, et al. Induction of late LTP-like plasticity in the human motor cortex by repeated non-invasive brain stimulation. *Brain Stimul.* 2013;6:424–432. <https://doi.org/10.1016/j.brs.2012.04.011>.
41. van 't Wout-Frank M, Shea MT, Larson VC, Greenberg BD, Philip NS. Combined transcranial direct current stimulation with virtual reality exposure for post-traumatic stress disorder: feasibility and pilot results. *Brain Stimul.* 2019;12:41–43. <https://doi.org/10.1016/j.brs.2018.09.011>.
42. Lee T-W, Xue S-W. Does emotion regulation engage the same neural circuit as working memory? A meta-analytical comparison between cognitive reappraisal of negative emotion and 2-back working memory task. *PLoS One.* 2018;13: e0203753. <https://doi.org/10.1371/journal.pone.0203753>.
43. Wischniewski M, Mantell KE, Opitz A. Identifying regions in prefrontal cortex related to working memory improvement: a novel meta-analytic method using electric field modeling. *Neurosci Biobehav Rev.* 2021;130:147–161. <https://doi.org/10.1016/J.NEUBIOREV.2021.08.017>.
44. Au J, Katz B, Buschkuehl M, et al. Enhancing working memory training with transcranial direct current stimulation. *J Cogn Neurosci.* 2016;28:1419–1432. https://doi.org/10.1162/jocn_a_00979.
45. Ruf SP, Fallgatter AJ, Plewnia C. Augmentation of working memory training by transcranial direct current stimulation (tDCS). *Sci Rep.* 2017;7:876. <https://doi.org/10.1038/s41598-017-01055-1>.
46. Ke Y, Wang N, Du J, et al. The effects of transcranial direct current stimulation (tDCS) on working memory training in healthy young adults. *Front Hum Neurosci.* 2019;13:19. <https://doi.org/10.3389/fnhum.2019.00019>.
47. Trumbo MC, Matzen LE, Coffman BA, et al. Enhanced working memory performance via transcranial direct current stimulation: the possibility of near and far transfer. *Neuropsychologia.* 2016;93:85–96. <https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2016.10.011>.
48. Au J, Katz B, Moon A, et al. Post-training stimulation of the right dorsolateral prefrontal cortex impairs working memory training performance. *J Neurosci Res.* 2021;99:2351–2363. <https://doi.org/10.1002/JNR.24784>.
49. Shires J, Carrasco C, Berryhill ME. No tDCS augmented working memory training benefit in undergraduates rewarded with course credit. *Brain Stimul.* 2020;13:1524–1526. <https://doi.org/10.1016/J.BRS.2020.08.015>.
50. Luque-Casado A, Rodríguez-Freiria R, Fogelson N, Iglesias-Soler E, Fernández-del-Olmo M. An integrative clustering approach to tDCS individual response variability in cognitive performance: Beyond a null effect on working memory. *Neuroscience.* 2020;443:120–130. <https://doi.org/10.1016/J.NEUROSCIENCE.2020.07.036>.
51. Perceval G, Flöel A, Meinzer M. Can transcranial direct current stimulation counteract age-associated functional impairment? *Neurosci Biobehav Rev.* 2016;65:157–172. <https://doi.org/10.1016/J.NEUBIOREV.2016.03.028>.
52. Leach RC, McCurdy MP, Trumbo MC, Matzen LE, Leshkar ED. Differential age effects of transcranial direct current stimulation on associative memory. *J Gerontol B Psychol Sci Soc Sci.* 2019;74:1163–1173. <https://doi.org/10.1093/GERONB/GBY003>.
53. Antonenko D, Nierhaus T, Meinzer M, et al. Age-dependent effects of brain stimulation on network centrality. *Neuroimage.* 2018;176:71–82. <https://doi.org/10.1016/J.NEUROIMAGE.2018.04.038>.
54. Schmieker M, Menze I, Schneider C, Taubert M, Zaehle T, Mueller NG. Making the rich richer: frontoparietal tDCS enhances transfer effects of a single-session distractor inhibition training on working memory in high capacity individuals but reduces them in low capacity individuals. *Neuroimage.* 2021;242:118438. <https://doi.org/10.1016/J.NEUROIMAGE.2021.118438>.
55. Emonson MRL, Fitzgerald PB, Rogasch NC, Hoy KE. Neurobiological effects of transcranial direct current stimulation in younger adults, older adults and mild cognitive impairment. *Neuropsychologia.* 2019;125:51–61. <https://doi.org/10.1016/j.neuropsychologia.2019.01.003>.
56. Brunoni AR, Tortella G, Benseñor IM, Lotufo PA, Carvalho AF, Fregni F. Cognitive effects of transcranial direct current stimulation in depression: results from the SELECT-TDCS trial and insights for further clinical trials. *J Affect Disord.* 2016;202:46–52. <https://doi.org/10.1016/J.JAD.2016.03.066>.

57. Splittgerber M, Salvador R, Brauer H, et al. Individual baseline performance and electrode montage impact on the effects of anodal tDCS Over the left dorso-lateral prefrontal cortex. *Front Hum Neurosci.* 2020;14:349. <https://doi.org/10.3389/fnhum.2020.00349>.
58. Tseng P, Hsu TY, Chang CF, et al. Unleashing potential: transcranial direct current stimulation over the right posterior parietal cortex improves change detection in low-performing individuals. *J Neurosci.* 2012;32:10554–10561. <https://doi.org/10.1523/JNEUROSCI.0362-12.2012>.
59. Berryhill ME, Jones KT. tDCS selectively improves working memory in older adults with more education. *Neurosci Lett.* 2012;521:148–151. <https://doi.org/10.1016/j.neulet.2012.05.074>.
60. Al-Kaysi AM, Al-Ani A, Loo CK, et al. Predicting tDCS treatment outcomes of patients with major depressive disorder using automated EEG classification. *J Affect Disord.* 2017;208:597–603. <https://doi.org/10.1016/j.jad.2016.10.021>.
61. Hordacre B, Moezzi B, Goldsworthy MR, Rogasch NC, Graetz LJ, Ridding MC. Resting state functional connectivity measures correlate with the response to anodal transcranial direct current stimulation. *Eur J Neurosci.* 2017;45:837–845. <https://doi.org/10.1111/EJN.13508>.
62. Thut G, Bergmann TO, Fröhlich F, et al. Guiding transcranial brain stimulation by EEG/MEG to interact with ongoing brain activity and associated functions: a position paper. *Clin Neurophysiol.* 2017;128:843–857. <https://doi.org/10.1016/j.clinph.2017.01.003>.
63. Schutter DJLG, Knyazev GG. Cross-frequency coupling of brain oscillations in studying motivation and emotion. *Motiv Emot.* 2012;36:46–54. <https://doi.org/10.1007/s11031-011-9237-6>.
64. Knyazev GG. Motivation, emotion, and their inhibitory control mirrored in brain oscillations. *Neurosci Biobehav Rev.* 2007;31:377–395. <https://doi.org/10.1016/j.neubiorev.2006.10.004>.
65. Tortella-Feliu M, Morillas-Romero A, Balle M, Llabrés J, Bornas X, Putman P. Spontaneous EEG activity and spontaneous emotion regulation. *Int J Psychophysiol.* 2014;94:365–372. <https://doi.org/10.1016/j.ijpsycho.2014.09.003>.
66. Putman P, van Peer J, Maimari I, van der Werff S. EEG theta/beta ratio in relation to fear-modulated response-inhibition, attentional control, and affective traits. *Biol Psychol.* 2010;83:73–78. <https://doi.org/10.1016/j.biopsycho.2009.10.008>.
67. Angelidis A, van der Does W, Schakel L, Putman P. Frontal EEG theta/beta ratio as an electrophysiological marker for attentional control and its test-retest reliability. *Biol Psychol.* 2016;121:49–52. <https://doi.org/10.1016/j.biopsycho.2016.09.008>.
68. Angelidis A, Hagenaars M, van Son D, van der Does W, Putman P. Do not look away! Spontaneous frontal EEG theta/beta ratio as a marker for cognitive control over attention to mild and high threat. *Biol Psychol.* 2018;135:8–17. <https://doi.org/10.1016/j.biopsycho.2018.03.002>.
69. Massar SAA, Kenemans JL, Schutter DJLG. Resting-state EEG theta activity and risk learning: sensitivity to reward or punishment? *Int J Psychophysiol.* 2014;91:172–177. <https://doi.org/10.1016/J.IJPSYCHO.2013.10.013>.
70. Schutte I, Kenemans JL, Schutter DJLG. Resting-state theta/beta EEG ratio is associated with reward- and punishment-related reversal learning. *Cogn Affect Behav Neurosci.* 2017;17:754–763. <https://doi.org/10.3758/s13415-017-0510-3>.
71. Wischniewski M, Zerr P, Schutter DJLG. Effects of theta transcranial alternating current stimulation over the frontal cortex on reversal learning. *Brain Stimul.* 2016;9:705–711. <https://doi.org/10.1016/j.brs.2016.04.011>.
72. Sari BA, Koster EHW, Pourtois G, Derakshan N. Training working memory to improve attentional control in anxiety: a proof-of-principle study using behavioral and electrophysiological measures. *Biol Psychol.* 2016;121:203–212. <https://doi.org/10.1016/j.biopsycho.2015.09.008>.
73. Harty S, Cohen Kadosh R. Suboptimal engagement of high-level cortical regions predicts random-noise-related gains in sustained attention. *Psychol Sci.* 2019;30:1318–1332. <https://doi.org/10.1177/0956797619856658>.
74. Sheffield JG, Raz G, Sella F, Kadosh RC. How can noise alter neurophysiology in order to improve human behaviour? A combined tRNS and EEG study. Preprint. Posted online January 9, 2020. <https://dx.doi.org/10.1101/2020.01.09.900118>.
75. Buhle JT, Silvers JA, Wager TD, et al. Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cereb Cortex.* 2014;24:2981–2990. <https://doi.org/10.1093/cercor/bht154>.
76. Thielscher A, Antunes A, Saturnino GB. Field modeling for transcranial magnetic stimulation: a useful tool to understand the physiological effects of TMS? *Annu Int Conf IEEE Eng Med Biol Soc.* 2015;2015:222–225. <https://doi.org/10.1109/EMBC.2015.7318340>.
77. Hanke M, Baumgartner FJ, Ibe P, et al. A high-resolution 7-Tesla fMRI dataset from complex natural stimulation with an audio movie. *Sci Data.* 2014;1:140003. <https://doi.org/10.1038/sdata.2014.3>.
78. Ma DS, Correll J, Wittenbrink B. The Chicago face database: a free stimulus set of faces and norming data. *Behav Res Methods.* 2015;47:1122–1135. <https://doi.org/10.3758/s13428-014-0532-5>.
79. Marteau TM, Bekker H. The development of a six-item short-form of the state scale of the Spielberger State-trait Anxiety Inventory (STAI). *Br J Clin Psychol.* 1992;31:301–306. Published correction appears in *Br J Clin Psychol.* 2020;59:276. <https://doi.org/10.1111/J.2044-8260.1992.TB00997.X>
80. Brunoni AR, Amadera J, Berbel B, Volz MS, Rizzerio BG, Fregni F. A systematic report on reporting and assessment of adverse effects associated with transcranial direct current stimulation. *Int J Neuropsychopharmacol.* 2011;14:1133–1145. <https://doi.org/10.1017/S1461145710001690>.
81. Dickerson SS, Kemeny ME. Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychol Bull.* 2004;130:355–391. <https://doi.org/10.1037/0033-2909.130.3.355>.
82. Schmitz A, Grillon C. Assessing fear and anxiety in humans using the threat of predictable and unpredictable aversive events (the NPU-threat test). *Nat Protoc.* 2012;7:527–532. <https://doi.org/10.1038/nprot.2012.001>.
83. Davis M. Neural systems involved in fear and anxiety measured with fear-potentiated startle. *Am Psychol.* 2006;61:741–756. <https://doi.org/10.1037/0003-066X.61.8.741>.
84. Blumenthal TD, Cuthbert BN, Filion DL, Hackley S, Lipp OV, Van Boxtel A. Committee report: guidelines for human startle eyeblink electromyographic studies. *Psychophysiology.* 2005;42:1–15. <https://doi.org/10.1111/j.1469-8986.2005.00271.x>.
85. Gross JJ, John OP. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol.* 2003;85:348–362. <https://doi.org/10.1037/0022-3514.85.2.348>.
86. Koole SL, Jostmann NB. Getting a grip on your feelings: effects of action orientation and external demands on intuitive affect regulation. *J Pers Soc Psychol.* 2004;87:974–990. <https://doi.org/10.1037/0022-3514.87.6.974>.
87. Reiss S, Peterson RA, Gursky DM, McNally RJ. Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behav Res Ther.* 1986;24:1–8. [https://doi.org/10.1016/0005-7967\(86\)90143-9](https://doi.org/10.1016/0005-7967(86)90143-9).
88. Taylor S, Zvolensky MJ, Cox BJ, et al. Robust dimensions of anxiety sensitivity: development and initial validation of the Anxiety Sensitivity Index-3. *Psychol Assess.* 2007;19:176–188. <https://doi.org/10.1037/1040-3590.19.2.176>.
89. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol.* 1988;54:1063–1070. <https://doi.org/10.1037/0022-3514.54.6.1063>.
90. Derryberry D, Reed MA. Anxiety-related attentional biases and their regulation by attentional control. *J Abnorm Psychol.* 2002;111:225–236. <https://doi.org/10.1037/0021-843x.111.2.225>.
91. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods.* 2004;134:9–21.
92. Lopez-Calderon J, Luck SJ. ERPLAB: an open-source toolbox for the analysis of event-related potentials. *Front Hum Neurosci.* 2014;8:213. <https://doi.org/10.3389/fnhum.2014.00213>.
93. Friehs MA, Frings C. Offline beats online: transcranial direct current stimulation timing influences on working memory. *Neuroreport.* 2019;30:795–799. <https://doi.org/10.1097/WNR.0000000000001272>.
94. Friehs MA, Frings C, Hartwigsen G. Effects of single-session transcranial direct current stimulation on reactive response inhibition. *Neurosci Biobehav Rev.* 2021;128:749–765. <https://doi.org/10.1016/J.NEUBIOREV.2021.07.013>.
95. Hill AT, Fitzgerald PB, Hoy KE. Effects of anodal transcranial direct current stimulation on working memory: a systematic review and meta-analysis of findings from healthy and neuropsychiatric populations. *Brain Stimul.* 2016;9:197–208. <https://doi.org/10.1016/j.brs.2015.10.006>.
96. R Core Team. *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing; 2022.
97. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 2015;67:1–48. <https://doi.org/10.18637/jss.v067.i01>.
98. Lenth RV. *Emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.7.3;* 2022. <https://CRAN.R-project.org/package=emmeans>.
99. Stanislaw H, Todorov N. Calculation of signal detection theory measures. *Behav Res Methods Instrum Comput.* 1999;31:137–149.
100. Balderston NL, Liu J, Roberson-Nay R, Ernst M, Grillon C. The relationship between dlPFC activity during unpredictable threat and CO₂-induced panic symptoms. *Transl Psychiatry.* 2017;7:1266. <https://doi.org/10.1038/s41398-017-0006-5>.
101. Ferry RA, Nelson BD. Differential impact of threat type on defensive motivation and attention during the NPU-threat task. *Motiv Emot.* 2020;44:670–685. <https://doi.org/10.1007/s11031-020-09835-5>.
102. Jones KT, Peterson DJ, Blacker KJ, Berryhill ME. Frontoparietal neurostimulation modulates working memory training benefits and oscillator synchronization. *Brain Res.* 2017;1667:28–40. <https://doi.org/10.1016/j.brainres.2017.05.005>.
103. Yamashita M, Kawato M, Imamizu H. Predicting learning plateau of working memory from whole-brain intrinsic network connectivity patterns. *Sci Rep.* 2015;5:7622. <https://doi.org/10.1038/srep07622>.
104. Talsma LJ, Kroese HA, Slagter HA. Boosting cognition: effects of multiple-session transcranial direct current stimulation on working memory. *J Cogn Neurosci.* 2017;29:755–768. https://doi.org/10.1162/jocn_a_01077.
105. Martin DM, Liu R, Alonzo A, et al. Can transcranial direct current stimulation enhance outcomes from cognitive training? A randomized controlled trial in healthy participants. *Int J Neuropsychopharmacol.* 2013;16:1927–1936. <https://doi.org/10.1017/S1461145713000539>.
106. Lally N, Nord CL, Walsh V, Roiser JP. Does excitatory fronto-extracerebral tDCS lead to improved working memory performance? *F1000Res.* 2013;2:219. <https://doi.org/10.12688/f1000research.2-219.v2>.
107. Fehring DJ, Illiparampil R, Acevedo N, Jaberzadeh S, Fitzgerald PB, Mansouri FA. Interaction of task-related learning and transcranial direct current stimulation of the prefrontal cortex in modulating executive functions. *Neuropsychologia.* 2019;131:148–159. <https://doi.org/10.1016/j.neuropsychologia.2019.05.011>.

108. Bortoletto M, Pellicciari MC, Rodella C, Miniussi C. The interaction with task-induced activity is more important than polarization: a tDCS study. *Brain Stimul.* 2015;8:269–276. <https://doi.org/10.1016/J.BRS.2014.11.006>.
109. Buch ER, Santarnecchi E, Antal A, et al. Effects of tDCS on motor learning and memory formation: a consensus and critical position paper. *Clin Neurophysiol.* 2017;128:589–603. <https://doi.org/10.1016/J.CLINPH.2017.01.004>.
110. Schutter DJLG, Hofman D, van Honk J. Fearful faces selectively increase corticospinal motor tract excitability: a transcranial magnetic stimulation study. *Psychophysiology.* 2008;45:345–348. <https://doi.org/10.1111/j.1469-8986.2007.00635.x>.
111. Coombes SA, Tandonnet C, Fujiyama H, Janelle CM, Cauraugh JH, Summers JJ. Emotion and motor preparation: a transcranial magnetic stimulation study of corticospinal motor tract excitability. *Cogn Affect Behav Neurosci.* 2009;9:380–388. <https://doi.org/10.3758/CABN.9.4.380>.
112. Ankri YLE, Braw Y, Luboshits G, Meiron O. The effects of stress and transcranial direct current stimulation (tDCS) on working memory: a randomized controlled trial. *Cogn Affect Behav Neurosci.* 2020;20:103–114. <https://doi.org/10.3758/s13415-019-00755-7>.
113. Esposito M, Ferrari C, Fracassi C, Miniussi C, Brignani D. Responsiveness to left-prefrontal tDCS varies according to arousal levels. *Eur J Neurosci.* 2022;55:762–777. <https://doi.org/10.1111/EJN.15584>.
114. Schutter DJLG, Van Honk J. Electrophysiological ratio markers for the balance between reward and punishment. *Brain Res Cogn Brain Res.* 2005;24:685–690. <https://doi.org/10.1016/J.COGBRAINRES.2005.04.002>.
115. Krawczyk DC, D'Esposito M. Modulation of working memory function by motivation through loss-aversion. *Hum Brain Mapp.* 2013;34:762–774. <https://doi.org/10.1002/HBM.21472>.
116. Di Rosa E, Brigadoi S, Cutini S, et al. Reward motivation and neurostimulation interact to improve working memory performance in healthy older adults: a simultaneous tDCS-fNIRS study. *Neuroimage.* 2019;202:116062. <https://doi.org/10.1016/J.NEUROIMAGE.2019.116062>.
117. Jones KT, Gözenman F, Berryhill ME. The strategy and motivational influences on the beneficial effect of neurostimulation: a tDCS and fNIRS study. *Neuroimage.* 2015;105:238–247. <https://doi.org/10.1016/J.NEUROIMAGE.2014.11.012>.
118. Jones KT, Berryhill ME. Parietal contributions to visual working memory depend on task difficulty. *Front Psychiatry.* 2012;3:81. <https://doi.org/10.3389/fpsy.2012.00081>.
119. Opitz A, Paulus W, Will S, Antunes A, Thielscher A. Determinants of the electric field during transcranial direct current stimulation. *Neuroimage.* 2015;109:140–150. <https://doi.org/10.1016/J.NEUROIMAGE.2015.01.033>.
120. Burke SN, Barnes CA. Neural plasticity in the ageing brain. *Nat Rev Neurosci.* 2006;7:30–40. <https://doi.org/10.1038/nrn1809>.
121. Cirillo G, di Pino G, Capone F, et al. Neurobiological after-effects of non-invasive brain stimulation. *Brain Stimul.* 2017;10:1–18. <https://doi.org/10.1016/j.brs.2016.11.009>.
122. Fonteneau C, Mondino M, Arns M, et al. Sham tDCS: a hidden source of variability? Reflections for further blinded, controlled trials. *Brain Stimul.* 2019;12:668–673. <https://doi.org/10.1016/j.brs.2018.12.977>.
123. Turi Z, Csifcsák G, Boayue NM, et al. Blinding is compromised for transcranial direct current stimulation at 1 mA for 20 min in young healthy adults. *Eur J Neurosci.* 2019;50:3261–3268. <https://doi.org/10.1111/EJN.14403>.
124. Greinacher R, Buhöt L, Möller L, Learmonth G. The time course of ineffective sham-blinding during low-intensity (1 mA) transcranial direct current stimulation. *Eur J Neurosci.* 2019;50:3380–3388. <https://doi.org/10.1111/ejn.14497>.
125. Ambrus GG, Al-Moyed H, Chaieb L, Sarp L, Antal A, Paulus W. The fade-in – short stimulation – fade out approach to sham tDCS – reliable at 1 mA for naïve and experienced subjects, but not investigators. *Brain Stimul.* 2012;5:499–504. <https://doi.org/10.1016/j.brs.2011.12.001>.

COMMENT

This negative study is an important addition to the literature, reminding the field of the importance to study the interaction between brain stimulation and various cognitive tasks. Although this intervention did not improve stress regulation, lessons learned from this work can help develop future intervention studies in patients.

Noah Philip, MD
Providence, RI, USA