

RESEARCH ARTICLE

Influence of different data-averaging methods on mean values of selected variables derived from preoperative cardiopulmonary exercise testing in patients scheduled for colorectal surgery

Ruud F. W. Franssen^{1,2,*}, Bart H. E. Sanders³, Tim Takken⁴, F. Jeroen Vogelaar², Maryska L. G. Janssen-Heijnen^{2,5}, Bart C. Bongers^{6,7}

1 Department of Clinical Physical Therapy, VieCuri Medical Center, Venlo, The Netherlands, **2** Department of Epidemiology, GROW School for Oncology and Reproduction, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands, **3** Department of Sports Medicine, VieCuri Medical Center, Venlo, The Netherlands, **4** Child Development and Exercise Center, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, The Netherlands, **5** Department of Epidemiology, VieCuri Medical Center, Venlo, The Netherlands, **6** Department of Nutrition and Movement Sciences, Nutrition and Translational Research in Metabolism (NUTRIM), Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands, **7** Department of Surgery, Nutrition and Translational Research in Metabolism (NUTRIM), Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands

* rfranssen@viecuri.nl



OPEN ACCESS

Citation: Franssen RFW, Sanders BHE, Takken T, Vogelaar FJ, Janssen-Heijnen MLG, Bongers BC (2023) Influence of different data-averaging methods on mean values of selected variables derived from preoperative cardiopulmonary exercise testing in patients scheduled for colorectal surgery. *PLoS ONE* 18(3): e0283129. <https://doi.org/10.1371/journal.pone.0283129>

Editor: Lindsay Bottoms, University of Hertfordshire, UNITED KINGDOM

Received: October 14, 2022

Accepted: March 3, 2023

Published: March 16, 2023

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0283129>

Copyright: © 2023 Franssen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction

Patients with a low cardiorespiratory fitness (CRF) undergoing colorectal cancer surgery have a high risk for postoperative complications. Cardiopulmonary exercise testing (CPET) to assess CRF is the gold standard for preoperative risk assessment. To aid interpretation of raw breath-by-breath data, different methods of data-averaging can be applied. This study aimed to investigate the influence of different data-averaging intervals on CPET variables used for preoperative risk assessment, as well as to evaluate whether different data-averaging intervals influence preoperative risk assessment.

Methods

A total of 21 preoperative CPETs were interpreted by two exercise physiologists using stationary time-based data-averaging intervals of 10, 20, and 30 seconds and rolling average intervals of 3 and 7 breaths. Mean values of CPET variables between different data averaging intervals were compared using repeated measures ANOVA. The variables of interest were oxygen uptake at peak exercise (VO_{2peak}), oxygen uptake at the ventilatory anaerobic threshold (VO_{2VAT}), oxygen uptake efficiency slope (OUES), the ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold (VE/VCO_{2VAT}), and the slope of the relationship between the minute ventilation and carbon dioxide production (VE/VCO_2 -slope).

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: This study was unconditionally financed by the Research and Innovation fund VieCuri Medical Center under reference number E.21.32.004-2. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Results

Between data-averaging intervals, no statistically significant differences were found in the mean values of CPET variables except for the ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold ($P = 0.001$). No statistically significant differences were found in the proportion of patients classified as high or low risk regardless of the used data-averaging interval.

Conclusion

There appears to be no significant or clinically relevant influence of the evaluated data-averaging intervals on the mean values of CPET outcomes used for preoperative risk assessment. Clinicians may choose a data-averaging interval that is appropriate for optimal interpretation and data visualization of the preoperative CPET. Nevertheless, caution should be taken as the chosen data-averaging interval might lead to substantial within-patient variation for individual patients.

Clinical trial registration

Prospectively registered at ClinicalTrials.gov ([NCT05353127](#)).

Introduction

Preoperative aerobic fitness is independently associated with postoperative outcomes following major abdominal surgery [1]. Consequently, cardiopulmonary exercise testing (CPET) is increasingly used within multimodal preoperative risk assessment [2], as it provides an objective, non-invasive, and accurate evaluation of a patient's aerobic fitness that represents the capacity to meet the increased oxygen demand following major abdominal surgery [3, 4]. The advantage of CPET over other risk assessment tools is that CPET encompasses an integrative evaluation of the cardiovascular, pulmonary, and muscular system [5]. In addition, CPET can be used to inform collaborative decision-making, to optimize comorbidities, to triage perioperative care (e.g., ward, intensive care), to advice on preoperative physical exercise training (e.g., risk assessment, contraindications), and to guide and personalize subsequent physical exercise training prescription [6].

During CPET, a patient exercises against a progressively increasing work rate until volitional exhaustion, while breath-by-breath respiratory gasses are analyzed. The large number of data-points that are collected by the breath-by-breath sampling rate can be a challenge for data visualization, as the signal can have high variability. Therefore, data-averaging is performed to optimize graphical data display and to aid CPET interpretation (see [Fig 1](#)). Although it is generally accepted that data-averaging methods influence the numerical value of CPET-derived variables, there is no consensus among existing guidelines on the best averaging method [7].

The most frequently used CPET-derived variables that are associated with postoperative complications in the current literature are the oxygen uptake at peak exercise (VO_{2peak}), the oxygen uptake at the ventilatory anaerobic threshold (VO_{2VAT}) [2, 6, 8], and the ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold (VE/VCO_{2VAT}) [9]. Measures that are less frequently used are the slope of the relationship between the minute ventilation and carbon dioxide production (VE/VCO_2 -slope), that can be used as an alternative for the VE/VCO_{2VAT} if the VAT is undeterminable [8], and the oxygen uptake efficiency slope (OUES) [10].




Fig 1. Visualization of the plot with oxygen uptake (VO_2) and carbon dioxide production (VCO_2) over time without data-averaging (graph A) and using the five different data-averaging intervals: a stationary time-based average of 10 seconds (graph B), 20 seconds (graph C), and 30 seconds (graph D), a rolling average interval of 3 breaths (graph E) and 7 breaths (graph F) in patient 21. See [S1 File](#) for a graphical display of the Wasserman plots of patient 21 with the different data-averaging intervals. Note that the number of data points is lower when stationary time-based averaging is used (and decreasing with longer data-averaging intervals) compared to when a rolling average is used. In addition, a lower number of data points leads to smoothing of the VO_2 and VCO_2 curves. Abbreviations: VAT = ventilatory anaerobic threshold; VCO_2 = carbon dioxide production; VO_2 = oxygen uptake; $\text{VO}_{2\text{peak}}$ = oxygen uptake at peak exercise. Vertical grey dotted lines represent start of the warm-up phase (W), test phase (T), and recovery phase (R).

<https://doi.org/10.1371/journal.pone.0283129.g001>

Although preoperative risk assessment should be multimodal, CPET-derived thresholds are often used to recognize patients with a low aerobic fitness who have a high risk for adverse surgical outcomes. In major abdominal surgery, often used thresholds to identify patients at high-risk for postoperative complications are a $\text{VO}_{2\text{peak}} < 18.2 \text{ mL/kg/min}$ and/or a $\text{VO}_{2\text{VAT}} < 11.1 \text{ mL/kg/min}$ [9]. Studies in healthy individuals have shown that the numerical value of the $\text{VO}_{2\text{peak}}$ can differ as much as $\sim 10\%$ depending on the data-averaging method [11–13], indicating that data-averaging might significantly influence threshold determination and subsequently might affect preoperative risk assessment. To date however, there are no studies quantifying the extent to which differences in data-averaging influence the numerical value of preoperative CPET-derived variables such as $\text{VO}_{2\text{peak}}$, $\text{VO}_{2\text{VAT}}$, OUES, $\text{VE}/\text{VCO}_{2\text{VAT}}$, and VE/VCO_2 -slope. Therefore, the primary aim of this study was to investigate the influence of different CPET data-averaging intervals on the numerical values of CPET-derived variables used for preoperative risk assessment in patients scheduled for elective colorectal cancer surgery. The secondary aim was to elucidate the impact of data-averaging intervals on the classification of patients into low or high risk for postoperative complications based on known risk assessment thresholds.

Methods

This observational cross-sectional study was performed at the VieCuri Medical Center, a large teaching hospital in Venlo, the Netherlands. The current study was executed as a secondary analysis of data collected in a study [14] that was approved by the Medical Ethics Review Committee–Zuyderland/Zuyd (Heerlen, the Netherlands) under reference number METCZ20190150. Reporting was done using the STROBE guidelines for reporting of cross-

sectional studies [15]. The study protocol was prospectively registered at ClinicalTrials.gov (NCT05353127).

Participants

Data from consecutive patients considered for colorectal cancer surgery who were ≥ 18 years of age, had a score ≤ 7 metabolic equivalents of task on the veterans-specific activity questionnaire, and therefore performed preoperative CPET as a part of a tele-prehabilitation study [14], were collected between July 2020 and September 2021. All patients signed informed consent. Preoperative CPET was conducted after diagnosis and before any intervention or treatment was initiated.

Preoperative cardiopulmonary exercise testing

Patients preoperatively performed incremental CPET up to volitional exertion in upright position on an electronically-braked cycle ergometer (Lode Corival, Lode BV, Groningen, the Netherlands). Prior to the test, patients were asked to refrain from vigorous physical activity, caffeine, and tobacco for 24 hours and meals for 2 hours, but to continue medication as usual. Seat height was adjusted to the participant's leg length. Before commencing CPET, forced vital capacity and forced expiratory volume in one second was obtained from maximal flow-volume curves (Ergostik, Geratherm Respiratory, Bad Kissingen, Germany) according to ATS/ERS standards [5]. Subsequently, baseline cardiopulmonary values were assessed during a three-minute rest period while seated at the cycle ergometer, thereafter a three-minute warm-up phase took place that consisted of unloaded cycling. After the warm-up, work rate was increased by constant increments of 5, 10, 15, 20, or 25 W/min in a ramp-like manner, depending on the subject's estimated physical fitness level and aimed at reaching a maximal effort within eight to twelve minutes. Throughout CPET, subjects maintained a pedaling frequency between 60 and 80 revolutions/min. The protocol continued until the patient's pedaling frequency fell definitely below 60 revolutions/min, despite strong verbal encouragement, or when the patient met the criteria for exercise termination before symptom limitation as proposed in the ATS/ACCP statement on cardiopulmonary exercise testing [5].

During CPET, subjects breathed through a facemask (Hans Rudolph, Kansas City, MO, USA) connected to an ergospirometry system (Ergostik, Geratherm Respiratory, Bad Kissingen, Germany). Before every test, calibration for respiratory gas analysis measurements (ambient air and a gas mixture of 16% oxygen and 5% carbon dioxide) and volume measurements (three-liter syringe) took place. Expired gas passed through a flow meter (triple V volume transducer), an oxygen analyzer, and a carbon dioxide analyzer. The flow meter and gas analyzers were connected to a computer that calculated breath-by-breath minute ventilation, oxygen uptake, carbon dioxide production, and the respiratory exchange ratio. Raw unfiltered breath-by-breath data was retrogradely averaged over five different data display intervals.

Procedures

Preoperative CPET patient data was anonymized and patient characteristics other than anthropometric measures were concealed. A medical and clinical exercise physiologist (BB) and a clinical exercise physiologist (RF) determined VO_{2peak} , VO_{2VAT} , OUES, VE/VCO_{2VAT} , and the VE/VCO_2 -slope in all CPETs by means of a predefined set of guidelines (see [S2 File](#)). A VO_{2peak} was conceived "valid" when objective criteria for maximal volitional exertion were reached defined as an RER ≥ 1.10 or reaching $\geq 95\%$ of the predicted maximal heart rate at peak exercise. CPET interpretation was performed using Blue Cherry software version 1.3.3.3 (Geratherm Respiratory GmbH, Bad Kissingen, Germany), in which observers interpreted the

CPET data together using TeamViewer software (TeamViewer GmbH, Göppingen, Germany). Final determination was based on consensus between the two observers. If the two observers were unable to reach consensus, a third observer (TT) was consulted. Data-averaging-intervals used were stationary time-based averages, calculated by averaging the breath-by-breath data over 10, 20, or 30 seconds and rolling averages calculated by averaging a fixed number of single breath measurements (i.e., 3 and 7), then discarding the first breath and adding a new breath to obtain a new breath averaging block. Determination of the aforementioned CPET variables was repeated for all five different data-averaging intervals.

Apart from the CPET data, the preoperative patient characteristics age, sex, body mass index, smoking status (never, former, current), age-adjusted Charlson comorbidity index, American Society of Anesthesiologists classification, veterans-specific activity questionnaire score, hemoglobin levels (mmol/L), and tumor location were recorded to characterize the study population.

Sample size

A sample size calculation was performed with G*Power [16] for F-test repeated measures within factors. Based on a mean \pm standard deviation (SD) value for VO_{2VAT} of 9.7 ± 2.3 mL/kg/min (based on preliminary analysis of the used data) for a mean difference between data-averaging methods of minimally 0.7 mL/kg/min, the estimated effect size is estimated at ~ 0.30 . With an α of 0.05 and a β of 0.80, a minimum of 15 CPETs are needed to detect the estimated effect size.

Statistical analysis

Continuous data were checked for normality using the Shapiro-Wilks test. To assess the difference between different CPET data-averaging intervals, differences in mean numerical values of VO_{2peak} , VO_{2VAT} , OUES, VE/VCO_{2VAT} , and the VE/VCO_2 -slope, between different data-averaging intervals were calculated and analyzed by means of within-factors repeated-measures analysis of variance (ANOVA). In case of a statistically significant difference between methods ($P < 0.05$), post-hoc testing was performed using the Bonferroni correction to identify exact differences. Effect sizes were estimated by calculating the eta squared (i.e., sum of squares of the effect divided by the total sum of squares). To evaluate the influence of data-averaging intervals on preoperative risk assessment, individual numerical values for VO_{2peak} , VO_{2VAT} , OUES, and VE/VCO_{2VAT} were compared with known preoperative risk assessment thresholds. Patients were classified as high-risk when having a $VO_{2peak} < 18.2$ mL/kg/min [9], $VO_{2VAT} < 11.1$ mL/kg/min [9], $OUES/kg < 20.6$ [10], and/or $VE/VCO_{2VAT} > 30.9$ [9]. Cochran's Q-test was used to determine whether differences in preoperative risk assessment exist between data-averaging methods. Differences between data-averaging methods were assumed statistically significant when $P < 0.05$.

Results

A total of 21 CPETs of patients with colorectal cancer (see Table 1 for patient characteristics) were re-assessed using five different data-averaging intervals. Thus, a total of 105 CPETs (five data-averaging intervals \times 21 CPETs) were evaluated. Mean \pm SD duration of the CPET ramp phase was 586 ± 174 seconds (9:46 \pm 2:54 min). A valid VO_{2peak} was reached in 70 (67.7%) of the evaluated CPETs. VO_{2VAT} and VE/VCO_{2VAT} were determinable in 104 out of 105 CPETs (99%). The OUES and $VE-VCO_2$ -slope were determinable in all 105 CPETs.

Mean values of the CPET-derived variables ranged from 14.5 mL/kg/min to 14.6 mL/kg/min for VO_{2peak} , from 9.3 mL/kg/min to 9.7 mL/kg/min for VO_{2VAT} , from 19.1 to 19.4 for

Table 1. Baseline characteristics of subjects.

Characteristics	n = 21
Age (years)	70.5 ± 12.5
Sex ratio (male; female)	12 (57%); 9 (43%)
Body mass index (kg/m ²)	28.6 ± 4.9
Age-adjusted Charlson comorbidity index	
≤3	10 (47.6%)
4–5	10 (47.6%)
6+	1 (4.8%)
ASA-classification	
I	4 (19.0%)
II	7 (33.3%)
III	9 (42.9%)
IV	1 (4.8%)
Hemoglobin level (mmol/L)	7.4 ± 1.2
Tumor location	
Colon	15 (71.4%)
Rectum	6 (28.6%)

Data are presented as mean ± standard deviation (SD) or as number (%).

<https://doi.org/10.1371/journal.pone.0283129.t001>

OUES/kg, from 31.2 to 31.9 for VE/VCO_{2VAT}, and from 33.6 to 35.3 for VE/VCO₂-slope, dependent on the different data-averaging intervals. There was a significant difference in mean values of VO_{2peak} between groups with different data averaging intervals, but this difference did not remain significant after post-hoc testing. For the variable VE/VCO_{2VAT}, the 3 breaths rolling average interval was statistically significant different from the time-based 20 seconds (P = 0.004) and 30 seconds (P = 0.005) data-averaging interval, as well as from the rolling average of 7 breaths (P = 0.021; see Table 2). The effect sizes for all variables were ≤0.009.

Fig 2 depicts within-patient variation in the numerical value of several CPET-derived variables using the five different data-averaging intervals. Although the numerical values for VO_{2peak} were consistent (maximal within patient difference, 0.4 mL/kg/min, or 5.6%), within patient variation could be as much as 4.0 mL/kg/min for VO_{2VAT} (40.8%), 5.7 for the OUES/kg (40.3%), 4.7 for VE/VCO_{2VAT} (13.4%), and 10.4 (37.3%) for VE/VCO₂-slope when using different data-averaging intervals (see Fig 2).

When dichotomizing patients into the high or low risk category for postoperative complications based on the numerical values of each CPET variable, the proportion of patients with a high risk based on their VO_{2peak} ranged from 76% to 81%, depending on the used data-averaging interval. Based on VO_{2VAT} the proportion of high-risk patients ranged from 67% to 76%, whereas this ranged from 57% to 67% for OUES/kg and from 76% to 86% for VE/VCO_{2VAT}. As depicted in Table 3, there were no statistically significant differences in the proportion of patients who were classified as at high risk between different data-averaging-intervals. As depicted in Fig 2, individual values of some patients crossed the risk threshold depending on the data-averaging interval that was used. Based on within-patient variation, the estimated risk could differ for 1 patient when based on VO_{2peak} (patient 15), for 5 patients based on VO_{2VAT} (patients 1, 6, 7, 12, and 19), for 2 patients based on OUES (patients 2 and 18), and for 4 patients based on VE/VCO_{2VAT} (patients 3, 7, 13, and 16), depending on the used data-averaging interval.

Table 2. Numerical values of CPET variables using different data-averaging intervals.

	Data-averaging interval					P-value ^a
	Stationary time-based average			Rolling average		
	10 seconds	20 seconds	30 seconds	3 breaths	7 breaths	
VO _{2peak} (mL/min)	1202 (1008–1396)	1194 (999–1389)	1193 (997–1390)	1201 (1008–1394)	1200 (1005–1396)	0.040^c
VO _{2peak} (mL/kg/min)	14.6 (12.5–16.7)	14.5 (12.4–16.6)	14.5 (12.4–16.6)	14.6 (12.5–16.7)	14.6 (12.5–16.7)	0.012^c
Valid VO _{2peak} (mL/kg/min) ^b	16.2 (13.7–18.7)	16.2 (13.6–18.7)	16.1 (13.5–18.8)	16.2 (13.6–18.8)	16.3 (13.7–18.7)	0.104
VO _{2VAT} (mL/min)	800 (684–916)	775 (656–895)	764 (669–859)	776 (668–884)	761 (669–851)	0.345
VO _{2VAT} (mL/kg/min)	9.7 (8.5–10.9)	9.4 (8.1–10.7)	9.3 (8.2–10.4)	9.5 (8.3–10.6)	9.3 (8.2–10.4)	0.435
OUES	1559 (1322–1795)	1559 (1338–1779)	1565 (1338–1792)	1582 (1354–1809)	1574 (1351–1798)	0.463
OUES/kg	19.1 (16.6–21.7)	19.2 (16.7–21.7)	19.3 (16.7–21.8)	19.5 (17.0–21.9)	19.4 (16.9–21.8)	0.479
VE/VCO _{2VAT}	34.2 (31.9–36.5)	34.6 (32.3–37.0) ^d	35.1 (32.6–37.5) ^d	33.6 (31.4–35.8) ^d	34.4 (32.0–36.8) ^d	0.001
VE/VCO ₂ -slope	31.4 (28.6–34.1)	31.8 (28.8–35.0)	31.2 (28.4–34.0)	31.8 (28.9–34.7)	31.8 (29.0–34.7)	0.608

Data are presented as mean and 95% confidence interval (CI), unless stated otherwise.

Abbreviations: OUES = oxygen uptake efficiency slope; VE/VCO₂-slope = the slope of the relationship between the minute ventilation and carbon dioxide production; VE/VCO_{2VAT} = ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold; VO_{2peak} = oxygen uptake at peak exercise; VO_{2VAT} = oxygen uptake at the ventilatory anaerobic threshold.

^a: as determined by repeated-measures ANOVA (within factors).

^b: as determined by a respiratory exchange ratio at peak exercise ≥ 1.10 and/or a heart rate at peak exercise $\geq 95\%$ of the predicted maximal heart rate based on the formula $208 - (0.8 \times \text{age in years})$.

^c: did not remain significant after post hoc testing with Bonferroni correction.

^d: the 3 breaths rolling average interval was statistically significant different from the stationary time-based interval of 20 seconds ($P = 0.004$) and 30 seconds ($P = 0.005$), as well as from the 7 breaths rolling average interval ($P = 0.021$).

<https://doi.org/10.1371/journal.pone.0283129.t002>

Discussion

To our knowledge, the current study was the first study that aimed to investigate whether the selection of different CPET data-averaging intervals would translate into differences in mean values of CPET-derived variables in patients with colorectal cancer who performed CPET for preoperative risk assessment. As CPET-derived variables are used to preoperatively classify patients into having a low or high risk for postoperative complications based on their CRF, the secondary aim of the current study was to investigate whether potential differences in the numerical values of CPET-derived variables would lead to differences in preoperative risk classification. Based on the mean values of the CPET-derived variables there were only statistically significant differences for the variables VO_{2peak} and VE/VCO_{2VAT} between different data-averaging intervals. For VO_{2peak}, the between-group difference did not remain significant after post-hoc analysis, whereas data-averaging group differences VE/VCO_{2VAT} were statistically significant between the 3 breaths moving average and the 20- and 30-second time-based interval, as well as the 7 breaths moving average.

For VO_{2peak}, the greatest observed difference between data-averaging groups was 0.1 mL/kg/min. Given that the coefficient of variation (a measure of reproducibility) for VO_{2peak} is estimated to be between ~5% and ~9% [5] (i.e., between ~0.7 mL/kg/min and ~1.3 mL/kg/min based on mean values of VO_{2peak} in the current study), the observed maximal difference of 0.1 mL/kg/min is not clinically relevant. The observation that this small difference in VO_{2peak} is not clinically relevant is further emphasized by the fact that no differences were found between the proportion of patients who were classified as low or high risk based on VO_{2peak} when using different data-averaging intervals in the current study. Provided that the critical difference of VE/VCO_{2VAT} in patients with colorectal cancer is assumed to be ~10% [17], the




Fig 2. Variation in the observed values of VO_{2peak} (graph A), VO_{2VAT} (graph B), OUES (graph C), VE/VCO_{2VAT} (graph D), and the VE/VCO_2 -slope (graph E) within individual patients. Dots represent individual numerical value with a unique color for each data-averaging interval throughout the graphs (red = 10 seconds; yellow = 20 seconds; green = 30 seconds; blue = 3 breaths; purple = 7 breaths). Error bars represent the mean values and 95% confidence intervals. Horizontal dotted lines represent known risk assessment thresholds defined as 18.2 mL/kg/min for VO_{2peak} (graph A), 11.1 mL/kg/min for VO_{2VAT} (graph B), <20.6 for OUES (graph C), and >30.9 for VE/VCO_{2VAT} (graph D). Note that individual values of patients often cross the risk threshold (dotted horizontal line). These patients might have a different risk estimation depending on the data-averaging interval. Abbreviations: OUES = oxygen uptake efficiency slope; VE/VCO_{2VAT} = ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold; VO_{2peak} = oxygen uptake at peak exercise; VO_{2VAT} = oxygen uptake at the ventilatory anaerobic threshold.

<https://doi.org/10.1371/journal.pone.0283129.g002>

maximal mean difference of 1.5 (5%) measured in the current study is not deemed clinically relevant. The observation that differences in the mean values of the VE/VCO_2 -slope are not clinically relevant is also supported by the very small effect size (0.009).

The main purpose of using data-averaging of CPET data is to reduce noise of breath-by-breath fluctuations and to aid CPET interpretation [5]. In the current study there seem to be

Table 3. Effect of different data-averaging intervals on classifying patients as having a high-risk for postoperative complications.

	Data-averaging interval					P-value ^a
	Stationary time-based average			Rolling average		
	10 seconds, n (%)	20 seconds n (%)	30 seconds (%) n (%)	3 breaths n (%)	7 breaths n (%)	
VO_{2peak}	17 (81%)	17 (81%)	17 (81%)	17 (81%)	16 (76%)	0.406
VO_{2VAT}	16 (76%)	16 (76%)	14 (67%)	14 (67%)	16 (76%)	0.615
OUES/kg	13 (62%)	13 (62%)	12 (57%)	14 (67%)	14 (67%)	0.231
VE/VCO_{2VAT}	16 (76%)	18 (86%)	17 (81%)	15 (71%)	16 (76%)	0.334

Data are presented as number (%).

Abbreviations: OUES = oxygen uptake efficiency slope; VE/VCO_{2VAT} = ventilatory equivalent for carbon dioxide at the ventilatory anaerobic threshold; VO_{2peak} = oxygen uptake at peak exercise; VO_{2VAT} = oxygen uptake at the ventilatory anaerobic threshold.

^a: determined by Cochran’s Q-test.

<https://doi.org/10.1371/journal.pone.0283129.t003>

no clinically relevant differences in CPET-derived variables between different data-averaging intervals. This is a reassuring observation that opens possibilities to be flexible in the use of data-averaging intervals as long as the interval is within certain boundaries. That is, the type and duration of CPET can be taken into consideration when determining the optimal data-averaging interval [7]. For example, using longer averaging intervals in longer tests, or using a rolling average for noisy data. On the other hand, longer intervals might mask dynamic pathophysiological processes such as oscillatory breathing. In these circumstances shorter time-based intervals might be optimal [7]. For preoperative exercise testing a stationary time-based average of 10 seconds, or a breath-based rolling average of 3 or 7 seconds might provide a good trade-off between the number of data-points and the duration of the test.

Although the literature is scarce with regard to the influence of data-averaging intervals on the determination of CPET-derived variables (and only available for VO_{2peak}), results of the current study are in line with a previous publication in which the effect of data-averaging intervals on VO_{2peak} in 22 healthy athletic subjects was investigated [18]. The authors found that only a stationary time-based data-averaging interval of 60 seconds was significantly different from all other data-averaging intervals (10, 15, 20, and 30 seconds) [18]. In a study evaluating VO_{2peak} values of 15 patients who were screened for heart transplant surgery (with comparable mean VO_{2peak} values as observed in the current study), no significant differences were found between stationary time-based data-averaging intervals of 15 and 30 seconds, and a 8 breaths rolling average interval [19]. Moreover, only a 60-second stationary time-based data-averaging interval was statistically significantly different from the aforementioned data-averaging intervals [19]. These long data-averaging intervals (of 60 seconds or more) are probably not used very often in preoperative CPETs and are not recommended by current preoperative CPET guidelines [8].

Based on the results of this study, the recommendation in the preoperative CPET guideline to use a breath-based data-averaging interval of 3–5 breaths or a time-based data-averaging interval of ~20 seconds seems plausible when evaluating the mean (group level) values. Nevertheless, caution should be taken when evaluating individual patients, as different data-averaging intervals caused substantial variation in the numerical values of CPET-derived variables within patients. As depicted in Fig 2, individual values of patients could differ as much as ~40%. In individual patients, the chosen data-averaging interval could induce a shift of that patient from low to high risk or vice versa. This is an important observation, as risk assessment could influence surgical planning for individual patients (e.g., enrollment in prehabilitation program, referring to a higher care unit postoperatively) and the shared decision-making process. It is recognized that preoperative risk assessment is not solely based on risk thresholds determined by CPET, but rather consists of a composite assessment, taking into account the full CPET in combination with other preoperative risk factors such as, but not limited to, malnutrition, comorbidities, and geriatric status. Nevertheless, the influence of the data-averaging interval could be taken into consideration, especially in patients in which the CPET values are close to the risk classification cut-off point. In addition, instead of rigid cut-off points inducing black and white risk assessment, grey zones (intermediate risk) could be introduced to account for individual differences [17].

A limitation of the current study was that VO_{2peak} was determined over a ~30 second interval [5] regardless of the data-averaging interval that was used. The use of the fixed 30 second interval might have masked some of the variability caused by the data-averaging interval, explaining the very small differences of VO_{2peak} values between data averaging intervals. A strength of the current study is that variation other than variation coming from the data-averaging interval was minimized. Firstly, by repeating interpretation of the 21 CPETs that were retrospectively formatted using 5 different data-averaging intervals, as opposed to repeated

testing of patients with different data averaging intervals. Secondly, to account for inter-observer variability, CPET interpretation was done by two clinical exercise physiologists, based on consensus, and by using a predefined set of guidelines (see [S2 File](#)). By doing so, the observed variation between groups of data-averaging intervals was exclusively caused by the used data-averaging interval and not by within-patient biological variation, measurement error, or inter-observer variability.

The current study opens possibilities for clinicians to be flexible in the data-averaging interval that is used for interpretation of the preoperative CPET. Current CPET literature does not provide clear and consistent guidance for clinicians about the choice of a data-averaging interval [7, 8]. As different (patho)physiological patterns might require different data-visualization, future research could focus on investigating optimal data-visualization methods that best fit the aim of the CPET, the properties of the CPET, and the (patho)physiological process the clinician is willing to evaluate.

Conclusion

On a group level there appear to be no clinically relevant differences in the mean values of VO_{2peak} , VO_{2VAT} , OUES, VE/VCO_{2VAT} , and VE/VCO_2 -slope between different data-averaging intervals used for interpretation of preoperative CPET in patients with colorectal cancer. In addition, the choice of data-averaging interval does not influence the proportion of patients classified as high or low risk for complications based on their exercise tolerance. Nevertheless, the chosen data-averaging interval might lead to substantial within patient variation for individual patients and should therefore be considered in patients in which the CPET values are close to the risk classification cut-off point.

Supporting information

S1 File. Graphical display of the Wasserman plots of patient 21.
(PDF)

S2 File. Guideline for systematic interpretation of preoperative cardiopulmonary exercise testing.
(PDF)

S1 Dataset.
(XLSX)

Acknowledgments

The authors like to thank Accuramed BVBA (Halen, Belgium) for providing a free unrestricted copy of the Blue Cherry software to support this study.

Author Contributions

Conceptualization: Ruud F. W. Franssen, Tim Takken, F. Jeroen Vogelaar, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

Data curation: Ruud F. W. Franssen, Bart H. E. Sanders.

Formal analysis: Ruud F. W. Franssen, Bart H. E. Sanders, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

Funding acquisition: Ruud F. W. Franssen, F. Jeroen Vogelaar, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

Investigation: Ruud F. W. Franssen, Bart C. Bongers.

Methodology: Ruud F. W. Franssen, Bart H. E. Sanders, Tim Takken, F. Jeroen Vogelaar, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

Project administration: Ruud F. W. Franssen, Bart H. E. Sanders.

Resources: Ruud F. W. Franssen.

Software: Ruud F. W. Franssen.

Supervision: F. Jeroen Vogelaar, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

Visualization: Ruud F. W. Franssen, Tim Takken, Bart C. Bongers.

Writing – original draft: Ruud F. W. Franssen, Bart H. E. Sanders, Bart C. Bongers.

Writing – review & editing: Ruud F. W. Franssen, Bart H. E. Sanders, Tim Takken, F. Jeroen Vogelaar, Maryska L. G. Janssen-Heijnen, Bart C. Bongers.

References

1. Moran J, Wilson F, Guinan E, McCormick P, Hussey J, Moriarty J. Role of cardiopulmonary exercise testing as a risk-assessment method in patients undergoing intra-abdominal surgery: a systematic review. *Br J Anaesth*. 2016; 116(2):177–91. <https://doi.org/10.1093/bja/aev454> PMID: 26787788
2. Steffens D, Ismail H, Denehy L, Beckenkamp PR, Solomon M, Koh C, et al. Preoperative Cardiopulmonary Exercise Test Associated with Postoperative Outcomes in Patients Undergoing Cancer Surgery: A Systematic Review and Meta-Analyses. *Ann Surg Oncol*. 2021; 28(12):7120–46. <https://doi.org/10.1245/s10434-021-10251-3> PMID: 34101066
3. Older P, Smith R, Courtney P, Hone R. Preoperative evaluation of cardiac failure and ischemia in elderly patients by cardiopulmonary exercise testing. *Chest*. 1993; 104(3):701–4. <https://doi.org/10.1378/chest.104.3.701> PMID: 8365279
4. Bongers BC, Dejong CHC, den Dulk M. Enhanced recovery after surgery programmes in older patients undergoing hepatopancreatobiliary surgery: what benefits might prehabilitation have? *Eur J Surg Oncol*. 2021; 47(3 Pt A):551–9. <https://doi.org/10.1016/j.ejso.2020.03.211> PMID: 32253075
5. American Thoracic S, American College of Chest P. ATS/ACCP Statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med*. 2003; 167(2):211–77.
6. Otto JM, Levett DZH, Grocott MPW. Cardiopulmonary Exercise Testing for Preoperative Evaluation: What Does the Future Hold? *Current Anesthesiology Reports*. 2020; 10(1):1–11.
7. Dumitrescu D, Rosenkranz S. Graphical Data Display for Clinical Cardiopulmonary Exercise Testing. *Ann Am Thorac Soc*. 2017; 14(Supplement_1):S12–S21. <https://doi.org/10.1513/AnnalsATS.201612-955FR> PMID: 28541745
8. Levett DZH, Jack S, Swart M, Carlisle J, Wilson J, Snowden C, et al. Perioperative cardiopulmonary exercise testing (CPET): consensus clinical guidelines on indications, organization, conduct, and physiological interpretation. *Br J Anaesth*. 2018; 120(3):484–500. <https://doi.org/10.1016/j.bja.2017.10.020> PMID: 29452805
9. West MA, Asher R, Browning M, Minto G, Swart M, Richardson K, et al. Validation of preoperative cardiopulmonary exercise testing-derived variables to predict in-hospital morbidity after major colorectal surgery. *Br J Surg*. 2016; 103(6):744–52. <https://doi.org/10.1002/bjs.10112> PMID: 26914526
10. Bongers BC, Berkel AE, Klaase JM, van Meeteren NL. An evaluation of the validity of the pre-operative oxygen uptake efficiency slope as an indicator of cardiorespiratory fitness in elderly patients scheduled for major colorectal surgery. *Anaesthesia*. 2017; 72(10):1206–16. <https://doi.org/10.1111/anae.14003> PMID: 28741667
11. Dideriksen K, Mikkelsen UR. Reproducibility of incremental maximal cycle ergometer tests in healthy recreationally active subjects. *Clin Physiol Funct Imaging*. 2017; 37(2):173–82. <https://doi.org/10.1111/cpf.12283> PMID: 26373336
12. Martin-Rincon M, Calbet JAL. Progress Update and Challenges on V. O₂max Testing and Interpretation. *Front Physiol*. 2020; 11:1070. <https://doi.org/10.3389/fphys.2020.01070> PMID: 33013459

13. Martin-Rincon M, Gonzalez-Henriquez JJ, Losa-Reyna J, Perez-Suarez I, Ponce-Gonzalez JG, de La Calle-Herrero J, et al. Impact of data averaging strategies on VO₂max assessment: Mathematical modeling and reliability. *Scand J Med Sci Sports*. 2019; 29(10):1473–88.
14. Franssen RFW, Bongers BC, Vogelaar FJ, Janssen-Heijnen MLG. Feasibility of a tele-prehabilitation program in high-risk patients with colon or rectal cancer undergoing elective surgery: a feasibility study. *Perioper Med (Lond)*. 2022; 11(1):28. <https://doi.org/10.1186/s13741-022-00260-5> PMID: 35879732
15. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007; 335(7624):806–8. <https://doi.org/10.1136/bmj.39335.541782.AD> PMID: 17947786
16. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007; 39(2):175–91. <https://doi.org/10.3758/bf03193146> PMID: 17695343
17. Rose GA, Davies RG, Davison GW, Adams RA, Williams IM, Lewis MH, et al. The cardiopulmonary exercise test grey zone; optimising fitness stratification by application of critical difference. *Br J Anaesth*. 2018; 120(6):1187–94. <https://doi.org/10.1016/j.bja.2018.02.062> PMID: 29793585
18. Midgley AW, McNaughton LR, Carroll S. Effect of the VO₂ time-averaging interval on the reproducibility of VO₂max in healthy athletic subjects. *Clin Physiol Funct Imaging*. 2007; 27(2):122–5. <https://doi.org/10.1111/j.1475-097X.2007.00725.x> PMID: 17309533
19. Johnson JS, Carlson JJ, VanderLaan RL, Langholz DE. Effects of sampling interval on peak oxygen consumption in patients evaluated for heart transplantation. *Chest*. 1998; 113(3):816–9. <https://doi.org/10.1378/chest.113.3.816> PMID: 9515862