

5-31-2024

## TENDINopathy severity assessment–achilles (TENDINS-A): Evaluation of reliability and validity in accordance with COSMIN recommendations

Myles C. Murphy  
*Edith Cowan University*

Fergus McCleary

Dana Hince

Ruth Chimenti

Paola Chivers

*See next page for additional authors*

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2022-2026>



Part of the [Sports Sciences Commons](#)

---

[10.1136/bjsports-2023-107741](https://doi.org/10.1136/bjsports-2023-107741)

This article has been accepted for publication in *British Journal of Sports Medicine*, 2024, following peer review, and the Version of Record can be accessed online at <https://doi.org/10.1136/bjsports-2023-107741>.

Murphy, M. C., McCleary, F., Hince, D., Chimenti, R., Chivers, P., Vosseller, J. T., . . . Rio, E. K. (2024). TENDINopathy severity assessment–achilles (TENDINS-A): Evaluation of reliability and validity in accordance with COSMIN recommendations. *British Journal of Sports Medicine*. Advance online publication. <https://doi.org/10.1136/bjsports-2023-107741>

This Journal Article is posted at Research Online.  
<https://ro.ecu.edu.au/ecuworks2022-2026/4080>

---

**Authors**

Myles C. Murphy, Fergus McCleary, Dana Hince, Ruth Chimenti, Paola Chivers, J. Turner Vosseller, Sophia Nimphius, Nonhlanhla S. Mkumbuzi, Peter Malliaras, Nicola Maffulli, Robert J. de Vos, and Ebonie K. Rio



1 **TENDINopathy Severity assessment – Achilles (TENDINS-A): Evaluation of reliability and validity in**  
2 **accordance with COSMIN recommendations.**

3

4 Myles C Murphy <sup>1,2</sup>, Fergus McCleary <sup>2</sup>, Dana Hince <sup>3</sup>, Ruth L Chimenti <sup>4</sup>, Paola T Chivers <sup>3,5</sup>, J. Turner  
5 Vosseller <sup>6</sup>, Sophia Nimphius <sup>5</sup>, Nonhlanhla Mkumbuzi <sup>7-10</sup>, Peter Malliaras <sup>11</sup>, Nicola Maffuli <sup>12-14</sup>, Robert-Jan  
6 De Vos <sup>15</sup>, Ebonie Rio <sup>16-18</sup>

7

8 <sup>1</sup> Nutrition and Health Innovation Research Institute, School of Medical and Health Sciences, Edith Cowan  
9 University, Joondalup, Western Australia, Australia

10 <sup>2</sup> School of Health Sciences and Physiotherapy, The University of Notre Dame Australia, Fremantle, Western  
11 Australia, Australia

12 <sup>3</sup> Institute for Health Research, The University of Notre Dame Australia, Fremantle, Western Australia, Australia

13 <sup>4</sup> Department of Physical Therapy and Rehabilitation Science, University of Iowa, Iowa City, Iowa, United  
14 States of America

15 <sup>5</sup> School of Medical and Health Sciences, Edith Cowan University, Joondalup, Western Australia, Australia

16 <sup>6</sup> Jacksonville Orthopaedic Institute, Jacksonville, Florida, United States of America

17 <sup>7</sup> Department of Sports, Exercise, and Rehabilitation, Northumbria University, Newcastle upon Tyne, United  
18 Kingdom

19 <sup>8</sup> Department of Rehabilitation, Midlands State University, Gweru, Zimbabwe

20 <sup>9</sup> Department of Human Movement Science, Nelson Mandela University, Summerstrand, Gqeberha, South  
21 Africa

22 <sup>10</sup> NtombiSport Pty Ltd, Cape Town, South Africa

23 <sup>11</sup> School of Primary and Allied Health Care, Faculty of Medicine Nursing and Health Science, Monash  
24 University, Victoria, Australia

25 <sup>12</sup> Department of Medicine, Surgery and Dentistry, University of Salerno, Baronissi, Salerno, Italy

26 <sup>13</sup> Faculty of Medicine, School of Pharmacy and Bioengineering, Keele University, Newcastle, United Kingdom

27 <sup>14</sup> Barts and the London School of Medicine and Dentistry, Centre for Sports and Exercise Medicine, Queen  
28 Mary University of London, London, United Kingdom

29 <sup>15</sup> Department of Orthopaedics and Sports Medicine, Erasmus MC University Medical Centre, Rotterdam, The  
30 Netherlands.

31 <sup>16</sup> La Trobe Sport and Exercise Medicine Research Centre, La Trobe University, Bundoora, Victoria, Australia

32 <sup>17</sup> Victorian Institute of Sport, Melbourne, Victoria, Australia

33 <sup>18</sup> The Australian Ballet, Melbourne, Victoria, Australia

34

35 **\*Corresponding author:** Dr Myles C Murphy. Nutrition and Health Innovation Research Institute, School of

36 Medical and Health Sciences, Edith Cowan University, Joondalup, Western Australia, Australia Email:

37 

38

39

40 **ABSTRACT**

41 **Objective:** To evaluate the construct validity (structural validity and hypothesis-testing), reliability (test-retest  
42 reliability, measurement error and internal consistency) and minimal important change (MIC) of the 13-item  
43 TENDINopathy Severity assessment–Achilles (TENDINS-A).

44 **Methods:** Participants with Achilles pain completed an online survey including: demographics, TENDINS-A,  
45 Foot and Ankle Outcome Score (FAOS), and Victorian Institute of Sport Assessment–Achilles (VISA-A).  
46 Exploratory factor analysis (EFA) assessed dimensionality. Confirmatory Factor Analysis (CFA) assessed  
47 structural validity [root-mean-squared error of approximation (RMSEA); Comparative Fit Index (CFI); Tucker-  
48 Lewis Index (TLI); Standardised Root Measure Square (SRMS)]. Correlations between TENDINS-A and the  
49 FAOS or VISA-A assessed hypothesis-testing. Intraclass correlation (ICC) assessed test-retest reliability.  
50 Cronbach’s  $\alpha$  assessed internal consistency. Standard error of the measurement (SEM) assessed measurement  
51 error. A distribution-based approach assessed MIC.

52 **Results:** Seventy-nine participants (51% female) with a mean (SD) age=42.6 (13.0) years, height=175.0 (11.7)  
53 cm and body mass=82.0 (19.1) kg were included. EFA identified three meaningful factors, proposed as pain,  
54 symptoms and function. The best model identified using CFA for TENDINS-A had structural validity  
55 (RMSEA= 0.101, CFI= 0.959, TLI= 0.947, SRMS=0.068), which included three factors (Pain, Symptoms, and  
56 Function), but excluded three items from the original TENDINS-A. TENDINS-A exhibited moderate positive  
57 correlation with FAOS ( $\rho=0.598, p<0.001$ ) and a moderate, negative correlation with VISA-A ( $r=-$   
58  $0.639, p<0.001$ ). Reliability of the TENDINS-A was excellent (ICC=0.930; Cronbach’s  $\alpha=0.808$ ; SEM=6.54  
59 units), with an MIC of 12 units.

60 **Conclusions:** Our evaluation of the revised 10-item TENDINS-A determined it has construct validity and  
61 excellent reliability, compared to the VISA-A and FAOS which lack content and construct validity. The  
62 TENDINS-A is recommended as the preferred patient-reported outcome measure to assess disability in people  
63 with Achilles tendinopathy.

64

65 **What is already known:** The TENDINopathy Severity assessment – Achilles (TENDINS-A) has been co-  
66 designed with patients, clinicians and researchers as a new patient-reported outcome measure (PROM) to assess  
67 the severity of disability in Achilles tendinopathy, but its psychometric properties have not been previously  
68 reported.

69 **What this study adds:** The TENDINS-A has content validity, construct validity and excellent reliability in the  
70 assessment of the severity of disability in Achilles tendinopathy.

71 **How this study might affect practice:** The TENDINS-A is recommended for use in both clinical practice and  
72 research to assess the severity of disability in Achilles tendinopathy.

73

## 74 INTRODUCTION

75 Achilles tendinopathy is characterised by focal Achilles tendon pain accompanied by impaired function with  
76 mechanical loading,<sup>1</sup> which is typically managed using predominantly exercise rehabilitation.<sup>2 3</sup> Achilles  
77 tendinopathy is common in the running population [prevalence of 4.2% (diagnosed using a pain mapping  
78 approach) in runners training for an event]<sup>4 5</sup> and also prevalent within the general population.<sup>6</sup> People with  
79 tendinopathy, expert clinicians and researchers have identified several core health domains as important in  
80 tendinopathy, including tendon-related disability.<sup>7</sup> However, multiple different measures to quantify tendon-  
81 related disability are used across studies, with no existing consensus.<sup>8 9</sup> In clinical practice measures of tendon-  
82 related disability are not commonly used,<sup>10</sup> in part due to inadequate content validity.<sup>11 12</sup> Thus, despite a  
83 consensus that tendon-related disability is an important aspect of health,<sup>7</sup> there remains no measure of tendon-  
84 related disability that is validated for research or clinical use.

85 The consensus-based standards for the selection of measurement instruments (COSMIN) guidelines represent  
86 international best-practice for the design and appraisal of Patient Reported Outcome Measures (PROMs).<sup>13 14</sup>  
87 However, the value of a PROM is not in its availability but in its quality, reflected by its clinometric  
88 properties.<sup>15</sup> Valid and reliable PROMs should be used and recommended, but this is not currently possible in  
89 Achilles tendinopathy research of the disability domain.<sup>16</sup> Existing PROMs that assess tendon-related disability,  
90 such as the Victorian Institute of Sport Assessment – Achilles (VISA-A),<sup>16 17</sup> have historically been  
91 recommended in the absence of other tools,<sup>18</sup> but do not satisfy the guidelines set out by COSMIN and suffer  
92 from substantial flaws in their design.

93 Using the VISA-A as an example, it has inadequate content and structural validity. VISA-A development did not  
94 involve patients<sup>11</sup> and methodological testing of the VISA-A scale using Rasch analysis has demonstrated it has  
95 insufficient structural validity<sup>19</sup> and hypothesis testing of the VISA-A shows it is significantly correlated to  
96 participant age and body mass index.<sup>20</sup> The recommendations from several reviews and original research studies  
97 highlight the VISA-A is not valid and should not be used in research or clinical practice.<sup>11 12 19 20</sup>

98 The newly developed TENDINopathy Severity assessment – Achilles (TENDINS-A) assesses the severity of the  
99 Tendinopathy Core Health Domain of Disability,<sup>7</sup> and consists of questions covering sub-domains of pain,  
100 symptoms, and physical function related to Achilles tendinopathy.<sup>21</sup> The TENDINS-A has adequate content  
101 validity, being co-designed by people with Achilles tendinopathy, expert clinicians and researchers.<sup>21</sup> Whilst the  
102 TENDINS-A has sufficient content validity, the structural validity and other measurement properties of this new  
103 PROM (e.g., reliability) remain unknown.

#### 104 **Objective**

105 The objective of this study was to evaluate the construct validity (including structural validity and hypothesis  
106 testing), reliability (including test-retest reliability, measurement error and internal consistency) and minimal  
107 important change of the TENDINS-A.

#### 108 **Hypotheses**

109 The TENDINS-A will have adequate criteria-based structural validity, demonstrate moderate significant  
110 correlations to commonly utilised PROMS for Achilles tendinopathy<sup>16</sup> [VISA-A and Foot and Ankle Outcome  
111 Score (FAOS)], no correlation to unrelated constructs such as baseline characteristics (e.g., age or body mass  
112 index) and excellent reliability.

#### 113 **METHODS**

##### 114 **Study Design**

115 Cross-sectional cohort study evaluating the psychometric properties of the TENDINS-A.

##### 116 **Participants and setting**

117 We used a network of greater than 20 clinicians known to the research team (exercise scientists, general  
118 practitioners, orthopaedic surgeons, physiotherapists, podiatrists, rheumatologists and sport and exercise  
119 physicians) and Achilles tendon researchers to identify participants with either mid-portion or insertional  
120 Achilles tendinopathy and provide them with our online survey using Qualtrics (convenience sampling). This  
121 was accessed via a quick response (QR) code or anonymous web link. Any person over the age of 18 years, who  
122 could read the English language, with self-reported Achilles tendinopathy was eligible.

##### 123 **Inclusion criteria**

124 Participants self-identified as having Achilles tendon pain and were provided with a pain map of established  
125 locations of Achilles tendon symptoms and asked to select all markings that corresponded to their region of pain  
126 (which is superior to both palpation and ultrasound tissue characterisation for pain localisation).<sup>22</sup> Participants



127 who selected the insertion of mid-portion Achilles tendon were included within this validation study and this  
128 methodology provides ‘near-perfect’ agreement to Sports Physician diagnosis of Achilles tendinopathy.<sup>23</sup>

### 129 **Outcome measures**

130 All outcome measures were completed by participants within a single survey and the survey forced responses to  
131 avoid missing data.

#### 132 *Participant characteristics*

133 Age (years), sex (male/female/intersex), height (cm), body mass (kg), ethnicity, country of residence, languages  
134 other than English spoken by the participant (self-reported), whether the participant performed moderate to  
135 vigorous physical activity (MVPA) most days (yes/no), highest level of education, work status and total  
136 household income were reported.

#### 137 *TENDINopathy Severity assessment – Achilles (TENDINS-A)*

138 The original TENDINS-A (which consists of 13 scoreable items numbered 3, 4, 6, 7, 8, 10, 11, 12, 13A, 13B,  
139 13C, 13D, 13E, and 4 non-scoreable items numbered 1, 2, 5, 9) was provided to participants as the first PROM  
140 within the survey and was scored between 0 and 100, with ‘0’ representing a perfect score (no disability) and  
141 ‘100’ representing complete disability.<sup>21</sup> If participants were unable to perform one of the pain-with-loading  
142 tests (e.g., single leg hop), they were instructed to leave it blank and a score of ‘10’ was provided.

143 Participants completed the TENDINS-A prior to the FAOS and the VISA-A. Directly after completing the  
144 VISA-A questionnaire, the TENDINS-A was immediately repeated. This ensured the clinical status of  
145 participants was unchanged,<sup>24</sup> whilst still allowing for any potential interference effects. Specifically, the  
146 participant would be unable to remember responses to the initial TENDINS-A as they have no option to go back  
147 in the survey and 50 questions (i.e., the VISA-A and FAOS) separated the initial and then repeated TENDINS-  
148 A.

#### 149 *Foot and Ankle Outcome Score*

150 The 42-item FAOS<sup>25</sup> was performed following the TENDINS-A, with a mean total score calculated for the  
151 purposes of this study and used for hypothesis testing. A score of ‘0’ represented a perfect score (no disability),  
152 with a score of ‘100’ representing complete disability. The FAOS is a reliable tool and has adequate internal  
153 consistency,<sup>25</sup> however it has inadequate content validity (as it was not developed for Achilles tendinopathy) and  
154 its construct validity in Achilles tendinopathy is unknown.

#### 155 *Victorian Institute of Sport Assessment – Achilles*

156 The 8-item VISA-A<sup>17</sup> was performed following the FAOS and used for hypothesis testing. The VISA-A is  
 157 inversely scored when compared to the TENDINS-A and FAOS with a score of '100' represented a perfect score  
 158 (no disability), and a score of '0' representing complete disability. Where participants selected multiple  
 159 responses in item 8 (related to how much physical activity can be performed before cessation due to symptoms  
 160 is required), the lowest score was retained for analysis. The VISA-A is a reliable tool and has adequate internal  
 161 consistency,<sup>17</sup> however it has inadequate content validity<sup>11</sup> and construct validity.<sup>19</sup>

162 **Power calculation**

163 Study size was informed by the recommendations of the COSMIN risk of bias checklist<sup>13 14</sup> rather than formal  
 164 power calculations. Questions 1, 2, 5 and 9 are not scored in the TENDINS-A; therefore the PROM was initially  
 165 considered to have 13 scorable items as one item has five secondary scales.<sup>21</sup> COSMIN guidelines suggest that  
 166 six-times the number of persons to items is adequate for assessing structural validity using classical test theory,<sup>13</sup>  
 167 <sup>14</sup> thus the minimum sample size for this study was determined to be 78 persons. We also required all  
 168 communalities in exploratory factor analysis (EFA) to be >0.3, thus samples of <100 participants are justified.<sup>26</sup>  
 169 Furthermore, a sample of >50 participants is considered adequate for reliability and internal consistency.<sup>13 14</sup>

170 **Statistical analysis**

171 The different statistical analyses, per each measurement property, are described below. All statistical analyses  
 172 were performed within IBM SPSS statistics (version 29.0), or IBM SPSS Amos (version 29.0) and we have  
 173 provided a glossary of statistical terminology used in Table 1. Where appropriate, confidence intervals are  
 174 presented, and statistical significance was considered when  $p < 0.05$ .

175 For the purposes of clarity in this manuscript, when referencing an Item (e.g., Item 7) these will always be in  
 176 reference to the original 13 items of the TENDINS-A. However, item numbers for the final TENDINS-A  
 177 (presented as an appendix) are different and have been re-ordered to avoid confusion (e.g., labelled 1-10).

178 **Table 1. Glossary of statistical terminology**

| Term                          | Definition  |
|-------------------------------|---|
| Bartlett's test of Sphericity | Assesses the probability that the correlation matrix has significant correlations amongst items in the dataset, which is a requirement for factor analysis. |
| Communality                   | Proportion of the common variance for an individual item, relative to all factors.  |
| Comparative Fit Index         | Assessment of model fit by quantifying discrepancies between the sample data and the model while adjusting for sample size.                                 |

|   |  |
|---|--|
| Eigenvalues                                     | Quantification of the proportion of the variance accounted for by an individual factor.  |
| Kaiser-Meyer-Olkin Measure of Sampling Adequacy | An assessment of whether data is appropriate for factor analysis by determining the proportion of variance among all items that might be from a common variance. |
| Root-mean-squared error of the approximation    | Estimate of the discrepancy the sample data and the model-implied data covariance matrices per degree of freedom.  |
| Scree Plot                                      | Graphical representation of eigenvalues to visually assess when new factors are unlikely to explain a sufficient level of the variance.                          |
| Standardised Root Measure Square                | Square root of the calculated discrepancy between the sample covariance matrix and the model covariance matrix.  |
| Tucker-Lewis Index                              | Measure of goodness of fit in the analysis of covariance structures.   |

179

180 *Dimensionality*

181 We performed EFA for all scoreable items using SPSS Statistics to assess dimensionality. To proceed to  
182 confirmatory factor analysis (CFA), we required all communalities to exceed 0.3,<sup>27</sup> required the Kaiser-Meyer-  
183 Olkin Measure of Sampling Adequacy to exceed 0.7<sup>28</sup> and for Bartlett's test of Sphericity to be significant  
184 ( $p < 0.01$ ),<sup>29</sup> which are all recognised criteria for progression to factor analysis.

185 Factors were eligible for inclusion within EFA where the Eigenvalues exceeded 1.0 (which are deemed  
186 substantial). The Scree plot was also visually inspected and additional models were performed to include more/  
187 less factors, based on author (MCM) judgement and with the TENDINS-A being proposed to measure three  
188 factors related to disability: pain, symptoms and function.<sup>21</sup> A principal axis factoring method was used to  
189 determine our initial factor matrix. As per standard practice, factors were considered meaningful when an item  
190 loaded at  $\geq 0.4$  for that factor.<sup>30</sup> As 'disability' is a domain that reflects pain, symptoms and function,<sup>7</sup> we  
191 hypothesised the TENDINS-A to have three factors identified by EFA.

192 *Structural Validity*

193 Confirmatory factor analysis was performed using SPSS Amos (22) to assess structural validity. We included all  
194 three factors (which were identified in the EFA and TENDINS-A content validity study),<sup>21</sup> within the initial  
195 model, and then removed variables as needed to achieve the best fit. Items with categorical (yes/ no only)

196 responses were not included within CFA (not appropriate for CFA but had been tested in EFA). Models were  
197 tested and selected based off the best root-mean-squared error of the approximation (RMSEA) and overall  
198 model chi-squared statistics, with a lower chi-squared statistic representing a better model fit. An overall  
199 acceptable model would require a Comparative Fit Index (CFI) OR Tucker-Lewis Index (TLI) of >0.95 AND  
200 RMSEA of <0.06 or Standardised Root Measure Square (SRMS) of <0.08.<sup>13</sup>

201 The CFA provided standardised factor loading and error variance per item. Composite reliability of >0.70 is  
202 considered adequate.<sup>31</sup> Composite reliability was calculated using the formula below when  $\lambda$  = standardised  
203 factor loading and  $\varepsilon$  = error variance:

204 
$$\text{Composite reliability} = \frac{(\sum \lambda_i)^2}{(\sum \lambda_i)^2 + (\sum \varepsilon_i)}$$

205 Convergent validity was assessed using the average variance extracted (AVE). An AVE of >0.50 is considered  
206 adequate.<sup>32</sup> The AVE was calculated using the formula below when  $\lambda$  = standardised factor loading and  $\varepsilon$  = error  
207 variance:

208 
$$\text{Average variance extracted} = \frac{\sum_{i=1}^n \lambda_i^2}{\sum_{i=1}^n \lambda_i^2 + \sum_{i=1}^n \text{Var}(\varepsilon_i)}$$

209  
210 The criteria for adequate structural validity of the final model were:

- 211 1. CFI or TLI >0.95,<sup>13</sup>
- 212 2. RMSEA of <0.06 or SRMS of <0.08,<sup>13</sup>
- 213 3. Composite Reliability >0.70,<sup>31</sup> and
- 214 4. AVE >0.50.<sup>32</sup>

#### 215 *Test-retest reliability*

216 Test-retest reliability of the TENDINS-A was calculated as a continuous scale. The absolute and relative test-  
217 retest reliability (which is a sub-type of reliability and demonstrates how closely someone will generate the  
218 same score on a PROM when repeated in a specified timeframe) were determined.<sup>33</sup> Relative test-retest  
219 reliability was reported as the intraclass correlation (ICC), with a two-way mixed approach used to assess  
220 absolute agreement of a single measure.<sup>34</sup> Absolute test-retest reliability is a sub-type of reliability and  
221 demonstrates the degree of uncertainty in a measurement. For example, measurements with more uncertainty

222 will have greater measurement error, hence it is systematic and random error that is not reflective of true  
223 change.<sup>34</sup> The standard error of the measurement<sup>35</sup> was calculated using the following equation:

224 
$$\text{Standard error of the measurement} = SD_{\text{baseline}}\sqrt{1-ICC}$$

225 *Minimal important change*

226 The Minimal important change (MIC) and 95% confidence intervals was calculated using a distribution-based  
227 method,<sup>36</sup> using the following equation:

228 
$$\text{Minimal important change} = \frac{SD_{\text{baseline}}}{2}$$

229 *Internal consistency*

230 Internal consistency is a sub-type of reliability that investigates whether different items that measure the same  
231 construct give comparable outcomes. The internal consistency for the entire 13-item TENDINS-A was  
232 calculated using the average inter-item correlation and a Cronbach's Alpha was reported with a positive rating  
233 given for values >0.7.<sup>13</sup>

234 *Convergent and divergent validity*

235 In addition to the convergent validity assessed using CFA, we also assessed convergent and divergent validity of  
236 the TENDINS-A scale using a correlation between scores of the TENDINS-A and another PROM theorised to  
237 measure the same overall core health domain of tendinopathy (disability)<sup>7 16</sup> or unrelated constructs, such as age  
238 and BMI (given they are continuous variables that should not be related to tendon-related disability). The  
239 TENDINS-A, FAOS and the VISA-A were assessed for normality using visual assessment and the one-sample  
240 Kolomogorov-Smirnov Test. For normally distributed data the Pearson's correlation coefficient was used and for  
241 non-normally distributed data the non-parametric Spearman rho correlation was used. Correlations were  
242 considered very weak (between 0-0.25), weak (0.26-0.49), moderate (0.5-0.69), strong (0.7-0.89) and very  
243 strong (0.9-1).

244 *Comparison between sub-groups*

245 To ensure that significant difference between sub-groups did not exist we calculated between-group differences  
246 for key baseline characteristics [sex (male/ female), MVPA most days (yes/no), multilingual status (yes/ no), and  
247 whether tendon loading activities are typically performed daily (yes/no)] using independent t-tests.

248 **Equity, diversity and inclusion**

249 The authorship group are of diverse gender, geographical location and research experience. Furthermore, the  
250 participants included in this study were diverse in relation to many demographic variables (e.g., sex, age, BMI,  
251 geographical location or education level).

252

## 253 RESULTS

### 254 Participants

255 Seventy-nine (n=79) participants (51% female, 49% male) provided survey data and were included within our  
256 analysis with no missing data. Participants had a mean (SD) age of 42.6 (13.0) years, height of 175.0 (11.7) cm  
257 and body mass of 82.0 (19.1) kg. Participants were predominantly of 'Australian' ethnicity (79.7%) and were  
258 not multi-lingual (64.6%) with those multilingual participants predominantly having English as their first  
259 language (53.6%). A significant number of participants reported that they performed MVPA most days (69.6%).  
260 Participant education level and employment varied, with most having completed tertiary studies (74.6%) and  
261 working full time (63.3%). Income levels ranged from less than \$30,000 Australian Dollars (AUD) (5.1%) to  
262 greater than \$200,000 AUD (27.8%), per annum. The activities that aggravated the pain of participants with  
263 Achilles tendinopathy varied: walking slow (15.2%), walking fast (44.3%), walking up and down stairs (43.0%),  
264 running up and down stairs (46.8%), running slow (60.8%), running fast (54.4%), hopping and jumping  
265 (62.0%), and rapidly changing direction while running (44.3%). The complete breakdown of participant  
266 characteristics is provided in Appendix A.

### 267 Uni-dimensionality

268 Bartlett's test of Sphericity suggested that the results of our correlation matrix were not random [ $\chi^2(78) = 708$ ,  
269  $p < 0.001$ ]. Our Kaiser-Meyer-Olkin Measure of Sampling Adequacy (0.811) far exceeded the minimum cut-off  
270 of 0.70. Thus, our correlation matrix was appropriate. The extracted communalities (Mean=0.67, SD= 0.16,  
271 Range= 0.18 to 0.93) for all but three items (items 6, 8 and 12) exceeded our cut-off score for CFA inclusion  
272 (Appendix B).

273 Principal axis factoring analysis was performed and the initial Eigen values of three factors exceeded 1.0 (see  
274 appendix C), explained 67.9% of the total variance and the first factor accounted for >20% of the variability.  
275 The three identified factors were included within the Factor Matrix, and associated factor loading per item, are  
276 presented within Table 2. However, no items were associated with a loading of >0.4 on Factor Three.

277 **Table 2. Factor Loading with Principal Axis Factoring Analysis for all items with loading >0.4 (n=79).**

| TENDINS- A | Factor One | Factor Two | Factor Three |
|------------|------------|------------|--------------|
| Item 3     | 0.412      |            | 0.400        |
| Item 4     | 0.496      | 0.479      |              |
| Item 6     |            |            |              |

|          |       |       |  |
|----------|-------|-------|--|
| Item 7   |       | 0.644 |  |
| Item 8   |       | 0.504 |  |
| Item 10  |       | 0.672 |  |
| Item 11  | 0.614 |       |  |
| Item 12  |       |       |  |
| Item 13A | 0.731 |       |  |
| Item 13B | 0.888 |       |  |
| Item 13C | 0.915 |       |  |
| Item 13D | 0.916 |       |  |
| Item 13E | 0.917 |       |  |

278 \*Blank cells represent loading of <0.4

279 **Structural Validity** Several Maximum Likelihood Estimates CFA models were trialled, informed by the factor  
280 loading from EFA, and subscales from our content validity study to generate our final CFA model. The best  
281 model used three factors [a) Pain, b) Symptoms, and c) Function] and eight items (Figure 1). Question 12 was  
282 excluded because of poor EFA initial communalities, non-meaningful factor loadings within EFA and being unable  
283 to load onto either of the three factors within CFA. Question 13A and 13B were removed as their exclusion  
284 resulted in substantially improved model fit (Appendix D). The scoring for the remaining 12 questions was  
285 amended to maintain a score range from 0 to 100, as presented within Appendix E.

286 In the final model, the chi-squared statistic was  $\chi^2(22) = 39.2$ , the CFI was 0.959, the TLI was 0.947, the  
287 RMSEA was 0.101 and the SRMS was 0.068, indicating sufficient structural validity. The model was also  
288 assumed accurate with CMIN/DF = 1.796, parsimony comparative fit index = 0.753 and parsimony normed fit  
289 index = 0.717.

290 The standardised factor loadings and error variance for the final model are included within Table 3. The  
291 correlation between 'Pain' and 'Symptoms' was estimated at 0.642 with significant covariances (Estimate =  
292 4.104, standard error = 1.101 and composite reliability = 3.728,  $p < 0.001$ ). The correlation between 'Pain' and  
293 'Function' was estimated at 0.668 with significant covariances (Estimate = 5.424, standard error = 1.300 and  
294 composite reliability = 4.171,  $p < 0.001$ ). The correlation between 'Symptoms' and 'Function' was estimated at  
295 0.275 with significant covariances (Estimate = 2.818, standard error = 1.353 and composite reliability = 2.082,  
296  $p = 0.037$ ).

297 **Table 3. Standardised Factor Loadings of the TENDINS-A (n=79)**

| TENDINS- A | Factor   | Standardised Factor Loading | Error Variance |
|------------|----------|-----------------------------|----------------|
| Item 11    | Pain     | 0.643                       | 0.587          |
| Item 3     | Pain     | 0.605                       | 0.634          |
| Item 4     | Pain     | 0.627                       | 0.607          |
| Item 7     | Symptoms | 0.741                       | 0.451          |
| Item 10    | Symptoms | 0.855                       | 0.269          |
| Item 13C   | Function | 0.921                       | 0.152          |
| Item 13D   | Function | 0.975                       | 0.049          |
| Item 13E   | Function | 0.967                       | 0.065          |

298 *Internal consistency.* For the 8-items assessed using CFA included in TENDINS-A theorised to measure  
 299 ‘disability’, the internal consistency was 0.842, hence displaying sufficient internal consistency.

300 *Composite reliability.* For the 8-items assessed using CFA included in TENDINS-A theorised to measure  
 301 ‘disability’, the composite reliability was 0.934, hence displaying sufficient composite reliability.

302 *Convergent validity.* For the 8-items assessed using CFA included in TENDINS-A theorised to measure  
 303 ‘disability’, the AVE was 0.648, hence displaying sufficient convergent validity.

304 Thus, our final CFA model could determine the contribution of different items in measuring ‘Pain’, ‘Symptoms’  
 305 and ‘Function’ and the TENDINS-A has sufficient structural validity. Consequently, the scores for the remaining  
 306 items were amended to ensure the TENDINS-A retained a maximal score of 100 (40 units for ‘Pain’, 30 units  
 307 for ‘symptoms’ and 30 units for ‘function’). Questions 6 and 8 were also excluded from CFA modelling being  
 308 dichotomous (Yes/No) items but were retained within the complete TENDINS-A in the ‘symptoms’ domain. All  
 309 subsequent analyses therefore relate to the revised TENDINS-A, presented in Appendix E.



310 **Distribution of data**

311 Whilst data for the TENDINS-A (Kolomogorov-Smirnov test statistic=0.09, p=0.097) and the VISA-A  
312 (Kolomogorov-Smirnov test statistic= 0.06, p=0.200) were normally distributed, the FAOS were not  
313 (Kolomogorov-Smirnov test statistic= 0.12, p=0.019). The TENDINS-A data had a minimum score of 0 and a  
314 maximum score of 100. The mean (SD) of the TENDINS-A was 47.89 (24.71) points, the FAOS was 23.60  
315 (18.63) percent, and the VISA was 53.58 (23.96) points. The mean (SD) of the repeated TENDINS-A was 44.34  
316 (24.16) points.

317 **Reliability**

318 *Intraclass correlation co-efficient*

319 The 10-item TENDINS-A mixed-effect ICC for absolute agreement of a single measure was ICC= 0.930 (95%  
320 CI = 0.881 to 0.959), p= < 0.001.

321 *Internal consistency*

322 The Cronbach's Alpha for the 10-items of the TENDINS-A was reported as  $\alpha=0.808$ , representing excellent  
323 internal consistency.

324 *Standard error of the measurement*

325 The TENDINS-A standard error of the measurement was calculated as 6.54 units.

326 **Minimal important change**

327 The MIC for the 10-item TENDINS-A was calculated as 12.36 (95% CI= 5.46 to 19.25) units of difference,  
328 representing 25.6% points of change from a mean TENDINS-A score of 47.89.

329 **Convergent validity**

330 The TENDINS-A exhibited a moderate positive correlation with the FAOS ( $\rho=0.598$ , 95%CI= 0.408 to 0.738,  
331  $p<0.001$ ), and a moderate, negative correlation with the VISA-A ( $r=-0.639$ , 95%CI= -0.764 to -0.467,  $p<0.001$ ).

332 **Divergent validity**

333 The TENDINS-A showed no evidence of a statistically significant association with age ( $p=0.426$ ) or BMI  
334 ( $p=0.189$ ).

335 **Comparison between sub-groups**

336 Between-group differences (Table 4) were observed for self-reported MVPA most days, with those performing  
337 moderate to vigorous physical activity most days having a lower TENDINS-A score ( $p=0.002$ ) and males  
338 reported a lower TENDINS-A score ( $p=0.044$ ). No differences in TENDINS-A score were observed for those  
339 performing specific tendon-loading exercise versus those who did not ( $p=0.079$ ), or whether participants were

340 multi-lingual ( $p=0.397$ ). As more males performed MVPA than females in our sample, we performed a *post hoc*  
 341 linear regression model to assess whether group differences due to sex were a result of higher levels of MVPA in  
 342 males than females. The model supported this idea, with no association between the TENDINS-A score and sex  
 343 ( $p=0.061$ ) after adjusting for MVPA [17.7 (95%CI= 6.46 to 28.97) unit increase in TENDINS-A score,  
 344  $p=0.002$ ].

345 **Table 4. Between group differences in TENDINS-A score.**

|   |                  | Mean (SD),<br>range    | Mean<br>Difference<br>(SED) | Test statistic<br>(Independent t-<br>test) | p-value |
|---|------------------|------------------------|-----------------------------|--|---------|
| Moderate to<br>vigorous<br>physical activity<br>most days | Yes (n=55)       | 37.69 (23.08),<br>0-91 | -18.81 (5.91)               | -3.18                                      | 0.002*  |
|   | No (n=24)        | 56.50 (26.54),<br>5-98 |                             |  |         |
| Sex   | Male (n=39)      | 37.56 (23.27),<br>5-98 | -11.54 (5.63)               | -2.048                                     | 0.044*  |
|   | Female<br>(n=40) | 49.10 (26.64),<br>0-91 |                             |  |         |
| Tendon-loading<br>exercise<br>performed most<br>days      | Yes (n=43)       | 38.79 (22.35),<br>5-86 | -10.13 (5.69)               | -1.779                                     | 0.079   |
|   | No (n=36)        | 48.92 (28.23),<br>0-98 |                             |  |         |
| Multilingual  | Yes (n=28)       | 48.79 (23.57),<br>5-91 | 8.34 (5.97)                 | 1.396                                      | 0.167   |
|   | No (n=51)        | 40.45 (26.32),<br>0-98 |                             |  |         |

346

347 Abbreviations: SD= standard deviation; SED= standard error of difference; n= number; range= min to max

348 \*  $p<0.05$

349 **DISCUSSION**

350 The present investigation evaluated the construct validity (consisting of structural validity and hypothesis  
351 testing), as well as reliability (consisting of internal consistency, test-retest reliability and measurement error) of  
352 the TENDINS-A. Based on this investigation, we present a revised TENDINS-A (Appendix E) that still includes  
353 all original questions, but only 10 of the original questions are scoreable.

354 The 10-item TENDINS-A, proposed to provide an overall measure of ‘disability’, demonstrated sufficient  
355 construct validity for three factors (pain; symptoms; function) and reliability, supporting its use in clinical and  
356 research settings. Furthermore, with normally distributed scores, there was no evidence of floor or ceiling  
357 effects within the sample. The TENDINS-A has a median completion time of ~8 minutes and is also likely to be  
358 meaningful to people with a lived experience of Achilles tendinopathy, as the research team included such  
359 individuals to inform its development.<sup>21</sup>

360 All continuous variables of the TENDINS-A, after the removal of item 12, 13A and 13B, were able to load onto  
361 one of three factors: pain, symptoms or function. The core health domain of ‘disability’ is defined as the  
362 “composite scores of a mix of patient-rated pain and disability due to the pain, usually relating to tendon-  
363 specific activities/tasks,” and included in tools such as the VISA-A.<sup>7 16</sup> Thus, the TENDINS-A is proposed to  
364 have adequate structural validity and measure the overall composite of disability, using scores from items  
365 related to ‘pain’, ‘symptoms’ and ‘function’. Therefore, the TENDINS-A is the only current PROM quantifying  
366 Achilles tendinopathy related disability with adequate content<sup>21</sup> and structural validity and should be used in  
367 place of all other PROMs. Researchers may still wish to use the VISA-A in addition to the TENDINS-A to allow  
368 for comparisons to previous research, but only the TENDINS-A should be used to report primary outcomes.

369 The reliability of the TENDINS-A is excellent ( $ICC > 0.90$ ). We opted to assess test-retest reliability on a single  
370 day as tendinopathy pain and symptoms are known to fluctuate daily.<sup>24</sup> The internal consistency of the  
371 TENDINS-A was also excellent ( $\alpha > 0.80$ ), ensuring all items were being scored consistently. The terminology  
372 surrounding the minimal detectable change, MIC and minimal clinically important difference (MCID) is  
373 confusing and for the purposes of discussion we have included all available comparisons. We calculated the  
374 minimal important change (MIC=12.4 units) of the TENDINS-A using a distribution-based method. This is  
375 larger than the MIC for the VISA-A, which is reported as between 7.8 to 8.2.<sup>37</sup> A study using an anchor-based  
376 method in the VISA-A reported the MCID as 14 units,<sup>38</sup> which is comparable to our estimated measure. Another  
377 study using the VISA-A estimated the MCID to be 6.5 units, however this was estimated from a selective  
378 sample of 15 participants with insertional Achilles tendinopathy, which may also explain the smaller MCID.<sup>39</sup>

379 To characterize the magnitude of change on the TENDINS-A that is meaningful to individuals with Achilles  
380 tendinopathy, future longitudinal studies are needed to estimate the minimal clinically-important difference  
381 (MCID).

382 The TENDINS-A demonstrated moderate correlations to both the VISA-A and FAOS, which is expected given  
383 they are proposed to measure the same overall health domain of disability.<sup>16</sup> We did not expect strong  
384 correlations to the VISA-A or FAOS given the limitations of the VISA-A,<sup>11 12 15</sup> which we theorised would  
385 extend to the FAOS as it is also lacking content validity. This lack of a strong correlation may be due to the  
386 TENDINS-A being a superior PROM to the FAOS and VISA-A (given the established content and structural  
387 validity of the TENDINS-A). The lack of a strong correlation can also result from the absence of a gold standard  
388 outcome measure for disability. Alternatively, we theorised the TENDINS-A should not have been associated  
389 with baseline characteristics such as age or BMI, and this was confirmed by our analysis. This differs from the  
390 VISA-A, whose total score is significantly associated with age and BMI,<sup>20</sup> reflecting a problem for previous  
391 studies using the VISA-A that did not adjust for age or BMI within the statistical analysis.

392 None of our specified sub-groups (sex, being multilingual, MVPA most days, tendon-loading exercise most  
393 days) showed a floor or ceiling effect. We expect this is likely due to items of the TENDINS-A being  
394 specifically linked to the function and aggravating tasks typically performed by the participant, making it  
395 suitable for both sedentary and physically active people with Achilles tendinopathy. Performing MVPA most  
396 days was associated with a lower TENDINS-A score, as expected. This is most likely a consequence of the  
397 numerous health benefits of physical activity.<sup>40</sup> Alternatively, the inactive group may undertake less tendon-  
398 specific loading exercise and aggravate the Achilles less. However, this seems unlikely, as performing tendon-  
399 loading activity was also assessed and no significant effect detected.

#### 400 **Limitations**

401 The sample for this study, whilst diverse in some respects, was heavily reliant on participants living in Australia.  
402 While other PROMS for Achilles tendinopathy showed acceptable cross-cultural validity in other languages,<sup>41 42</sup>  
403 future research on cross-cultural adaptation of the TENDINS-A is needed. Cross-cultural adaptation of the  
404 TENDINS-A is currently being performed by several different research groups internationally and these data  
405 can also be used to assess differential item functioning.

406 Self-report may have resulted in a misdiagnosis of Achilles tendinopathy. However in this study, we enhanced  
407 diagnostic accuracy by implementing a standardized pain map.<sup>22</sup> A prior investigation demonstrated a 93%  
408 concordance between patient-reported Achilles tendinopathy when using a pain map and diagnoses made by a  
409 physician.<sup>23</sup> These findings suggest that, in the majority of cases, self-reported Achilles tendinopathy aligns with  
410 the clinical diagnosis.

411 Finally, further details related to medical co-morbidities and medication usage<sup>43</sup> would have allowed analysis of  
412 their influence on total score and would be recommended in future research trials.

### 413 **CONCLUSION**

414 Our evaluation of the revised TENDINS-A (Appendix E), which includes 10-scoreable items across three sub-  
415 domains of pain, symptoms and physical function has adequate construct validity and reliability. These findings,  
416 and previously established content validity, ensure that the revised TENDINS-A with 10-scoreable items can be  
417 recommended for immediate use in both research and clinical practice, being the preferred tool over the VISA-A  
418 and FAOS to assess disability in individuals with Achilles tendinopathy.

419

420 **Author Contributions:** Myles C Murphy, Dana Hince and Ebonie K Rio conceived the study and designed the  
421 methods. Fergus McCleary collected all data. Myles C Murphy, Dana Hince, Paola T Chivers and Sophia  
422 Nimphius performed all analysis. All authors contributed to interpretation and manuscript preparation.

423 **Competing Interests and Funding:** No authors have any competing interests, and this research was not  
424 funded.

425 **Acknowledgements:** Dr Myles C Murphy is supported by a Western Australian Department of Health  
426 Innovation Fellowship (G1006615) and a Near-Miss Award (G1006605).

427 **Data Sharing:** The data from this research study will be considered to be shared upon reasonable request of the  
428 corresponding author.

429 **Ethical Approvals:** This research was approved by the University of Notre Dame Australia Human Research  
430 Ethics Committee (ID: 2022-175F) and all participants gave informed, electronic consent.

431 **Patient involvement:** The TENDINS-A was co-designed with people having a lived experience of Achilles  
432 tendinopathy and was found to have adequate content validity, ensuring that the TENDINS-A is relevant to  
433 patients.

434 **Equity, diversity and inclusion:** The authorship group are of diverse gender, geographical location and  
435 research experience. Furthermore, the participants included in this study were diverse in relation to many  
436 demographic variables (e.g., sex, age, BMI, geographical location or education level).

437

438 **REFERENCES**

- 439 1. Scott A, Squier K, Alfredson H, et al. ICON 2019: International Scientific Tendinopathy Symposium  
440 Consensus: Clinical Terminology. *British journal of sports medicine* 2020;54(5):260-62. doi:  
441 10.1136/bjsports-2019-100885
- 442 2. Murphy M, Travers MJ, Gibson W, et al. The rate of improvement of pain and function in mid-portion  
443 Achilles tendinopathy with loading protocols: A Systematic Review and Longitudinal Meta-Analysis.  
444 *Sports medicine (Auckland, NZ)* 2018;48(8):1875-91. doi: 10.1007/s40279-018-0932-2.
- 445 3. Murphy MC, Travers MJ, Chivers P, et al. Efficacy of heavy eccentric calf training for treating mid-portion  
446 Achilles tendinopathy: a systematic review and meta-analysis. *British journal of sports medicine*  
447 2019;53(17):1070-77. doi: 10.1136/bjsports-2018-099934
- 448 4. Chen W, Cloosterman KLA, Bierma-Zeinstra SMA, et al. Epidemiology of insertional and midportion  
449 Achilles tendinopathy in runners: A prospective cohort study. *Journal of sport and health science* 2023  
450 doi: 10.1016/j.jshs.2023.03.007 [published Online First: 20230323]
- 451 5. Lopes AD, Hespanhol LC, Yeung SS, et al. What are the Main Running-Related Musculoskeletal Injuries?  
452 *Sports medicine (Auckland, NZ)* 2012;42(10):891-905. doi: 10.1007/bf03262301
- 453 6. Albers IS, Zwerver J, Diercks RL, et al. Incidence and prevalence of lower extremity tendinopathy in a Dutch  
454 general practice population: a cross sectional study. *BMC Musculoskelet Disord* 2016;17(1):16. doi:  
455 10.1186/s12891-016-0885-2
- 456 7. Vicenzino B, de Vos R-J, Alfredson H, et al. ICON 2019—International Scientific Tendinopathy Symposium  
457 Consensus: There are nine core health-related domains for tendinopathy (CORE DOMAINS): Delphi  
458 study of healthcare professionals and patients. *British journal of sports medicine* 2019;54(8):444-51.  
459 doi: 10.1136/bjsports-2019-100894
- 460 8. Murphy M, Rio E, Debenham J, et al. Evaluating the progress of mid-portion Achilles Tendinopathy during  
461 rehabilitation. A Review of Outcome Measures for muscle structure and function, tendon structure and  
462 neural and pain associated mechanisms. *Int J Sports Phys Ther* 2018;13(3):537-51.
- 463 9. Murphy M, Rio E, Debenham J, et al. Evaluating the progress of mid-portion Achilles Tendinopathy during  
464 rehabilitation. A Review of Outcome Measures for Self- Reported Pain and Function. *Int J Sports Phys*  
465 *Ther* 2018;13(2):283-92. doi: 10.26603/ijsp20180283

- 466 10. Murphy M, Debenham J, Bulsara C, et al. Assessment and monitoring of Achilles Tendinopathy in clinical  
467 practice: A qualitative descriptive exploration of the barrier's clinicians face. *BMJ Open Sport Exerc*  
468 *Med* 2022;ePub ahead of print
- 469 11. Korakakis V, Kotsifaki A, Stefanakis M, et al. Evaluating lower limb tendinopathy with Victorian Institute of  
470 Sport Assessment (VISA) questionnaires: a systematic review shows very-low-quality evidence for  
471 their content and structural validity-part I. *Knee Surg Sports Traumatol Arthrosc* 2021;29(9):2749-64.  
472 doi: 10.1007/s00167-021-06598-5 [published Online First: 20210521]
- 473 12. Korakakis V, Whiteley R, Kotsifaki A, et al. A systematic review evaluating the clinimetric properties of the  
474 Victorian Institute of Sport Assessment (VISA) questionnaires for lower limb tendinopathy shows  
475 moderate to high-quality evidence for sufficient reliability, validity and responsiveness—part II. *Knee*  
476 *Surg Sports Traumatol Arthrosc* 2021 doi: 10.1007/s00167-021-06557-0
- 477 13. Gagnier JJ, Lai J, Mokkink LB, et al. COSMIN reporting guideline for studies on measurement properties of  
478 patient-reported outcome measures. *Qual Life Res* 2021;30(8):2197-218. doi: 10.1007/s11136-021-  
479 02822-4 [published Online First: 20210405]
- 480 14. Terwee CB, Prinsen CAC, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of  
481 patient-reported outcome measures: a Delphi study. *Qual Life Res* 2018;27(5):1159-70. doi:  
482 10.1007/s11136-018-1829-0 [published Online First: 20180317]
- 483 15. Korakakis V, Whiteley R, Kotsifaki A, et al. Tendinopathy VISAs have expired—is it time for outcome  
484 renewals? *Knee Surg Sports Traumatol Arthrosc* 2021 doi: 10.1007/s00167-021-06596-7
- 485 16. Grävare Silbernagel K, Malliaras P, de Vos R-J, et al. ICON 2020—International Scientific Tendinopathy  
486 Symposium Consensus: A Systematic Review of Outcome Measures Reported in Clinical Trials of  
487 Achilles Tendinopathy. *Sports medicine (Auckland, NZ)* 2021 doi: 10.1007/s40279-021-01588-6
- 488 17. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical  
489 severity of Achilles tendinopathy. *British journal of sports medicine* 2001;35(5):335-41. [published  
490 Online First: 2001/10/02]
- 491 18. Murphy MC. Exercise rehabilitation for mid-portion Achilles tendinopathy: a critique of evidence and  
492 assumptions (PhD Academy Award). *British journal of sports medicine* 2022 doi: 10.1136/bjsports-  
493 2022-106024 [published Online First: 20221124]



- 494 19. Comins J, Siersma V, Coupe C, et al. Assessment of content validity and psychometric properties of VISA-  
495 A for Achilles tendinopathy. *PloS one* 2021;16(3):e0247152. doi: 10.1371/journal.pone.0247152  
496 [published Online First: 20210311]
- 497 20. Travers N, Murphy MC, Wand B, et al. Does the Victorian Institute of Sport Assessment – Achilles assess  
498 disability in people with Achilles tendinopathy: a Rasch Measurement Theory analysis. *Under Review*  
499 2023
- 500 21. Murphy MC, Newsham-West RJ, Cook J, et al. TENDINopathy Severity assessment – Achilles (TENDINS-  
501 A): Development and Content Validity assessment of a new Patient-Reported Outcome Measure for  
502 Achilles Tendinopathy. . *The Journal of orthopaedic and sports physical therapy* 2023;Epub Ahead of  
503 Print
- 504 22. Rio EK, Rabusin C, Munteanu S, et al. Where is your pain? Pain location on loading is different to  
505 palpation, imaging and recall location: a study of Achilles tendinopathy research participants. . *The*  
506 *Journal of orthopaedic and sports physical therapy* 2023;Under Review
- 507 23. Sleswijk Visser TSO, van Es EM, Meuffels DE, et al. Standardized pain mapping for diagnosing Achilles  
508 tendinopathy. *Journal of science and medicine in sport* 2022;25(3):204-08. doi:  
509 10.1016/j.jsams.2021.10.010 [published Online First: 20211021]
- 510 24. Tavakkoli Oskouei S, Malliaras P, Hill KD, et al. Evaluating daily physical activity and biomechanical  
511 measures using wearable technology in people with Achilles tendinopathy: A descriptive exploratory  
512 study. *Musculoskeletal science & practice* 2022;58:102534. doi: 10.1016/j.msksp.2022.102534  
513 [published Online First: 20220222]
- 514 25. Golightly YM, Devellis RF, Nelson AE, et al. Psychometric properties of the foot and ankle outcome score  
515 in a community-based study of adults with and without osteoarthritis. *Arthritis Care Res (Hoboken)*  
516 2014;66(3):395-403. doi: 10.1002/acr.22162
- 517 26. MacCallum RC, Widaman KF, Zhang S, et al. Sample size in factor analysis. *Psychological Methods*  
518 1999;4(1):84-99. doi: 10.1037/1082-989X.4.1.84
- 519 27. Child D. The essentials of factor analysis: A&C Black 2006.
- 520 28. Kaiser HF. An index of factorial simplicity. *Psychometrika* 1974;39(1):31-36. doi: 10.1007/BF02291575
- 521 29. Bartlett MS. A Note on the Multiplying Factors for Various  $\chi^2$  Approximations. *Journal of the*  
522 *Royal Statistical Society Series B (Methodological)* 1954;16(2):296-98.
- 523 30. Stevens JP. Applied multivariate statistics for the social sciences: Routledge 2012.

- 524 31. Psychometric theory : Nunnally and Bernstein, 3rd edition; 2007.
- 525 32. Bagozzi RP, Yi Y. On the evaluation of structural equation models. *Journal of the Academy of Marketing*  
526 *Science* 1988;16(1):74-94. doi: 10.1007/BF02723327
- 527 33. Mokkink LB, Terwee CB, Patrick DL, et al. The COSMIN study reached international consensus on  
528 taxonomy, terminology, and definitions of measurement properties for health-related patient-reported  
529 outcomes. *J Clin Epidemiol* 2010;63(7):737-45. doi: 10.1016/j.jclinepi.2010.02.006
- 530 34. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability  
531 Research. *J Chiropr Med* 2016;15(2):155-63. doi: 10.1016/j.jcm.2016.02.012 [published Online First:  
532 20160331]
- 533 35. de Vet HCW, Terwee CB, Mokkink LB, et al. *Measurement in Medicine: A Practical Guide*. Cambridge:  
534 Cambridge University Press 2011.
- 535 36. Mouelhi Y, Jouve E, Castelli C, et al. How is the minimal clinically important difference established in  
536 health-related quality of life instruments? Review of anchors and methods. *Health Qual Life Outcomes*  
537 2020;18(1):136. doi: 10.1186/s12955-020-01344-w [published Online First: 20200512]
- 538 37. Challoumas D, Zouvani A, Creavin K, et al. Determining minimal important differences for patient-reported  
539 outcome measures in shoulder, lateral elbow, patellar and Achilles tendinopathies using distribution-  
540 based methods. *BMC Musculoskelet Disord* 2023;24(1):158. doi: 10.1186/s12891-023-06261-9  
541 [published Online First: 20230302]
- 542 38. Lagas IF, van der Vlist AC, van Oosterom RF, et al. Victorian Institute of Sport Assessment-Achilles (VISA-  
543 A) Questionnaire-Minimal Clinically Important Difference for Active People With Midportion Achilles  
544 Tendinopathy: A Prospective Cohort Study. *The Journal of orthopaedic and sports physical therapy*  
545 2021;51(10):510-16. doi: 10.2519/jospt.2021.10040
- 546 39. McCormack J, Underwood F, Slaven E, et al. The minimum clinically important difference on the VISA-  
547 Aand LEFS for patients with insertional Achilles tendinopathy. *Int J Sports Phys Ther* 2015;10(5):639-  
548 44.
- 549 40. Warburton DE, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *CMAJ : Canadian*  
550 *Medical Association journal = journal de l'Association medicale canadienne* 2006;174(6):801-9. doi:  
551 10.1503/cmaj.051351

- 552 41. Tu X, Tu Z, Lin W, et al. The Victorian Sports Assessment Institute-Achilles Tendinopathy Questionnaire  
553 (VISA-A): Chinese cross-cultural adaptation and psychometric validation. *Health Qual Life Outcomes*  
554 2022;20(1):111. doi: 10.1186/s12955-022-02025-6 [published Online First: 20220723]
- 555 42. Bahari M, Hadadi M, Vosoughi AR, et al. Cross-cultural adaptation, reliability and validity of the Persian  
556 version of the Victorian Institute of Sport Assessment-Achilles questionnaire (VISA-A). *Disability and*  
557 *rehabilitation* 2022;44(6):983-91. doi: 10.1080/09638288.2020.1781268 [published Online First:  
558 20200714]
- 559 43. Rio EK, Mc Auliffe S, Kuipers I, et al. ICON PART-T 2019–International Scientific Tendinopathy  
560 Symposium Consensus: recommended standards for reporting participant characteristics in  
561 tendinopathy research (PART-T). *British journal of sports medicine* 2019;54(11):627-30. doi:  
562 10.1136/bjsports-2019-100957
- 563  
564

565 **Figure 1. Factor Loading within Confirmatory Factor Analysis**

566 *Legend: Small circles to the left of the diagram (e.g., E11) represent error related to each item. Medium sized*  
567 *circles (e.g., Item 11) represent scores for each TENDINS-A item (known as indicators). Large circles to the*  
568 *right of the diagram (e.g., PAIN) represent factors causing disability (known as unobserved or latent variables).*  
569 *Arrows linking factors and items represent the standardised factor loading for each item. Arrows linking factors*  
570 *represent covariance between factors.*

571