



Correction

Correction: Shekarkar Azgomi et al. A Rapid and Simple Multiparameter Assay to Quantify Spike-Specific CD4 and CD8 T Cells after SARS-CoV-2 Vaccination: A Preliminary Report. *Biomedicines* 2021, 9, 1576

Mojtaba Shekarkar Azgomi ^{1,2}, Marco Pio La Manna ^{1,2}, Giusto Davide Badami ^{1,2}, Paolo Ragonese ², Antonino Trizzino ³, Francesco Dieli ^{1,2} and Nadia Caccamo ^{1,2,*}

¹ Central Laboratory of Advanced Diagnostic and Biomedical Research (CLADIBIOR), University of Palermo, 90127 Palermo, Italy; mojtaba.shekarkarazgomi@unipa.it (M.S.A.); marcopio.lamanna@unipa.it (M.P.L.M.); giustodavide.badami@unipa.it (G.D.B.); francesco.dieli@unipa.it (F.D.)

² Department of Biomedicine, Neurosciences and Advanced Diagnostic (Bi.N.D.), University of Palermo, 90127 Palermo, Italy; paolo.ragonese@unipa.it

³ Department of Pediatric Hematology and Oncology, A.R.N.A.S. Civico Di Cristina and Benfratelli Hospital, 90127 Palermo, Italy; triznino@hotmail.com

* Correspondence: nadia.caccamo@unipa.it

In the original publication, there was a mistake in Figure 1A, as published [1]. A FACS plot for IL-2 production was duplicated during the editing of the Figure through mere error, between non-stimulated and stimulated conditions. In detail, we duplicated the sixth plot of the upper row from the third plot of the middle row. The corrected Figure 1 appears below.



Citation: Shekarkar Azgomi, M.; La Manna, M.P.; Badami, G.D.; Ragonese, P.; Trizzino, A.; Dieli, F.; Caccamo, N. Correction: Shekarkar Azgomi et al. A Rapid and Simple Multiparameter Assay to Quantify Spike-Specific CD4 and CD8 T Cells after SARS-CoV-2 Vaccination: A Preliminary Report. *Biomedicines* 2021, 9, 1576. *Biomedicines* 2023, 11, 2378. <https://doi.org/10.3390/biomedicines11092378>

Received: 20 July 2023

Accepted: 25 July 2023

Published: 25 August 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. Quantification of spike-specific CD4⁺ and CD8⁺ T cells after SARS-CoV-2 vaccination. Gating strategy used to identify spike-specific CD4⁺ and CD8⁺ T cells and to detect their cytokine expression in response to spike-derived peptides. (A) Cumulative frequency and distribution of spike-specific CD4⁺ and CD8⁺ T cells (B) in SARS-CoV-2-vaccinated individuals (*n* = 40). Cut-off for positivity was set at <0.02 for CD4⁺ T cells and <0.05 for CD8⁺ T cells. Analysis of distinct cytokine expression by spike-specific CD4⁺ (C) and CD8⁺ (D) T cells using a Kruskal–Wallis test with Dunn’s correction. ** *p* < 0.01; *** *p* < 0.001; ns: not significant.

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Shekarkar Azgomi, M.; La Manna, M.P.; Badami, G.D.; Ragonese, P.; Trizzino, A.; Dieli, F.; Caccamo, N. A Rapid and Simple Multiparameter Assay to Quantify Spike-Specific CD4 and CD8 T Cells after SARS-CoV-2 Vaccination: A Preliminary Report. *Biomedicines* **2021**, *9*, 1576. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.