

Compounds identified on hexane and dichloromethane extracts of *Salicornia ramosissima*

 Vera M.S. Isca,^[a] Ana M.L. Seca,^[a,b] Diana C.G.A. Pinto,^[a] Artur M. S. Silva^[a] and Helena Silva^[c]
^[a]Department of Chemistry & QOPNA, University of Aveiro, 3810-193 Aveiro, Portugal; ^[b]DCTD, University of Azores, Rua Mãe de Deus, 9501-801 Ponta Delgada, Portugal; ^[c]Department of Biology, University of Aveiro, 3810-193 Aveiro, Portugal. e-mail: veraisca@ua.pt

Salicornia ramosissima J. Woods [4]

Aliphatic profile


Analysis of hexane extracts by GC-MS revealed the presence of 27 fatty acids of which 6 are unsaturated, 15 alcohols, 7 sterols, 5 sugars and 3 cinnamic acids. The fatty acids, alcohols and sterols are the most abundant ones. Table 1 shows some of the identified compounds. Palmitic acid, tetracosan-1-ol and Docosan-1-ol represent the main components of the extracts.

Table 1 – Identified compounds by GC-MS

Identified Compounds	NE (%)	NE-H (%)	E (%)	E-H (%)
Fatty acids				
Palmitic acid C16:0	2.54	5.32	3.32	6.55
Tetracosanoic acid C24:0	0.63	1.02	0.56	1.17
Docosanoic acid C22:0	0.56	0.95	0.65	1.39
Octadec-11-enoic acid C18:1 (11E)	0.15	0.41	0.18	0.57
Tricosanoic acid C23:0	-	0.31	-	0.40
2-Hydroxy-heptanoic acid COOHCH(OH)(CH ₂) ₄ CH ₃	-	0.03	-	0.04
Fatty alcohols				
Tetracosan-1-ol C24-OH	2.50	2.33	2.42	3.26
Dososan-1-ol C22-OH	1.55	2.70	1.87	3.17
Hexacosan-1-ol C26-OH	1.45	1.13	1.28	1.69
tridecan-2-ol C13-OH	-	0.05	-	0.10
Pentacosan-1-ol C25-OH	-	-	0.14	0.20
Sterols				
Stigmasterol	0.78	0.60	0.67	0.92
β-sitosterol	0.66	0.77	0.74	1.13
Sitostanol	0.46	0.64	0.60	0.87
Aromatic acid				
Isoferulic acid	-	0.51	-	0.23


The lipid profile among *Salicornia* species is quite distinct however the fatty acid content in *Salicornia ramosissima* is similar to the *Salicornia fruticosa* [5], the saturated fatty acids are present in higher amount (87%) than unsaturated (13%) and palmitic acid (41,9%) represent the main component of the fatty acids fraction.

The analysis of the dichloromethane extract from *S. ramosissima* aerial parts allowed the isolation of the triacylglycerol (1) and the 2,2'-((oxybis(3-(tert-butyl)-6,1-phenylene))bis(propane-2,2-diyl))bis(4-(tert-butyl)phenol) (2). Here we present their unequivocal structure elucidation.



The spectroscopic characterization of compound 2 led to an unique structure with uncommon *t*-butyl groups. Is not the first time that a compound like this is discovered in nature [5]. The structure's was confirmed by mass spectroscopy (Fig 3).

Fig 3



Acknowledgments

Thanks are due to the University of Aveiro, Fundação para a Ciência e a Tecnologia (FCT) and FEDER for funding the Organic Chemistry Research Unit (project PEst-C/UI0062/2011) and Portuguese National NMR Network (RNRMN).

References

- [1] Silva, H.; Freitas, H.; Caldeira, G., *Rev. Biol.* **1999**, *17*, 193.
- [2] Isca, V. M. S. Seca, A. M. L.; Pinto, D. C. G. A.; Silva, A. M. S. in: *Natural Products: Research Review*, Vol 5. M/S Daya Publishing House, New Delhi. 2012 (accepted for publication).
- [3] Isca, V. M. S.; Seca, A. M. L.; Pinto, D. C. G. A.; Silva, A. M. S.; Silva, H. *6thSPJOCs, Book abstract*, Lisbon **2012**, p. 146.
- [4] http://www.species.ie/burren/species.php?species_group=Burren&menuentry=soorten&id=491&tab=beschrijving, access in 20/10/2011.
- [5] Radwan, H.M.; Nazif, N.M.; Abou-Setta, L.M., *Res. J. Med. Med. Sci.* **2007**, *2*, 72.
- [6] Bisel, P.; Al-Momani, L.; Müller, M., *Org. Biomol. Chem.*, **2008**, *6*, 2655.

Material and Methods

Plant collected in Ria de Aveiro (Portugal) from distinct growth conditions:
 NE – growth with natural irrigation
 E - growth with additional irrigation with canal water from 15 to 15 days

Plant air dried and aerial parts extract with dichloromethane or hexane at room temperature


Dichloromethane extract

Silica gel column chromatography (gradient mixtures of hexane and ethyl acetate)

Silica gel TLC (different polarity eluent mixtures)

Compounds 1 and 2

Identified compounds family's in each extract



Prior to the hydrolysis of long chain alcohols are the most abundant family of compounds. This will change significantly after hydrolysis since fatty acids become majority (Fig 1).

The compounds tricosanoic, 2-hydroxy-heptanoic, isoferulic acids and the tridecan-2-ol exist exclusively in ester form, since they appears only after hydrolysis.

In the plant irrigated in excess (E), is observed an increase in the production of esterified compounds (Fig 2). The main alteration is the production of pentacosan-1-ol, nonexistent in NE.

Free compounds





Fig 2

Esterified compounds



Spectroscopical characterization

In order to elucidate the fatty acid linked to the glycerol, the compound 1 was submitted to an alkaline hydrolysis and the result analyzed by GC-MS. The tetracosanoic acid was identified.

