

2-7-2024

## Expert Opinion, Real-World Classification, and Decision-Making in Thoracolumbar Burst Fractures Without Neurologic Deficits?

Gaston Camino-Willhuber

Sebastian Bigdon

Charlotte Dandurand

Marcel F. Dvorak

Cumhur F. Öner

Follow this and additional works at: [https://jdc.jefferson.edu/rothman\\_institute](https://jdc.jefferson.edu/rothman_institute)  
See next page for additional authors



Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Camino-Willhuber, Gaston; Bigdon, Sebastian; Dandurand, Charlotte; Dvorak, Marcel F.; Öner, Cumhur F.; Schnake, Klaus; Muijs, Sander; Benneker, Lorin M.; Vialle, Emiliano; Tee, Jin W.; Keynan, Ory; Chhabra, Harvinder S.; Joaquim, Andrei F.; Popescu, Eugen C.; Canseco, Jose A; Holas, Martin; Kanna, Rishi M.; Aly, Mohamed M.; Fallah, Nader; Schroeder, Gregory D.; Spiegl, Ulrich; El-Skarkawi, Mohammad; Bransford, Richard J.; Rajasekaran, Shanmuganathan; and Vaccaro, Alexander R., "Expert Opinion, Real-World Classification, and Decision-Making in Thoracolumbar Burst Fractures Without Neurologic Deficits?" (2024). *Rothman Institute Faculty Papers*. Paper 247.  
[https://jdc.jefferson.edu/rothman\\_institute/247](https://jdc.jefferson.edu/rothman_institute/247)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

## Authors

Gaston Camino-Willhuber, Sebastian Bigdon, Charlotte Dandurand, Marcel F. Dvorak, Cumhur F. Öner, Klaus Schnake, Sander Muijs, Lorin M. Benneker, Emiliano Vialle, Jin W. Tee, Ory Keynan, Harvinder S. Chhabra, Andrei F. Joaquim, Eugen C. Popescu, Jose A Canseco, Martin Holas, Rishi M. Kanna, Mohamed M. Aly, Nader Fallah, Gregory D. Schroeder, Ulrich Spiegl, Mohammad El-Skarkawi, Richard J. Bransford, Shanmuganathan Rajasekaran, and Alexander R. Vaccaro

# Expert Opinion, Real-World Classification, and Decision-Making in Thoracolumbar Burst Fractures Without Neurologic Deficits?

Global Spine Journal  
2024, Vol. 14(1S) 49S–55S  
© The Author(s) 2023  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/21925682231194456  
[journals.sagepub.com/home/gsj](https://journals.sagepub.com/home/gsj)



Gaston Camino-Willhuber, MD<sup>1,\*</sup> , Sebastian Bigdon, MD<sup>2,\*</sup> ,  
Charlotte Dandurand, MD, MSc, FRCSC<sup>3</sup> , Marcel F. Dvorak, MD, MBA<sup>3</sup>,  
Cumhur F. Öner, MD, PhD<sup>4</sup>, Klaus Schnake, MD<sup>5,6</sup>, Sander Muijs, MD, PhD<sup>4</sup>,  
Lorin M. Benneker, MD<sup>7</sup>, Emiliano Vialle, MD<sup>8</sup> , Jin W. Tee, MD<sup>9</sup>, Ory Keynan, MD<sup>10</sup>,  
Harvinder S. Chhabra, MBBS, MS<sup>11</sup>, Andrei F. Joaquim, MD, PhD<sup>12</sup> ,  
Eugen C. Popescu, MD, PhD<sup>13</sup> , Jose A. Canseco, MD, PhD<sup>14</sup> , Martin Holas, MD<sup>15</sup> ,  
Rishi M. Kanna, MS, MRCS, FNB<sup>16</sup> , Mohamed M. Aly, MD, PhD<sup>17,18</sup>, Nader Fallah, PhD<sup>3</sup>,  
Gregory D. Schroeder, MD<sup>14</sup>, Ulrich Spiegl, MD<sup>19</sup> , Mohammad El-Skarkawi, MD<sup>20</sup> ,  
Richard J. Bransford, MD<sup>21</sup>, Shanmuganathan Rajasekaran, MD, PhD<sup>16</sup> , and  
Alexander R. Vaccaro, MD, PhD, MBA<sup>14</sup>

## Abstract

**Study Design:** Retrospective analysis of prospectively collected data.

- 
- <sup>1</sup> Institute of Orthopedics “Carlos E. Ottolenghi”, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina  
<sup>2</sup> Department of Orthopaedic Surgery and Traumatology, Inselspital, University Hospital, University of Bern, Bern, Switzerland  
<sup>3</sup> Combined Neurosurgical and Orthopaedic Spine Program, Department of Orthopaedic Surgery, University of British Columbia, Vancouver, BC, Canada  
<sup>4</sup> University Medical Centers, Utrecht, The Netherlands  
<sup>5</sup> Center for Spinal and Scoliosis Surgery, Malteser Waldkrankenhaus St. Marien, Erlangen, Germany  
<sup>6</sup> Department of Orthopedics and Traumatology, Paracelsus Private Medical University Nuremberg, Nuremberg, Germany  
<sup>7</sup> Spine Unit, Sonnenhof Spital, University of Bern, Bern, Switzerland  
<sup>8</sup> Cajuru Hospital, Catholic University of Paraná, Curitiba, Brazil  
<sup>9</sup> Department of Neurosurgery, National Trauma Research Institute (NTRI), Melbourne, VIC, Australia  
<sup>10</sup> Rambam Health Care Campus, Haifa, Israel  
<sup>11</sup> Department of Spine Service, Indian Spinal Injuries Centre, Sector C, Vasant Kunj, New Delhi, India  
<sup>12</sup> Department of Neurology, State University of Campinas, Campinas-Sao Paulo, Brazil  
<sup>13</sup> “Prof. Dr. N. Oblu” Emergency Hospital, Iasi, Romania  
<sup>14</sup> Department of Orthopaedic Surgery, Rothman Institute, Thomas Jefferson University Hospital, Philadelphia, PA, USA  
<sup>15</sup> Klinika Úrazovej Chirurgie SZU a FN sP F.D. Roosevelta, Banská Bystrica, Slovakia  
<sup>16</sup> Department of Orthopaedics and Spine Surgery, Ganga Hospital, Coimbatore, Tamil Nadu, India  
<sup>17</sup> Department of Neurosurgery, Prince Mohammed Bin Abdulaziz Hospital, Riyadh, Saudi Arabi  
<sup>18</sup> Department of Neurosurgery, Mansoura University, Mansoura, Egypt  
<sup>19</sup> Klinik für Orthopädie, Unfallchirurgie und plastische Chirurgie, Universitätsklinik Leipzig, Leipzig, Germany  
<sup>20</sup> Department of Orthopaedic and Trauma Surgery, Assiut University Medical School, Assiut, Egypt  
<sup>21</sup> Department of Orthopaedics and Sports Medicine, University of Washington Harborview Medical Center, Seattle, WA, USA

\*Gaston Camino-Willhuber and Sebastian Bigdon Contributed equally.

## Corresponding Author:

Gaston Camino-Willhuber, MD, Institute of Orthopedics “Carlos E. Ottolenghi,” Hospital Italiano de Buenos Aires, Buenos Aires, Potosi 4215, Argentina.  
Email: [gaston.camino@hospitalitaliano.org.ar](mailto:gaston.camino@hospitalitaliano.org.ar)



Creative Commons Non Commercial No Derivs CC BY-NC-ND: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

**Objectives:** To compare decision-making between an expert panel and real-world spine surgeons in thoracolumbar burst fractures (TLBFs) without neurological deficits and analyze which factors influence surgical decision-making.

**Methods:** This study is a sub-analysis of a prospective observational study in TL fractures. Twenty two experts were asked to review 183 CT scans and recommend treatment for each fracture. The expert recommendation was based on radiographic review.

**Results:** Overall agreement between the expert panel and real-world surgeons regarding surgery was 63.2%. In 36.8% of cases, the expert panel recommended surgery that was not performed in real-world scenarios. Conversely, in cases where the expert panel recommended non-surgical treatment, only 38.6% received non-surgical treatment, while 61.4% underwent surgery. A separate analysis of A3 and A4 fractures revealed that expert panel recommended surgery for 30% of A3 injuries and 68% of A4 injuries. However, 61% of patients with both A3 and A4 fractures received surgery in the real world. Multivariate analysis demonstrated that a 1% increase in certainty of PLC injury led to a 4% increase in surgery recommendation among the expert panel, while a .2% increase in the likelihood of receiving surgery in the real world.

**Conclusion:** Surgical decision-making varied between the expert panel and real-world treating surgeons. Differences appear to be less evident in A3/A4 burst fractures making this specific group of fractures a real challenge independent of the level of expertise.

### Keywords

thoracolumbar burst fracture, surgical decision making, surgical treatment

## Introduction

The thoracolumbar junction is most susceptible for injuries. About two thirds of all fractures occur at this site. The most common mechanism is falling from height, during sports or within a motor vehicle accident.<sup>1</sup> The axial impact to the spine with or without a concomitant bending moment may lead to the fracture of the vertebral column.<sup>2</sup> In addition to the high impact on the quality of life, there are high socioeconomic costs due to treatment, loss of work, short- and long-term disability.<sup>2,3</sup> Therapeutic decision-making includes non-surgical and surgical treatment in fractures without neurological compromise while surgical treatment is more commonly recommended in cases with neurological compromise or refractory pain. Despite the frequency and severe sequelae of the fracture, treatment of thoracolumbar burst fractures (TLBF) in patients who are neurologically intact (Type A3/A4 according to AO Spine Thoracolumbar Injury Classification System) remains controversial.

The AO Spine classification for thoracolumbar fractures has shown substantial inter- and intraobserver reliability ( $k = .62$ ).<sup>4</sup> A proper classification allows the comparison of different therapeutic approaches among injuries with similar morphological characteristics.

Despite the substantial reliability of the AO Spine classification in general, thoracolumbar fractures may present a heterogeneous radiological pattern, and misclassification is possible. Interestingly, the lowest agreement ( $\kappa = .19$ ) within the validation study was found for A4 fractures. There is an ongoing debate in literature for the best treatment of A3 and A4 fractures.<sup>5,6</sup> Failing to recognize a specific fracture pattern might be one of the factors that contributes to a wide disparity in treatment recommendations.

The AO spine classification provides modifiers for more precise fracture analysis, arguing for or against surgical treatment. However, these modifiers are used inconsistently in both clinical and research settings. Other than the exact fracture pattern of the bone and potential modifiers, the integrity of the posterior ligamentous complex (PLC) is thought to be very important to guide surgeons in making therapeutic decisions.<sup>7,8</sup> Injury to the PLC affects the static of the spine and may increase the risk for fracture subsidence.

In addition to fracture classification and related factors, the decision-making process for treatment may also be affected by variables such as the surgeon's educational background, individual experience, regional treatment guidelines, resource availability, and patient preferences. Consequently, the extent to which radiological characteristics, such as fracture morphology, and fracture classification are influenced by a surgeon's expertise remains uncertain. This equipoise study offers the unique opportunity to compare how the radiographic assessment influences decision making and how it differs from the actual decision making where other factors are considered.

The aims of this study are:

- (1) To analyze the degree to which the classification of fractures and the treatment recommendations of a panel of expert surgeons correlate with the actual classification and treatment that patients receive in a real-world setting.
- (2) To evaluate the influence of radiological characteristics such as the probability of PLC injury and M1 case-specific modifier in decision making.

## Methods

This study was conducted by the AO Spine Knowledge Forum Trauma, an expert research study group supported by the not-for-profit organization AO Foundation whose members are generally recognized for their academic expertise in spinal trauma. The study is part of a prospective observational study of TL Fractures, the Spine TL A3/A4 study.<sup>9</sup>

The detailed methodology is available in the article of Dandurand et al “Understanding Decision Making as it Influences Treatment in Thoracolumbar Burst Fractures Without Neurological Deficit: Conceptual Framework and Methodology” in this focus issue. The AO Spine Knowledge Forum Trauma completed consent and recruitment for a multicenter prospective observational study of TL Fractures; the Spine A3/A4 study. Each enrolling center obtained local approval from their local institutional review board. The baseline CT scans and conventional radiographs of 183 patients were available for this study. All patients were neurologically intact and had injuries between T11 and L2.

In the analysis presented here, we compare the expert group’s classification and treatment recommendations with those of the treating surgeons’ actual decision on-site (Aim 1). Twenty two Spine Trauma experts from different parts of the world were asked to recommend a treatment for 183 TLBFs. The 22 experts, all with extensive experience in management of spinal trauma were recruited from the AO Spine Knowledge Forum Trauma (KF Trauma). Each member of the expert panel independently reviewed the DICOM images of the 183 TL fracture cases and were asked to classify each injury based on the latest AO TL Injury Classification System, assess the degree of certainty of PLC disruption and the degree of comminution. Finally, they were asked to recommend treatment – either surgical or non-operative, which specific type of treatment and finally asked how confident they were in this recommendation. These experts were agnostic to the actual treatment that the patient received within the Spine TL A3/A4 Study and were also agnostic to any results of the TL Spine A3/A4 study.

Each expert recommendation was compared to the treatment actually received in real-world. We use descriptive statistics and crosstabulation to compare the overall therapy recommendation of the expert group with the actual on-site treatment. In addition, we consider the treatment decision or recommendation concerning the fracture pattern according to the AO Spine TL injury classification system. To test validity, we used Pearson Chi-Square Test.

To evaluate the influence of radiologic characteristics, such as the presence of PLC injury and case-specific M1 modifier-indeterminate injury to the tension band based on MRI, on decision-making (Aim 2), we performed a multivariate logistic regression analysis. To assess the impact of radiologic features on decision-making (Aim 2), we focused on factors like the presence of PLC injury and case specific M1 modifier. We conducted a multivariate logistic regression analysis to

evaluate these influences systematically. The experts were asked to assess the probability (0%–100%) of the presence of a PLC injury based on the available images. PLC values and the M1 modifier were used as independent variables for logistic regression model. SPSS, R and SAS were used for data analysis.

## Results

### Overall Recommendations

We included 183 cases of TLBFs without neurologic deficit. Twenty-two experts were asked to give a treatment recommendation after reviewing all available radiological images without knowing the actual treatment these patients have received. We therefore received 3929 aggregated recommendations.

### Correlation Between the Expert Panel and Real-World Treating Surgeons

Regarding surgical treatment, the overall agreement between expert panel and real-world surgeons was observed in 63.2%. In contrast, in 36.8% of cases, the expert panel recommended surgery that was not performed in the real-world scenario.

Vice versa, in all the cases where the expert panel recommended non-surgical treatment after reviewing the imaging, only 38.6% actually received non-surgical treatment. The remaining 61.4% of patients received surgery.

### Comparison Between Expert Panel Surgeons and Real-World Treating Surgeons Based on Injury Fracture Pattern

Surgery indication recommendation between the expert panel (Table 1) and real-world treating surgeons (Table 2) is presented as well as comparison based on the distinction between A3 and A4 patterns (Table 3).

The expert Panel was as likely to recommend surgery as non-surgical care (51% vs 49%) when looking at A3/A4 fractures together while in the real world almost 2/3 of patients were treated surgically (62%). When A3 and A4 was explored separately, the expert panel recommended surgical care in 30% of A3 injuries and 68% in A4 fractures. In the real world an identical proportion (61%) of patients with A3 and A4 fractures actually received surgery.

### Correlation of PLC Injury Identification and M1-Modifier in Surgical Recommendation Between Expert Panel and Real-World Treating Surgeons

Multivariate analysis showed an Odds ratio for surgery recommendation based on PLC identification of 1.039 ( $P$ .001), meaning that for each 1% of PLC injury in certainty, there was a 4% increase in surgery recommendation among the expert

**Table 1.** Surgery Recommendations by Expert Panel Based on Injury Pattern Groups.

Injury Groups	Surgery Recommendation (%/N)		Total
	No	Yes	
A3-A4 subtypes	51.1 (1811)	48.9 (1733)	100 (3544)
B1-B2-C subtypes	2.6 (10)	97.4 (375)	100 (385)
All fracture types combined	46.3 (1821)	53.7 (2108)	100 (3929)

**Table 2.** Surgery Actually Performed by Real-World Treating Surgeons Based on Injury Pattern Groups.

Injury Group	Surgery Actually Performed in Real-World (%/N)		Total
	No	Yes	
A3-A4	38.5 (1364)	61.5 (2180)	100 (3544)
B1-B2-C	29.6 (114)	70.4 (271)	100 (385)
All fracture types combined	37.6 (1478)	62.4 (2451)	100 (3929)

**Table 3.** Comparison Between Expert Panel Surgeons and Real-World Treating Surgeons Based on Distinction Between A3 and A4 Fracture Pattern.

Fracture Type	Expert Panel Recommendation			Real World Treatment Received		
	Surgery n (%)	Non-Surgical Management n (%)	P Value	Surgery n (%)	Non-Surgical Management n (%)	P Value
A3 vs A4 fractures						
A3	565 (30.9)	1266 (69.1)	<.001	1130 (61.7)	701 (38.3)	.412
A4	1168 (68.2)	545 (31.8)		1050 (61.3)	663 (38.7)	

**Table 4.** Influence of Surgery Recommendation Based on PLC and Modifier Likelihood Among the Panel of Experts and Real-World Treating Surgeons.

Variable	Panel of Experts		Real World Treating Surgeons	
	OR (95% CI)	P	OR (95% CI)	P
Certainty of PLC injury percentage	1.039 (1.036-1.043)	<.001	1.002 (1.000-1.004)	.039
Presence of MI-modifier	4.055 (3.535-4652)	<.001	1.052 (.93-1.199)	.450

panel. On the other hand, of each 1% increase in the certainty of PLC injury, there was .2% increase in the chance of receiving surgery in the real world. Additionally, surgery recommendation significantly increased based on the presence of modifiers (OR = 4.05,  $P = .0001$ ) among the expert panel but was not statistically significant among the real-world treating surgeons. (Table 4).

## Discussion

Treatment decision-making in TL fractures without neurological compromise is controversial among spine surgeons.<sup>10-12</sup> The recommendations may vary according to the surgeon's experience and background, the patient's status, comorbidities, and proper classification based upon imaging findings.<sup>10-12</sup> In this study, we compared treatment recommendation from international experts reviewing available

imaging data to on-site spine surgeons treating patients with these fractures in a real-world setting.

Interesting differences can be found in the therapeutic recommendations of the expert panel compared to the actual real-world decisions. If the expert group considered surgical treatment based on imaging findings, 63.2% of the patients actually underwent surgical treatment in the real-world, compared with 36.8% who were managed non-surgically.

Particularly interesting are the patients for whom the experts did not consider initial surgical treatment. Almost two thirds (61.3%) of these patients nevertheless received surgical treatment from the treating surgeon on site, while only 38.7% were treated non-surgically.

Investigating the B1, B2 and C type subgroups of injuries, while the expert panel strongly favored surgery (97%), only 70% of these patients actually were treated with surgery in the real world. This strongly suggest that there are variables other than the



morphological appearance of the fracture pattern on CT scan that are directing actual real world treatment decisions. These variables could include strong local preferences for non-surgical and surgical care among specific surgeons that over-ride any considerations based upon morphological features and subsequent classification categories. It may be that patient preference has a stronger influence in the real-world and guides treatment. Finally, there may be issues such as individual patient characteristics (body habitus, etc) as well as resource availability that influence treatment decisions.

Interestingly, when comparing surgical indications in the A3/A4 fracture patterns, the differences between the expert panel and real-world treating surgeons were less evident with almost identical proportion of surgical and non-surgical recommendations from the expert panel (49% and 51%). In the real world, 61% of A3 and A4 fractures were both treated surgically.

The distinction between A4 and A3 fractures becomes particularly intriguing when analyzed separately. The expert panel differentiates these two injury types by recommending surgery for 70% of A4 cases and non-surgical care for 70% of A3 cases. However, this distinction does not seem to fit real-world treatment decisions, as both A3 and A4 fractures undergo surgical treatment in 61% of cases, irrespective of the morphological differences between A4 and A3 injuries, as outlined in the AO classification.

There are several radiographic characteristics that are thought to influence the categorization of TLBF's and guide treatment recommendations. These include: degree of vertebral body comminution, local kyphosis, PLC integrity, and canal compromise. Add to this the influence of each surgeons' experience, local resources availability, patient preference and deeply held local beliefs regarding what the best treatment approach is for these fractures. Which of these factors have a strong influence on the decision-making remains currently unclear although we have shown that radiographic analysis by a group of experts is not adequate on its own to identify which patients actually receive surgery in the real-world setting.

In a prospective randomized trial, Wood et al compared non-operative vs operative management of TLBF without neurological compromise. They reported that patients managed non-operatively had less pain and better function in the conservative group (ODI of 20 in the operative group vs 2 in the non-operative group –  $P < .001$ ) without significant differences in local kyphosis ( $13^\circ$  for those who had surgery vs  $19^\circ$  for those treated non-surgically,  $P = .003$ ) in long-term follow-up (16 to 22 years).<sup>14</sup>

On the other hand, Siebenga et al demonstrated in a multicenter randomized trial that there was some patient benefit when treated surgically with a short segment fusion, with better functional outcome scores (VAS pain, VAS Spine Score and RMDQ-24).<sup>15</sup> Finally, a Meta-analysis performed by Gnanenthiran et al comparing non-operative vs operative treatment for burst fractures without neurological compromise, surgical treatment showed a slight improvement in residual kyphosis ( $12.8^\circ$  to  $11^\circ$  in the operative group) without differences between conservative and surgical groups in terms of pain and function. However, surgery had higher

complication rates and costs.<sup>16</sup> In this regard, our results are consistent with the literature showing the lack of consensus on the best treatment options for these injuries.

In our study, the expert panel of surgeons recommended treatment only based on radiological imaging characteristics. Many authors have analyzed the value of imaging studies when classifying thoracolumbar fractures. Rajasekaran et al evaluated the classification process of 30 thoracolumbar injuries among 41 spine surgeons. They reported that x-rays alone were insufficient to classify fractures correctly (only 43.4% of the cases were correctly classified), and ct scan added significant accuracy to the classification process (further 18.2%) with a modest gain in sensitivity when adding an MRI.<sup>17</sup> They concluded that CT was mandatory for proper classification.

When analyzing the factors that increased the likelihood for surgical recommendation from the expert panel of surgeons, we observed that PLC injury and the presence of M1 case-specific modifier were the most important. We did not analyze the M2 (patient-specific comorbidity) modifier as the expert panel was unaware of baseline clinical information. Expert surgeons were asked how certain they were about PLC injury from 0%–100% regarding suspected PLC injury. We found that the likelihood to indicate surgery increases by 4% for each 1% increase of suspicion in PLC injury among the expert panel, this result contrasted with the real-world surgeon's group, in which the indication for surgery was not as clearly influenced by the radiographic evidence of PLC disruption only increasing 1% by each 1% increase in PLC injury. CT-scan suspected PLC injury has an essential role in expert decision-making. Reliability of identifying PLC integrity based on CT scan findings is also subject to faults and variations,<sup>18-20</sup> and this could affect decision-making even among experts. On the other hand, the presence of M1-modifier significantly increased the potential for a recommendation for surgery by the expert panel 4-fold, meaning that these two factors were significant radiological predictors for surgical recommendations from the experts. Regarding the real-world surgeons, the presence of M1-modifier did not appear to influence treatment. We found no statistically significance. Therefore, this modifier probably did not influence surgical decision-making. We believe that in the real world, there must be other factors influencing surgical decision-making in addition to the morphology of the fracture that was made available to the expert panel.

This study has several strengths. Firstly, the retrospective analysis of prospectively collected data lends credibility to the findings. The comparison of decision-making between an expert panel and real-world spine surgeons in TLBFs without neurological deficits provides valuable insights into the factors that influence surgical decision-making. Additionally, the use of a diverse group of 22 experts to review 183 acute, baseline injury CT scans further strengthens the study.

However, there are also limitations to this study. As a sub-analysis of a prospective observational study in thoracolumbar fractures, the results might not be generalizable to other types of fractures or different patient populations. Moreover, the study only

assessed the impact of radiographic review on the experts' decision-making and did not consider other factors that may have influenced the actual treatment received by patients, such as resource availability, regional treatment protocols, or patient preferences.

Despite these limitations, the study offers valuable insights into the differences in surgical decision-making between expert panels and real-world treating surgeons. It highlights the challenges associated with A3/A4 burst fractures, emphasizing the need for further research to understand the factors that contribute to the discrepancies in surgical recommendations and actual treatments.

## Conclusion

Radiologic factors are essential in decision-making for TLBFs. International experts who assessed only the available imaging agreed with the treating surgeons on site in 63% of cases when deciding on surgical treatment. On the other hand, 61% of patients for whom the experts suggested nonoperative management underwent surgical treatment, indicating the possible influence of other factors. While the expert assessors differentiated between A3 and A4 fractures, this was not the case in the real-world tendency to treat patients surgically or non-surgically. As the likelihood of PLC injury increased and modifiers were present, the likelihood of the experts recommending surgery also increased, suggesting that better identification of PLC injury may improve the capacity of the AO Classification system to predict final treatment.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was organized and funded by AO Spine through the AO Spine Knowledge Forum Trauma, a focused group of international Trauma experts. AO Spine is a clinical division of the AO Foundation, which is an independent medically-guided not-for-profit organization. Study support was provided directly through AO Network Clinical Research.

## ORCID iDs

Gaston Camino-Willhuber  <https://orcid.org/0000-0002-5684-7679>  
 Sebastian Bigdon  <https://orcid.org/0000-0002-4649-0610>  
 Charlotte Dandurand  <https://orcid.org/0000-0002-4520-6454>  
 Emiliano Vialle  <https://orcid.org/0000-0003-1157-4889>  
 Andrei F. Joaquim  <https://orcid.org/0000-0003-2645-0483>  
 Eugen C. Popescu  <https://orcid.org/0000-0001-5732-5402>  
 Jose A. Canseco  <https://orcid.org/0000-0002-2152-5725>  
 Martin Holas  <https://orcid.org/0000-0002-3763-8767>  
 Rishi M. Kanna  <https://orcid.org/0000-0001-5817-4909>  
 Ulrich Spiegl  <https://orcid.org/0000-0002-5179-4192>  
 Mohammad El-Skarkawi  <https://orcid.org/0000-0001-6177-7145>  
 Shanmuganathan Rajasekaran  <https://orcid.org/0000-0001-6043-006X>

## References

- Bigdon SF, Hecht V, Fairhurst PG, Deml MC, Exadaktylos AK, Albers CE. Injuries in alpine summer sports - types, frequency and prevention: a systematic review. *BMC Sports Sci Med Rehabil.* 2022;14(1):79. doi:10.1186/s13102-022-00468-4.
- Bigdon SF, Saldarriaga Y, Oswald KAC, et al. Epidemiologic analysis of 8000 acute vertebral fractures: evolution of treatment and complications at 10-year follow-up. *J Orthop Surg Res.* 2022;17(1):270. doi:10.1186/s13018-022-03147-9.
- Zileli M, Sharif S, Fornari M. Incidence and epidemiology of thoracolumbar spine fractures: WFNS spine committee recommendations. *Neurospine.* 2021;18(4):704-712. doi:10.14245/ns.2142418.209.
- Urrutia J, Zamora T, Yurac R, et al. An independent interobserver reliability and intraobserver reproducibility evaluation of the new AOSpine thoracolumbar spine injury classification system. *Spine (Phila Pa 1976).* 2015;40(1):E54-E58. doi:10.1097/BRS.0000000000000656.
- Abudou M, Chen X, Kong X, Wu T. Surgical versus non-surgical treatment for thoracolumbar burst fractures without neurological deficit. *Cochrane Database Syst Rev.* 2013;6:CD005079. doi:10.1002/14651858.CD005079.pub3.
- Pehlivanoglu T, Akgul T, Bayram S, et al. Conservative versus operative treatment of stable thoracolumbar burst fractures in neurologically intact patients: Is there any difference regarding the clinical and radiographic outcomes? *Spine (Phila Pa 1976).* 2020;45(7):452-458. doi:10.1097/BRS.0000000000003295.
- Vaccaro AR, Oner C, Kepler CK, et al. AOSpine thoracolumbar spine injury classification system: fracture description, neurological status, and key modifiers. *Spine (Phila Pa 1976).* 2013;38:2028-2037.
- Vaccaro AR, Lehman RA, Hurlbert RJ, et al. A new classification of thoracolumbar injuries: The importance of injury morphology, the integrity of the posterior ligamentous complex, and neurologic status. *Spine (Phila Pa 1976).* 2005;30:2325-2333.
- AO Innovation Translation Center. Thoracolumbar burst fractures (AOSpine A3, A4) in neurologically intact patients: An observational, multicenter cohort study comparing surgical versus non-surgical treatment. 2021. clinicaltrials.gov. <https://clinicaltrials.gov/ct2/show/NCT02827214>. Accessed January 20, 2022.
- Rajasekaran S, Kanna RM, Schroeder GD, et al. Does the spine surgeon's experience affect fracture classification, assessment of stability, and treatment plan in thoracolumbar injuries? *Global Spine J.* 2017;7(4):309-316. doi:10.1177/2192568217699209.
- Sadiqi S, Oner FC, Dvorak MF, Aarabi B, Schroeder GD, Vaccaro AR. The influence of spine surgeons' experience on the classification and intraobserver reliability of the novel AOSpine thoracolumbar spine injury classification system-an international study. *Spine (Phila Pa 1976).* 2015;40(23):E1250-E1256. doi:10.1097/BRS.0000000000001042.
- Kepler CK, Vaccaro AR, Koerner JD, et al. Reliability analysis of the AOSpine thoracolumbar spine injury classification system by a worldwide group of naïve spinal surgeons. *Eur Spine J.* 2016;25(4):1082-1086. doi:10.1007/s00586-015-3765-9.



13. Yacoub AR, Joaquim AF, Ghizoni E, Tedeschi H, Patel AA. Evaluation of the safety and reliability of the newly-proposed AO spine injury classification system. *J Spinal Cord Med.* 2017;40(1):70-75. doi:[10.1179/2045772315Y.0000000042](https://doi.org/10.1179/2045772315Y.0000000042).
14. Wood KB, Buttermann GR, Phukan R, et al. Operative compared with non-operative treatment of a thoracolumbar burst fracture without neurological deficit: A prospective randomized study with follow-up at sixteen to twenty-two years. *J Bone Joint Surg Am.* 2015;97(1):3-9. doi:[10.2106/JBJS.N.00226](https://doi.org/10.2106/JBJS.N.00226).
15. Siebenga J, Leferink VJ, Segers MJ, Elzinga MJ, Bakker FC, Haarmanal HJ. Treatment of traumatic thoracolumbar spine fractures: a multicenter prospective randomized study of operative versus non-surgical treatment. *Spine (Phila Pa 1976).* 2006;31(25):2881-2890.
16. Gnanenthiran SR, Adie S, Harris IA. Non-operative versus operative treatment for thoracolumbar burst fractures without neurologic deficit: A meta-analysis. *Clin Orthop Relat Res.* 2012;470(2):567-577. doi:[10.1007/s11999-011-2157-7](https://doi.org/10.1007/s11999-011-2157-7).
17. Rajasekaran S, Vaccaro AR, Kanna RM, et al. The value of CT and MRI in the classification and surgical decision-making among spine surgeons in thoracolumbar spinal injuries. *Eur Spine J.* 2017;26(5):1463-1469. doi:[10.1007/s00586-016-4623-0](https://doi.org/10.1007/s00586-016-4623-0).
18. Maheswaran A, Aiyer SN, Farouk O, et al. Assessment of interobserver variability for CT scan-based evaluation of posterior ligament complex injury in thoracolumbar fractures: An international multicenter pilot study. *Global Spine J.* 2020;10(2): 118-129. doi:[10.1177/2192568219839414](https://doi.org/10.1177/2192568219839414).
19. Aly MM, Al-Shoaibi AM, Abduraba Ali S, Al Fattani A, Eldawoody H. How often would MRI change the thoracolumbar fracture classification or decision-making compared to CT alone? *Global Spine J.* 2022:21925682221089579. doi:[10.1177/21925682221089579](https://doi.org/10.1177/21925682221089579).
20. Karamian BA, Schroeder GD, Lambrechts MJ, et al. The influence of regional differences on the reliability of the AO spine sacral injury classification system. *Global Spine J.* 2022: 21925682211068419. doi:[10.1177/21925682211068419](https://doi.org/10.1177/21925682211068419).