

Assessing kidney transplantation using ECMO-supported donors within a KDPI-based allocation system

Supplementary Material

Supplementary Table S1: Kidney transplant outcomes in ECMO-supported donors stratified by KDPI

		KDPI 0-20%	KDPI 21-34%	KDPI 35-85%	KDPI 86-100%
Number		21 (VA: 10, VV: 11)	21 (VA: 18, VV: 3)	79 (VA: 47, VV: 32)	7 (VA: 5, VV: 2)
Delayed Graft Function		8 (38.10%)	10 (47.62%)	40 (50.63%)	5 (71.43%)
Creatinine at 1-year follow-up (mg/dL)		1.20 (0.92-1.46)	1.21 (1.10-1.46)	1.40 (1.17-1.87)	2.90 (1.85-3.15)
<i>Graft Survival</i>					
	1-year	100.00%	90.00%	84.87%	68.57%
	3-year	100.00%	90.00%	79.97%	45.71%
	5-year	88.89%	70.13%	58.03%	45.71%
<i>Patient Survival</i>					
	1-year	100.00%	100.00%	87.64%	83.33%
	3-year	100.00%	94.74%	87.64%	83.33%
	5-year	94.44%	73.82%	82.54%	83.33%

Supplementary Figure S1: Kidney Transplant Graft Survival by KDPI Category in ECMO-Supported Donors

