



11-15-2023

Psychostimulant Misuse Among American Indian, Alaskan Native, or Native Hawaiian College Students in the U.S. From 2015 to 2019

Fares Qeadan

Sydney Ross

William A. Barbeau

Erin F. Madden

Kamilla L. Venner

See next page for additional authors

Follow this and additional works at: https://ecommons.luc.edu/publichealth_facpubs



Part of the [Community Health and Preventive Medicine Commons](#), and the [Public Health Education and Promotion Commons](#)

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Parkinson School of Health Sciences and Public Health by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

© The Author(s), 2023.

Authors

Fares Qeadan, Sydney Ross, William A. Barbeau, Erin F. Madden, Kamilla L. Venner, and Kevin English

Psychostimulant Misuse Among American Indian, Alaskan Native, or Native Hawaiian College Students in the U.S. From 2015 to 2019

Fares Qeadan¹, Sydney Ross¹, William A Barbeau¹, Erin F Madden², Kamilla L Venner³ and Kevin English⁴

¹Parkinson School of Health Sciences and Public Health, Loyola University Chicago, Maywood, IL, USA. ²Department of Family Medicine and Public Health Sciences, Wayne State University, Detroit, MI, USA. ³Department of Psychology, Center on Alcohol, Substance Use and Addictions (CASAA), University of New Mexico, Albuquerque, NM, USA. ⁴Albuquerque Area Southwest Tribal Epidemiology Center, Albuquerque, NM, USA.

Substance Abuse: Research and Treatment
Volume 17: 1–14
© The Author(s) 2023
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11782218231209667



ABSTRACT

INTRODUCTION: This study examines factors associated with psychostimulant misuse, including polysubstance use and social factors, among the understudied American Indian/Alaska Native/Native Hawaiian (AI/AN/NH) college student population.

METHODS: Data were from the 2015 to 2019 American College Health Association-National College Health Assessment IIc (ACHA-NCHA IIc) survey. Multivariable logistic regression models and odds ratios were used to estimate associations between psychostimulant misuse and potential risk and protective factors among AI/AN/NH college students, including licit and illicit substance use, social support, relationship factors, exposure to violence or abuse, mental health symptoms, drug and alcohol education, and sample demographics.

RESULTS: Opioid misuse among AI/AN/NH college students significantly increased the odds of using psychostimulants. Specifically, for cocaine use, the adjusted odds ratio (aOR) was 3.17 with a 95% confidence interval (C.I.) of 2.17 to 4.63; for methamphetamine use, the aOR was 38.87 (95% C.I. 19.24-78.52). For amphetamine misuse among non-Tobacco users, the aOR was 5.47 (95% C.I. 3.49-8.55), while among Tobacco users, the aOR was 2.65 (95% C.I. 2.07-3.41). For cocaine and other stimulant misuse, the aOR was 3.64 (95% C.I. 2.30-5.67). Additionally, the use of other types of licit and illicit substances was associated with greater odds of psychostimulant use and misuse. Conversely, factors such as age, living on campus, and residing in parental/guardian housing were linked with lower odds of psychostimulant use and misuse.

CONCLUSION: Substance use prevention and treatment interventions targeting AI/AN/NH college students should address polysubstance use, including the combined use of opioids and psychostimulants. Substance use interventions should not be siloed to focus narrowly on single substances but rather should leverage potential protective factors against substance use, such as promoting supportive campus and family living conditions and other social support networks, in broad efforts to reduce multiple forms of substance use among AI/AN/NH students.

KEYWORDS: Psychostimulant misuse, American Indian/Alaska Native/Native Hawaiian, potential risk factors, potential protective factors, opioid misuse

RECEIVED: June 22, 2023. **ACCEPTED:** October 4, 2023.

TYPE: Original Research

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Fares Qeadan, Parkinson School of Health Sciences and Public Health, Loyola University Chicago, 2160 S 1st Ave, Maywood, IL 60153, USA.
Email: fqeadan@luc.edu

Introduction

In the United States (U.S.), psychostimulants, including cocaine, methamphetamine, and amphetamine, are most commonly used among individuals aged 18 to 25.¹ Young adults enrolled in college have particularly high rates of misuse for some types of stimulants, with estimates suggesting they are 1.3 times more likely to engage in prescription stimulant misuse than their non-college peers.² Although Native Hawaiian and Pacific Islander (69%) and American Indian/Alaska Native (AI/AN) populations (62.4%) respectively have the highest and third highest rates of alcohol abstinence in the past month compared to each other racial/ethnic group, some measures

suggest that AI/AN populations are disproportionately affected by forms of psychostimulant misuse.³ For example, AI/AN populations had higher rates of past year central nervous system stimulant misuse in the 2021 National Survey on Drug Use and Health (NSDUH) (6.3% among AI/AN adults), compared to national rates (3.5% across all races/ethnicities).³ And fatal overdoses involving psychostimulants other than cocaine increased from 6.6 to 16.5 per 100 000 from 2018 to 2021 among AI/AN populations, compared to a rise of 3.9 to 9.8 deaths per 100 000 across the whole of the U.S. population during the same period.⁴ Recent spikes in overdose deaths are increasingly driven by combinations of psychostimulants and



opioids.^{5,6} While Black or African American populations suffered the highest overdose mortality rates from combining any kind of psychostimulant with opioids in 2021 (17.3 per 100 000), AI/AN populations experienced the greatest percent change increase (238.9%) in overdose due to the combination of these substances between 2018 through 2021 (3.6–12.2 per 100 000) compared to other racial groups.⁴ Yet, when restricting many sources of substance use survey data or vital records data to specific AI/AN or other indigenous subpopulations (eg, Native Hawaiians (NH)), or within those groups to only young adults, sample sizes are small, and data are often unavailable or suppressed, as in the case of both the NSDUH³ and CDC multiple causes of death databases.⁴ Thus, it is unclear how psychostimulant use may be related to other kinds of drug use or what potential protective factors may be leveraged in substance use interventions targeting indigenous young adults, especially college students.

Despite these data limitations, several studies have worked to identify likely risk and protective factors associated with psychostimulant use and misuse among AI/AN populations, as well as separately among diverse cohorts of young adults, including college students.^{7–11} However, few studies have contextualized the complex factors that may be related to psychostimulant use within a coherent conceptual model, and such studies have not specifically evaluated potential risk and protective factors among indigenous college students. Eitle and Eitle's stress process model suggests several significant factors contribute to methamphetamine use, including identifying as male, being in a romantic relationship, and facing stressful life events.¹² Several other empirical studies outline additional factors that may contribute to a fuller model of relevant factors contributing to or preventing psychostimulant use. In particular, studies note the use of licit and illicit substances, especially opioids, is associated with greater odds of engaging in psychostimulant misuse.^{10,13,14} In fact, Native individuals who engage in opioid misuse have been found to have 10.2 times greater odds of engaging in methamphetamine compared to those who do not engage in opioid misuse.¹⁴ Studies also note measures of relationships and social support, specifically multiple sexual partners and fraternity/sorority membership, are also associated with a greater likelihood of psychostimulant use among emerging adults.^{13,15–18} Furthermore, empirical findings have found individuals who identify as a sexual or gender minority, face anxiety, depression, or stress, or have been exposed to violence or abuse are more likely to engage in psychostimulant uses than their peers.^{11,14,19,20} Additionally, evidence indicates students with low GPAs or who live in rural areas use psychostimulants more than their peers.^{13,16,21,22} Conversely, empirical literature indicates that living with a parent/guardian, living in on-campus housing, or receiving education on the dangers of drug misuse may prevent college students from engaging in psychostimulant misuse.^{7,13,23–25} A modified stress process model that incorporates these additional factors provides a

framework to guide analyses of potential risk and protective factors that may relate to psychostimulant use among indigenous college students (Figure 1). Because risk and protective factors do not exist in isolation but rather function in a complex, cumulative manner, prevention and treatment programs for substance use are only effective if they account for multiple contextual spheres that influence health and behavior.²⁶

With a sizable population of indigenous young adults enrolled in post-secondary institutions (121 000 in 2020)²⁷ and empirical literature denoting high rates of stimulant misuse among college population samples and AI/AN populations 12 and older, it is likely that psychostimulant misuse and the health risks associated with them may be a public health concern for AI/AN/NH college students.^{2,13,28,29} This study hypothesizes that opioid misuse operates within a complex web of social factors as a likely risk factor for stimulant misuse among AI/AN/NH college students. Since little empirical literature exists on risk and protective factors associated with psychostimulant misuse among this subpopulation, evaluation of opioid misuse and the other variables outlined in this study's conceptual model (Figure 1) as potential risk and protective factors associated with psychostimulant use among AI/AN/NH college students offer key public health information that may inform subsequent interventions.

Methods

Data source and sample

This study utilized aggregated data from the American College Health Association–National College Health Assessment IIc (ACHA–NCHA IIc) survey administered between the fall of 2015 through the spring of 2019, as detailed on the ACHA website.³⁰ The ACHA–NCHA IIc survey is a nationally recognized research survey that captures a broad array of health statuses and behaviors among college students. The survey encompasses questions on health and safety, health education, substance use, sexual and reproductive health, physical and mental health, and impediments to academic performance. Data were collected for a total of 426 425 participants between the fall of 2015 and the spring of 2019. In this study, the sample was restricted to students identifying as American Indian, Alaskan Native, or Native Hawaiian (AI/AN/NH; $n = 8103$).

Measures

Psychostimulant misuse. The primary outcomes of interest in this study were 4 measures of psychostimulant use: (a) cocaine use, (b) methamphetamine use, (c) amphetamine misuse excluding methamphetamine, and (d) cocaine and other stimulant misuse. All 4 primary outcomes were binary variables, defined by a “Yes” or “No” response. As seen in Supplemental Table 1 (Table S1), the answers to the ACHA–NCHA IIc survey were utilized in the variable construction of all 4 primary outcomes.




Figure 1. Conceptual model on psychostimulant misuse among American Indian, Native Alaskan, or Native Hawaiian college students; adapted from Eitle and Eitle¹².

Opioid misuse. Opioid misuse was the primary exposure of interest in this study. As outlined in Supplemental Table 1 (Table S1), opioid misuse was a binary variable and was identified by a positive response to either of the following survey questions: “Within the last 30 days, on how many days did you use opiates (heroin, smack)?” and “In the last 12 months, have you taken any prescription pain killers (eg, OxyContin, Vicodin, Codeine) that were not prescribed to you?”

Control variables. Guided by the adapted stress process model illustrated in Figure 1, analysis models considered and adjusted for other behaviors and characteristics, including other (non-opioid) forms of licit and illicit substance use (ie, tobacco,¹³ alcohol,^{13,17} marijuana,^{9,13} and other substance use⁹). Analysis models also adjusted for measures of relationships and social support (ie, multiple past-year sexual partners³¹ and fraternity/sorority membership^{16,17}), exposure to violence or abuse,³² mental health symptoms (ie, anxiety¹⁹ and depression^{13,19}), stress,¹⁹ living/housing situation,^{7,24} and receipt of drug and alcohol education.¹³ Furthermore, demographic variables, including gender identity,³³ sexual orientation,²⁰ and educational achievement,¹⁶ were incorporated into the analysis. The detailed construction of these variables is outlined in Table S1.

Statistical analysis

Descriptive and inferential statistics were used to estimate the prevalence of psychostimulant misuse among AI/AN/NH college students and assess the association between opioid misuse

and cocaine, methamphetamine, amphetamine, and cocaine and other stimulant misuse among this cohort. Frequencies (n) and percentages (%) were used to summarize the marginal distributions of demographic and individual characteristics across opioid misuse. Furthermore, frequencies (n) and column percentages (%) were used to assess the distribution of the primary outcomes across opioid misuse.

Multivariable logistic regression models were constructed and used to evaluate the association between opioid misuse and each of the 4 measures of psychostimulant misuse, while controlling for other individual and societal factors identified in the conceptual framework (Figure 1). In addition to the primary analyses, we conducted an in-depth examination of interaction effects across all models. Our analysis explored interactions between opioid misuse and other substance use, substance use and demographic factors, substance use and social factors, and opioid misuse and mental health symptoms. From this examination, significant interactions were identified only in the model for amphetamine misuse, excluding methamphetamine.

To prevent overfitting, forward selection was implemented for each of the 4 multivariable logistic regression models. A significance level of 0.1 was used for entry into the models while forcing the variable “opioid misuse” to be included in each model. Additionally, a stop criterion of 3 was implemented during forward selection for the multivariable logistic regression model on methamphetamine in order to follow the “7 events per variable” rule of thumb.³⁴ Accounting for bias due to

rare events, all logistic regression models were run using Firth's regression.^{35,36} Unadjusted and adjusted odds ratios (uOR and aOR) and their corresponding 95% confidence intervals were reported along with *P*-values from Firth's penalized maximum likelihood estimates to discern associations. The Hosmer Lemeshow test and area under the receiver operating characteristic curve (AUC) were used to assess the goodness-of-fit of all 4 regression models. All statistical tests in this study are 2 sided with a 5% significance level, and all analyses were carried out through SAS version 9.4.

Results

This study consisted of 8103 AI/AN/NH college students, of which 2.1% (*n*=166) reported using cocaine, 0.8% (*n*=60) noted using methamphetamine, 8.7% (*n*=700) disclosed misuse of amphetamines excluding methamphetamine, and 1.5% (*n*=120) engaged in cocaine and other stimulant misuse (Table 2). The majority of participants did not engage in opioid misuse, with 91.3% reporting having not engaged in opioid misuse and only 8.8% reporting having misused opioids (Table 1). On average, the participants in this study were 23.4 years old (SD=7.5 years; Table 1). Most participants identified as cisgender female (64.9%), and heterosexuality (75.9%) was the predominant sexual orientation in this sample population (Table 1).

Opioid misuse

Opioid misuse was associated with increased psychostimulant use and misuse, including cocaine use, methamphetamine use, amphetamine misuse, and cocaine and other stimulant use. Frequencies and percentages revealed a crude association between opioid misuse and all 4 of the psychostimulant outcomes, with a mean difference in psychostimulant misuse of 15.1% across those who reported having engaged in opioid misuse and those who didn't (Table 2). Unadjusted logistic regression models found AI/AN/NH college students who engaged in opioid misuse to have 14.3, 61.1, 9.2, and 21.5 times the odds of using cocaine, methamphetamine, amphetamine, and cocaine and other stimulants, respectively than those who did not misuse opioids (Table 3).

After considering confounding factors in the adjusted models, the association between opioid misuse and psychostimulant use decreased but remained statistically significant (Tables 4–7). Adjusted logistic regression models revealed AI/AN/NH college students who engaged in opioid misuse to have 38.9, 3.6, and 3.2 times the odds of engaging in methamphetamine use, cocaine and other stimulant misuse, and cocaine use, respectively, than those who did not misuse opioids (Tables 4, 5 and 7). For amphetamine misuse, the association was modified by tobacco use such that the adjusted odds ratio among non-Tobacco users was 5.47 (95% C.I. 3.49–8.55), while among Tobacco users, it was 2.65 (95% C.I. 2.07–3.41) (Table 6).

Other substance use

In addition to opioid misuse, substance use (ie, tobacco and marijuana use) was identified with a significant increase in the odds of engaging in psychostimulant misuse among AI/AN/NH college students. Tobacco use was found to be associated with increased odds of engaging in psychostimulant misuse, with an adjusted odds ratio of 2.4 for cocaine and other stimulant misuse (Table 7) and 2.2 for cocaine (Table 4). Marijuana was also identified with an increased odds of stimulant misuse, with an adjusted odds ratio of 6.0 for cocaine use (Table 4) and 6.7 for cocaine and other stimulant misuse (Table 7). Furthermore, those who engaged in other substance use (ie, hallucinogens or club drugs) were 3.8 times more likely to engage in amphetamine misuse (Table 6) and 7.9 times more likely to use cocaine and other stimulants (Table 7), than those who did not engage in other substance use. Additionally, the use of alcohol was found to be associated with 51% greater odds of engaging in amphetamine misuse (Table 6).

Age, on-campus housing, and parental/guardian housing

Amongst the potential risk factors identified, 2 socio-demographic characteristics were found to be associated with a decreased odds of engagement in psychostimulant misuse amongst AI/AN/NH college students. Age was identified with a significant decrease in the odds of engagement in cocaine use and amphetamine misuse. Every 1-year increase in age was associated with roughly a 5% decrease in the odds of using cocaine among all users and 7.5% in the odds of misusing amphetamine among tobacco users (Tables 4 and 6). Age was also found to be associated with a decreased odds of engagement in cocaine and other stimulant misuse but this association was on the boundary of statistical significance (aOR: 0.968; 95% C.I. 0.935–1.002; *P*-value = .0626; Table 7). Living in on-campus housing or parental/guardian housing may also serve as a potential protective factor against psychostimulant misuse. Individuals living on-campus were found to have odds of engaging in amphetamine misuse 25.0% lower than students living off-campus without a parent/guardian (Table 6). Living on campus was also found to be associated with a decrease in odds of engagement in cocaine use, but this association was on the boundary of statistical significance (aOR: 0.656; 95% C.I. 0.430–1.003; *P*-value = .0518; Table 4). Living in parental/guardian housing was found to be associated with a greater decrease in odds of engagement in stimulant misuse than living on campus. Living in parental/guardian housing was found to be associated with 43.4% and 78.0% lower odds of engaging in amphetamine and cocaine use, respectively (Tables 4 and 6).

Discussion

This study offers foundational estimates on the prevalence of psychostimulant use and misuse among AI/AN/NH college

Table 1. Demographic characteristics of American Indian, Alaskan Native, or Native Hawaiian college students in the U.S. by opioid misuse: ACHA-NCHA IIc data from 2015 to 2019.

DEMOGRAPHIC CHARACTERISTICS	OPIOID MISUSE ^a		P-VALUE ^b	TOTAL
	NO X (SD)	YES X (SD)		X (SD)
Age	23.3 (7.5)	24.0 (8.1)	0.0268	23.4 (7.5)
	n (% ^c)	n (% ^c) ^a	P-value ^d	n (% ^c)
Total	7386 (91.3)	708 (8.8)		
Gender identity			<0.0001	
Cisgender female	4821 (65.5)	414 (59.0)		5235 (64.9)
Cisgender male	2158 (29.3)	211 (30.1)		2369 (29.4)
Gender diverse	203 (2.8)	34 (4.8)		237 (2.9)
Transgender	177 (2.4)	43 (6.1)		220 (2.7)
Sexual orientation			<0.0001	
Heterosexual	5644 (77.0)	455 (65.1)		6099 (75.9)
Gay/Lesbian	239 (3.3)	31 (4.4)		270 (3.4)
Bisexual	632 (8.6)	88 (12.6)		720 (9.0)
Unsure/Other	818 (11.2)	125 (17.9)		943 (11.7)
Tobacco use			<0.0001	
No	4572 (62.1)	191 (27.1)		4763 (59.0)
Yes	2791 (37.9)	515 (73.0)		3306 (41.0)
Alcohol use			<0.0001	
No	1729 (23.5)	69 (9.8)		1798 (22.3)
Yes	5627 (76.5)	636 (90.2)		6263 (77.7)
Marijuana use			<0.0001	
No	4341 (59.1)	156 (22.2)		4497 (55.8)
Yes	3008 (40.9)	548 (77.8)		3556 (44.2)
Other substance use ^e			<0.0001	
No	6313 (85.5)	208 (29.4)		6521 (80.6)
Yes	1073 (14.5)	500 (70.6)		1573 (19.4)
Multiple past-year sexual partners			<0.0001	
No	5394 (74.8)	383 (55.8)		5777 (73.1)
Yes	1818 (25.2)	303 (44.2)		2122 (26.9)
Fraternity or sorority member			0.0004	
No	6776 (92.0)	619 (88.1)		7395 (91.6)
Yes	593 (8.1)	84 (12.0)		677 (8.4)

(Continued)

Table 1. (Continued)

DEMOGRAPHIC CHARACTERISTICS	OPIOID MISUSE ^a		P-VALUE ^b	TOTAL
	NO	YES		\bar{X} (SD)
	\bar{X} (SD)	\bar{X} (SD)		
Low educational achievement ^f			<0.0001	
No	7090 (96.4)	659 (93.3)		7749 (96.1)
Yes	69 (0.9)	29 (4.1)		98 (1.2)
N/A	196 (2.7)	18 (2.6)		214 (2.7)
Depression			<0.0001	
No	5261 (71.4)	340 (48.1)		5601 (69.4)
Yes	2106 (28.6)	367 (51.9)		2473 (30.63)
Anxiety			<0.0001	
No	5653 (76.5)	434 (61.3)		6087 (75.2)
Yes	1733 (23.5)	274 (38.7)		2007 (24.8)
Stress			<0.0001	
No	2444 (33.1)	171 (24.2)		2615 (32.4)
Yes	4930 (66.9)	536 (75.8)		5466 (67.6)
Exposure to violence or abuse			<0.0001	
No	4729 (64.0)	252 (35.6)		4981 (61.6)
Yes	2656 (36.0)	456 (64.4)		3112 (38.5)
Living in a rural environment			0.4020	
No	7307 (99.1)	702 (99.4)		8009 (99.2)
Yes	64 (0.9)	4 (0.6)		68 (0.8)
Housing/living situation				
Off-campus without parent(s)/guardian(s)	2898 (39.3)	291 (41.2)	<0.0001	3189 (39.5)
Fraternity or sorority housing	81 (1.1)	20 (2.8)		101 (1.3)
On-campus	2718 (36.9)	220 (31.2)		2938 (36.4)
With parent(s)/guardian(s)	1132 (15.4)	110 (15.6)		1242 (15.4)
Other	537 (7.3)	65 (9.2)		602 (7.5)
Receipt of drug and alcohol education			0.0017	
No	2065 (28.1)	237 (33.7)		2302 (28.6)
Yes	5288 (71.9)	467 (66.3)		5755 (71.4)

Abbreviations: \bar{x} , sample mean; SD, sample standard deviation; n, sample size.

Bold P-values indicates statistical significance at the 5% significance level.

^aOpioid misuse includes illicit/non-prescription and prescription opioids.

^bSatterthwaite P-value from 2 sample t-test.

^cColumn percentage.

^dP-value from Chi-Square test for independence.

^eOther substance misuse includes use of sedatives, hallucinogens, anabolic steroids, inhalants, other club drugs, other illegal drugs, or misuse of prescription antidepressants, erectile dysfunction drugs, or sedatives.

^fLow educational achievement is denoted by an approximate cumulative grade average of D/F.

Table 2. Psychostimulant misuse among American Indian, Alaskan Native, or Native Hawaiian college students in the U.S. by opioid misuse: ACHA-NCHA IIc data from 2015 to 2019.

PRIMARY OUTCOMES	OPIOID MISUSE ^a		P-VALUE ^c	TOTAL
	NO	YES		N (% ^b)
	N (% ^b)	N (% ^b)		
Cocaine use			<0.0001	
No	7262 (99.0)	613 (87.1)		7875 (97.9)
Yes	75 (1.0)	91 (12.9)		166 (2.1)
Methamphetamine use			<0.0001	
No	7341 (99.9)	651 (92.7)		7992 (99.3)
Yes	9 (0.1)	51 (7.3)		60 (0.8)
Amphetamine misuse, excluding methamphetamine			<0.0001	
No	6945 (94.1)	447 (63.1)		7392 (91.4)
Yes	439 (6.0)	261 (36.9)		700 (8.7)
Cocaine and other stimulant use ^d			<0.0001	
No	7295 (99.4)	626 (89.0)		7921 (98.5)
Yes	42 (0.6)	78 (11.0)		120 (1.5)

n=Sample size. Bold P-values indicates statistical significance at the 5% significance level.

^aOpioid misuse includes illicit/non-prescription and prescription opioids.

^bColumn percentage.

^cP-value from Chi-Square test for independence.

^dCocaine and other stimulant use includes cocaine, methamphetamine, other amphetamines, and misuse of prescription stimulants.

Table 3. Unadjusted odds ratios and confidence intervals (C.I.) for psychostimulant misuse among American Indian, Alaskan Native, or Native Hawaiian college students in the U.S. who misuse opioids: ACHA-NCHA IIc survey, 2015 to 2019.

	COCAINE USE	METHAMPHETAMINE USE	AMPHETAMINE MISUSE, EXCLUDING METHAMPHETAMINE	COCAINE AND OTHER STIMULANT MISUSE ^a
	UOR ^b (95% C.I.)	UOR ^b (95% C.I.)	UOR ^b (95% C.I.)	UOR ^b (95% C.I.)
	P ^c	P ^c	P ^c	P ^c
Opioid misuse ^d				
No	Ref = 1	Ref = 1	Ref = 1	Ref = 1
Yes	14.345 (10.461-19.673) <0.0001	61.088 (30.432-122.624) <0.0001	9.236 (7.710-11.063) <0.0001	21.509 (14.677-31.521) <0.0001

Bold indicates statistical significance at the 5% significance level.

^aCocaine and other stimulant use includes cocaine, methamphetamine, other amphetamines, and misuse of prescription stimulants.

^bUnadjusted odds ratio.

^cP-value from Firth's Penalized Maximum Likelihood Estimate.

^dOpioid misuse includes illicit/non-prescription and prescription opioids.

students who are routinely excluded from publicly available data due to small sample sizes. Adjusted analyses in this study also identified behaviors, characteristics, and other factors associated with both increased and decreased odds of psychostimulant use and misuse. Statistical tests indicate that opioid misuse is associated with a significant increase in the odds of engaging in cocaine use, methamphetamine use, amphetamine misuse, and cocaine and other stimulant misuse. A pivotal observation, however, was the modifying effect of tobacco use on the

relationship between opioid misuse and amphetamine misuse. Specifically, the risk associated with opioid misuse on amphetamine misuse was notably higher among non-tobacco users compared to those who use tobacco. This interaction features the intricate relationship between various substances and their combined effects on psychostimulant misuse. Additionally, adjustment for covariates revealed several other significant factors that impact the odds of engaging in psychostimulant use, including other forms of substance use (licit and illicit),

Table 4. Adjusted odds ratios and confidence intervals (C.I.) of cocaine use for opioid misuse and confounding variables, among American Indian, Alaskan Native, or Native Hawaiian college students in the U.S.: ACHA-NCHA IIc survey, 2015 to 2019.

		COCAINE USE
		AOR ^a (95% C.I.) P ^b
Opioid misuse ^c	No	Ref= 1
	Yes	3.172 (2.171-4.634) <0.0001
Age		0.953 (0.922-0.986) 0.0051
Gender	Cisgender female	Ref= 1
	Cisgender male	1.627 (1.098-2.411) 0.0153
	Gender diverse	2.482 (1.140-5.403) 0.0220
	Transgender	1.815 (0.848-3.883) 0.1246
Tobacco use	No	Ref= 1
	Yes	2.160 (1.204-3.876) 0.0098
Marijuana use	No	Ref= 1
	Yes	6.019 (2.454-14.765) <0.0001
Other substance use ^d	No	Ref= 1
	Yes	4.582 (2.825-7.431) <0.0001
Multiple past-year sexual partners	No	Ref= 1
	Yes	2.335 (1.590-3.428) <0.0001
Fraternity or sorority member	No	Ref= 1
	Yes	2.814 (1.814-4.366) <0.0001
Anxiety	No	Ref= 1
	Yes	1.365 (0.942-1.977) 0.1001
Exposure to violence or abuse	No	Ref= 1
	Yes	1.466 (0.971-2.215) 0.0689
Housing/living situation	Off-campus without parent(s)/guardian(s)	Ref= 1
	Fraternity or sorority housing	1.705 (0.763-3.815) 0.1937
	On-campus	0.656 (0.430-1.003) 0.0518
	With parent(s)/guardian(s)	0.220 (0.090-0.535) 0.0009
	Other	0.721 (0.336-1.544) 0.3992

Bold indicates statistical significance at the 5% significance level and italic indicates being on the boundary of statistical significance ($.05 < P < .10$). The Hosmer-Lemeshow test showed adequate fit ($P = .4904$), and the AUC (0.928) showed excellent fit and classification ability of the multivariable model.

^aAdjusted odds ratio.

^bP-value from Firth's Penalized Maximum Likelihood Estimate.

^cOpioid misuse includes illicit/non-prescription and prescription opioids.

^dOther substance misuse includes use of sedatives, hallucinogens, anabolic steroids, inhalants, other club drugs, other illegal drugs, or misuse of prescription antidepressants, erectile dysfunction drugs, or sedatives.

Table 5. Adjusted odds ratios and confidence intervals (C.I.) of methamphetamine use for opioid misuse and confounding variables, among American Indian, Alaskan Native, or Native Hawaiian college students in the U.S.: ACHA-NCHA IIc survey, 2015 to 2019.

		METHAMPHETAMINE USE
		AOR ^a (95% C.I.)
		P ^b
Opioid misuse ^c	No	Ref = 1
	Yes	38.865 (19.236-78.522) <0.0001
Gender	Cisgender female	Ref = 1
	Cisgender male	6.763 (3.112-14.696) <0.0001
	Gender diverse	21.652 (8.448-55.490) <0.0001
	Transgender	19.479 (8.090-46.900) <0.0001
Anxiety	No	Ref = 1
	Yes	4.259 (2.327-7.795) <0.0001

Bold indicates statistical significance at the 5% significance level. The Hosmer-Lemeshow test showed adequate fit ($P = .6800$), and the AUC (0.953) showed excellent fit and classification ability of the multivariable model.

^aAdjusted odds ratio.

^bP-value from Firth's Penalized Maximum Likelihood Estimate.

^cOpioid misuse includes illicit/non-prescription and prescription opioids.

relationships, social support, and living context. These factors may be useful for public health interventions seeking to support the health and wellbeing of indigenous post-secondary students.

Adjusted analyses revealed psychostimulant misuse to be significantly higher among AI/AN/NH college students who also reported tobacco use, marijuana use, and other substance use. The study found indigenous college students who engaged in tobacco, marijuana, or other substance use to be more likely to also engage in psychostimulant misuse. This aligns with existing literature denoting increased odd of psychostimulant misuse among AI/AN individuals who use nicotine (OR: 31.6; 95% C.I. 1.87-5.35), cannabis (OR: 7.39; 95% C.I. 2.28-23.96), or engage in prescription tranquilizers or sedative misuse (OR: 13.0; 95% C.I. 5.72-29.6), compared to AI/AN individuals who do not use these substances.¹⁴ Amongst all college students, the odds of co-use of psychostimulants and other substances may be substantially higher among AI/AN/NH college students, with a recent study finding college students who misuse stimulants to be 0.34 times less likely to use tobacco ($P < .001$) and 0.17 times less like to use marijuana.³⁷ The high prevalence of psychostimulant misuse among AI/AN/NH college students who engage in other forms of drug use, may be related to higher estimated rates of tobacco, marijuana, and illicit substance use among indigenous populations compared to other races and ethnicities.^{28,38} With evidence illustrating a significant association between licit and illicit substance use and psychostimulant misuse, interventions may

benefit from including education about the risks of polysubstance use, skills training to resist peer pressure to engage in polysubstance use, and counseling to address the underlying issues that may contribute to polysubstance use.

The observed interaction between tobacco use and opioid misuse in relation to amphetamine misuse is particularly intriguing. One possible explanation for this interaction could be the underlying neurobiological mechanisms. Both opioids and nicotine (from tobacco) act on the brain's reward system, albeit through different pathways.^{39,40} Chronic use of either substance can lead to alterations in this system, potentially affecting an individual's susceptibility to the effects of other drugs, such as amphetamines.⁴¹ For tobacco users, the concurrent use of opioids might not amplify the risk of amphetamine misuse to the same extent as in non-tobacco users because their reward system is already modulated by nicotine.^{40,42} Additionally, behavioral factors might play a role. Individuals who use tobacco might have different patterns of drug-seeking behavior or might be part of social networks where polysubstance use is less prevalent or less normalized. It is also possible that some unmeasured confounding factors, such as specific cultural or community norms around drug use, could influence this interaction. While our study sheds light on this interaction, further research is needed to fully understand the underlying mechanisms and implications.

Social support and relationship factors also showed significant associations with psychostimulant use in the analyses. Namely, AI/AN/NH college students who had multiple

Table 6. Adjusted odds ratios and confidence intervals (C.I.) of amphetamine misuse for opioid misuse and confounding variables, among American Indian, Alaskan Native, or Native Hawaiian college students in the U.S.: ACHA-NCHA IIc survey, 2015 to 2019.

		AMPHETAMINE MISUSE, EXCLUDING METHAMPHETAMINE
		AOR ^a (95% C.I.)
		P ^b
Opioid misuse ^c among non- Tobacco users	No	Ref= 1
	Yes	5.467 (3.494-8.554) <0.0001
Opioid misuse ^c among Tobacco users	No	Ref= 1
	Yes	2.654 (2.066-3.408) <0.0001
Age among non- Tobacco users		1.002 (0.978 1.026) 0.9681
Age among Tobacco users		0.925 (0.904-0.948) <0.0001
Sexual orientation	Heterosexual	Ref= 1
	Gay/Lesbian	0.615 (0.363-1.041) 0.0621
	Bisexual	0.645 (0.477-0.874) 0.0042
	Unsure/Other	1.005 (0.976-1.306) 0.9941
Alcohol use	No	Ref= 1
	Yes	1.506 (0.976-2.322) 0.0589
Marijuana use	No	Ref= 1
	Yes	2.090 (1.589-2.748) <0.0001
Other substance use ^d	No	Ref= 1
	Yes	3.753 (3.041-4.633) <0.0001
Multiple past-year sexual partners	No	Ref= 1
	Yes	2.014 (1.661-2.442) <0.0001
Fraternity or sorority member	No	Ref= 1
	Yes	1.907 (1.453-2.503) <0.0001
Anxiety	No	Ref= 1
	Yes	1.323 (1.088-1.610) 0.0052
Exposure to violence or abuse	No	Ref= 1
	Yes	1.567 (1.286-1.911) <0.0001
Housing/living situation	Off-campus without parent(s)/guardian(s)	Ref= 1
	Fraternity or sorority housing	1.493 (0.819-2.720) 0.2014
	On-campus	0.746 (0.597-0.933) 0.0093
	With parent(s)/guardian(s)	0.566 (0.411-0.779) 0.0004
	Other	0.725 (0.485-1.084) 0.1092

Bold indicates statistical significance at the 5% significance level and italic indicates being on the boundary of statistical significance ($.05 < P < .10$). The Hosmer-Lemeshow test showed adequate fit ($P = .1832$), and the AUC (0.871) showed good fit and classification ability of the multivariable model.

^aAdjusted odds ratio.

^bP-value from Firth's Penalized Maximum Likelihood Estimate.

^cOpioid misuse includes illicit/non-prescription and prescription opioids.

^dOther substance misuse includes use of sedatives, hallucinogens, anabolic steroids, inhalants, other club drugs, other illegal drugs, or misuse of prescription antidepressants, erectile dysfunction drugs, or sedatives.

Table 7. Adjusted odds ratios and confidence intervals (C.I.) of cocaine and other stimulant misuse for opioid misuse and confounding variables, among American Indian, Alaskan Native, and Native Hawaiian college students in the U.S.: ACHA-NCHA IIc survey, 2015 to 2019.

		COCAINE AND OTHER STIMULANT MISUSE
		AOR ^a (95% C.I.)
		P ^b
Opioid misuse ^c	No	Ref = 1
	Yes	3.636 (2.330-5.672) <0.0001
Age		0.968 (0.935-1.002) 0.0626
Gender	Cisgender female	Ref = 1
	Cisgender male	2.348 (1.467-3.758) 0.0004
	Gender diverse	3.844 (1.623-9.105) 0.0022
	Transgender	1.757 (0.722-4.272) 0.2140
Tobacco use	No	Ref = 1
	Yes	2.420 (1.105-5.301) 0.0271
Marijuana	No	Ref = 1
	Yes	6.658 (1.823-24.323) 0.0041
Other substance use ^d	No	Ref = 1
	Yes	7.893 (3.788-16.445) <0.0001
Multiple past-year sexual partners	No	Ref = 1
	Yes	2.816 (1.735-4.569) <0.0001
Fraternity or sorority member	No	Ref = 1
	Yes	3.620 (2.255-5.809) <0.0001
Anxiety	No	Ref = 1
	Yes	1.914 (1.230-2.979) 0.0040
Exposure to violence or abuse	No	Ref = 1
	Yes	1.986 (1.155-3.417) 0.0131

Bold indicates statistical significance at the 5% significance level and italic indicates being on the boundary of statistical significance ($.05 < P < .10$). The Hosmer-Lemeshow test showed adequate fit ($P = .9858$), and the AUC (0.955) showed excellent fit and classification ability of the multivariable model.

^aAdjusted odds ratio.

^bP-value from Firth's Penalized Maximum Likelihood Estimate.

^cOpioid misuse includes illicit/non-prescription and prescription opioids.

^dOther substance misuse includes use of sedatives, hallucinogens, anabolic steroids, inhalants, other club drugs, other illegal drugs, or misuse of prescription antidepressants, erectile dysfunction drugs, or sedatives.

past-year sexual partners or who were a member of a fraternity or sorority were more than 2 times more likely to engage in psychostimulant misuse. These findings are consistent with other empirical findings on Greek life involvement and

multiple sexual partners.^{13,15-17,43-46} These results support that stimulant misuse is heightened among college students with multiple past-year sexual partners or involved in a fraternity/sorority. In turn, such findings suggest that stimulant misuse

interventions should consider peer relationships and student organization factors in addressing substance use risks and psychosocial factors that may impact individual coping skills surrounding substance misuse.^{44,46} Interventions targeting these social factors may include providing education and support to students in fraternities and sororities and promotion of healthy relationship behaviors.

Analogous with other empirical literature, this study identified age, living on campus, and living with parent(s)/guardian(s), to be associated with significant decreases in the odds of engaging in cocaine use and amphetamine misuse.^{7,24} In a recent prevention study, social network analysis revealed close contact and sharing of information with peers and cousins, which could support youth in making prosocial decisions.⁴⁷ Living on campus or with parent(s)/guardian(s) may provide support, clear disapproval and sanctions, and monitoring/supervision that reduces individuals' likelihood of engaging in stimulant use and misuse. One study showed parental monitoring to be a stronger protective factor for female AI youth in terms of the onset of alcohol and marijuana use.⁴⁸ Importantly, the quality of parent-child relationships has been shown to be associated with lower rates of substance use in Bahamian youth, such that open communication and rules about curfew were protective⁴⁹ and decreasing communication and parental monitoring were associated with higher rates of alcohol and marijuana use in general population urban eighth graders.⁵⁰ Furthermore, a qualitative study of AI youth found immediate, extended family, and cousins to serve as both risk and protective factors for AI youth engagement in substance use.⁵¹ Specifically, AI youth reported lack of parental supervision as a risk factor for using substances.⁵¹ In contrast, perspectives that parents were strict were seen as protective by these youth.⁵¹ The impact of these nuances in parent-child relationships on substance use highlights the importance of incorporating these dynamics in substance use prevention programs. In the context of post-secondary students, benefits could be yielded from interventions that target younger students and those living off-campus or without a parent/guardian. Moreover, interventions aim to promote supportive campus and family living conditions may serve as potential protective factors against psychostimulant use among college students.

Furthermore, anxiety was found to be associated with an increased odds of psychostimulant use and misuse among AI/AN/NH college students. These findings parallel existing literature on the association between mental health problems and engagement in substance use.^{13,52} One study found college students with a generalized anxiety disorder to have 2.8 (95% C.I. 1.9–4.0), 1.4 (95% C.I. 1.1–1.9), and 1.8 (95% C.I. 1.2–2.6) times the odds of engaging in any cigarette smoking, any binge drinking, and any frequent binge drinking, respectively.⁵² Research has identified that individuals with mental health symptoms, such as social

anxiety, may engage in substance use as a means of managing and coping with mental health symptoms.⁵³ Furthermore, it has been found that neuroinflammatory dysfunction caused by stress and substance use can work synergistically and may contribute to anxiety and substance misuse comorbidities.⁵⁴ These findings emphasize the importance of mental health support, such as counseling and therapy, to address mental health symptoms and psychostimulant use among college students.

There are several limitations to this study and study design that must be taken into account when interpreting the results. For one, as a self-report assessment, the NCHA-ACHA IIc survey may be adversely impacted by respondent recall bias and non-response. Furthermore, while the NCHA-ACHA IIc survey covers over 100 post-secondary institutions, including tribal colleges and universities, this data may not be representative of all AI/AN/NH U.S. college students. The study's cross-sectional design prevents inferences of causality between opioid misuse and all other adjusted variables. Additionally, the study is limited to variables covered in the NCHA-ACHA IIc survey and there is no distinction between commercial and traditional tobacco use. Despite these limitations, the study's results fill in a gap in empirical literature on the prevalence and associated risk factors of stimulant use and misuse among AI/AN/NH college students.

Conclusion

The results of this study demonstrate high co-use of opioid misuse and psychostimulant misuse, along with other forms of licit and illicit substance use, among AI/AN/NH college students. A notable interaction was observed between opioid misuse and tobacco use, highlighting the nuanced effects of polysubstance use on amphetamine misuse. These findings suggest that substance use prevention and treatment interventions targeting AI/AN/NH college students should address polysubstance use, including combined use of opioids and psychostimulants, because these are increasingly driving overdose deaths nationally. Substance use interventions should not be siloed to focus narrowly on single substances and should instead leverage potential protective factors, such as promoting supportive campus and family living conditions, and address risk factors, such as participation in Greek life, in broad efforts to reduce multiple forms of substance use among AI/AN/NH students.

Acknowledgements

This study uses data from the American College Health Association National College Health Assessment (ACHA-NCHA). The opinions, findings, and conclusions reported in this article are those of the authors and are in no way meant to represent the corporate opinions, views, or policies of the ACHA. ACHA does not warrant nor assume any liability or responsibility for the accuracy, completeness, or usefulness of

any information presented in this article. We thank Rona Bern for working on the initial literature review.

Author Contributions

Each author contributed to the submission in the following manner. Study Concept: FQ Data Management: SR, Statistical Analysis: FQ, WAB, SR Interpretation of data: FQ, WAB, SR Initial drafting of manuscript: FQ, SR Editing/revision of manuscript: FQ, SR, WAB, EFM, KLV, KE. All authors have made significant contributions to this manuscript and have approved this version for submission.

Author Disclosures

FQ, EFM, KLV, and KE are partially funded from NIH grants (1) 5R61DA049382: Leveraging CDC opioid overdose surveillance funding from the Albuquerque area southwest tribal epidemiology center to create tribal data and culturally center medications for opioid use disorder; and (2) R01DA057658: Weighting Longitudinal Data to Access Opioid Analgesia Tapering Outcomes among Patients with Co-occurring Chronic Pain and Substance Use Disorder.

Supplemental Material

Supplemental material for this article is available online.

REFERENCES

- Substance Abuse and Mental Health Services Administration. *Key Substance Use and Mental Health Indicators in the United States: Results from the 2020 National Survey on Drug Use and Health*, Center for Behavioral Health Statistics and Quality & Substance Abuse Mental Health Services Administration; 2021. Accessed August 3, 2023. <https://www.samhsa.gov/data/sites/default/files/reports/rpt35325/NSDUHFRRPDFWHHTMLFiles2020/2020NSDUHFRR1PDFW102121.pdf>
- Ford JA, Pomykacz C. Non-Medical use of prescription stimulants: A comparison of college students and their Same-Age peers who do not attend college. *J Psychoactive Drugs*. 2016;48:253-260.
- Substance Abuse and Mental Health Services Administration. 2021 NSDUH Detailed Tables. 2021. Accessed August 3, 2023. <https://www.samhsa.gov/data/report/2021-nsduh-detailed-tables>
- CDC WONDER. Mortality 2018-2021. 2023. Accessed August 3, 2023. <http://wonder.cdc.gov/mcd-icd10-expanded.html>
- Kariisa M, Seth P, Scholl L, Wilson N, Davis NL. Drug overdose deaths involving cocaine and psychostimulants with abuse potential among racial and ethnic groups - United States, 2004-2019. *Drug Alcohol Depend*. 2021;227:109001.
- Mattson CL, Tanz LJ, Quinn K, et al. Trends and geographic patterns in drug and synthetic opioid overdose deaths — United States, 2013-2019. *MMWR Morb Mortal Wkly Rep*. 2021;70:202-207.
- DiBello AM, Benz MB, Miller MB, Merrill JE, Carey KB. Examining residence status as a risk factor for health risk behaviors among college students. *J Am Coll Health*. 2018;66:187-193.
- Herman-Stahl MA, Krebs CP, Kroutil LA, Heller DC. Risk and protective factors for nonmedical use of prescription stimulants and methamphetamine among adolescents. *J Adolesc Health*. 2006;39:374-380.
- Herman-Stahl MA, Krebs CP, Kroutil LA, Heller DC. Risk and protective factors for methamphetamine use and nonmedical use of prescription stimulants among young adults aged 18 to 25. *Addict Behav*. 2007;32:1003-1015.
- Russell K, Dryden DM, Liang Y, et al. Risk factors for methamphetamine use in youth: a systematic review. *BMC Pediatr*. 2008;8:48.
- Stone AL, Becker LG, Huber AM, Catalano RF. Review of risk and protective factors of substance use and problem use in emerging adulthood. *Addict Behav*. 2012;37:747-775.
- Eitle DJ, Eitle TM. Methamphetamine use among rural white and Native American adolescents: an application of the stress process model. *J Drug Educ*. 2013;43:203-221.
- Benson K, Flory K, Humphreys KL, Lee SS. Misuse of stimulant medication among college students: a comprehensive review and meta-analysis. *Clin Child Fam Psychol Rev*. 2015;18:50-76.
- Coughlin LN, Lin LA, Jannausch M, Ilgen MA, Bonar EE. Methamphetamine use among American Indians and Alaska Natives in the United States. *Drug Alcohol Depend*. 2021;227:108921.
- DeSantis AD, Webb EM, Noar SM. Illicit use of prescription ADHD medications on a college campus: A Multimethodological Approach. *J Am Coll Health*. 2008;57:315-324.
- McCabe SE, Teter CJ, Boyd CJ. Medical use, illicit use and diversion of prescription stimulant medication. *J Psychoactive Drugs*. 2006;38:43-56.
- Norman L, Ford J. Undergraduate prescription stimulant misuse: the impact of academic strain. *Subst Use Misuse*. 2018;53:1482-1491.
- Yen CF. Relationship between methamphetamine use and risky sexual behavior in adolescents. *Kaohsiung J Med Sci*. 2004;20:160-165.
- Lueck JA, Costantini R, Knobloch M. The Making of an addiction: examining psychological determinants of prescription stimulant abuse among college students. *Health Commun*. 2020;35:946-954.
- Philbin MM, Greene ER, Martins SS, LaBossier NJ, Mauro PM. Medical, nonmedical, and illegal stimulant use by sexual identity and Gender. *Am J Prev Med*. 2020;59:686-696.
- Lambert D, Gale JA, Hartley D. Substance Abuse by youth and Young Adults in rural America. *J Rural Health*. 2008;24:221-228.
- Rabiner DL, Anastopoulos AD, Costello EJ, et al. Motives and perceived consequences of nonmedical ADHD medication use by college students: are students treating themselves for attention problems? *J Atten Disord*. 2009;13:259-270.
- Arria AM, Caldeira KM, Vincent KB, O'Grady KE, Wish ED. Perceived harmfulness predicts nonmedical use of prescription drugs among college students: Interactions with sensation-seeking. *Prev Sci*. 2008;9:191-201.
- Bavarian N, Flay BR, Ketcham PL, Smit E. Illicit use of prescription stimulants in a college student sample: A theory-guided analysis. *Drug Alcohol Depend*. 2013;132:665-673.
- Dussault CL, Weyandt LL. An examination of prescription stimulant misuse and psychological variables among sorority and fraternity college populations. *J Atten Disord*. 2013;17:87-97.
- Substance Abuse and Mental Health Services Administration. Risk and Protective Factors. 2019. <https://www.samhsa.gov/sites/default/files/20190718-samhsa-risk-protective-factors.pdf>
- U.S. Department of Education. *Higher Education General Information Survey (HEGIS): Fall Enrollment in Colleges and Universities*. 2021. Retrieved August 3, 2023, from <https://nces.ed.gov/programs/digest/d21/>
- Center for Behavioral Health Statistics and Quality. Racial/Ethnic Differences in Substance Use, Substance Use Disorder, and Substance Use Treatment Utilization Among People Aged 12 or Older (2015-2019). 2021. Retrieved August 3, 2023, from <https://www.samhsa.gov/data/sites/default/files/reports/rpt35326/2021NSDUHSUChartbook102221B.pdf>
- National Center for Educational Statistics. College Enrollment Rates. 2022. Retrieved August 3, 2023, from <https://nces.ed.gov/programs/coe/indicator/cpb/college-enrollment-rate#suggested-citation>
- American College Health Association (ACHA). National College Health Assessment: Survey. 2015-2019. Accessed Sep 30, 2023. https://www.acha.org/NCHA/About_ACHA_NCHA/Survey/NCHA/About/Survey.aspx?hkey=7e9f6752-2b47-4671-8ce7-ba7a529c9934
- Benotsch EG, Koester S, Luckman D, Martin AM, Cejka A. Non-medical use of prescription drugs and sexual risk behavior in young adults. *Addict Behav*. 2011;36:152-155.
- Ogden SN, Dichter ME, Bazzi AR. Intimate partner violence as a predictor of substance use outcomes among women: A systematic review. *Addict Behav*. 2022;127:107214.
- De Pedro KT, Gilreath TD, Jackson C, Esqueda MC. Substance use among transgender students in California public middle and High Schools. *J Sch Health*. 2017;87:303-309.
- Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in logistic and Cox regression. *Am J Epidemiol*. 2007;165:710-718.
- Firth D. Bias reduction of maximum likelihood estimates. *Biometrika*. 1993;80:27-38.
- King G, Zeng L. Logistic regression in rare events data. *Polit Anal*. 2001;9:137-163.
- Cole VT, Hussong AM. Psychosocial functioning among college students who misuse stimulants versus other drugs. *Addict Behav*. 2020;105:106290.
- American Lung Association. Tobacco Use in Racial and Ethnic Populations. 2022. Accessed August 3, 2023. <https://www.lung.org/quit-smoking/smoking-facts/impact-of-tobacco-use/tobacco-use-racial-and-ethnic>
- Nestler EJ. Is there a common molecular pathway for addiction? *Nat Neurosci*. 2005;8:1445-1449.

40. Kohut SJ. Interactions between nicotine and drugs of abuse: a review of pre-clinical findings. *Am J Drug Alcohol Abuse*. 2017;43:155-170.
41. Koob GF, Volkow ND. Neurobiology of addiction: a neurocircuitry analysis. *Lancet Psychiatry*. 2016;3:760-773.
42. D'Souza MS, Markou A. Neuronal mechanisms underlying development of nicotine dependence: implications for novel smoking-cessation treatments. *Addict Sci Clin Pract*. 2011;6:4.
43. Bonar EE, Cunningham RM, Chermack ST, et al. Prescription Drug Misuse and sexual risk behaviors among adolescents and emerging adults. *J Stud Alcohol Drugs*. 2014;75:259-268.
44. Okumu M, Ombayo BK, Small E, Ansong D. Psychosocial syndemics and sexual risk practices among U.S. Adolescents: Findings from the 2017 U.S. Youth Behavioral Survey. *Int J Behav Med*. 2019;26:297-305.
45. Rabiner DL, Anastopoulos AD, Costello EJ, Hoyle RH, Swartzwelder HS. Predictors of nonmedical ADHD medication use by college students. *J Atten Disord*. 2010;13:640-648.
46. Tapert SF, Aarons GA, Sedlar GR, Brown SA. Adolescent substance use and sexual risk-taking behavior. *J Adolesc Health*. 2001;28:181-189.
47. Mason WA, Rentschler JK, Habecker P, Whitbeck LB. Social network analysis of diffusion among American Indian Youth in a culturally adapted, family-focused prevention program. *Prev Sci*. 2023;24:728-738.
48. Rusby JC, Light JM, Crowley R, Westling E. Influence of parent-youth relationship, parental monitoring, and parent substance use on adolescent substance use onset. *J Fam Psychol*. 2018;32:310-320.
49. Wang B, Stanton B, Li X, et al. The influence of parental monitoring and parent-adolescent communication on Bahamian adolescent risk involvement: a three-year longitudinal examination. *Soc Sci Med*. 2013;97:161-169.
50. Tobler AL, Komro KA. Trajectories of parental monitoring and communication and effects on drug use among urban young adolescents. *J Adolesc Health*. 2010;46:560-568.
51. Hurdle DE, Okamoto SK, Miles B. Family influences on alcohol and drug use by American Indian Youth: Implications for Prevention. *J Fam Soc Work*. 2003;7:53-68.
52. Cranford JA, Eisenberg D, Serras AM. Substance use behaviors, mental health problems, and use of mental health services in a probability sample of college students. *Addict Behav*. 2009;34:134-145.
53. Buckner JD, Morris PE, Abarno CN, Glover NI, Lewis EM. Biopsychosocial model social anxiety and substance use revised. *Curr Psychiatry Rep*. 2021;23:35.
54. Smiley CE, Wood SK. Stress- and drug-induced neuroimmune signaling as a therapeutic target for comorbid anxiety and substance use disorders. *Pharmacol Ther*. 2022;239:108212.