

NON-LINEAR TESTS OF WEAKLY EFFICIENT MARKETS: EVIDENCE FROM PORTUGAL^(*)

ANTÓNIO AFONSO^(**)

JOÃO TEIXEIRA^(***)

This version: 1998

JEL classification: C14, G14.

Keywords: market efficiency; non-linearity tests; stock market; Portugal.

Note: Correspondence should be addressed to António Afonso, Instituto Superior de Economia e Gestão, R. Miguel Lúpi 20, P - 1200 Lisbon, Portugal

^(*) The authors are grateful for helpful comments from Miguel St. Aubyn, and participants in conferences at ISEG, namely in the “II Conference on Complexity in Economics” and in the “1st CIEF Workshop on Financial Economics.” The usual disclaimer applies.

^(**) Instituto Superior de Economia e Gestão, Universidade Técnica de Lisboa and Instituto de Gestão do Crédito Público. Corresponding author (e-mail: aafonso@iseg.utl.pt).

^(***) Comissão do Mercado de Valores Mobiliários (e-mail: joaoteixeira@cmvm.pt).

Abstract: This paper investigates the existence of non-linear dependence in Portuguese financial time series namely stock exchange indexes returns. Non-linear dependence may exist in a series even if we have already concluded for the lack of linear dependence. If present, non-linear dependence would contradict the random walk model and the financial markets weak form efficiency hypothesis. Using daily observations for the period 1990-1997 some so-called non-linearity tests are performed in order to decide whether we can accept the weak form efficiency hypothesis. The results seem to confirm the existence of exploitable excess profit opportunities in the Portuguese stock market.

Resumo: Este trabalho investiga a existência de dependência não linear em séries financeiras portuguesas, concretamente para as rentabilidades de índices de acções. Uma série pode apresentar dependência não linear mesmo que se tenha concluído que não existe dependência linear. A existência de dependência não linear iria contrariar o modelo do passeio aleatório e a hipótese da eficiência de forma fraca dos mercados financeiros. Utilizando observações diárias para o período entre 1990 e 1997 foram levados a cabo “testes de não linearidade” no sentido de averiguar se é possível aceitar a hipótese de eficiência de forma fraca. Os resultados parecem confirmar a existência de oportunidades de lucros anormais no mercado accionista português.

No matter how we try to make it so, the world is not orderly: nature is not orderly, nor are the human creations called institutions. Economies and the capital markets are particularly lacking in orderliness. The capital markets are our own creation; yet we do not understand how they work. (...) current capital market theory is based on a linear view of the society. (...) However, people, and nature in general are nonlinear.

EDGAR PETERS

Chaos and Order in the Capital Markets (1996)

1 - Introduction

An efficient financial market can be described as one for which no deterministic pattern can be detected. Market efficiency implies the absence of pure arbitrage opportunities and denies the profitability by the use of historical data. Efficiency validation is sometimes reduced to test whether the returns generating process of a certain asset is deterministic (evidence against market efficiency) or stochastic (evidence for market efficiency).

Most papers concerning the issue of financial market efficiency used to test the weak form efficiency hypothesis by performing runs tests or autocorrelations tests. Usually we accept the existence of linear independence for a series P (for instance stock prices) when it is generated by a (logarithmic) random walk model, given by

$$\log P_t = C + \log P_{t-1} + u_t \quad (1)$$

where u_t is an independent and identically distributed (iid) random variable with zero

mean and finite variance (often called “white noise”) and C is a constant drift. Evidence that $\log P$ follows a random walk is a sign to accept the weak form efficiency hypothesis then returns, the log changes in prices, are unpredictable.¹

If the logarithmic price process follows a random walk then the present and past returns R_t , where

$$R_t = \log P_t - \log P_{t-1}, \quad (2)$$

are not associated with future returns and hence they have no predictability for future values of R . Therefore, R_t 's independence implies the existence of an efficient market and univariate time series methods won't succeed in capturing any returns process patterns.

However, even if there is no linear dependence that does not rule out non-linear dependence. In fact, non-linear dependence may exist in a series even if we have already concluded for the lack of linear dependence. If present, non-linear dependence would contradict the random walk model and the financial market weak form efficiency hypothesis. Checking if autocorrelation coefficients are not statistically different from zero is not enough. It is therefore necessary to test for the non-linearity of returns.

Brock, Hsieh and LeBaron (1991) mention several tests for non-linearity.² In our attempt to validate the weak form efficiency hypothesis, we are going to use the Engle test, the Tsay test, the Hinich bispectrum test, the Lyapunov exponent test and the BDS statistic.³

In the next section we introduce the data and describe the series main statistical features. Section three offers a summary description of the non-linearity tests used in the paper, section four presents the empirical results and section five concludes.

¹ For a comprehensive review on the efficient capital markets theory and test results see for instance Fama (1991).

² Other useful tests for detecting the presence of non-linearity are used by Lee, White and Granger (1993) and Brooks (1996).

³ Al-Lougahni and Chappell (1996) call this a new test in opposition to runs tests and autocorrelation tests.

2 - Preliminary Data Exploration

The assumption often assumed, in finance theory, that prices move randomly and that returns are identically distributed following a normal distribution is very handy even if quite unrealistic.⁴ The main reason for that assumption seems to be related to the fact that the normal distribution is characterised by only two parameters: its mean and its standard deviation. Actually, there is a large consensus among researchers on several stylised facts about the statistical properties of financial time series returns. For instance it is a well-known fact that returns distributions are not normal but rather thick-tailed (leptokurtic). Taylor (1986) presents a review of this topic.

In this study we analyse three financial time series: the BVL (Bolsa de Valores de Lisboa) General Index, the PSI20 (Portuguese Stock Index 20) Index and the BVL30 Index. These are composite stock market indexes that reflect the aggregate movements of the Portuguese stock market. Since the Portuguese financial market is a very thin one, individual stock prices don't seem to be a good choice.

The available data for those indexes allow us to use more than 1700 daily observations for BVL and around 1200 observations for the PSI20 and BVL30 series.⁵ The accurate samples and its main statistical features are described below on Table 1 and the three series are plotted on Figures 1, 2, and 3 (all figures are presented on the Annex).

Table 1
Descriptive Statistics for BVLG, PSI20 and BVL30

INDEX	SAMPLE		N	Mean	S. d.	Minimum	Maximum
	From	To					
BVLG	1990-01-10	1998-01-06	1793	909.17	333.88	537.20	2014.53
PSI20	1993-01-04	1998-01-06	1239	4764.17	1405.29	2917.56	9294.05
BVL30	1993-01-05	1998-01-06	1238	1922.99	672.32	980.14	3992.42

Notes: N is the number of observations in sample.

⁴ Afonso (1997) presents results for the absence of normality on Portuguese stock indexes.

⁵ The reader must be aware that until November 1997 Portuguese capital markets belonged to the so-called emerging markets category. Since December 2, 1997, the Portuguese Stock Exchange was included in the Morgan Stanley Capital Index for developed markets.

Table 2 gives a selection of descriptive statistics for the daily log returns. The log returns are also represented on Figures 4, 5, and 6. All three series returns show strong departure from normality, as the coefficients of skewness and kurtosis are statistically different from those of a normal distribution.⁶ All three series are leptokurtic and have asymmetric tails: PSI20 is skewed to the left and both BVLG and BVL30 are skewed to the right.

Table 2
Descriptive Statistics for BVLG, PSI20 and BVL30 Returns

INDEX	N	Mean	S. d.	Minimum	Maximum	Skewness	Kurtosis
BVLG	1792	0.00059692	0.0067877	-0.059529	0.075718	0.62951	21.2378
PSI20	1238	0.00091338	0.0080675	-0.070627	0.069413	-0.35051	14.6280
BVL30	1237	0.00111920	0.0076165	-0.068987	0.063209	0.28656	15.3008

Notes: N is the number of observations in sample.

3 - Non-linear Tests

A brief theoretical presentation of the non-linear tests is given in this section. The first two tests presented in this section are intended to test for neglected non-linearity and will be applied to the residuals of an AR(p) model. The number of lags was selected using the Schwarz Bayesian Information Criterion (SBIC).

3.1 - Engle's Test

Autoregressive conditional heteroscedasticity (ARCH) models were developed by Engle (1982) who also proposed a test that explicitly examines for non-linearity in the second moment. In its simplest form an ARCH (p) process can be written formally as

$$Y_t = \mathbf{b}_1 + \mathbf{b}_2 X_{2t} + \dots + \mathbf{b}_k X_{kt} + \mathbf{e}_t \quad (3)$$

$$\mathbf{e}_t \sim N(0, \mathbf{s}_t) \quad (4)$$

$$\mathbf{s}_t^2 = \mathbf{a}_0 + \mathbf{a}_1 \mathbf{e}_{t-1}^2 + \mathbf{a}_2 \mathbf{e}_{t-2}^2 + \dots + \mathbf{a}_p \mathbf{e}_{t-p}^2 \quad (5)$$

⁶ A normal distribution should have a zero skewness statistic and a kurtosis statistic of three.

The null hypothesis of no autocorrelation in the error variance is $H_0: \mathbf{a}_1 = \mathbf{a}_2 = \dots = \mathbf{a}_p = 0$, which if accepted would lead us to deny the existence of an ARCH (p) model. The procedure to test that hypothesis is as follows:

1. Regress Y_t linearly on X_t (if the information set, X_t , is restricted to the past observations of Y_t then we simply estimate an AR (p) process) and save the estimated residuals $\hat{\mathbf{e}}_t$.

2. Regress the squares of the estimated standard residuals $\hat{\mathbf{e}}_t^2$ on an intercept and p lagged values of $\hat{\mathbf{e}}_t^2$ as

$$\hat{\mathbf{e}}_t^2 = \mathbf{a}_0 + \mathbf{a}_1 \hat{\mathbf{e}}_{t-1}^2 + \mathbf{a}_2 \hat{\mathbf{e}}_{t-2}^2 + \dots + \mathbf{a}_p \hat{\mathbf{e}}_{t-p}^2 + \hat{\mathbf{h}}_t \quad (6)$$

and save the estimated residuals $\hat{\mathbf{h}}_t$.

3. Calculate the R^2 from the second regression and test the null hypothesis using the nR^2 statistic that follows a $X^2(p)$ distribution under the null of no ARCH dependence.

3.2 - Tsay's test

While the Engle test examines evidence for non-linearity in the variance the Tsay test checks for non-linearity on the mean.⁷ The procedure to compute the test proposed by Tsay (1986) is as follows:

1. Regress Y_t linearly on X_t (in practice we estimate an AR model if the information set, X_t , is restricted to the past observations of Y_t) and keep the estimated residuals $\hat{\mathbf{e}}_t$.

2. For each observation of Y_t build the vector Z_t of the cross products of past observations in other words, $Y_{t-i}Y_{t-j}$ for $i, j = 1, \dots, p$ where $i \geq j$. For example if $p = 2$ then $Z_t = [Y_{t-1}^2, Y_{t-1}Y_{t-2}, Y_{t-2}^2]^T$ (notice that our vector Z_t has $p(p+1)/2$ elements).

3. Regress the vector Z_t on the explanatory variables and save the estimated residuals $\hat{\mathbf{h}}_t$.

4. Regress the estimated residuals $\hat{\mathbf{e}}_t$ on \mathbf{h}_t

$$\hat{\mathbf{e}}_t = \mathbf{d}_0 + \mathbf{d}_1 \mathbf{h}_{t-1} + \mathbf{d}_2 \mathbf{h}_{t-2} + \dots + \mathbf{d}_p \mathbf{h}_{t-p} + \hat{\mathbf{x}}_t \quad (7)$$

and save the estimated residuals $\hat{\mathbf{x}}_t$.

5. Calculate the Tsay test statistic

$$\hat{F} = \frac{(\hat{\mathbf{e}}^T \mathbf{h})^T (\mathbf{h}^T \mathbf{h})^{-1} (\mathbf{h}^T \hat{\mathbf{e}}) / m}{(\hat{\mathbf{x}}^T \hat{\mathbf{x}}) / (n - p - m - 1)}, \quad (8)$$

where $m = p(p+1)/2$ and test the null hypothesis

$$H_0: \mathbf{d}_1 = \mathbf{d}_2 = \dots = \mathbf{d}_p = 0.$$

Tsay (1986) showed that the statistic (8) has an $F(m, n-p-m-1)$ distribution under the null hypothesis and is sensitive to departures from linearity in the mean.

3.3 - Hinich Bispectrum Test

The Hinich bispectrum test is used to estimate the bispectrum of a stationary time series and provides a direct test for non-linearity and also a direct test for Gaussianity.⁸ If the process generating the data (in our case the rates of return) is linear then the skewness of the bispectrum will be constant. If the test rejects constant skewness then a non-linear process is implied.

Linearity and Gaussianity can be tested using a sample estimator of the skewness function $\Gamma(w_1, w_2)$ with

$$\Gamma^2(w_1, w_2) = \frac{|B_{xxx}(w_1, w_2)|^2}{S_{xx}(w_1)S_{xx}(w_2)S_{xx}(w_1 + w_2)}. \quad (9)$$

where $S_{xx}(w)$ is the spectrum of $x(t)$ at frequency w . The bispectrum at frequency pairs (w_1, w_2) is defined as

⁷ Guarda and Salmon (1996) present several other tests that focus on non-linearity in the mean.

⁸ Details of the bispectrum estimation method can be found in Hinich (1982), Hinich and Patterson (1985), Ashley, Patterson and Hinich (1986) and also in Barnett, Gallant, Hinich, Jungeilges, Kaplan and Jensen (1996).

$$B_{.xxx}(w_1, w_2) = \sum_{r=-\infty}^{\infty} \sum_{s=-\infty}^{\infty} C_{.xxx}(r, s) e^{[-i2p(w_1r+w_2s)]} \quad (10)$$

in the principal domain given by $\Omega = \{(w_1, w_2): 0 < 0.5, w_2 < w_1, 2w_1 + w_2 < 1\}$.

The null of the Hinich “linearity” test is actually given by

H₀: flat skewness function, absence of third order non-linear dependence;

H₁: non-linear dependence, absence of efficiency,

and H₀ is rejected if the standard normal test statistic Z is large, over 2 or 3. When the null is Gaussianity the related test statistic is denoted by H and is also a standard normal random variate under the null.

3.4 - Lyapunov Exponent

Lyapunov exponents measure the exponential rate at which two nearby orbits are moving apart. They provide an estimate of sensitive dependence on initial conditions, a defining feature of chaos. It basically means that if we allow for small changes in the state of a system it will grow at an exponential rate.

Consider two points, x_0 and $x_0+\epsilon$, apart from each other by only the infinitesimal difference ϵ and apply a map function to each of the two points n times.⁹ The difference between the results is given by

$$d_n = e^{nI(x_0)} \epsilon \quad (11)$$

and after solving for the convergence (or divergence) rate λ we have the Lyapunov exponent

⁹ On this topic our notation is similar to the one used by Barnett, Medio and Serletis (1997).

$$I = \lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \frac{d_n}{e} \right|. \quad (12)$$

If a system has at least one positive Lyapunov exponent then the system is chaotic and trajectories, which start at two similar states, will diverge exponentially. The larger the dominant positive exponent the more chaotic the system and the shorter the time span of system predictability. A positive Lyapunov exponent is therefore viewed as “an operational definition of chaotic behavior”.¹⁰

In an earlier paper Wolf, Swift and Vastano (1985) estimated the Lyapunov exponent by averaging the observed orbits divergence rates. Some authors however argue that this method is less reasonable when we are dealing with noisy systems. Therefore McCaffrey, Ellner, Gallant and Nychka (1992) and Nychka, Ellner, McCaffrey and Gallant (1992) use an alternative approach based on Jacobian methods in order to avoid upward bias when estimating Lyapunov exponents.¹¹

3.5 - BDS Test¹²

The BDS test can only be used to produce indirect evidence about non-linearity because the sampling distribution of the test statistic is not known. This statistic is useful to test for patterns that occur more (or less) frequently than would be expected in independent data. The BDS test can be used to test for the remaining non-linear dependence in the residuals of an ARIMA process. The null hypothesis may be formulated as

H_0 : pure whiteness, independent data, data generated by an iid stochastic process, efficiency;

H_1 : non-linear dependence, absence of efficiency,

¹⁰ As stated by Abhyankar, Copeland and Wong (1995).

¹¹ S. Ellner, D. W. Nychka and A. R. Gallant developed the LENNS (Lyapunov Exponent of Noisy Nonlinear Systems) software, a program that also estimates the dominant Lyapunov exponent and tests for chaos. Unfortunately we were no able to get an executable version of this software.

¹² The BDS test is named after William A. Brock, W. Davis Dechert and J. A. Scheinkman.

and when the BDS test statistic is large (which means larger than 2 or perhaps 3), H_0 is rejected.

Before looking at our results we briefly present the BDS test of independence and identical distribution or the so-called BDS test for non-linearity.

The Brock-Dechert-Scheinkman (BDS) statistic is a non-parametric test to assess the null hypothesis that a univariate series $\{x_t, t = 1, \dots, n\}$ is independently and identically distributed against an unspecified alternative. This test is performed by examining the underlying probability structure of $\{x_t\}$ in order to search for any kind of dependence. Often referred as a non-linear test, the BDS statistic can be used to detect any deviation from independence even if due to the presence of non-linear dependence in the data.

Let us define the correlation integral, given by equation (13), as a measure of the fraction of pairs of points (x_t^m, x_s^m) in the series that are within a distance of ϵ (metric bound) from each other:

$$C_m^n(\mathbf{e}) = \frac{2}{N(N-1)} \sum_{t=1}^{N-1} \sum_{s=t+1}^N I(x_t^m, x_s^m) \quad (13)$$

with,

$$I(a, b) = \begin{cases} 1 & \text{if } \|a - b\| \leq \mathbf{e} \\ 0 & \text{otherwise} \end{cases}, \quad (14)$$

where $\|\cdot\|$ is the supremum norm ($L_\infty = \max\{|a|\}$) and

$$N = n - m + 1;$$

n - number of observations and

m - embedding dimension.

In the literature, x_t^m is commonly referred to as the m -history and can generally be written as $x_t^m = (x_t, x_{t+1}, \dots, x_{t+m-1})$. For instance, with $m = 2$, the first 3 histories will be: (x_t, x_{t+1}) , (x_{t+1}, x_{t+2}) and (x_{t+2}, x_{t+3}) . Observe that since the first and third history do not have any repeated element, they are said to be non-overlapping histories.

If the data is generated by a strictly stationary stochastic process, which is absolutely regular, then equation (13) can also be written as

$$C_m(\varepsilon) = \lim_{n \rightarrow +\infty} C_m^n(\varepsilon) = \int_{r^m} \int_{r^m} I(a, b) dF_m(a) dF_m(b) \quad (15)$$

and Brock, Dechert, Sheinkman and LeBaron (1991) show that if $\{x_t\}$ is iid then we have $C_m(\varepsilon) = C_1(\varepsilon)^m$.

The BDS statistic is then given by

$$W_m^n(\mathbf{e}) = \frac{\sqrt{N} (C_m^n(\mathbf{e}) - (C_1^n(\mathbf{e}))^m)}{s_m(\mathbf{e})}, \quad (16)$$

where $W_m^n(\mathbf{e})$ converges in distribution to a standard normal, $N(0,1)$, as n approaches infinity.¹³ Brock, Hsieh and LeBaron (1991) show that the normal distribution was found to be, asymptotically, a good approximation for the distribution of the BDS statistic when there are more than 500 observations.

One should be aware that the BDS statistic depends, to a great extent, on the choice of values for ε and m . With large (small) ε the spatial correlation between the data points will tend to be high (low). The greater is the embedding dimension the smaller will be the number of non-overlapping histories and, as a consequence, the points defined by the embedded vectors will become “closer” and the value of the BDS statistic will tend to be higher. For a large absolute value of the test statistic we reject the null hypothesis

¹³ The statistical properties of the BDS test were studied by, for instance, Brock and Baek (1991), Brock, Dechert and Sheinkman (1991), Brock, Hsieh and LeBaron (1991), Hsieh (1991) and

of iid (randomness) since this provides evidence that the data are non-linear.

4 – Empirical Results

Non-linear dependence may not be absent from the stock exchange returns. In order to validate that hypothesis we now proceed our work by testing for neglected non-linearity.

Engle and Tsay test results

After selecting an AR(1), for the log returns of the three stock exchange indexes (using the SBIC), we applied the two tests to the residuals of the AR(1) model. The results are reported in Table 3 and allow us to reject the null hypothesis in both cases. This means that we can neither conclude for the absence of an ARCH (1) model using the Engle test (there may be non-linearity in the variance) or for the lack of non-linearity on the mean using the Tsay test.¹⁴

Table 3
Engle and Tsay tests

	ARCH			Tsay		
	Test statistic	<i>p</i> -value	df	Test statistic	<i>p</i> -value	df
BVLG	146.99	0.0000	1	8.82	0.0030	(1, 1789)
PSI20	207.80	0.0000	1	14.71	0.0001	(1, 1235)
BVL30	228.94	0.0000	1	12.49	0.0004	(1, 1234)

Note: df – degrees of freedom.

Hinich Test Results

Without prewhitening and using the first differences of the natural logarithms of our stock exchange indexes we performed the Hinich bispectrum test. We did not use previously any linear filtering of the data since Ashley, Patterson and Hinich (1986)

Scheinkman and LeBaron (1989).

¹⁴ Teixeira (1997) produced similar results for the BVLG index using an AR(1) and an AR(2)

proved that the Hinich test is invariant to linear filtering. If the original series were non-linear then the non-linearity would pass through the linear filter to the residuals.

The evidence from Table 4, rejection of linearity with the Hinich test, supplies support for the conclusion on non-linearity in Portuguese stock indexes returns and therefore for the absence of weak form efficiency.¹⁵

Table 4
Hinich Bispectral Test

Series	Gaussianity test statistic (H)	Linearity test statistic (Z)	Conclusion
BVLG	9,14	3,97	Reject Gaussianity and linearity
PSI20	8,96	2,71	Reject Gaussianity and linearity
BVL30	8,77	2,35	Reject Gaussianity and linearity

Note: the tests are one sided and the null hypothesis is rejected in both cases if the test statistic exceeds 2 or 3. As suggested by Hinich (1982) the bispectral smoothing parameter was set to approximately the square root of the number of observations.

However, even if we did not reject the null of linearity, one should be aware, as reminded by, Barnett, Gallant, Hinich, Jungeiles and Jensen (1997), that “acceptance of the null of linearity (...) provides only weak support for the linearity, since the [Hinich] test, as currently constructed, actually tests the broader null of absence of third order nonlinear dependence”.¹⁶

processes for the Engle test, and with an AR(2) and an AR(24) for the Tsay test.

¹⁵ Our calculations were obtained with software developed by D. M. Patterson (Bispectrum Estimation Program, Version 5.11, September 1992).

¹⁶ The interesting paper of Barnett, Gallant *et al.* (1997) concerning non-linearity and tests for chaos may be found in <http://wuecon.wustl.edu/~barnett/Papers.html>.

Estimates of Lyapunov Exponents

Table 5 presents estimates of the maximum Lyapunov exponents, of the daily log returns series, using the estimation method of Wolf *et al.* (1985).¹⁷ We computed the Lyapunov Exponents with embedding dimensions up to five which, considering the available data, is already more than the value suggested by Wolf (1991).

Table 5
Lyapunov Exponents for log returns

Ndim	max λ		
	BVLG	PSI20	BVL30
1	0.0144	0.0216	0.0120
2	0.0155	0.0130	0.0282
3	0.0204	0.0135	0.0205
4	0.0291	0.0168	0.0144
5	0.0136	0.0099	0.0167

Notes: max λ - maximum estimated value of Lyapunov Exponent;
ndim - embedding dimension.

The results, all λ are positive, seem to point to the above stated operational definition of chaos: trajectories that start at two almost identical state vectors diverge exponentially as time passes, allowing us to accept chaos for our three financial time series.

If chaos exist one implication is that profitable non-linearity based trading rules exist at least in the short-run. The problem would then be to find out the generating mechanism in order to take advantage of the financial markets inefficiencies.

¹⁷ We computed the Lyapunov exponents with a software developed and distributed by A. Wolf (programs BASGEN and FET, 1991). These programs may be obtained from A. Wolf's Web site, <http://www.users.interport.net/~wolf/>, in a self-decompressing executable file. Wolf (1991) provides valuable indications on how to use those programs.

The BDS Test

The BDS test will be applied to the residuals of a fitted linear model, in our case an ARIMA model, which we presumed has extracted as much linear structure as possible from the data.

Using the Box-Jenkins methodology we tried to adjust an ARIMA (autoregressive integrating moving average) process for each of our chosen stock exchange indexes log returns. Briefly, we fitted an ARIMA (p, d, q) model to our time series, where p denotes the number of autoregressive terms, d the number of times the time series has to be differenced before it becomes stationary, and q is the number of moving average terms.¹⁸

Visual inspection may be used (see Figures 4, 5 and 6 on the Annex) to assess, as a preliminary step, whether the series are stationary. In fact, the Box-Jenkins methodology applies only to stationary data series, which means that the time series has a mean and a variance essentially constant through time. Together with the analysis of the correlograms of the autocorrelation and partial autocorrelation functions we tentatively identified an ARIMA (0, 1, 3) model for the returns of the three original time series. Using the lag operator L, that is,

$$LX_t = X_{t-1}, \quad (17)$$

our ARIMA models assumed the form

$$(1 - L)Y_t = (1 - \mathbf{q}_1 L - \mathbf{q}_2 L^2 - \mathbf{q}_3 L^3)u_t \quad (18)$$

where Y is the first difference of the natural logarithm of the original time series, \mathbf{q}_1 , \mathbf{q}_2 and \mathbf{q}_3 are the moving average coefficients and the random shock u_t is assumed to be an iid random variable with a mean of zero and a constant variance.

¹⁸ We did not restrict our linear specifications to an autoregressive form, solution used for instance by Brooks (1996).

Table 6 presents the main results of the fitted models, which satisfy the stationarity and invertibility conditions.¹⁹ The residuals of the ARIMA models will now be used to perform the BDS test.

Table 6
ARIMA models for Stock Exchange Indexes daily returns

Index	Model		q_1	q_2	q_3
BVLG	ARIMA(0,1,3)	Coefficient	0.802075	0.137213	0.050019
		T-Statistic	33.97	4.55	2.12
PSI20	ARIMA(0,1,3)	Coefficient	0.791847	0.133191	0.067637
		T-Statistic	27.91	3.69	2.38
BVL30	ARIMA(0,1,3)	Coefficient	0.799907	0.106930	0.08518
		T-Statistic	23.18	2.94	2.99

Notes: q_i is the parameter for an MA at lag i .

To compute the BDS statistic Brock, Hsieh and LeBaron (1991) recommend using ε between one-half to two times the standard deviation of the raw data ($0,5s \leq \varepsilon \leq 2s$) while suggesting that $m = 2$. For samples with less than 500 observations m should be set less or equal to 5.

Table 7 presents the results of the BDS test for the ARIMA residuals of log returns.²⁰ It can be seen that the null hypothesis of R_t being iid is rejected, at the 5 per cent level, for all indexes.

¹⁹ These models are always stationary since they have no autoregressive coefficients. The invertibility conditions require that $|q_i| < 1$ for all i .

²⁰ Our BDS statistics were obtained with software developed by William Dechert (BDS STATS, version 8.20). The interested reader may try to get the BDS software either from <http://www.ssc.wisc.edu/~lebaron/software/index.html> or from the alternative web site gopher://gopher.ssc.wisc.edu/00/econgopher/software/bds/dos.

Table 7
BDS Statistics
Residuals from the ARIMA to log returns

INDEX	m	ϵ			
		$0,5\sigma$	σ	$1,5\sigma$	2σ
BVLG	2	14,719	15.292	13.438	12.064
	3	18.322	17.763	14.765	12.933
	4	20.876	19.319	15.644	13.424
	5	24.284	21.014	16.187	13.527
PSI20	2	12.178	10.729	10.045	10.450
	3	15.219	12.513	10.693	10.550
	4	18.407	14.128	11.404	10.834
	5	22.656	15.792	12.065	10.929
BVL30	2	11.306	10.615	9.765	9.687
	3	14.518	12.297	10.268	9.793
	4	17.503	13.905	10.076	10.256
	5	21.379	15.583	11.735	10.376

Notes: m - embedding dimension; ϵ - distance between points, measured in terms of number of standard deviations of the raw data; σ - standard deviation. All statistics significant at the 5 per cent level.

In other words, non-linear dependence is not absent from the series returns and we must therefore conclude that the weak form efficiency hypothesis is not validated for all of the Portuguese stock exchange indexes in our data set.²¹

5 - Conclusions

The hypothesis of linearity is rejected by almost all the tests we computed. In fact, the results allow us to reject the null hypothesis of daily returns being iid, non-linear dependence is present on those returns therefore contradicting the random walk model supposition. Since the test results indicate that non-linear structure is present in the data, it is possible that exploitable excess profit opportunities may exist in the Portuguese stock market.²²

These conclusions are consistent with the results of Omran (1997) from the UK

²¹ These conclusions confirm the results from previous work by Afonso and Teixeira (1997) concerning namely stock exchange indexes for several activity sectors.

²² Obviously, and as Michael Brennan pointed out to us, non-linearity does not imply predictability.

and of de Lima (1995) from the US since those authors argue that non-linear dependence is present in stock returns after the 1987 crash.

Our results seem therefore to challenge the belief that daily rates of return can be viewed as independent random variables. In fact, those returns may be potentially forecastable. Nevertheless, the option of concluding from so-called non-linear tests that financial markets are inefficient implies always a strong assumption. Abhyankar, Copeland and Wong (1995) argue that non-linear structure may be consistent with lack of efficiency given risk neutrality and/or negligible transaction costs.

References

Abhyankar, A.; Copeland, L. S. and Wong, W. (1995). “Nonlinear Dynamics in Real-Time Equity Market Indices: Evidence from the United Kingdom,” *The Economic Journal*, 105, (431), July, 864-880.

Afonso, A. (1997). “Normality and Efficiency in Portuguese Stock Exchange Indexes,” March, *Estudos de Economia*, XVI-XVII, Fall, 101-106.

Afonso, A. and Teixeira, J. (1997). “Efficiency in Portuguese Stock Exchange Indexes: Runs Tests and BDS Statistics,” Working Paper N° 2/97, Department of Economics, Instituto Superior de Economia e Gestão, Universidade Técnica de Lisboa.

Al-Loughani, N. and Chappell, D. (1996). “On the Validity of the Weak Form Efficient Markets Hypothesis Applied to the London Stock Exchange,” *Applied Financial Economics*, 7 (2), April, 173-176.

Ashley, R. A.; Patterson, D. M. and Hinich, M. J. (1986). “A Diagnostic Test for Nonlinear Serial Dependence in Time Series Fitting Errors,” *Journal of Time Series Analysis*, 7 (3), 165-178.

Barnett, W. A.; Gallant, A. R.; Hinich, M. J.; Jungeilges, J.; Kaplan, D. and Jensen, M. J. (1996). “An Experimental Design to Compare Tests of Nonlinearity and Chaos,” in *Nonlinear Dynamics and Economics*. Eds. William Barnett, Alan Kirman and Mark Salmon. Cambridge. Cambridge U. Press.

Barnett, W. A.; Medio, A. and Serletis, A. (1997). “Nonlinear and Complex Dynamics in Economics,” *mimeo*.

Barnett, W. A.; Gallant, A. R.; Hinich, M. J.; Jungeilges, J.; Kaplan, D. and Jensen, M. J. (1997). “A Single-Blind Controlled Competition Among Tests for Nonlinearity and Chaos,” *Journal of Econometrics* (forthcoming).

Brock, W. A. and Baek, E. G. (1991). “Some Theory of Statistical Inference for Nonlinear Science,” *Review of Economic Studies*, 58, 697-716.

Brock, W. A.; Hsieh, D. A. and LeBaron, B. (1991). *Nonlinear Dynamics, Chaos, and Instability: Statistical Theory and Economic Evidence*, The MIT Press.

Brock, W. A.; Dechert, W. D.; Scheinkman, J. A. and LeBaron, B. (1991). *A Test for Independence Based on the Correlation Dimension*, Social Systems Research Institute, University of Wisconsin, March.

Brooks, C. (1996). “Testing For Non-linearity in Daily Sterling Exchange Rates,” *Applied Financial Economics*, 6 (4), August, 307-317.

de Lima, P. F. (1995). “Nonlinearities and Nonstationarities in Stock Returns,” Working Papers in Economics, The John Hopkins University, Department of Economics.

Engle, R. F. (1982). “Autoregressive Conditional Heteroskedasticity with Estimates of the Variance of United Kingdom Inflation,” *Econometrica*, 50 (4), 987-1007.

Fama, E. (1991). “Efficient Capital Markets II,” *Journal of Finance*, XLVI (5), December, 1575-1617.

Guarda, P. and Salmon, M. (1996). “Detection of nonlinearity in foreign-exchange data,” in *Nonlinear Dynamics and Economics*. Eds. William Barnett, Alan Kirman and Mark Salmon. Cambridge. Cambridge U. Press.

Hinich, M. J. (1982). “Testing for Gaussianity and Linearity in a Stationary Time Series,” *Journal of Time Series Analysis*, 3 (3), 169-176.

Hinich, M. J. and Patterson, D. M. (1985). “Evidence of Nonlinearity in Daily Stock Returns,” *Journal of Business and Economic Statistics*, 3 (1), 69-77.

Hinich, M. J. and Patterson, D. M. (1985). “Evidence of nonlinearity in the trade-by-trade stock market return generating process,” in Barnett, W. A.; Geweke, J. and Shell, K. (eds.), *Economic Complexity: Chaos, Sunspots, Bubbles, and Nonlinearity*, Cambridge University Press.

Hsieh, D. A. (1991). “Chaos and Nonlinear Dynamics: Application to Financial Markets,” *Journal of Finance*, XLVI (5), December, 1839-1877.

Lee, T. H.; White, H. and Granger, C. (1993). “Testing for Neglected Nonlinearity in Time Series Models: A Comparison of Neural Networks Methods and Alternative Tests,” *Journal of Econometrics*, N° 56, 269-290.

McCaffrey, D.; Ellner, S.; Gallant, R. and Nychka, D. (1992). “Estimating the Lyapunov Exponent of a Chaotic system With Nonparametric Regression,” *Journal of the American Statistical Association*, September, 87 (419), 682-695.

Nychka, D.; Ellner, S.; McCaffrey, D. and Gallant, R. (1992). “Finding Chaos in Noisy Systems,” *Journal of the Royal Statistical Society, Series B*, 54 (2), 399-426.

Omran, M. F. (1997). “Nonlinear Dependence and Conditional Heteroscedasticity in Stock Returns: UK Evidence,” *Applied Economics Letters*, 4 (10), 647-650.

Scheinkman, J. A. and LeBaron, B. (1989). “Nonlinear Dynamics and Stock Returns,” *Journal of Business*, 62 (3), 311-337.

Taylor, S. (1986). *Modelling Financial Time Series*, John Wiley & Sons.

Teixeira, J. (1997). *Previsão Baseada em Modelos Neurais e de Regressão Dinâmica*, October, Masters dissertation, Faculdade de Ciências de Lisboa.

Tsay, R. S. (1986). “Nonlinearity Tests for Time Series,” *Biometrika*, 73 (2), 461-466.

Wolf, A. (1991). *Lyapu-News: documentation for FET, a program that quantifies chaos in time series, mimeo.*

Wolf, A.; Swift, J. B.; Swinney, H. L. and Vastano, J. A. (1985). “Determining Lyapunov Exponents from a Time Series,” *Physica*, 16D, 285-317.

Annex

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

