

Review

Ophthalmologica

Ophthalmologica
DOI: 10.1159/000331277

Received: July 28, 2011
Accepted: July 29, 2011
Published online: September 15, 2011

Pseudophakic Cystoid Macular Edema

Conceição Lobo

Association for Innovation and Biomedical Research on Light and Image, Department of Ophthalmology, University Hospital of Coimbra, Institute of Biomedical Research on Light and Image, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

Key Words

Cataract surgery · Cystoid macular edema · Inflammatory mediators · Management

Abstract

Cataract surgery is an efficient procedure, and is generally associated with good visual results. Nevertheless, cystoid macular edema (CME) may develop, and this can result in suboptimal postoperative vision. Many factors are considered to contribute to its development, and although the treatment options depend upon the underlying cause of CME, the usual therapeutic approach for prophylaxis and treatment of CME is directed towards blocking the inflammatory mediators. This article provides a review of possible risk factors, pathogeneses, incidence rates, and methods of diagnosis, as well as the current guidelines for managing CME.

Copyright © 2011 S. Karger AG, Basel

Introduction

Cataract extraction is one of the most commonly performed surgeries. Over recent years, it has benefited from consistent innovations in instrumentation, lens design, and surgical technique that have led to improved outcomes following cataract surgery [1]. Phacoemulsification using small incisions and implantation of a foldable

intraocular lens (IOL) is currently the preferred technique among cataract surgeons [2]. The procedure is efficient, and uneventful surgery is generally associated with good visual results [3–5]. Nevertheless, cystoid macular edema (CME) may develop, and this can result in suboptimal postoperative vision [6–8]. It can occur after uncomplicated surgery in patients with otherwise healthy eyes, after complicated surgery, or after surgery in patients with ocular diseases such as uveitis or diabetic retinopathy [9].

CME following cataract surgery was initially reported by Irvine in 1953 and demonstrated angiographically by Gass and Norton in 1966, so it is known as the Irvine-Gass syndrome [10–12].

Angiographic CME is diagnosed in patients who are otherwise asymptomatic with respect to visual acuity, but have detectable leakage from the perifoveal capillaries on fluorescein angiography (FA). Clinical CME is diagnosed in those patients who have detectable visual impairment as well as angiographic and/or biomicroscopic findings.

The clinical diagnosis is confirmed using optical coherence tomography (OCT) and FA.

The incidence of pseudophakic CME depends on the methodology used in its detection.

The prophylactic use of non-steroidal anti-inflammatory drugs (NSAIDs) preoperatively, and the combination of steroids and NSAIDs in the postoperative period, is recommended to reduce the incidence of pseudophakic CME.

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2011 S. Karger AG, Basel
0030-3755/11/0000-0000\$38.00/0

Accessible online at:
www.karger.com/oph

Conceição Lobo
AIBILI, Faculty of Medicine, University of Coimbra-IBILI, Depart. Ophthalmology
University Hospital of Coimbra, Azinhaga Sta Comba, Celas
PT- 3000-548 Coimbra (Portugal)
Tel. +351 239 480 100, E-Mail clobofonseca@gmail.com

Etiology and Risk Factors

The specific etiology of aphakic and pseudophakic macular edema is not fully understood. Many factors have been considered to contribute to its development, such as the type of cataract surgery, light toxicity, vitreomacular traction, inflammatory mediators, use of adrenergic drugs, age, vitreous loss, integrity of the posterior capsule, hypertension, and diabetes [7, 13].

The type of cataract surgery used is associated with different outcomes and complications, such as CME. The change in procedure from large-incision intracapsular cataract extraction to small-incision extracapsular phacoemulsification was associated with a clear decrease in the incidence of this complication [7, 13–18]. This has been explained by less blood-aqueous barrier damage after phacoemulsification with an intact continuous curvilinear capsulorhexis than after extracapsular cataract extraction [19, 20].

Although microscope light toxicity has been indicated to be a possible contributor to CME, a prospective randomized study did not support this finding, and showed no statistically significant difference in the rate of angiographic CME from using a pupillary light occluder [21]. However, IOLs that filter UV light may reduce the rate of angiographic CME [22].

The age of the patients is another factor that needs to be considered. Some authors have indicated a positive correlation with age, demonstrating an increased incidence of CME in older patients [7, 23].

Changes occurring in the vitreous body during surgery are another pathogenic mechanism that has been proposed as a cause of CME development. Although CME can occur after uneventful cataract surgery, certain surgical complications raise the risk of CME. Rupture of the posterior capsule as well as secondary capsulotomy (including YAG laser capsulotomy) are associated with a higher rate of CME [24]. Vitreous loss increases the prevalence of CME by 10–20% [7, 25, 26]. Different studies have confirmed that the rate of clinical CME associated with vitreous loss is higher compared to cases without this complication after either extracapsular cataract extraction or phacoemulsification [25, 27–29]. The lower rate of CME occurring when there is vitreous loss during phacoemulsification has also been explained by the small wound construction and greater stability in this type of surgery compared with large-incision cataract surgery [17, 30–32]. Vitreous to the wound prolongs CME, and can be associated with a poorer prognosis [26]. Iris incarceration, an additional risk factor for CME, may have a more impor-

tant association with poor vision in patients with chronic postsurgical CME than with other intraoperative complications [24, 33]. Retained lens fragments, even when they are removed by pars plana vitrectomy, are another complication associated with an increased rate of CME and more severe visual loss, in eyes with a sulcus fixated posterior chamber IOL, anterior chamber IOL, or aphakic eyes [34–37]. Retained lens fragments cause inflammation that is often severe, and these eyes have longer cataract surgery times than eyes with uneventful cataract extraction. Prolonged exposure to light from the operating microscope during the initial surgery combined with possible pars plana vitrectomy, as a second surgery, may also contribute to a higher incidence of CME [38, 39].

There are multiple reports about the association of CME after cataract surgery with the topical use of latanoprost, an ocular hypotensive lipid that is a prostaglandin analogue agent [40–43]. Other similar agents (unoprostone, travoprost and bimatoprost) were also associated with an increased incidence of CME [44]. The explanation of the increased incidence of CME associated with the prostaglandin analogues is based on the associated increase in blood-aqueous barrier disruption and inflammatory activity. Considering this possibility, it is recommended to avoid or discontinue this medication and substitute it for another type of hypotensive drug before surgery.

In diabetic patients, CME shows an increase risk, especially in patients with pre-existing diabetic retinopathy [45]. The incidence of CME in diabetic patients, even in the absence of diabetic retinopathy, is higher than in patients without this pathology [46]. Although the frequency of this problem has decreased with the widespread use of smaller incisions and phacoemulsification, it remains an important issue to consider [47]. The poor visual acuity after cataract surgery in these patients due to macular edema is common, and the two clinical forms (diabetic macular edema and edema caused by Irvine-Gass syndrome) probably co-exist [45, 48, 49]. These are difficult to differentiate, although some authors have suggested that if there is post-operative hyperfluorescence of the optic disc in the angiograms, it probably corresponds to a CME which would be likely to resolve spontaneously [50]. It is very important to take into consideration that if the patient has already some degree of macular edema when the cataract surgery is considered, it should be treated before the surgery and – in cases when this is not possible – intravitreal anti-inflammatory medication at the surgical time may be used.

Certain patients appear to have a higher risk of CME after cataract surgery. Patients with uveitis frequently develop CME, and it is the most common reason for poor

outcomes of cataract surgery in these patients [51]. After cataract surgery in patients with juvenile rheumatoid arthritis or pars planitis, the occurrence of CME can reach an incidence of about 50% [52]. Foster et al. [53] reported similar incidences in a retrospective study with uveitis patients. Other studies have reported lower incidences of CME. It is important to note that there is a wide variation in severity of uveitis, and surgeons should consider these patients at risk of postoperative CME. Taking this into account, it is imperative to control preoperative inflammation and introduce postoperative medication according to the expected risk.

In conclusion, the most frequent risk factors associated with pseudophakic CME are iris trauma, rupture of posterior capsule, vitreous loss or incarceration, dislocated IOL, use of iris fixed lenses, active uveitis, and diabetes.

Pathogenesis

The pathogenesis of CME following cataract surgery remains uncertain, but clinical observations and experimental studies indicate that the pathophysiology of this postoperative problem may be multifactorial [26, 39]. Among the factors of potential clinical importance are vitreous traction [10, 54], functional disturbance of the blood-retinal barrier (BRB) [13, 55–57], and inflammation mediated by prostaglandins [7, 39, 58–62]. Most investigators consider inflammation as the major etiologic factor in the development of CME following cataract surgery. It is noteworthy that inflammation is directly associated with a breakdown of the BRB. It is probable that after surgery the inflammation mediators (prostaglandins, cytokines, and other vascular permeability factors) are released from the anterior segment of the eye and diffuse to the vitreous cavity and retina, stimulating the breakdown of the BRB and subsequent leakage of fluids across the retinal vessel wall and through the retinal pigment epithelium (RPE) into perifoveal retinal tissues resulting in macular edema [13, 63, 64]. Surgical manipulation, which happens during the cataract surgery, always causes trauma to the iris. It is known that the iris is a metabolically active tissue that releases inflammatory mediators when traumatized. After surgery, the physiological healing process is sufficient to slowly, but progressively, suppress the inflammation [65]. In about 90% of patients with macular edema following cataract surgery, a spontaneous resolution of the edema and a recovery of visual acuity can be observed. In specific situations, an excessive leakage occurs, which can lead to a severe and irreversible impair-

ment of visual acuity. The different degrees of leakage explain the higher incidence of angiographically detectable CME compared to clinically identified CME [66].

The inflammatory mediators probably play the essential initiating role in the development of inflammatory CME, but the exact factors and events responsible for further CME development and its chronicity have not yet been clearly identified.

Incidence

Though the rate of CME has been declining with decreasing surgical incision and the use of microsurgical instrumentation, the high volume of cataract surgeries performed each year makes CME a common cause of poor vision in pseudophakic patients [32].

Determining the overall incidence of CME has been difficult because of variations in the patient populations evaluated (with varying risk factors) and the use of different methods for evaluating macular thickening. Another factor that contributes to the uncertain incidence is the use of different prophylactic medications before and after cataract surgery. There are multiple reports comparing the incidence of clinical and angiographic CME under specific anti-inflammatory drugs [67–69]. The multiple reports and reviews that mention or discuss the incidence of CME show different numbers and the explanations are, in certain cases, unclear [26, 70]. However, it is important to recognize the many variables involved and the difficulty in identifying their potential impact on the interpretation of results from therapeutic trials that evaluated potential treatments for CME.

The true current incidence of clinical CME using modern techniques appears to be very low in uncomplicated cases. It is likely to be in the range of 0.2–2%, and the rate of angiographic CME is at least ten-fold higher in magnitude [26, 32, 71]. These numbers increase if another large-incision technique is required, instead of small-incision phacoemulsification with implantation of a foldable IOL; they also increase if we have a complicated surgery with iris trauma, or capsule rupture with vitreous loss, or in patients at high risk (such as those with uveitis or diabetes).

Diagnosis

The diagnosis of CME can generally be made on clinical examination with evidence of perifoveal cystic spaces, and has traditionally been confirmed with use of FA

to document the classic petaloid pattern of leakage induced by inflammatory mediators that lead to breakdown of the BRB. Today, the most definite evidence is obtained from OCT examination.

Signs and symptoms of clinically significant CME typically develop 4–12 weeks after surgery and reach a peak at 4–6 weeks postoperatively. The patient may complain of impaired vision after an initial period of improved vision [24]. Generally, this responds to treatment with topical anti-inflammatory medications and usually resolves within 6 months. However, some cases respond poorly to conservative treatment, persist for more than 6 months, and may develop permanent visual loss. Although the incidence of chronic CME is much less frequent, being reported at 1–2% of uncomplicated cases and about 8% after complicated cataract surgery, the associated vision loss makes it a serious complication.

In ophthalmoscopy and biomicroscopy, the loss of foveal depression is the most frequent sign of CME. Intraretinal cystoid spaces can be detected and the perifoveal area can appear with a yellowish coloration [7, 24, 32]. A swelling of the optic nerve head can be detectable [26]. In cases of chronic CME, the cystoid spaces fuse to foveal cysts.

Angiographic CME is diagnosed in patients who are otherwise asymptomatic with respect to visual acuity, but have detectable leakage from perifoveal capillaries on FA demonstrated by hyperfluorescence in the central macula and optic disc. Clinical CME is diagnosed in those patients who have detectable visual impairment as well as angiographic and/or biomicroscopic findings [32, 72].

In the early phase of the FA, capillary dilatation and leakage from small perifoveal capillaries are visible. In later phases, pooling in the outer plexiform layer results in the classic perifoveal ‘petaloid’ staining pattern. Another common sign is late leakage and staining of the optic nerve due to capillary leakage [73]. In severe CME, the cystoid spaces may have a ‘honeycomb’ appearance in FA, corresponding to large cystoid spaces, extending outside the immediate perifoveal region [24]. Vitreous fluorometry, a more sensitive technique for measuring alterations in the BRB using fluorescein, shows an even higher incidence of alterations in the period immediately after cataract surgery [55]. The retinal leakage analyzer is a methodology that can be used to objectively measure the disruption of the blood-retinal barrier, leading to a quantitative evaluation of the macular edema [74].

With OCT, it is possible to observe the cystic spaces in the outer nuclear layer of the central macula and, in correlation with histological findings, measure the edema

which should correspond to the value of the observed retinal thickening [75–77]. The foveal thickness can increase significantly, and it correlates moderately with the decrease in visual acuity, but clearly better than the correlation between FA and visual acuity.

Although the FA has been considered the diagnostic gold standard for pseudophakic CME, OCT is now the method of choice, being a non-invasive technique to evaluate and follow CME after cataract surgery [78]. An additional advantage of the use of OCT is that the thickening of macular area effectively measures the edema, and therefore correlates better with vision than the angiogram grading [26].

Classification

Angiographic CME is defined as the presence of fluorescein leakage on FA. It is mostly asymptomatic. A decrease in visual acuity does not correlate with an extension of leakage. Clinically significant CME presents with decreased vision, and is generally diagnosed by biomicroscopy and OCT.

Usually the CME appears within 4 months of surgery, and is designed as acute. If the onset occurs after more than 4 months postoperatively, it is designed late-onset CME.

The CME is classified as chronic if it lasts more than 6 months.

Management

Angiographic CME after cataract surgery is not necessarily associated with poor visual outcome. Usually it resolves spontaneously and only about 1–3% of cases persist, and this corresponds to clinical CME with persistent symptoms [79].

It is now accepted that in cataract surgery it is important to perform a therapeutic intervention both for prophylaxis and for treatment of CME based on the existing knowledge of its pathogenesis. Many of the multiple studies performed to test the efficacy of different drugs in pseudophakic CME have been poorly designed. Therefore, the results are not consistent and controversy persists. However, it is now widely accepted that it is important to introduce anti-inflammatory drugs not only to treat but also specifically to prevent the development of CME after cataract surgery [80, 81].

Although the treatment options depend on the underlying cause of CME, the usual therapeutic approach for

the prophylaxis and treatment of CME is directed towards blocking the inflammatory mediators, mainly the prostaglandins in the anterior segment of the eye, using topical steroids and NSAIDs.

Topical NSAIDs have demonstrated effectiveness in preventing both angiographic and clinical CME [68]. There are multiple reports concerning the use of different NSAIDs in the prevention and treatment of CME. In general, studies have proven the effectiveness of use of this type of medication, and without clear statistical differences between the different NSAIDs.

Corticosteroids are effective, but they can cause an increase in intraocular pressure in a small percentage of patients. Usually the medical treatment of uncomplicated cases can first include NSAIDs with, in some cases, the addition of topical steroids. Clinical evidence suggests that the combined use of NSAIDs and steroids is synergistic [82, 83].

In patients with evidence of anterior segment inflammation or those with chronic CME following posterior capsule trauma, steroids have been shown to be effective, and a stepwise plan may be undertaken starting with topical administration, followed by local injection, reserving high-dose intra-vitreous administration for severe refractory cases. This can reduce macular edema and improve vision in eyes with CME that persists or recurs despite previous medical treatment [84–86]. However, CME may recur in some cases, even after more than one intravitreal injection of triamcinolone acetonide.

A carbonic anhydrase inhibitor (acetazolamide) may help reduce the edematous component. These have the ability to stimulate the RPE to pump excess fluid out of the macula. Furthermore, carbonic anhydrase inhibitors induce acidification of the subretinal space, and thereby increase fluid resorption from the retina through the RPE into the choroid [24, 87]. There are some reports documenting the positive effect of acetazolamide in post-surgical CME [88].

The use of antiangiogenic agents has also been proposed as an alternative therapeutic approach in some cases of refractory CME. VEGF, a potent inducer of alterations in the BRB, may be increased in postoperative CME. However, bevacizumab, a monoclonal antibody against all isoforms of VEGF, which has been used to treat some neovascular diseases, was used in patients with postoperative CME without positive effects [89].

Vitreolysis using the Nd:YAG laser has shown positive effects in cases of vitreous incarceration in the cataract incision wound [90].

Surgical treatment, namely vitrectomy, is indicated in some specific cases of chronic CME and appears to be effective. The rationale for performing vitrectomy in CME following surgery includes the removal of vitreous adhesions and inflammatory mediators and improved access of topical medication to the posterior pole.

The current guidelines for managing post-cataract surgery inflammation state that the prevention of inflammation should be the main goal, with good patient selection, correct eye/patient preparation, taking special care during surgery not to cause iris trauma, appropriate resolution of intraoperative complications, and timely treatment of postoperative inflammation [91].

It should be noted that diabetic patients with pre-surgical macular edema have a higher risk of worsening their macular edema after cataract surgery. In these patients, it is recommended to first treat the macular edema and only perform the surgery afterwards. In these cases, the use of intravitreal triamcinolone acetonide (4 mg) injected at the end of cataract surgery is indicated.

Conclusions/Recommendations for Managing Pseudophakic CME

- First, it is very important to perform a correct preoperative evaluation of the patient. Patients can be divided into normal and high-risk cataract patients.
- If s/he is a high-risk patient, the potential for improvement or correction should be considered, and, if indicated, medication should be introduced and the therapeutic schema adapted.
- If s/he is a normal patient, it is recommended (as prophylaxis) to administer topical NSAIDs during the first month in combination with topical steroids, using decreasing doses during the first 2 weeks.
- If there is a recognized CME, first the topical NSAIDs and steroids should be re-introduced for 1 month. After this, visual acuity should be evaluated and an OCT should be performed to estimate any possible improvement in the CME.
- If the CME has not improved, the introduction of acetazolamide (for 1 month or more) should be considered, or, as an alternative, a periocular corticosteroid or intravitreal triamcinolone.
- If there is vitreous incarceration or a persistent inflammatory reaction, surgery it (vitrectomy) should be considered.

References

- 1 DeCross FC, Afshari NA: Perioperative antibiotics and anti-inflammatory agents in cataract surgery. *Curr Opin Ophthalmol* 2008; 19:22–26.
- 2 Panchapakesan J, Rohtchina E, Mitchell P: Five-year change in visual acuity following cataract surgery in an older community: the Blue Mountains Eye Study. *Eye* 2004;18:278–282.
- 3 Linebarger EJ, Hardten DR, Shah GK, et al: Phacoemulsification and modern cataract surgery. *Surv Ophthalmol* 1999;44:123–147.
- 4 Gogate PM, Kulkarni SR, Krishnaiah S, et al: Safety and efficacy of phacoemulsification compared with manual small-incision cataract surgery by a randomized controlled clinical trial: six-week results. *Ophthalmology* 2005;112:869–874.
- 5 Riaz Y, Mehta JS, Wormald R, et al: Surgical interventions for age related cataract. *Cochrane Database Syst Rev* 2006;(4): CD001323.
- 6 O' Brien TP: Emerging guidelines for use of NSAD therapy to optimize cataract surgery patient care. *Curr Med Res Opin* 2005;21: 1131–1137, correction 2005;21:1431–1432.
- 7 Rossetti L, Autelitano A: Cystoid macular edema following cataract surgery. *Curr Opin Ophthalmol* 2000;11:65–72.
- 8 Mohammadpour M, Jafarinasab MR, Javadi MA: Outcomes of acute postoperative inflammation after cataract surgery. *Eur J Ophthalmol* 2007;17:20–28.
- 9 Nelson ML, Martidis A: Managing cystoid macular edema after cataract surgery. *Curr Opin Ophthalmol* 2003;14:39–43.
- 10 Irvine AR: A newly defined vitreous syndrome following cataract surgery, interpreted according to recent concepts of the structure of the vitreous. *Am J Ophthalmol* 1953; 36:599–619.
- 11 Gass JD, Norton EW: Cystoid Macular edema and papilledema following cataract extraction: a fluorescein fundoscopic and angiographic study. *Arch Ophthalmol* 1966;76: 646–661.
- 12 Irvine AR: Cystoid maculopathy. *Surv Ophthalmol* 1976;21:1–17.
- 13 Ursell PG, Spalton DJ, Whitcup SM, Nussenblatt RB: Cystoid macular edema after phacoemulsification: relationship to blood-aqueous barrier damage and visual acuity. *J Cataract Refract Surg* 1999;25:1492–1497.
- 14 Hitchings RA, Chisholm IH, Bird AC: Aphakic macular edema: incidence and pathogenesis. *Invest Ophthalmol Soc* 1975;14:68–72.
- 15 Jaffe NS, Clayman HM, Jaffe MS: Cystoid macular edema after intracapsular and extracapsular extraction with and without an intraocular lens. *Ophthalmology* 1982;89: 25–29.
- 16 Kraff MC, Sanders DR, Jampol LM, Lieberman HL: Effect of primary capsulotomy with extracapsular surgery on the incidence of pseudophakic cystoid macular edema. *Am J Ophthalmol* 1984;98:166–170.
- 17 Koch PS: Anterior vitrectomy; in Nordan LT, Maxwell WA, Davidson JA (eds): *Surgical Rehabilitation of Vision: An Integrated Approach to Anterior Segment Surgery*. London, Gower, 1992, chapter 17.
- 18 Powe NR, Schein OD, Gieser SC, et al: Synthesis of the literature on visual acuity and complications following cataract extraction with intraocular lens implantation. *Cataract Patient Outcome Research Team. Arch Ophthalmol* 1994;112:239–252.
- 19 Oshika T, Yoshimura K, Miyata N: Postsurgical inflammation after phacoemulsification and extracapsular extraction with soft or conventional intraocular lens implantation. *J Cataract Refract Surg* 1992;18:356–361.
- 20 Pande MV, Spalton DJ, Kerr-Muir MG, Marshall J: Postoperative inflammatory response to phacoemulsification and extracapsular cataract surgery: aqueous flare and cells. *J Cataract Refract Surg* 1996;22:770–774.
- 21 Kraff MC, Lieberman HL, Jampol LM, Sanders DR: Effect of a pupillary light occluder on cystoid macular edema. *J Cataract Refract Surg* 1989;15:658–660.
- 22 Nagpal M, Nagpal K, Nagpal PN: Postcataract cystoid macular edema. *Ophthalmol Clin North Am* 2001;14:651–659.
- 23 Stern AL, Taylor DM, Dalburg LA, et al: Pseudophakic cystoid maculopathy: a study of 50 cases. *Ophthalmology* 1981;88:942–946.
- 24 Loewenstein A, Zur D: Postsurgical cystoid macular edema; in Coscas G (ed): *Macular Edema*. Dev Ophthalm. Basel, Karger, 2010; vol 47, pp 148–159.
- 25 Frost NA, Sparrow JM, Strong NP, Rosenthal AR: Vitreous loss in planned extracapsular cataract extraction does lead to a poorer visual outcome. *Eye* 1995;9:446–451.
- 26 Flach AJ: The incidence, pathogenesis and treatment of cystoid macular edema following cataract surgery. *Trans Am Ophthalmol Soc* 1998;96:557–634.
- 27 Balent A, Civerchia LL, Mohamadi P: Visual outcomes of cataract extraction and lens implantation complicated by vitreous loss. *J Cataract Refract Surg* 1988;14:158–160.
- 28 Bergman M, Laatikainen L: Cystoid macular oedema after complicated cataract surgery and implantation of an anterior chamber lens. *Acta Ophthalmol (Copenh)* 1994;72: 178–180.
- 29 Blomquist PH, Rugwani RM: Visual outcomes after vitreous loss during cataract surgery performed by residents. *J Cataract Refract Surg* 2002;28:847–852.
- 30 Ah-Fat FG, Sharma MK, Majid MA, Yang YC: Vitreous loss during conversion from conventional extracapsular cataract extraction to phacoemulsification. *J Cataract Refract Surg* 1998;24:801–805.
- 31 Morel C, Roman S, Metge F, et al: Surgery of intravitreal nuclear luxations post-phacoemulsification (in French). *J Fr Ophthalmol* 1998;21:170–175.
- 32 Ray S, Amico DJ: Pseudophakic cystoid macular edema. *Semin Ophthalmol* 2002;17: 167–180.
- 33 Spaide RF, Yannuzzi LA, Sisco LJ: Chronic cystoid macular edema and predictors of visual acuity. *Ophthalmic Surg* 1993;24:262–267.
- 34 Borne MJ, Tasman W, Regillo C, et al: Outcomes of vitrectomy for retained lens fragments. *Ophthalmology* 1996;103:971–976.
- 35 Kapusta MA, Chen JC, Lam WC: Outcomes of dropped nucleous during phacoemulsification. *Ophthalmology* 1996;103:1184–1187.
- 36 Rossetti A, Doro D: Retained intravitreal lens fragments after phacoemulsification: complications and visual outcome in vitrectomized and nonvitrectomized eyes. *J Cataract Refract Surg* 2002;28:310–315.
- 37 Cohen SM, Davis A, Cukrowski C: Cystoid macular edema and pars plana vitrectomy for retained lens fragments. *J Cataract Refract Surg* 2006;32:1521–1526.
- 38 Sheidow TG, Gonder JR: Cystoid macular edema following combined phacoemulsification and vitrectomy in macular hole. *Retina* 1998;18:510–514.
- 39 Jampol LM: Aphakic cystoid macular edema: a hypothesis. *Arch Ophthalmol* 1985; 103:1134–1135.
- 40 Callanan D, Fellman RL, Savage JA: Latanoprost-associated cystoid macular edema. *Am J Ophthalmol* 1998;126:134–135.
- 41 Miyake K, Ota I, Maekubo K, et al: Latanoprost accelerates disruption of the blood-aqueous barrier and the incidence of angiographic cystoid macular edema in early postoperative pseudophakias. *Arch Ophthalmol* 1999;117:34–40.
- 42 Moroi SE, Gottfredsdottir MS, Scheingart MT, et al: Cystoid macular edema associated with latanoprost therapy in a case series of patients with glaucoma and ocular hypertension. *Ophthalmology* 1999;106:1024–1029.
- 43 Ayyala RS, Cruz DA, Margo CE, et al: Cystoid macular edema associated with latanoprost in aphakic and pseudophakic eyes. *Am J Ophthalmol* 1998;126:602–604.
- 44 Wand M, Gaudio AR: Cystoid macular edema associated with ocular hypotensive lipids. *Am J Ophthalmol* 2002;133:403–405.

- 45 Pollack A, Leiba H, Bukelman A, Oliver M: Cystoid macular edema following cataract extraction in patients with diabetes. *Br J Ophthalmol* 1992;76:221–224.
- 46 Schmier JK, Halpern MT, Covert DW, et al: Evaluation of cost for cystoid macular edema among patients after cataract surgery. *Retina* 2007;27:621–628.
- 47 Henricsson M, Heijl A, Janzon L: Diabetic retinopathy before and after cataract surgery. *Br J Ophthalmol* 1996;80:789–793.
- 48 Dowler JGF, Hykin PG, Lightman SL, et al: Visual acuity following extracapsular cataract extraction in diabetes: a meta-analysis. *Eye* 1995;9:313–317.
- 49 Schatz H, Atienza D, McDonald HR, et al: Severe diabetic retinopathy after cataract surgery. *Am J Ophthalmol* 1994;117:314–321.
- 50 Dowler JGF, Sehmi KS, Hykin PG, et al: The natural history of macular edema after cataract surgery in diabetes. *Ophthalmology* 1999;106:663–668.
- 51 Okhravi N, Lightman SL, Towler HM: Assessment of visual outcome after cataract surgery in patients with uveitis. *Ophthalmology* 1999;106:710–722.
- 52 Hooper PL, Rao NA, Smith RE: Cataract extraction in uveitis patients. *Surv Ophthalmology* 1990;35:120–144.
- 53 Foster RE, Lowder CY, Meisler OM, Zakov ZN: Extracapsular cataract extraction and posterior chamber intraocular lens implantation in uveitis patients. *Ophthalmology* 1992;99:1234–1241.
- 54 Schepens CL, Avila MP, Jalkh AE, et al: The role of vitreous in cystoid macular edema. *Surv Ophthalmol* 1984;28:499–504.
- 55 Cunha-Vaz JG, Travassos A: Breakdown of the blood-retinal barriers and cystoid macular edema. *Survey Ophthalmol* 1984;28:485–492.
- 56 Miyake K: Vitreous fluorophotometry in aphakic or pseudophakic eyes with persistent cystoid macular edema. *Jpn J Ophthalmol* 1985;29:146–152.
- 57 Smith RT, Campbell CJ, Koester CJ, et al: The barrier function in extracapsular cataract surgery. *Ophthalmology* 1990;97:90–95.
- 58 Miyake K: Prevention of cystoid macular edema after lens extraction by topical indomethacin. I. A preliminary report. *Albrecht von Graefes Arch Klin Exp Ophthalmol* 1977;203:81–88.
- 59 Miyake K: Prevention of cystoid macular edema after lens extraction by topical indomethacin. II. A control study in bilateral extractions. *Jpn J Ophthalmol* 1978;22:80–94.
- 60 Jampol LM: Cystoid macular edema following cataract surgery. *Arch Ophthalmol* 1988;106:894–895.
- 61 Stark WJ, Maumenee AE, Fagadau W, et al: Cystoid macular edema in pseudophakia. *Surv Ophthalmol* 1984;28:442–445.
- 62 Bitó LZ: Prostaglandins: old concepts and new perspectives. *Arch Ophthalmol* 1987;105:1036–1039.
- 63 Tso MOM: Pathophysiology of cystoid macular edema. *Ophthalmology* 1982;89:902–915.
- 64 Miyake K, Ibaraki N: Prostaglandins and cystoid macular edema. *Surv Ophthalmol* 2002;47(suppl 1):S203–S218.
- 65 Augustin A, Loewenstein A, Kuppermann BD: General pathophysiology; in Coscas G (eds): *Macular Edema*. Dev Ophthalmol. Basel, Karger, 2010, vol 47, pp 10–26.
- 66 Montes J, Erakgun T, Afrashi F, et al: Incidence of cystoid macular edema after uncomplicated phacoemulsification. *Ophthalmologica* 2003;217:408–412.
- 67 Asano S, Kensaku M, Ota I, et al: Reducing angiographic cystoid macular edema and blood-aqueous barrier disruption after small-incision phacoemulsification and foldable intraocular lens implantation. Multicenter prospective randomized comparison of topical diclofenac 0.1% and betamethasone 0.1%. *J Cataract Refract Surg* 2008;34:57–63.
- 68 Rossetti L, Chaudhwi H, Dickersin K: Medical prophylaxis and treatment of cystoid macular edema after cataract surgery: the results of a meta-analysis. *Ophthalmology* 1998;105:397–405.
- 69 Sivaprasad S, Bunce C, Wormald R: Nonsteroidal anti-inflammatory agents for cystoid macular oedema following cataract surgery: a systematic review. *Br J Ophthalmol* 2005;89:1420–1422.
- 70 Kraff MC, Sanders DR, Jampol LM, et al: Factors affecting pseudophakic cystoid macular edema: five randomized trials. *J Am Intraoc Implant Soc* 1985;11:380–385.
- 71 Henderson BA, Kim JY, Ament CS, et al: Clinical pseudophakic cystoid macular edema: risk factors for development and duration after treatment. *J Cataract Refract Surg* 2007;33:1550–1558.
- 72 Kim A, Stark WJ: Are topical NSAIDs needed for routine cataract surgery? *Am J Ophthalmol* 2008;146:554–560.
- 73 Arciere ES, Santana A, Rocha FN: Blood-aqueous barrier changes after the use of prostaglandins analogues in patients with pseudophakia and aphakia. *Arch Ophthalmol* 2005;123:186–192.
- 74 Lobo CL, Faria PM, Soares MA, et al: Macular alterations after small incision cataract surgery. *J Cataract Refract Surg* 2004;30:752–760.
- 75 Hee MR, Izatt JA, Swanson EA, et al: Optical coherence tomography of the human retina. *Arch Ophthalmol* 1995;113:325–332.
- 76 Perente I, Utine CA, Ozturker C, et al: Evaluation of macular changes after uncomplicated phacoemulsification surgery by optical coherence tomography. *Curr Eye Res* 2007;32:241–247.
- 77 Biro Z, Balla Z, Kovacs B: Change of foveal and perifoveal thickness measured by OCT after phacoemulsification and IOL implantation. *Eye* 2008;22:8–12.
- 78 Almeida DR, Johnson D, Hollands H, et al: Effect of prophylactic nonsteroidal anti-inflammatory drugs on cystoid macular edema assessed using optical coherence tomography quantification of total macular volume after cataract surgery. *J Cataract Refract Surg* 2008;34:64–69.
- 79 Salomon LD: Efficacy of topical flurbiprofen and indomethacin in preventing pseudophakic cystoid macular edema. Flurbiprofen – CME study group I. *J Cataract Refract Surg* 1995;21:73–81.
- 80 Cervantes-Costes G, Sánchez-Castro Y, Mendoza-Schuster E, et al: Inhibition of surgically induced miosis and prevention of postoperative macular edema with nepafenac. *Clin Ophthalmol* 2009;3:219–226.
- 81 Wolf EJ, Braunstein A, Shih C, et al: Incidence of visually significant pseudophakic macular edema after uneventful phacoemulsification in patients treated with nepafenac. *J Cataract Refract Surg* 2007;33:1546–1549.
- 82 Heier JS, Topping TM, Baumann W, et al: Ketorolac versus prednisolone versus combination therapy in treatment of acute pseudophakic cystoid macular edema. *Am Acad Ophthalmology* 2000;107:2034–2038.
- 83 Flach AJ: Nonsteroidal anti-inflammatory drugs; in Tasman W (ed): *Duane's Foundations of Clinical Ophthalmology*. Philadelphia, Lippincott, 1994, pp 1–32.
- 84 Jonas JB, Kreissing I, Segenring RF: Intravitreal triancinolone acetamide for pseudophakic cystoid macular edema. *Am J Ophthalmol* 2003;136:384–386.
- 85 Boscia F, Furino C, Dammaco R, et al: Intravitreal triancinolone acetamide in refractory pseudophakic cystoid macular edema: functional and anatomic results. *Eur J Ophthalmol* 2005;15:89–95.
- 86 Conway MD, Canakis C, Livir-Rallatos C, et al: Intravitreal triancinolone acetamide for refractory chronic pseudophakic cystoid macular edema. *J Cataract Refract Surg* 2003;29:27–33.
- 87 Tranos PG, Wickremasinghe SS, Stangos NT, et al: Macular edema. *Surv Ophthalmol* 2004;5:470–490.
- 88 Catiar A, Tadayoni R, Massin P, et al: Advantages of acetazolamide associated with anti-inflammatory medications in postoperative treatment of macular edema (in French). *J Fr Ophthalmol* 2005;28:1027–1031.
- 89 Spitzer MS, Ziemssen F, Yoeruek E, et al: Efficacy of intravitreal bevacizumab in treating postoperative pseudophakic cystoid macular edema. *J Cataract Refract Surg* 2008;34:70–75.
- 90 Steinert RF, Wasson PI: Neodymium: YAG laser anterior vitreolysis for Irvine-Gass cystoid macular edema. *J Cataract Refract Surg* 1989;15:304–307.
- 91 Alió J, Bodaghi B, Tassignon MJ: Guidelines for managing post-cataract surgery inflammation. Can we reach a consensus? *Ophthalmol Times Europe* 2008;(suppl Nov):1–11.