

## COMPOSITION AND BIOLOGICAL ACTIVITY OF THE ESSENTIAL OIL OF PERUVIAN *LANTANA CAMARA*

JULIO BENITES<sup>1</sup>, CRISTINA MOITEIRO\*<sup>2</sup>, GRAÇA MIGUEL<sup>3</sup>, LEONEL ROJO<sup>1</sup>, JOSÉ LÓPEZ<sup>1</sup>, FLORÊNCIA VENÂNCIO<sup>2</sup>, LUÍS RAMALHO<sup>2</sup>, SÓNIA FEIO<sup>2</sup>, SUSANA DANDLEN<sup>3</sup>, HUGO CASANOVA<sup>4</sup>, AND IVÁN TORRES<sup>4</sup>

<sup>1</sup>Departamento de Ciencias Químicas y Farmacéuticas, Universidad Arturo Prat, Casilla 121, Iquique, Chile.

<sup>2</sup>INETI, Instituto Nacional de Engenharia Tecnologia e Inovação I.P., Departamento de Tecnologia de Indústrias Químicas, Estrada do Paço do Lumiar, 1649-038 Lisboa, Portugal.

<sup>3</sup>Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal.

<sup>4</sup>Facultad de Farmacia y Bioquímica, Universidad Privada Antonio Guillermo Urrelo, Cajamarca, Perú.

(Received: February 23, 2009 - Accepted: June 24, 2009)

### ABSTRACT

The composition of the essential oil from *Lantana camara* L. (Verbenaceae) obtained by hydrodistillation of the aerial parts was examined by GC, GC/MS, and <sup>13</sup>C-NMR. The GC analysis showed that carvone is the most abundant monoterpene 75.9%, together with limonene 16.9%, accounting for 92.8% of the oil. The major components were also tested by <sup>13</sup>C-NMR analysis of the essential oil. The *L. camara* oil was assayed against several microorganisms, showing moderate antibacterial activity against the human pathogen *Staphylococcus aureus* (MIC 200 µg/ml). High antioxidant activity evaluated by the Trolox equivalent antioxidant capacity assay (TEAC) was found (29.0 mmol Trolox/kg) and relative low anti-inflammatory activity due to its weak ability for inhibiting lipoxigenase (IC<sub>50</sub> = 81.5 µg/ml).

**Keywords:** Antibacterial, antioxidant, anti-inflammatory, carvone, <sup>13</sup>C-NMR, essential oil composition, GC-MS, *Lantana camara*, limonene, Verbenaceae.

### INTRODUCTION

*Lantana camara* L. (Verbenaceae) is a hairy shrub native to Central and South America. This plant has been introduced in many countries as an ornamental or hedge plant, and is now a highly invasive weed in many places around the world. In Peru, *L. camara* (locally named as “yerba de la maestranza” and “siete colores”)<sup>1</sup> has been completely naturalized throughout the coast. Different parts of the plant are used in folk and traditional medicine systems for the treatment of health disorders including tetanus, malaria, tumors, and rheumatism, as well as also symptoms such as itches, dermatitis, ulcers, swellings, catarrh, dysentery, bilious fever, and eczema<sup>1-6</sup>.

The essential oils (EOs) derived from leaves and flowers of different colors of *L. camara* isolated by diverse methods and collected in different locations and seasons have been previously investigated, showing great variation in

their chemical composition (Table 1). In general, β-caryophyllene was the sesquiterpene identified in all essential oils from different origins (6-35%). The EOs of *L. camara* showed a wide spectrum of antibacterial and antifungal activities<sup>5,7</sup>.

Antioxidant and (Rosas-Romero & Saavedra, 2005) and anti-inflammatory activities<sup>9-11</sup> of several extracts of *L. camara* from different origins have been reported, but to the best of our knowledge this is the first report on the composition and biological activity of EO obtained from the Peruvian specimens of *L. camara*.

Here we report the analysis of EO from aerial parts of *L. camara* (yellow flowers) collected in Trujillo, Peru, and its biological activity against human pathogenic bacteria, as well as its antioxidant and anti-inflammatory activity. Our previous studies have demonstrated that Peru EO is enriched in carvone and limonene.

**Table 1.** Main constituents of the essential oils of *Lantana camara* from different geographical origins previously reported

Retention Indices Apolar <sup>a</sup>	Polar <sup>a</sup>	Main components	Percentage (%)	Origin	Part of plant	Type of isolation	Reference
1612 1780 2004		$\beta$ -Caryophyllene <i>ar</i> -Curcumene Caryophyllene epoxide	13 25 7	Cameroon (West Africa)	Leaves + flowers	Hydrodistillation	19
1123 1612 1724 1780 2031		Sabinene $\beta$ -Caryophyllene Zingiberene <i>ar</i> -Curcumene Nerolidol	7 11 19 16 11	Cameroon	Leaves	Hydrodistillation	19
1612 1780 2031		$\beta$ -Caryophyllene <i>ar</i> -Curcumene Nerolidol	8 27 13	Cameroon	Flowers	Hydrodistillation	19
1612 1724 1780		$\beta$ -Caryophyllene Zingiberene <i>ar</i> -Curcumene	20 11 27	Cameroon	Leaves	Headspace by dynamic method	19
1612 2040		$\beta$ -Caryophyllene Davanone	12 15	Madagascar	Leaves + flowers	Hydrodistillation	19
1118 1582		$\Delta^3$ -Carene $\beta$ -Caryophyllene	10 19	Madagascar	Not referred	Not referred	20
Not determined		Davanone $\beta$ -Caryophyllene Sabinene Linalool 1,8-Cineole $\alpha$ -Humulene $\beta$ -Bisabolene <i>ar</i> -Curcumene Caryophyllene oxide $\gamma$ -Cadinene	23-26 11-14 9-11 5-6 4-5 4-5 2 1-2 1 <0.5	Madagascar	Aerial part (pink-violet flowers)	Steam distillation	21
Not determined		$\beta$ -Caryophyllene $\beta$ -Bisabolene Sabinene $\gamma$ -Cadinene $\alpha$ -Humulene <i>ar</i> -Curcumene 1,8-Cineole Linalool Davanone Caryophyllene oxide	26-31 14-15 9-14 1-5 2-3 1-3 1 0.4-1 0-1 <05	Madagascar	Aerial part (yellow-orange flowers)	Steam distillation	21
Not determined		$\beta$ -Caryophyllene Davanone Linalool 1,8-Cineole $\beta$ -Bisabolene $\gamma$ -Muuroolene	16 12 5 4 2 1	Madagascar	Commercial	Not referred	21
1427 1460 1487 1500 1550 1560 1577 1640 1643		$\beta$ -Caryophyllene $\alpha$ -Humulene Germacrene D Bicyclogermacrene Elemol Nerolidol Caryophyllene oxide Cubenol $\alpha$ -Cadinol	21 11 4 9 3 4 4 2 2	Brazzaville (Congo)	Leaves	Hydrodistillation	22
968 1020 1419 1451 1495		Sabinene 1,8-Cineole $\beta$ -Caryophyllene $\alpha$ -Humulene $\delta$ -Guayene	13 9 19 10 5	Republic of Benin (Western Africa)	Leaves	Hydrodistillation	7

1474 1474 1658 1746	<i>ar</i> -Curcumene $\gamma$ -Curcumene <i>epi</i> - $\beta$ -bisabolol + others (-)- $\gamma$ -Curcumen-15-al	10 8 14 15	Brazil	Commercial (not referred)	Distillation	23	
974 1008 1028 1419 1480	Sabinene $\alpha$ -Phellandrene Limonene $\beta$ -Caryophyllene Germacrene D	9 16 17 11 13	State of Amapá (North Brazil)	Leaves + thin branches	Hydrodistillation	24	
1419 1480 1555	$\beta$ -Caryophyllene Germacrene D Germacrene B	6 28 9	State of Roraima (North Brazil)	Leaves + thin branches	Hydrodistillation	24	
1453 1484 1500	$\alpha$ -Humulene $\gamma$ -Curcumene + <i>ar</i> - curcumene $\alpha$ -Zingiberene	11 28 19	State of Pará (North Brazil)	Leaves + thin branches	Hydrodistillation	24	
1453 1484 1500	$\alpha$ -Humulene $\gamma$ -Curcumene + <i>ar</i> - curcumene $\alpha$ -Zingiberene	10 32 16	State of Pará (North Brazil)	Flowers	Hydrodistillation	24	
982 993 1425 1490 1494 1509 1515 1581	1124 1157 1628 1760 1726 1740 1758 1966	$\beta$ -Pinene Myrcene <i>trans</i> - $\beta$ -Caryophyllene Bicyclosesquiphellandrene $\alpha$ -Zingiberene <i>E,E</i> - $\alpha$ -Farnesene $\gamma$ -Bisabolene <i>trans</i> - $\beta$ -Caryophyllene oxide	0,2-3 2-4 14-15 16-18 5-6 3-4 3-4 4-9	Colombia	Leaves + branches	Hydrodistillation; Microwaves hydrodistillation; SFE; Dynamic headspace extraction	25
1020 975 1419 1450 1491	1,8-Cineole Sabinene $\beta$ -Caryophyllene $\alpha$ -Humulene Bicyclogermacrene	10 17 14 6 8	Iran	Leaves + flowers	Hydrodistillation	25	
1020 975 1419 1450 1491	1,8-Cineole Sabinene $\beta$ -Caryophyllene $\alpha$ -Humulene Bicyclogermacrene	6 7 23 11 18	Iran	Leaves + flowers	Steam distillation	26	
Not determined	$\beta$ -Caryophyllene Geranyl acetate Terpinyl acetate Bornyl acetate D-limonene	35 22 6 4 2	India	Aerial Parts	Not referred	5	
1380 1391 1425 1434 1484 1498	$\alpha$ -Copaene $\beta$ -Elemene $\beta$ -Caryophyllene $\alpha$ -Cadinene Germacrene D $\gamma$ -Elemene	5 7 9 3 21 10	Northern India	Leaves		27	
1380 1391 1425 1434 1484 1498	$\alpha$ -Copaene $\beta$ -Elemene $\beta$ -Caryophyllene $\alpha$ -Cadinene Germacrene D $\gamma$ -Elemene	11 15 7 7 11 7	Northern India	Flowers		27	
864 877 901 905	Germacrene D $\alpha$ -Humulene $\beta$ -Caryophyllene Germacrene B	16 9 12 6	South China	Aerial parts	Steam distillation	6	
	$\beta$ -Caryophyllene (10-30%)/ sabinene (30%)/bisabolenes (15%)/ $\beta$ -sesquiphellandrene (>10%)		Egypt	Leaves + flowers		28	

\*GC Column

## MATERIAL AND METHODS

### Plant material

*L. camara* plants with yellow flowers were collected in January 2007 from Trujillo at 32 m.a.s.l., in the region of La Libertad, Perú, and identified by Prof. Fredy Pelaéz from the "Herbario Trujillense de la Facultad de Ciencias Biológicas de la Universidad Nacional de Trujillo." A voucher sample under accession N° 932 was deposited in this herbarium.

### Oil essential isolation

Air dried aerial parts, 40 g, were subjected to hydrodistillation for 5 h using a Clevenger-type apparatus. The oil yield on a moisture-free basis was 4.5% (w/w).

### GC and GC-MS analysis

Chromatographic analysis of the oil of *L. camara* was carried out on a Hewlett Packard 5890 series II GC equipped with an Automatic Sampler HP 6890 series injector linked to two injector modules, two flame ionization detectors and two columns, together with a computer station for data treatment. The columns were an OV-101 fused silica (50 m x 0.25 mm, 0.25 µm film thickness) and Supelco wax10 (30 m x 0.25 mm, 0.25 µm film thickness). Oven temperature was held at 70°C for 5 min and then programmed to 220°C at 2°C/min. The detector and injector were kept at 250°C and 240°C, respectively. Helium was used as carrier gas, at constant pressure 55 kPa for the OV-101 and 45 kPa for the Supelco wax10. A 1 µl of oil was injected. The identification of the essential oil components is given in Table 2 and was made by comparison of their retention indices (RI) obtained from calculated values relative to C<sub>8</sub>-C<sub>20</sub> *n*-alkanes. The intensity of each peak was integrated. The sample was analyzed three times. The average peak areas of all GC signals were added together and the percentage of each component peak was calculated by comparing its average area to the total area.

The GC-MS analysis for identification of compounds was carried out in a Carlo Erba HRGC-MS gas chromatography equipped with a KRATOS mass detector model MS25RF (sector instrument) and a HP-5MS column (30 m x 0.25 mm, 0.25 µm film thickness), carrier gas helium, constant pressure 90 kPa, split 1:20. The oven was programmed initially from 70°C with 2 min hold up time to the final temperature of 250°C with 5°C/min ramp. The final temperature hold time was 5 min. The inlet and GC/MS interface temperatures were kept at 250°C and 280°C, respectively. The temperature of EI 70 eV source was 200°C with full scan (25-450m/z), scan time 0.3 s. The mass spectra of essential oil components were identified by comparing the mass spectra of the analytes with those of authentic standards from the mass spectra of Wiley 6.0 and Mass Spectra Library (NIST 98), and with corresponding data of components from reference oils analyzed in our laboratory.

### NMR Spectroscopy

The <sup>13</sup>C and DEPT spectra were recorded on a Bruker AMX300 spectrometer operating at 75.468 MHz equipped with a 5 mm QNP probe at room temperature. The EO (150 mg) was dissolved in deuteriochloroform (0.5 ml), with all chemical shift values (δ) refer to tetramethylsilane (TMS) as internal standard and are given in ppm. <sup>13</sup>C-NMR spectra were recorded with the following parameters: spectral width of 27777 Hz under low-power proton decoupling and <sup>13</sup>C pulse width of 7.5 µs, 45° excitation pulse, 2 s relaxation delay (D1) between scans, acquisition time of 3.867 s, line broadening of 1.0 Hz, for 64K data table. The number of accumulated scans was 1 K (1024) and the plot limits +230 to -6 ppm. Chemical shifts and peaks attribution of <sup>13</sup>C-NMR spectra were made according to those in the literature<sup>12</sup> and carvone and limonene standards. Multiplicities were obtained by 45, 90, and 135° DEPT experiments while relative intensity was determined as relative percentage of NMR peak height.

### Screening for antimicrobial activity

All microorganism strains were obtained from the Culture Collection of Industrial Microorganisms (CCMI) Laboratório de Microbiologia Industrial, Lisbon, Portugal. Bacteria: *Escherichia coli* CCMI 270, *Listeria monocytogenes* CCMI 1106 *Micrococcus luteus* CCMI 322, *Pseudomonas aeruginosa* CCMI 331, *Streptococcus faecium* CCMI 338, *Staphylococcus aureus* CCMI 335, *Streptococcus faecium* CCMI 338 were used in a quantitative bioassay. Brain Heart Infusion (Merck, Darmstadt, Germany) was used as culture medium. The temperature and incubation time were 37°C at 24 h, except for *M. luteus* which was incubated at 30°C for 48 h. The antimicrobial activity of essential oil was

determined by the broth dilution method<sup>13</sup>. The following concentrations were tested: 200, 100, 50, 25, and 12.5 µg/ml. After incubation the microbial growth was examined. The results are expressed in Minimal Inhibitory Concentration (MIC), the weakest concentration of essential oil yielding no visible growth. The bactericidal/bacteriostatic activity was determined by a sub-cultivation of the samples in normal culture media at appropriate temperature and incubation times. The MIC of each compound was determined at least twice. The essential oil was dissolved in dimethylsulphoxide (DMSO) (Merck, Darmstadt, Germany).

### Antioxidant activity

Green ABTS<sup>•+</sup> radicals were generated by oxidizing 7 mM aqueous solution of colorless ABTS [2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid)] with 2.45 mM potassium persulfate and allowing the mixture to stand in the dark at room temperature for 12-16 h before use. Radical scavenging capacity was measured against ABTS<sup>•+</sup> using 10 µl of oil samples or Trolox diluted with 900 µl of ABTS<sup>•+</sup> solution, and the absorbance reading was taken 1 min after initial mixing at 734 nm. The results were calculated using a standard curve prepared with Trolox and expressed as mM of Trolox equivalents per Kg of essential oil of *L. camara*.

### 5-Lipoxygenase assay

The assay mixture contained 10 µl of oil dissolved in DMSO and Tween 20, 0.1 M potassium phosphate buffer (pH 6.3, 2.95 ml) and 100 µM linoleic acid. The reaction is initiated with the addition of 100 U 5-lipoxygenase diluted with phosphate buffer kept at 4°C. The increase in absorbance at 234 nm was recorded for 10 min. The percentage inhibition of enzyme activity was calculated by comparison with the negative control constituted by DMSO and Tween 20. The percentage of enzyme activity was plotted against concentration of the oil. The IC<sub>50</sub> value is the concentration of the *L. camara* oil that caused 50% enzyme inhibition. Nordihydroguaiaretic acid (NDGA) represented the positive control.

## RESULTS AND DISCUSSION

The results of the GC and GC-MS analyses are listed in Table 2. Identification of the components was based on the comparison of the GC retention indices (RI) on polar and non-polar columns, determined relative to the retention time of a series of *n*-alkanes with linear interpolation with those of standards and with our essential oils database. The GC analysis of these EO showed that carvone is the most abundant monoterpene 75.9%, together with limonene 16.9% making up 92.8% of the total oil. Peru oil also contains traces of sabinene, *p*-cymene, terpinene, β-caryophyllene and germacrene D. The presence of the two major components was confirmed by MS, <sup>13</sup>C and DEPT NMR analysis (Table 3). The chemical shift values of the all carbons were compared with those reported for other oils<sup>12</sup> and with carvone and limonene standards. <sup>13</sup>C-NMR provided qualitative information on the total oil content of *L. camara* aerial parts.

**Table 2.** Percentage composition of essential oil components identified in *Lantana camara* collected in Trujillo, Peru.

Compound	RI <sup>1</sup>	Relative content (%)	Mode of identification
$\alpha$ -Thujene	916	0.4	RI, MS
$\alpha$ -Pinene	925	0.1	RI, MS
Camphene	941	0.1	RI, MS
Sabinene	964	0.2	RI, MS
Myrcene	979	0.4	RI, MS
$\alpha$ -Phellandrene	1000	0.1	RI, MS
<i>p</i> -Cymene	1014	0.2	RI, MS
Limonene	1025	16.9	RI, MS, NMR
<i>cis</i> -Ocimene	1028	0.6	RI, MS
<i>trans</i> -Ocimene	1037	0.1	RI, MS
Linalool	1081	0.7	RI, MS
<i>cis</i> -Limonene oxide	1118	0.1	RI, MS
Menthone	1136	0.2	RI, MS
Isomenthone	1153	0.2	RI, MS
<i>cis</i> -Dihydrocarvone	1175	0.1	RI, MS
<i>trans</i> -Dihydrocarvone	1182	0.2	RI, MS
<i>trans</i> -Carveol	1206	0.2	RI, MS
Carvone	1228	75.9	RI, MS, NMR
Piperitone	1235	0.8	RI, MS
Carvone oxide	1251	0.3	RI, MS
Piperitenone	1317	0.5	RI, MS
$\beta$ -Bourbonene	1391	0.5	RI, MS
$\beta$ -Elemene	1394	0.2	RI, MS
$\beta$ -Caryophyllene	1426	0.1	RI, MS
Germacrene D	1486	0.9	RI, MS
Nerolidol	1546	0.1	RI, MS

<sup>1</sup>Retention Indices identical to bibliography calculated through Kovats Indices on the OV-101 column.

**Table 3.** Spectral assignments of main monoterpenes detected in <sup>13</sup>C-NMR spectrum of *Lantana camara* essential oil

Peak	Compound	$\delta$ (ppm)	% Intensity	Assignment
1	Carvone	15.29	79.8	CH <sub>3</sub> (C-7)
2	Carvone	20.09	74.9	CH <sub>3</sub> (C-10)
3	Limonene	20.40	15.9	CH <sub>3</sub> (C-10)
4	Limonene	23.08	17.8	CH <sub>3</sub> (C-7)
5	Limonene	27.50	20.5	CH <sub>2</sub> (C-3)
6	Limonene	30.18	19.2	CH <sub>2</sub> (C-5)
7	Limonene	30.40	19.9	CH <sub>2</sub> (C-2)
8	Carvone	30.89	97.6	CH <sub>2</sub> (C-5)
9	Limonene	40.66	15.9	CH (C-4)
10	Carvone	42.08	100.0	CH (C-4)
11	Carvone	42.71	85.6	CH <sub>2</sub> (C-3)
12	Limonene	108.05	22.8	CH <sub>2</sub> (C-9)
13	Carvone	110.07	94.0	CH <sub>2</sub> (C-9)
14	Limonene	120.26	13.5	C=C (C-6)
15	Limonene	132.90	2.0	C=C (C-1)
16	Carvone	134.94	22.3	C=C (C-1)
17	Carvone	144.20	53.9	CH (C-6)
18	Carvone	146.21	28.8	C=C (C-8)
19	Limonene	149.00	<i>t.</i>	C=C (C-8)
20	Carvone	199.09	7.93	C=O (C-2)

*t.* = traces.

Antibacterial activity of *L. camara* EO was evaluated against the human pathogens *E. coli*, *P. aeruginosa*, *S. aureus*, *S. faecium* and the food spoilages *M. luteus* and *L. monocytogenes*. The EO inhibited the growth of *S. aureus* with bacteriostatic activity at 200  $\mu$ g/ml but did not exhibit any antibacterial activity against other microorganisms tested (MIC were >200  $\mu$ g/ml). Some authors<sup>5</sup> reported the activity of *L. camara* oil from India against *S. aureus*, *E. coli*, and *P. aeruginosa*. Peru oil did not show activity against the last two microorganisms, but the results are not comparable because the methodologies used and the chemical composition of the oils is different. The antibacterial activity may be related to the chemical proportion of the main compounds, carvone and limonene, as well to the minor components present in our EO. Some authors<sup>14</sup> detected that isolated (*R*)-(+)-isomers of limonene and carvone appeared to be much more potent against a range of bacteria when compared to the essential oils where they occur. However, these authors also observed that corresponding (*S*)-(-)-isomers may exhibit different activities which could be due to the synergism of the minor components present in EOs. Indeed, chemically the EOs are quite complex mixtures of a large variety of constituents, and their antimicrobial activity can be attributed to several active compounds as well as to the synergetic effects between them and to the different proportions in which they are present in the mixtures. Thus, this complexity makes it often difficult to explain the activities.

Several assays have been frequently used to estimate antioxidant capacities in essential oils, including 2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS)<sup>15,16</sup>. This method is based on the ability of antioxidants to quench the ABTS radical cation relative to that of Trolox (water-soluble vitamin E analogue). The activity found for *L. camara* was 29.0 mmol Trolox/kg. This value is similar to that obtained by some authors<sup>16</sup> for *Origanum vulgare* L. oil (25.1 mmol Trolox/kg) and higher than that of *Lippia alba* Mill. oil (14.4 mmol Trolox/kg) in which carvacrol and carvone predominated in the respective oils. The authors attributed the antioxidant activity of *O. vulgare* oil to the relatively large amount of carvacrol, but in the same work other essential oils also showed values close to that observed for oregano oil, and they were constituted by *trans*-anethole (*Pimpinella anisum* L.) (24.3 mmol Trolox/kg), ledol (*Lepechinia schiedeana*) (25.7 mmol Trolox/kg), and linalool (*Cananga odorata*) (38.6 mmol Trolox/kg). Comparing these results with ours, we can consider that the carvone-rich oil of *L. camara* from Peru possesses a particularly high antioxidant capacity. Such results may be partially explained by the presence of carvone, a  $\alpha,\beta$ -unsaturated ketone, which is more active than ketone monoterpenes according some authors<sup>17</sup>. Until now, only extracts of *L. camara* from Bolivia were reported as good antioxidants when evaluated by two other methods:  $\beta$ -carotene bleaching technique and DPPH free radical technique.

Inflammation is a normal pathophysiological response to the attack of infectious agents or to physical, chemical or traumatic injuries. The mechanism of inflammation involves complex cascades of events in which the pro-inflammatory leukotrienes, resulting from the metabolism of arachidonic acid have an important role. The enzyme 5-lipoxygenase, which promotes the oxidation of arachidonic acid, is involved in the formation of this class of messengers<sup>18</sup>. The concentration of *L. camara* oil that produced a 50% inhibition (IC<sub>50</sub>) of lipoxygenase was 81.5  $\mu$ g/ml, not as good as the positive control (49.5  $\mu$ g/ml). According to an arbitrary relative *in vitro* anti-inflammatory activity scale established by Baylac and Racine<sup>18</sup>, we can consider that the *L. camara* oil from Peru possesses a weak activity (51 < IC<sub>50</sub> £ 100 mg L/1) when assessed by this method. Extracts of *L. camara* have revealed anti-inflammatory activities<sup>9,11</sup>, but such results cannot be directly compared to ours since the methods used for anti-inflammatory evaluations were quite different. Some authors<sup>10</sup> also reported anti-inflammatory activity in extracts of *L. camara* in which saponins, tannins and flavonoids were present. The anti-inflammatory activity of *L. camara* has been attributed to the presence of the pentacyclic triterpenes, oleanolic acid, and ursolic acid, which act as inhibitors of human leucocyte elastase as well as COX-2 (cyclooxygenase isoenzyme)<sup>2</sup>. The high antioxidant activity of *L. camara* oil from Peru, evaluated by the Trolox equivalent antioxidant capacity assay (TEAC) was observed (29.0 mmol Trolox/kg). We suggest that this effect may be related to the presence of carvone and limonene, which are the major components the EO from *L. camara* (92.8%) and also considerable antioxidants. Since carvone had not yet been identified in *L. camara* oils (see Table 1) the Peru oil seems to be a new chemotype, which may be confirmed through the chemical composition analysis of plants recollected in different Peru regions and seasons in order to compare the chemical evolution of the main constituents.

## ACKNOWLEDGEMENTS

We acknowledge with thanks the financial support of the Dirección de Investigación, Universidad Arturo Prat in Chile, and Departamento de Tecnología de Industrias Químicas-INETI and Centro de Desenvolvimento de Ciências e Técnicas de Produção Vegetal (CDCTPV), in Portugal.

## REFERENCES

1. A. Brick, Diccionario Enciclopédico de Plantas útiles del Perú. Programa de las Naciones Unidas para el Desarrollo. Centro de Estudios Regionales Andinos "Bartolomé de Las Casas" Cusco, Perú, 1999.
2. E. L. Ghisalberti, *Fitoterapia*. **71**, 467, (2000).
3. O. P. Sharma, *Vet. Hum. Toxicol.* **26**, 488, (1984).
4. O. P. Sharma, H. P. S. Makkar, R. K. Dawra, S. S. Negi, *Toxicol.* **26**, 975, (1988).
5. M. J. Deena, J. E. Thoppil, *Fitoterapia*. **71**, 453, (2000).
6. A. J. Sundufú, H. Shoushan, *Flavour. Fragr. J.* **19**, 229, (2004).
7. G. Alitonou, F. Avlessi, I. Bokossa, E. Ahoussi, J. Dangou, D. C. K. Sohounhloué, *C.R. Chim.* **7**, 1101, (2004).
8. A. Rosas-Romero, G. Saavedra, *Pharm. Biol.* **43**, 79, (2005).
9. A. M. Forestieri, M. T. Monforte, S. Ragusa, A. Trovato, *Phytother. Res.* **10**, 100, (1996).
10. O. O. Oyedapo, F. C. Sab, J. A. Olagunju, *Biomed. Lett.* **59**, 175, (1999).
11. S. Basu, B. Hazra, *Phytother. Res.* **20**, 896, (2006).
12. K.-H. Kubeczka, V. Formáček, *Essential Oils Analysis by Capillary Gas Chromatography and Carbon-13 NMR Spectroscopy*, 2nd Edition, John Wiley & Sons, Ltd, England, 2002.
13. P. A. Wayne, Performance standards for antimicrobial susceptibility testing. Ninth informational supplement NCCLS document M100-S9. National Committee for Clinical Laboratory Standards, 2008.
14. K. K. Aggarwal, S. P. S. Khanuja, A. Ahmad, T. R. Santha Kumar, V. K. Gupta, S. Kumar, *Flavour. Fragr. J.* **17**: 59, (2002).
15. D. Mantle, J. G. Anderton, G. Falkous, M. Barnes, P. Jones, E. K. Perry, *Comp. Biochem. Physiol. (Part B)*. **121**, 385, (1998).
16. M. Puertas-Mejía, S. Hillebrand, E. Stashenko, P. Winterhalter, *Flavour. Fragr. J.* **17**, 380, (2002).
17. G. Ruberto, M. T. Baratta, *Food Chem.* **69**, 167, (2000).
18. S. Baylac, P. Racine, *Int. J. Aromather.* **23**, 138, (2003).
19. M. B. Ngassoum, S. Yonkeu, L. Jirovetz, G. Buchbauer, G. Schmaus, F. J. Hammerschmidt, *Flavour. Fragr. J.* **14**, 245, (1999).
20. S. Mollenbeck, T. König, P. Schreier, W. Schwab, J. Rajaonarivony, L. Ranarivelo, *Flavour. Fragr. J.* **12**, 63, (1997).
21. J.-A. Randrianalijaona, P. Ramanoelina, J. Rasoarahona, E. Gaydou, *Anal. Chim. Acta.* **545**, 46, (2005).
22. J.-M. Ouamba, A. Ouabonzi, A. Ekouya, J.-M. Bessière, C. Menut, A. A. Abena, J.-T. Banzouzi, *Flavour. Fragr. J.* **21**, 158, (2006).
23. P. Weyerstahl, H. Marschall, A. Eckhardt, C. Christiansen, *Flavour. Fragr. J.* **14**, 15, (1999).
24. M. H. da Silva, E. H. Andrade, M. G. Zoghbi, A. I. Luz, J. D. Silva, J. G. Maia, *Flavour. Fragr. J.* **14**, 208, (1999).
25. E. E. Stasbenko, B. F. Jaramillo, J. R. Matinez, *Rev. Acad. Coloma Cienc.* **27**, 579, (2003).
26. F. Sefidkon, *Flavour. Fragr. J.* **17**, 78, (2002).
27. M. Khan, S. K. Srivastava, K. V. Syamasundar, M. Singh, A. A. Naqvi, *Flavour. Fragr. J.* **17**, 75, (2002).
28. M. M. Fathy, *Bull. Fac. Pharm. Cairo Univ.* **38**, 105, (2000).