


Morphological and molecular identification for four new wood-inhabiting species of *Trechispora* (Basidiomycota) from China

Kai-Yue Luo¹, Jiang-Qing Su^{1,2}, Chang-Lin Zhao^{1,2}

¹ The Key Laboratory of Forest Resources Conservation and Utilization in the Southwest Mountains of China Ministry of Education, Key Laboratory of National Forestry and Grassland Administration on Biodiversity Conservation in Southwest China, Yunnan Provincial Key Laboratory for Conservation and Utilization of In-forest Resource, Southwest Forestry University, Kunming 650224, China

² College of Forestry, Southwest Forestry University, Kunming 650224, China

Corresponding author: Chang-Lin Zhao (fungi@swfu.edu.cn)

Abstract

Four new wood-inhabiting fungi, *Trechispora albofarinosa*, *T. bisterigmata*, *T. pileata* and *T. wenshanensis* **spp. nov.**, are proposed based on a combination of morphological features and molecular evidence. *Trechispora albofarinosa* is characterized by the farinose basidiomata with flocculence hymenial surface, a monomitic hyphal system with clamped generative hyphae, and ellipsoid, warted basidiospores. *Trechispora bisterigmata* is characterized by the membranous basidiomata with odontoid hymenial surface, rhizomorphic sterile margin, barrelled basidia and subglobose to broad ellipsoid, smooth basidiospores. *Trechispora pileata* is characterized by the laterally contracted base, solitary or imbricate basidiomata, fan shaped pileus, radially striate-covered surface with appressed scales, odontoid hymenophore surface, and subglobose to broad ellipsoid, thin-walled, smooth basidiospores. *Trechispora wenshanensis* is characterized by a cottony basidiomata with a smooth hymenial surface, and ellipsoid, thin-walled, warted basidiospores. Sequences of ITS and LSU marker of the studied samples were generated, and phylogenetic analyses were performed with the maximum likelihood, maximum parsimony, and Bayesian inference methods. The phylogenetic tree inferred from the ITS+nLSU sequences highlighted that four new species were grouped into the genus *Trechispora*.

Key words: East Asia, macrofungi, molecular systematics, taxonomy, 4 new taxa



Academic editor: Bao-Kai Cui

Received: 7 February 2024

Accepted: 29 April 2024

Published: 15 May 2024

Citation: Luo K-Y, Su J-Q, Zhao C-L (2024) Morphological and molecular identification for four new wood-inhabiting species of *Trechispora* (Basidiomycota) from China. MycoKeys 105: 155–178. <https://doi.org/10.3897/mycokeys.105.120438>

Copyright: © Kai-Yue Luo et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Fungi represent one of the most diverse groups of organisms on earth, with an indispensable role in the processes and functioning of ecosystems (Wu et al. 2020, 2021; Wang et al. 2021; Hyde 2022; Guan and Zhao 2023). Wood-inhabiting fungi play an important role in the carbon cycle (Dai et al. 2015; Wu et al. 2017; Chen and Zhao 2020; Guan et al. 2021; Liu et al. 2022a, b; Luo and Zhao 2022; Mardones et al. 2023; Yuan et al. 2023; Zhang et al. 2023; Zhao et al. 2023a, b; Zhou et al. 2023). The wood-inhabiting fungal order Trechisporales

K.H. Larss. is a species-poor order, compared with most other orders within Agaricomycetes, Basidiomycota (Wijayawardene et al. 2022).

Trechispora P. Karst. (Hydnodontaceae) typified by *T. onusta* P. Karst., which is characterized by resupinate to effused basidiomata; a smooth to hydroid to poroid hymenophore; ampullaceous septa; short cylindrical basidia; and smooth to verrucose or aculeate basidiospores (Karsten 1890; Bernicchia and Gorjón 2010). Currently, MycoBank and Index Fungorum have registered 163 recorded and 156 recorded intraspecific names in *Trechispora*, respectively. About 100 species are currently accepted in *Trechispora* worldwide (Karsten 1890; Bondartsev and Singer 1941; Rogers and Jackson 1943; Rogers 1944; Bondartsev 1953; Liberta 1966, 1973; Parmasto 1968; Burdsall and Gilbertson 1982; Gilbertson and Budington 1970; Jülich 1975, 1976; Ryvarden 1975; Ryvarden and Liberta 1978; Hallenberg 1978, 1980; Jülich and Stalpers 1980; Rauschert 1987; Vries 1987; Larsson 1992, 1994, 1995, 1996; Hjortstam and Larsson 1995; Ryvarden 2002; Trichies and Schultheis 2002; Ryvarden et al. 2003; Miettinen and Larsson 2006; Dai 2011; Yuan and Dai 2012; Ordynets et al. 2015; Phookamsak et al. 2019; Xu et al. 2019; Chikowski et al. 2020; Haelewaters et al. 2020; Crous et al. 2021; de Meiras-Ottoni et al. 2021; Zhao and Zhao 2021; Liu et al. 2022a, b; Luo and Zhao 2022; Sommai et al. 2023), of which 38 species of the genus have been found in China (Dai 2011; Yuan and Dai 2012; Xu et al. 2019; Dai et al. 2021; Zhao and Zhao 2021; He et al. 2022; Liu et al. 2022a, b; Luo and Zhao 2022; Deng et al. 2023; Liu et al. 2024a, b).

There have been many studies on the phylogeny of this genus in recent years. A high phylogenetic diversity on the corticioid Agaricomycetes based on two genes, 5.8S and 28S showed that nine taxa of *Trechispora* nested into trechisporoid clade (Larsson et al. 2004). The molecular systematics suggested that *Trechispora* belonged to Hydnodontaceae and was related to genera *Brevicellium* K.H. Larss. & Hjortstam, *Porpomyces* Jülich, *Sistotremastrum* J. Erikss., and *Subulicystidium* Parmasto (Telleria et al. 2013). Based on the ITS and nLSU datasets, the phylogenetic study of *Trechispora* reported two new *Trechispora* species as *T. cyatheae* Ordynets, Langer & K.H. Larss. and *T. echinocrystallina* Ordynets, Langer & K.H. Larss., on La Réunion Island (Ordynets et al. 2015). The phylogeny of Trechisporales was inferred from a combined ITS-nLSU sequences, which revealed that two related genera *Porpomyces*, *Scytinopogon* Singer, grouped closely together with *Trechispora* and all of them nested within Hydnodontaceae (Liu et al. 2019). Based on ITS dataset, the three new species of *Trechispora* were described and used to evaluate the phylogenetic relationship with other species of this genus, in which *T. murina* was retrieved as a sister to *T. bambusicola* with moderate supports, and *T. odontioidea* formed a single lineage and then grouped with *T. fimbriata* and *T. nivea*, while *T. olivacea* formed a monophyletic lineage with *T. farinacea*, *T. hondurensis*, and *T. mollis* (Luo and Zhao 2022). Recently, based on the morphological features and molecular evidence, three new species of *Trechispora* have been reported from Northern and Northeastern Thailand (Sommai et al. 2023).

During investigations into the wood-inhabiting fungi in the Yunnan-Guizhou Plateau of China, samples representing four additional species belonging to genus *Trechispora* were collected. To clarify the placement and relationships of the four species, we carried out a phylogenetic and taxonomic study on *Trechispora*, based on the ITS+nLSU.

Materials and methods

Morphology

The specimens studied were deposited at the herbarium of Southwest Forestry University (SWFC), Kunming, Yunnan Province, China. The macromorphological descriptions were based on field notes and photos captured in the field and laboratory. Color, texture, taste and odor of basidiomata were mostly based on authors' field trips. Color terminology followed Kornerup and Wanscher (1978). All materials were examined under a Nikon 80i microscope. Drawings were made with the aid of a drawing tube. The measurements and drawings of the microscopic structures were made (Wu et al. 2022). The following abbreviations were used: KOH = 5% potassium hydroxide water solution, CB = cotton blue, CB- = acyanophilous, IKI = Melzer's reagent, IKI- = both inamyloid and indextrinoid, L = spore length (arithmetic average for all spores), W = spore width (arithmetic average for all spores), Q = L/W ratios of the specimens studied, and n = a/b (a = total number of spores measured, from b = number of specimens).

Molecular phylogeny

The CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd, Beijing) was used to obtain genomic DNA from the dried specimens following the manufacturer's instructions (Zhao and Wu 2017). The nuclear ribosomal ITS region was amplified with the primers ITS5 and ITS4 (White et al. 1990). The nuclear ribosomal LSU gene was amplified with the primers LR0R and LR7 (Vilgalys and Hester 1990; Rehner and Samuels 1994). The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min, followed by 35 cycles at 94 °C for 40 s, 58 °C for 45 s and 72 °C for 1 min, and a final extension of 72 °C for 10 min. The PCR procedure for nLSU was as follows: initial denaturation at 94 °C for 1 min, followed by 35 cycles at 94 °C for 30 s, 48 °C for 1 min and 72 °C for 1.5 min, and a final extension of 72 °C for 10 min. The PCR products were purified and directly sequenced at Kunming Tsingke Biological Technology Limited Company, Yunnan Province, China. All newly-generated sequences were deposited in NCBI GenBank (Table 1).

The sequences were aligned in MAFFT version 7 (Katoh et al. 2019) using the G-INS-i strategy. The alignment was adjusted manually using AliView version 1.27 (Larsson 2014). Each dataset was aligned separately at first and then the ITS+nLSU regions were combined with Mesquite version 3.51. The combined dataset was deposited in TreeBASE (submission ID 31349). Sequences of *Fibrodontia alba* Yurchenko & Sheng H. Wu and *F. brevidens* (Pat.) Hjortstam & Ryvar den retrieved from GenBank were used as an outgroup in the ITS analysis (Luo and Zhao 2022).

Maximum parsimony analysis in PAUP* version 4.0a169 (<http://phylosolutions.com/paup-test/>) was applied to ITS+nLSU following a previous study (Zhao and Wu 2017). All characters were equally weighted and gaps were treated as missing data. Trees were inferred using the heuristic search option with TBR branch swapping and 1,000 random sequence additions. Max-trees were set to 5,000, branches of zero length were collapsed and all parsimonious trees were saved. Clade robustness was assessed using bootstrap (BT) analysis with 1,000 pseudo replicates (Felsenstein 1985).

Table 1. List of species, specimens and GenBank accession numbers of sequences used in this study.

Species name	Specimen No.	GenBank accession No.		References
		ITS	LSU	
<i>Fibrodontia alba</i>	TNM F24944	NR153983	NG060401	Yurchenko and Wu 2014
<i>F. brevidens</i>	Wu 9807-16	KC928276	KC928277	Yurchenko and Wu 2014
<i>Trechispora alba</i>	CH21384	OR557258	–	Liu et al. 2024a
<i>T. albofarinosa</i>	CLZhao 4356	OQ241383	OQ282703	This study
<i>T. amianthina</i>	CBS 202.54	–	MH868822	Vu et al. 2019
<i>T. araneosa</i>	KHL 8570	AF347084	–	Larsson et al. 2004
<i>T. bambusicola</i>	CLZhao 3302	MW544021	MW520171	Zhao and Zhao 2021
<i>T. bambusicola</i>	CLZhao 3305	MW544022	MW520172	Zhao and Zhao 2021
<i>T. bispora</i>	CBS 142.63	MH858241	MH869842	Larsson et al. 2004
<i>T. bisterigmata</i>	CLZhao 2522	OQ241386	–	This study
<i>T. bisterigmata</i>	CLZhao 7870	OQ241387	–	This study
<i>T. byssinella</i>	UC 2023068	KP814481	–	Unpublished
<i>T. chartacea</i>	FLOR56185	MK458775	–	Liu et al. 2022a
<i>T. clancularis</i>	FRDBI 4426619	MW487976	–	Unpublished
<i>T. cohaerens</i>	HHB-19445	MW740327	–	Unpublished
<i>T. copiosa</i>	AMO427	MN701015	MN687973	de Meiras-Otoni et al. 2021
<i>T. copiosa</i>	AMO450	MN701017	MN687974	de Meiras-Otoni et al. 2021
<i>T. crystallina</i>	LWZ 20170729-2	OM523419	OM339238	Liu et al. 2022a
<i>T. cyatheae</i>	FR0219443	UDB024016	UDB024017	Ordynets et al. 2015
<i>T. cyatheae</i>	FR0219446	UDB024020	UDB024021	Ordynets et al. 2015
<i>T. dentata</i>	Dai 22565	OK298491	OM049408	Liu et al. 2022b
<i>T. dimitiella</i>	Dai 21181	OK298493	OK298949	Liu et al. 2022b
<i>T. dimitiella</i>	Dai 21931	OK298492	OK298948	Liu et al. 2022b
<i>T. echinospora</i>	E11/37-10	JX392850	JX392851	Telleria et al. 2013
<i>T. echinospora</i>	E11/37-12	JX392853	JX392854	Telleria et al. 2013
<i>T. farinacea</i>	356	AF347089	–	Larsson et al. 2004
<i>T. farinacea</i>	MA-Fungi 79474	JX392855	JX392856	Telleria et al. 2013
<i>T. fimbriata</i>	CLZhao 7969	MW544024	MW520174	Zhao and Zhao 2021
<i>T. fimbriata</i>	CLZhao 9006	MW544025	MW520175	Zhao and Zhao 2021
<i>T. foetida</i>	FLOR 56315	MK458769	–	Liu et al. 2022a
<i>T. fragilis</i>	Dai 20535	OK298494	OK298950	Liu et al. 2022b
<i>T. gelatinosa</i>	AMO824	MN701020	MN687977	de Meiras-Otoni et al. 2021
<i>T. gelatinosa</i>	AMO1139	MN701021	MN687978	de Meiras-Otoni et al. 2021
<i>T. gracilis</i>	LWZ 20170814-17	OM523435	OM339253	Liu et al. 2022a
<i>T. havencampii</i>	DED8300	NR154418	NG059993	Desjardin and Perry 2015
<i>T. hondurensis</i>	HONDURAS19-F016	NR178152	NG081479	Haelewaters et al. 2020
<i>T. hondurensis</i>	HONDURAS19-F016a	MT571523	MT636540	Haelewaters et al. 2020
<i>T. hymenocystis</i>	KHL 8795	AF347090	–	Unpublished
<i>T. hymenocystis</i>	KHL 16444	MT816397	–	Unpublished
<i>T. incisa</i>	GB0090521	KU747093	KU747086	Unpublished
<i>T. incisa</i>	GB0090648	KU747095	KU747087	Unpublished
<i>T. invisitata</i>	5425_537	ON963772	–	Unpublished
<i>T. invisitata</i>	UC2023088	KP814425	–	Unpublished

Species name	Specimen No.	GenBank accession No.		References
		ITS	LSU	
<i>T. kavinioides</i>	KG 981002	AF347086	–	Unpublished
<i>T. laevispora</i>	Dai 21655	OK298495	OM108710	Liu et al. 2022b
<i>T. larssonii</i>	LWZ 20190817-11a	OM523442	OM339259	Liu et al. 2022a
<i>T. longiramosa</i>	HG 140168	OM523448	OM339264	Liu et al. 2022a
<i>T. mellina</i>	URM85756	–	MH280000	Unpublished
<i>T. microspora</i>	FRDBI 18772216	OL828778	–	Unpublished
<i>T. mollis</i>	URM85884	MK514945	MK514945	Unpublished
<i>T. mollis</i>	URM85885	–	MT423667	Unpublished
<i>T. mollusca</i>	iNAT 30809943	MZ269232	–	Unpublished
<i>T. mollusca</i>	CFMR:DLL2011-186	KJ140681	–	Unpublished
<i>T. nivea</i>	MA-Fungi 76238	JX392824	JX392825	Telleria et al. 2013
<i>T. nivea</i>	MA-Fungi 76257	JX392826	JX392827	Telleria et al. 2013
<i>T. pallescens</i>	FLOR56184	MK458767	–	Unpublished
<i>T. pallescens</i>	FLOR56188	MK458774	–	Unpublished
<i>T. papillosa</i>	AMO713	MN701022	MN687979	de Meiras-Otoni et al. 2021
<i>T. papillosa</i>	AMO795	MN701023	MN687981	de Meiras-Otoni et al. 2021
<i>T. patawaensis</i>	VPapp-GF1901	OL314550	OL314546	Unpublished
<i>T. perminispora</i>	LWZ2019081639a	OM523525	OM339329	Liu et al. 2024a
<i>T. pileata</i>	CLZhao 4456	OQ241388	OQ282715	This study
<i>T. praefocata</i>	FRDBI 18819116	OL828784	–	Unpublished
<i>T. regularis</i>	KHL 10881	AF347087	–	Unpublished
<i>T. rigida</i>	URM85754	MT406381	MH279999	Unpublished
<i>T. sinensis</i>	LWZ 20170816-35	OM523479	OM339287	Liu et al. 2022a
<i>T. stellulata</i>	14153	MW023104	–	Unpublished
<i>T. stellulata</i>	33962903	ON364078	–	Unpublished
<i>T. stellulata</i>	UC2023099	KP814451	–	Unpublished
<i>T. stellulata</i>	UC2023230	KP814491	–	Unpublished
<i>T. stevensonii</i>	MA-Fungi 70669	JX392841	JX392842	Telleria et al. 2013
<i>T. stevensonii</i>	MA-Fungi 70645	JX392843	JX392844	Telleria et al. 2013
<i>T. subfarinacea</i>	LWZ2020092133a	OM523528	OM339331	Liu et al. 2024a
<i>T. subhelvetica</i>	7089	JN710601	–	Unpublished
<i>T. subhymenocystis</i>	LWZ 20190818-29b	OM523492	OM339299	Liu et al. 2022a
<i>T. subregularis</i>	VPapp-GF2103	OL331097	OL314548	Unpublished
<i>T. subsinensis</i>	LWZ 20190611-9	OM523497	OM339304	Liu et al. 2022a
<i>T. subsphaerospora</i>	KHL 8511	AF347080	–	Unpublished
<i>T. termitophila</i>	AMO396	MN701025	MN687983	de Meiras-Otoni et al. 2021
<i>T. termitophila</i>	AMO893	MN701026	MN687984	de Meiras-Otoni et al. 2021
<i>T. torrendii</i>	URM85886	MK515148	MH280004	Unpublished
<i>T. tropica</i>	LWZ 20170613-16	OM523503	OM339311	Liu et al. 2022a
<i>T. tuberculata</i>	Dai17433	OM523507	OM339314	Liu et al. 2024a
<i>T. wenshanensis</i>	CLZhao 11649	OQ241389	OQ282716	This study
<i>T. wenshanensis</i>	CLZhao 11715	PP712100	–	This study
<i>T. wenshanensis</i>	CLZhao 22940	PP712101	–	This study
<i>T. yunnanensis</i>	CLZhao 210	NR177488	MN654918	Xu et al. 2019
<i>T. yunnanensis</i>	CLZhao 214	MN654922	MN654919	Xu et al. 2019

Descriptive tree statistics - tree length (TL), composite consistency index (CI), composite retention index (RI), composite rescaled consistency index (RC) and composite homoplasy index (HI) - were calculated for each maximum parsimonious tree generated. The combined dataset was also analysed using Maximum Likelihood (ML) in RAxML-HPC2 through the CIPRES Science Gateway (Miller et al. 2012). Branch support (BS) for the ML analysis was determined by 1000 bootstrap pseudoreplicates.

MrModeltest 2.3 (Nylander 2004) was used to determine the best-fit evolution model for each dataset for the purposes of Bayesian inference (BI), Bayesian inference was performed using MrBayes 3.2.7a with a GTR+I+G model of DNA substitution and a gamma distribution rate variation across sites (Ronquist et al. 2012). A total of four Markov chains were run for two runs from random starting trees for 1.7 million generations for ITS+nLSU with tree and parameters sampled every 1,000 generations. The first quarter of all of the generations were discarded as burn-ins. A majority rule consensus tree was computed from the remaining trees. Branches were considered as significantly supported if they received a maximum likelihood bootstrap support value (BS) of > 70%, a maximum parsimony bootstrap support value (BT) of > 70% or a Bayesian posterior probability (BPP) of > 0.95.

Results

Molecular phylogeny

The ITS+nLSU dataset comprised sequences from 88 fungal specimens representing 64 taxa. The dataset had an aligned length of 2271 characters, of which 1376 characters were constant, 190 were variable and parsimony-uninformative and 705 were parsimony-informative. Maximum parsimony analysis yielded 300 equally parsimonious tree (TL = 5543, CI = 0.2979, HI = 0.7021, RI = 0.5278 and RC = 0.1572). The best model of nucleotide evolution for the ITS+nLSU dataset estimated and applied in the Bayesian analysis was found to be GTR+I+G. Bayesian analysis and ML analysis resulted in a similar topology as in the MP analysis. The Bayesian analysis had an average standard deviation of split frequencies = 0.012925 (BI) and the effective sample size (ESS) across the two runs is double the average ESS (avg. ESS) = 389. The phylogenetic tree inferred from the ITS+nLSU sequences highlighted that four new species were grouped into the genus *Trechispora* (Fig. 1).

Taxonomy

***Trechispora albofarinosa* K.Y. Luo & C.L. Zhao, sp. nov.**

MycoBank No: 849463

Figs 2, 3

Holotype. CHINA. Yunnan Province, Pu'er, Jingdong County, Huangcaoling, Wuliangshan National Nature Reserve, 24°23'N, 100°45'E, altitude 2350 m a.s.l., on the fallen branch of *Pinus*, leg. C.L. Zhao, 5 October 2017, CLZhao 4356 (SWFC).

Etymology. *Albofarinosa* (Lat.): referring to the farinose basidiomata with white hymenial surface.




Figure 1. Maximum parsimony strict consensus tree illustrating the phylogeny of the four new species and related species in *Trechispora*, based on ITS+nLSU sequences. Branches are labelled with maximum likelihood bootstrap values > 70%, parsimony bootstrap values > 50% and Bayesian posterior probabilities > 0.95, respectively.




Figure 2. Basidiomata of *Trechispora albofarinosa* (holotype) **A** the front of the basidiomata **B** characteristic hymenophore. Scale bars: 1 cm (**A**); 1 mm (**B**).




Figure 3. Microscopic structures of *Trechispora albofarinosa* (holotype) **A** basidiospores **B** basidia and basidioles **C** a cross section of basidiomata. Scale bars: 5 μm (**A**); 10 μm (**B**, **C**).

Description. Basidiomata annual, resupinate, farinose, without odor or taste when fresh, up to 3.5 cm long, 1.5 cm wide, and 300–500 μm thick. Hymenial surface flocculence, white when fresh, white to cream on drying. Sterile margin indistinct, white, and up to 0.5 mm wide.

Hyphal system monomitic, generative hyphae with clamp connections with ampullaceous septa, colorless, thick-walled, frequently branched, interwoven, 2–3.5 μm in diameter; IKI–, CB–, tissues unchanged in KOH.

Cystidia and cystidioles absent; basidia clavate, with four sterigmata and a basal clamp connection, 6.5–10 \times 3.5–5 μm .

Basidiospores ellipsoid, colorless, thin-walled, aculeate, IKI–, CB–, 2.5–3.5 (–4) \times 2–2.5 (–3.5) μm , L = 3.18 μm , W = 2.44 μm , Q = 1.3 (n = 30/1).

***Trechispora bisterigmata* K.Y. Luo & C.L. Zhao, sp. nov.**

MycoBank No: 849464

Figs 4, 5

Holotype. CHINA. Yunnan Province, Yuxi, Xinping County, Mopanshan National Forestry Park, 23°56'N, 101°29'E, altitude 2200 m a.s.l., on the trunk of *Albizia julibrissin*, leg. C.L. Zhao, 20 August 2017, CLZhao 2522 (SWFC).

Etymology. *Bisterigmata* (Lat.): referring to the basidia mainly with two sterigmata.

Description. Basidiomata annual, resupinate, adnate, membranous, without odor or taste when fresh, up to 2.5 cm long, 1.5 cm wide, and 4 mm thick. Hymenial surface odontoid, cream. Sterile margin indistinct, white, rhizomorphic, up to 0.5 mm wide.

Hyphal system monomitic, generative hyphae with clamp connections, colorless, slightly thick-walled, ampullate septa frequently present in subiculum and hymenium with crystals, up to 6 μm wide, branched, interwoven, 2.5–4 μm in diameter; IKI–, CB–, tissues unchanged in KOH.

Cystidia and cystidioles are absent; basidia barrelled, slightly constricted, with two or four sterigmata and a basal clamp connection, 6.5–14.5 \times 3.5–5.5 μm .




Figure 4. Basidiomata of *Trechispora bisterigmata* (holotype) **A** the front of the basidiomata **B** characteristic hymenophore. Scale bars: 1 cm (**A**); 1 mm (**B**).

Basidiospores subglobose to broad ellipsoid, colorless, slightly thick-walled, smooth, IKI-, CB-, (2-) 2.5–4 × 2–3.5 μm, L = 3.03 μm, W = 2.41 μm, Q = 1.23–1.28 (n = 60/2).

Additional specimen examined (paratype). China. Yunnan Province, Yuxi, Xiping County, Mopanshan National Forestry Park, 23°56'N, 101°29'E, altitude 2200 m a.s.l., on the living angiosperm tree, leg. C.L. Zhao, 19 August 2018, CLZhao 7870 (SWFC).




Figure 5. Microscopic structures of *Trechispora bisterigmata* (holotype) **A** basidiospores **B** basidia and basidioles **C** a cross section of basidiomata. Scale bars: 5 μm (**A**); 10 μm (**B**, **C**).

***Trechispora pileata* K.Y. Luo & C.L. Zhao, sp. nov.**

MycoBank No: 849465

Figs 6, 7

Holotype. CHINA. Yunnan Province, Pu'er, Jingdong County, Wuliangshan National Nature Reserve, 24°23'N, 100°45'E, altitude 2350 m a.s.l., on the angiosperm trunk, leg. C.L. Zhao, 6 October 2017, CLZhao 4456 (SWFC).

Etymology. *Pileata* (Lat.): referring to the pileate basidiomata.

Description. Basidiomata annual, with a laterally contracted base, solitary or imbricate. Pileus fan shaped, cortical to corky, up to 1.5 cm long, 1 cm wide, and 2 mm thick, yellowish to yellowish brown, the surface radially striate covered with appressed scales, azonate; the hymenophore surface odontoid, yellowish brown, up to 1 mm long. Context cream, 1 mm thick. Sterile margin indistinct, slightly buff, and 0.5 mm wide.

Hyphal system monomitic, generative hyphae with clamp connections, colorless, thick-walled, frequently branched, interwoven, hyphae in spines 2.5–4 μ m in diameter, IKI–, CB–, tissues unchanged in KOH. Hyphae in context colorless, thin- to thick-walled, unbranched, interwoven, 4.5–6 μ m in diameter, IKI–, CB–, tissues unchanged in KOH.

Cystidia and cystidioles absent; basidia subcylindrical, constricted, with four sterigmata and a basal clamp connection, 5–7 \times 2.5–4 μ m.




Figure 6. Basidiomata of *Trechispora pileata* (holotype) **A, B** the front of the basidiomata **C, D** the back of the basidiomata. Scale bars: 0.5 cm (**A**); 1 mm (**B**); 0.5 cm (**C**); 1 mm (**D**).




Figure 7. Microscopic structures of *Trechispora pileata* (holotype) **A** basidiospores **B** basidia and basidioles **C** hyphae of context of pileus **D** a spine trama of basidiomata. Scale bars: 5 μ m (**A**); 10 μ m (**B**, **C**).

Basidiospores subglobose to broad ellipsoid, colorless, thin-walled, smooth, IKI–, CB–, (2.5–) 2.8–5 (–5.5) × (2.5–) 3–4.7 μm, L = 4 μm, W = 3.56 μm, Q = 1.12 (n = 30/1).

***Trechispora wenshanensis* K.Y. Luo & C.L. Zhao, sp. nov.**

MycoBank No: 849466

Figs 8, 9

Holotype. CHINA. Yunnan Province, Wenshan, Babao Town, Balao battle site, 23°22'N, 104°15'E, altitude 1300 m a.s.l., on the fallen angiosperm branch, leg. C.L. Zhao, 19 January 2019, CLZhao 11649 (SWFC).

Etymology. *Wenshanensis* (Lat.): referring to the locality (Wenshan) of the type specimen.

Description. Basidiomata annual, resupinate, adnate, cottony, easily to separate from substrate, without odor or taste when fresh, up to 5.5 cm long, 4 cm wide, and 200–400 μm thick. Hymenial surface smooth, slightly cream when fresh, cream to buff on drying. Sterile margin indistinct, cream, and 1–2 mm wide.

Hyphal system monomitic, generative hyphae with clamp connections, colorless, thin- to thick-walled, branched, interwoven, 1–2 μm in diameter; IKI–, CB–, tissues unchanged in KOH.

Cystidia and cystidioles are absent; basidia barrelled, with four sterigmata and a basal clamp connection, 7–10 × 3–5 μm.

Basidiospores ellipsoid, colorless, thin-walled, warted, IKI–, CB–, (2–) 2.5–3.7 (–4) × (1.5–) 2–3 μm, L = 3.02 μm, W = 2.37 μm, Q = 1.25–1.30 (n = 90/3).

Additional specimens examined (paratypes). China. Yunnan Province, Wenshan, Funing county, Guying village, 23°42'N, 105°53'E, altitude 1000 m a.s.l., on the fallen angiosperm branch, leg. C.L. Zhao, 20 January 2019, CLZhao 11715; Yunnan Province, Lincang, Lancangjiang Forestry Region, 25°37'N, 97°30'E, altitude 1750 m a.s.l., on the fallen angiosperm branch, leg. C.L. Zhao, 21 July 2022, CLZhao 22940 (SWFC).




Figure 8. Basidiomata of *Trechispora wenshanensis* (holotype) **A** the front of the basidiomata **B** characteristic hymenophore. Scale bars: 1 cm (**A**); 1 mm (**B**).




Figure 9. Microscopic structures of *Trechispora wenshanensis* (holotype) **A** basidiospores **B** basidia and basidioles **C** a cross section of basidiomata. Scale bars: 5 μm (**A**); 10 μm (**B, C**).

Discussion

Many recently described wood-inhabiting fungal taxa have been reported in the subtropics and tropics, including in the genus *Trechispora* (Ordynets et al. 2015; Phookamsak et al. 2019; Xu et al. 2019; Chikowski et al. 2020; Haelewaters et al. 2020; Crous et al. 2021; de Meiras-Otoni et al. 2021; Zhao and Zhao 2021; Liu et al. 2022a, b; Luo and Zhao 2022; Deng et al. 2023; Sommai et al. 2023). The present study reports four new species in *Trechispora*, based on a combination of morphological features and molecular evidence.

Based on ITS+nLSU topology (Fig. 1), four new species were grouped into the genus *Trechispora*, in which *T. albofarinosa* was sister to *T. araneosa* (Höhn. & Litsch.) K.H. Larss., However, morphologically, *T. araneosa* can be delimited from *T. albofarinosa* by its odontoid to poroid hymenial surface and larger ba-

basidiospores ($5\text{--}6.5 \times 4\text{--}5 \mu\text{m}$; Larsson 1995). The second new species *T. bisterigmata* grouped closely with *T. cohaerens* (Schwein.) Jülich & Stalpers and *T. laevispora* Z.B. Liu, Y.D. Wu & Yuan Yuan. However, morphologically, *T. cohaerens* is different from *T. bisterigmata* by its thin-walled hyphal (Jülich and Stalpers 1980); *T. laevispora* can be delimited from *T. bisterigmata* by having the smooth hymenial surface, and thin-walled basidiospores (Liu et al. 2022b). The third species *T. pileata* formed a monophyletic lineage. The species *T. wenshanensis* grouped closely with *T. mellina* (Bres.) K.H. Larss. However, morphologically, *T. mellina* can be delimited from *T. wenshanensis* by having the longer basidia ($15\text{--}20 \times 4.5\text{--}5 \mu\text{m}$) and smooth basidiospores (Chikowski et al. 2020).

Morphologically, *Trechispora albofarinosa* resembles *T. olivacea* K.Y. Luo & C.L. Zhao and *T. yunnanensis* C.L. Zhao by sharing the farinosa basidiomata. However, *T. olivacea* differs from *T. albofarinosa* by olivaceous hymenial surface and thick-walled basidiospores (Luo and Zhao 2022); *T. yunnanensis* can be delimited from *T. albofarinosa* due to its thick-walled, larger basidiospores ($7\text{--}8.5 \times 5\text{--}5.5 \mu\text{m}$; Xu et al. 2019). The new species *T. albofarinosa* is similar to *T. bambusicola* C.L. Zhao, *T. fimbriata* C.L. Zhao, *T. fissurata* C.L. Zhao and *T. murina* K.Y. Luo & C.L. Zhao in its presence of ellipsoid basidiospores. *T. bambusicola* can be delimited from *T. albofarinosa* by odontoid hymenial surface with aculei cylindrical to conical (0.3–0.5 mm long), and thick-walled basidiospores (Zhao and Zhao 2021); *T. fimbriata* can be delimited from *T. albofarinosa* due to its hydroid hymenial surface, and thick-walled basidiospores (Zhao and Zhao 2021); *T. fissurata* is different from *T. albofarinosa* by hydroid hymenial surface and thick-walled, broadly basidiospores ($3.3\text{--}4 \times 2.8\text{--}3.5 \mu\text{m}$; Zhao and Zhao 2021); *T. murina* can be delimited from *T. albofarinosa* due to its grandinoid hymenial surface and thick-walled basidiospores (Luo and Zhao 2022).

Trechispora bisterigmata is similar to *T. fastidiosa* (Pers.) Liberta by sharing the membranous basidiomata. However, *T. fastidiosa* differs from *T. bisterigmata* by smooth hymenial surface and larger basidiospores ($6\text{--}7 \times 4.5\text{--}5.5 \mu\text{m}$; Bernicchia and Gorjón 2010). *T. bisterigmata* resembles *T. bambusicola* C.L. Zhao, *T. canariensis* Ryvarde & Liberta and *T. christiansenii* (Parmasto) Liberta in its monomitic hyphal system and presence of the crystals. However, *T. bambusicola* differs from *T. bisterigmata* by its odontoid hymenial surface and ornamented basidiospores (Zhao and Zhao 2021); *T. canariensis* differs from *T. bisterigmata* due to its larger basidia ($15\text{--}20 \times 5\text{--}6 \mu\text{m}$) and thin-walled, larger basidiospores ($5\text{--}7 \times 3\text{--}3.5 \mu\text{m}$; Ryvarde and Liberta 1978); *T. christiansenii* can be delimited from *T. bisterigmata* by its larger basidia ($15\text{--}20 \times 6\text{--}7 \mu\text{m}$) and larger basidiospores ($5.5\text{--}7 \times 4\text{--}4.5 \mu\text{m}$; Liberta 1966).

Trechispora pileata is similar to *T. byssinella* (Bourdot) Liberta, *T. kavinioides* B. de Vries, *T. silvae-ryae* (J. Erikss. & Ryvarde) K.H. Larss. and *T. subsphaerospora* (Litsch.) Liberta by sharing smooth basidiospores. However, *T. byssinella* differs from *T. pileata* by having narrower ellipsoid basidiospores (Bernicchia and Gorjón 2010); *T. kavinioides* can be delimited from *T. pileata* by its odontoid hymenial surface, and narrower ellipsoid to lacrymiform basidiospores (Bernicchia and Gorjón 2010); *T. silvae-ryae* is different from *T. pileata* by dimitic hyphal system (Bernicchia and Gorjón 2010); *T. subsphaerospora* differs from *T. pileata* by having angular basidiospores (Bernicchia and Gorjón 2010). In addition, the delimitation characteristics of the genus have full resupinate basidiomata, but this new species has the pileate basidiomata with a laterally contracted base. Based on the phylogenetic

analyses, this new species groups with *Trechispora* species, therefore, we propose that the genus *Trechispora* accommodate this new species in the present study.

Trechispora wenshanensis resembles *T. fastidiosa* and *T. laevispora* Z.B. Liu, Y.D. Wu & Yuan Yuan by sharing a smooth hymenial surface. However, *T. fastidiosa* differs from *T. wenshanensis* by larger basidiospores ($6-7 \times 4.5-5.5 \mu\text{m}$; Bernicchia and Gorjón 2010); *T. laevispora* differs from *T. wenshanensis* by fimbriate margin of the basidiomata and smooth basidiospores (Liu et al. 2022b). The new species *T. wenshanensis* is similar to *T. bambusicola* C.L. Zhao, *T. fimbriata* C.L. Zhao, *T. fissurata* C.L. Zhao, *T. murina* K.Y. Luo & C.L. Zhao and *T. yunnanensis* C.L. Zhao due to its ellipsoid basidiospores. However, *T. bambusicola* can be delimited from *T. wenshanensis* by odontoid hymenial surface, and thick-walled basidiospores (Zhao and Zhao 2021); *T. fimbriata* differs from *T. wenshanensis* due to its hydroid hymenial surface, and thick-walled basidiospores (Zhao and Zhao 2021); *T. fissurata* is different from *T. wenshanensis* by hydroid hymenial surface, and thick-walled, broadly basidiospores ($3.3-4 \times 2.8-3.5 \mu\text{m}$; Zhao and Zhao 2021); *T. murina* can be delimited from *T. wenshanensis* due to its grandinoid hymenial surface, and thick-walled basidiospores (Luo and Zhao 2022); *T. yunnanensis* is different from *T. wenshanensis* by farinaceous hymenial surface and thick-walled, larger basidiospores ($7-8.5 \times 5-5.5 \mu\text{m}$; Xu et al. 2019).

Key to 42 accepted species of *Trechispora* in China

- | | | |
|----|--|---------------------------|
| 1 | Basidiomata with clavarioid | 2 |
| – | Basidiomata without clavarioid | 6 |
| 2 | Basidiomata grayish brown to pale purple | 3 |
| – | Basidiomata pure white to pale yellow | 4 |
| 3 | Basidiomata with dense branches and long terminal branches | |
| | <i>T. longiramosa</i> | |
| – | Basidiomata with loose branches | <i>T. laxa</i> |
| 4 | Basidiomata with flattened branches | 5 |
| – | Basidiomata without flattened branches | <i>T. tongdaoensis</i> |
| 5 | Basidiomata branches polychotomous | <i>T. alba</i> |
| – | Basidiomata branches dichotomous | <i>T. khokpasiensis</i> |
| 6 | Basidiomata pileate | <i>T. pileata</i> |
| – | Basidiomata resupinate to effused | 7 |
| 7 | Hymenophore poroid | 8 |
| – | Hymenophore smooth, colliculose, irpicoid, grandinoid, odontoid, hydroid | 13 |
| 8 | Hyphal system dimitic | <i>T. dimitiella</i> |
| – | Hyphal system monomitic | 9 |
| 9 | Subicular hyphae thick-walled | 10 |
| – | Subicular hyphae thin-walled | 11 |
| 10 | Ampullate septa present on subicular hyphae | <i>T. mollusca</i> |
| – | Ampullate septa absent on subicular hyphae | <i>T. suberosa</i> |
| 11 | Crystals in subiculum as numerous rodlets | <i>T. candidissima</i> |
| – | Crystals in subiculum as rhomboidal plates or various shapes | 12 |
| 12 | Sphaerocysts present in cords and the adjacent part of subiculum | |
| | <i>T. hymenocystis</i> | |
| – | Sphaerocysts absent | <i>T. subhymenocystis</i> |

13	Basidiospores smooth	14
–	Basidiospores ornamented.....	16
14	Basidiomata with rhizomorph	<i>T. bisterigmata</i>
–	Basidiomata without rhizomorph.....	15
15	Basidiospores subglobose, angular to turbinate.....	<i>T. confinis</i>
–	Basidiospores ellipsoid.....	<i>T. laevispora</i>
16	Basidiomata < 50 µm thick	17
–	Basidiomata > 50 µm thick.....	19
17	Crystals absent.....	<i>T. gracilis</i>
–	Crystals present.....	18
18	Crystals aggregated, rhomboidal fakes	<i>T. perminispora</i>
–	Crystals butterfly-like, easily broken into irregular shapes	<i>T. subaraneosa</i>
19	Hymenophore smooth	20
–	Hymenophore colliculose, irpicoid, grandinioid, odontoid, hydroid.....	27
20	Basidiospores slightly cyanophilous.....	<i>T. incisa</i>
–	Basidiospores acyanophilous.....	21
21	Basidiospores > 6.5 µm long	<i>T. yunnanensis</i>
–	Basidiospores < 6.5 µm long	22
22	Generative hyphae < 2 µm in diameter.....	<i>T. wenshanensis</i>
–	Generative hyphae > 2 µm in diameter	23
23	Generative hyphae thin-walled.....	24
–	Generative hyphae thick-walled.....	25
24	Hymenophore farinaceous	<i>T. larssonii</i>
–	Hymenophore arachnoid	<i>T. subfarinacea</i>
25	Generative hyphae > 3.5 µm in diameter	<i>T. latehypha</i>
–	Generative hyphae < 3.5 µm in diameter	26
26	Basidiospores ellipsoid, thin-walled.....	<i>T. albofarinosa</i>
–	Basidiospores broadly ellipsoid to globose, thick-walled.....	<i>T. olivacea</i>
27	Hymenial surface colliculose, irpicoid or grandinioid	28
–	Hymenial surface odontoid or hydroid.....	30
28	Generative hyphae thick-walled.....	<i>T. murina</i>
–	Generative hyphae thin-walled.....	29
29	Growth on bamboo.....	<i>T. taiwanensis</i>
–	Growth on other plant	<i>T. crystallina</i>
30	Tramal hyphae thin-walled or slightly thick-walled	31
–	Tramal hyphae distinctly thick-walled.....	35
31	Crystals absent in trama.....	<i>T. tropica</i>
–	Crystals present in trama.....	32
32	Basidiospores subglobose to globose.....	<i>T. odontioidea</i>
–	Basidiospores ellipsoid or broadly ellipsoid	33
33	Tramal hyphae 3–6 µm wide, spines of basidiospores constricted	<i>T. constricta</i>
–	Tramal hyphae 2–4 µm wide, spines of basidiospores not constricted...	34
34	Cystidia present.....	<i>T. chaibuxiensis</i>
–	Cystidia absent.....	<i>T. nivea</i>
35	Hymenophore aculei > 0.4 mm long	36
–	Hymenophore aculei < 0.4 mm long	39
36	Margin smooth	<i>T. fissurata</i>
–	Margin fimbriate	37

37	Basidiomata irpicoid	<i>T. dentata</i>
–	Basidiomata odontoid or hydroid	38
38	Hymenophore aculei sparse, cream to buff-yellow when fresh...	<i>T. fimbriata</i>
–	Hymenophore aculei dense, white when fresh.....	<i>T. fragilis</i>
39	Generative hyphae ampullate septa absent	<i>T. bambusicola</i>
–	Generative hyphae ampullate septa present	40
40	Basidiospores with sharp spines	<i>T. subfissurata</i>
–	Basidiospores without sharp spines.....	41
41	Spines of basidiospores constricted	<i>T. subsinensis</i>
–	Spines of basidiospores not constricted	<i>T. sinensis</i>

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The research was supported by the National Natural Science Foundation of China (Project Nos: 32170004, U2102220), Forestry Innovation Programs of Southwest Forestry University (Project No: LXXK-2023Z07), and the Yunnan Province College Students Innovation and Entrepreneurship Training Program (Project no. s202310677034), and the High-level Talents Program of Yunnan Province (YNQR-QNRC-2018-111).

Author contributions

Conceptualization, C.-L.Z.; methodology, C.-L.Z. and K.-Y.L.; software, C.-L.Z. and K.-Y.L.; validation, C.-L.Z. and K.-Y.L.; formal analysis, C.-L.Z., K.-Y.L. and J.-Q.S.; investigation, C.-L.Z., K.-Y.L. and J.-Q.S.; resources, C.-L.Z.; writing–original draft preparation, C.-L.Z. and K.-Y.L.; writing–review and editing, C.-L.Z. and K.-Y.L.; visualization, C.-L.Z. and K.-Y.L.; supervision, C.-L.Z.; project administration, C.-L.Z.; funding acquisition, C.-L.Z. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Jiang-Qing Su  <https://orcid.org/0009-0008-5480-4502>

Chang-Lin Zhao  <https://orcid.org/0000-0002-8668-1075>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bernicchia A, Gorjón SP (2010) Fungi Europaei 12: Corticiaceae s.l. Edizioni Candusso, Alassio, Italy.
- Bondartsev AS (1953) The Polyporaceae of the European USSR and Caucasia. Israel Program for Scientific Translations, Jerusalem, 1–896.

- Bondartsev AS, Singer R (1941) Zur Systematik der Polyporaceen. *Annales Mycologici* 39(1): 43–65.
- Burdsall Jr HH, Gilbertson RL (1982) New species of Corticiaceae (Basidiomycotina, Aphyllophorales) from Arizona. *Mycotaxon* 15: 333–340.
- Chen JZ, Zhao CL (2020) Morphological and molecular identification of four new resupinate species of *Lyomyces* (Hymenochaetales) from southern China. *MycKeys* 65: 101–118. <https://doi.org/10.3897/mycokeys.65.48660>
- Chikowski RS, Larsson KH, Gibertoni TP (2020) Taxonomic novelties in *Trechispora* (Trechisporales, Basidiomycota) from Brazil. *Mycological Progress* 19(12): 1403–1414. <https://doi.org/10.1007/s11557-020-01635-y>
- Crous PW, Osieck ER, Jurjević Ž (2021) Fungal Planet description sheets: 1284–1382. *Persoonia* 47: 178–374. <https://doi.org/10.3767/persoonia.2021.47.06>
- Dai YC (2011) A revised checklist of corticioid and hydroid fungi in China for 2010. *Mycoscience* 52(1): 69–79. <https://doi.org/10.1007/S10267-010-0068-1>
- Dai YC, Cui BK, Si J, He SH, Hyde KD, Yuan HS, Liu XY, Zhou LW (2015) Dynamics of the worldwide number of fungi with emphasis on fungal diversity in China. *Mycological Progress* 14(8): 62. <https://doi.org/10.1007/s11557-015-1084-5>
- Dai YC, Yang ZL, Cui BK, Wu G, Yuan HS, Zhou LW, et al. (2021) Diversity and systematics of the important macrofungi in Chinese forests. *Junwu Xuebao* 40: 770–805.
- de Meiras-Ottoni A, Larsson KH, Gibertoni TB (2021) Additions to *Trechispora* and the status of *Scytinopogon* (Trechisporales, Basidiomycota). *Mycological Progress* 20(2): 203–222. <https://doi.org/10.1007/s11557-021-01667-y>
- Deng PT, Yan J, Liu XF, He ZM, Lin Y, Lu MX, Zhang P (2023) Three coraloid species of the genus *Trechispora* (Trechisporales, Basidiomycota) in China: Two newly discovered taxa and one reported for the first time. *MycKeys* 99: 153–170. <https://doi.org/10.3897/mycokeys.99.109375>
- Desjardin DE, Perry BA (2015) A new species of *Scytinopogon* from the island of Principe, Republic of Sao Tome and Principe, West Africa. *Mycosphere* 6(4): 433–440. <https://doi.org/10.5943/mycosphere/6/4/5>
- Felsenstein J (1985) Confidence intervals on phylogenetics: An approach using bootstrap. *Evolution; International Journal of Organic Evolution* 39(4): 783–791. <https://doi.org/10.2307/2408678>
- Gilbertson RL, Budington AB (1970) Three new species of wood-rotting fungi in the Corticiaceae. *Mycologia* 62(4): 673–678. <https://doi.org/10.1080/00275514.1970.12019011>
- Guan QX, Zhao CL (2023) Five new species of Schizoporaceae (Basidiomycota, Hymenochaetales) from East Asia. *MycKeys* 96: 25–56. <https://doi.org/10.3897/mycokeys.96.99327>
- Guan QX, Li YF, Zhao CL (2021) Morphological and phylogenetic evidence for recognition of two new species of *Hyphoderma* (Basidiomycota) from southern China, with a key to all Chinese *Hyphoderma*. *MycKeys* 83: 145–160. <https://doi.org/10.3897/mycokeys.83.69909>
- Haelewaters D, Dima B, Abdel-Hafiz All, Abdel-Wahab M, Abul-Ezz SR, Acar I, Aguirre-Acosta E, Aime MC, Aldemir S, Ali M, Ayala-Vásquez O, Bakhit MS, Bashir H, Battistin E, Bendiksen E, Castro-Rivera R, Çolak ÖF, De Kesel A, de la Fuente JI, Dizkırırcı A, Hussain S, Jansen GM, Kaygusuz O, Khalid AN, Khan J, Kiyashko AA, Larsson E, MartínezGonzález CR, Morozova OV, Niazi AR, Noordeloos ME, Pham THG, Popov ES, Psurtseva NV, Schoutteten N, Sher H, Türkekul I, Verbeken A, Ahmad H, Afshan NS, Christe P, Fiaz M, Glazot O, Liu J, Majeed J, Markotter W, Nagy A, Nawaz H,

- Papp V, Péter Á, Pfliegler WP, Qasim T, Riaz M, Sándor AD, Szentiványi T, Voglmayr H, Yousaf N, Krisai-Greilhuber I (2020) Fungal systematics and evolution: FUSE 6. *Sydowia* 72: 231–356.
- Hallenberg N (1978) Wood-fungi (Corticaceae, Coniophoraceae, Lachnocladiaceae, Thelephoraceae) in N. Iran. I. *Iranian Journal of Plant Pathology* 14: 38–87.
- Hallenberg N (1980) New taxa of Corticiaceae from N. Iran (Basidiomycetes). *Mycotaxon* 11(2): 447–475.
- He MQ, Zhu XY, Li TH, Cui BK, Wang K, Bau Tolgor, Zhao RL (2022) Macrofungal classification system and information platform <http://www.nmdc.cn/macrofungi/> is launched. *Mycosystema* 41(6): 899–905.
- Hjortstam K, Larsson KH (1995) Annotated check-list to genera and species of corticioid fungi (Aphylliphorales, Basidiomycotina) with special regards to tropical and subtropical areas. *Windahlia* 21: 1–75.
- Hyde KD (2022) The numbers of fungi. *Fungal Diversity* 114(1): 1. <https://doi.org/10.1007/s13225-022-00507-y>
- Jülich W (1975) Studies in resupinate basidiomycetes - III. *Persoonia* 8(3): 291–305.
- Jülich W (1976) Studies in resupinate basidiomycetes - IV. *Persoonia* 8(4): 431–442.
- Jülich W, Stalpers JA (1980) The resupinate non-poroid Aphylliphorales of the temperate northern hemisphere. *Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen* 74: 1–335.
- Karsten PA (1890) *Fragmenta mycologica XXIX*. *Hedwigia* 29: 147–149.
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kornerup A, Wanscher J (1978) *Metheun Hand Book of Colour*, 3rd edn. Metheun, London, 144–148.
- Larsson KH (1992) The genus *Trechispora* (Corticaceae, Basidiomycetes). Göteborg: Dept. of Systematic Botany.
- Larsson KH (1994) Poroid species in *Trechispora* and the use of calcium oxalate crystals for species identification. *Mycological Research* 98(10): 1153–1172. [https://doi.org/10.1016/S0953-7562\(09\)80200-1](https://doi.org/10.1016/S0953-7562(09)80200-1)
- Larsson KH (1995) Taxonomy of *Trechispora farinacea* and proposed synonyms I. Species with a grandinoid or hydroid hymenophore. *Symbolae Botanicae Upsalienses* 30(3): 101–118.
- Larsson KH (1996) New species and combinations in *Trechispora* (Corticaceae, Basidiomycotina). *Nordic Journal of Botany* 16(1): 83–98. <https://doi.org/10.1111/j.1756-1051.1996.tb00218.x>
- Larsson A (2014) AliView: A fast and lightweight alignment viewer and editor for large data sets. *Bioinformatics (Oxford, England)* 30(22): 3276–3278. <https://doi.org/10.1093/bioinformatics/btu531>
- Larsson KH, Larsson E, Kõljalg U (2004) High phylogenetic diversity among corticioid homobasidiomycetes. *Mycological Research* 108(9): 983–1002. <https://doi.org/10.1017/S0953756204000851>
- Liberta AE (1966) On *Trechispora*. *Taxon* 15(8): 317–319. <https://doi.org/10.2307/1216118>
- Liberta AE (1973) The genus *Trechispora* (Basidiomycetes, Corticaceae). *Canadian Journal of Botany* 51(10): 1871–1892. <https://doi.org/10.1139/b73-240>
- Liu SL, Ma HX, He SH, Dai YC (2019) Four new corticioid species in *Trechisporales* (Basidiomycota) from East Asia and notes on phylogeny of the order. *MycKeys* 48: 97–113. <https://doi.org/10.3897/mycokeys.48.31956>

- Liu SL, He SH, Wang XW, May TW, He G, Chen SL, Zhou LW (2022a) Trechisporales emended with a segregation of Sistotrematrales ord. nov. (Basidiomycota). *Mycosphere* 13(1): 862–954. <https://doi.org/10.5943/mycosphere/13/1/11>
- Liu ZB, Wu YD, Zhao H, Lian YP, Wang YR, Wang CG, Mao W-L, Yuan Y (2022b) Outline, divergence times, and phylogenetic analyses of *Trechisporales* (Agaricomycetes, Basidiomycota). *Frontiers in Microbiology* 13: 818358. <https://doi.org/10.3389/fmicb.2022.818358>
- Liu SL, Wang XW, Li GJ, Deng CY, Rossi W, Leonardi M, Liimatainen K, Kekki T, Niskanen T, Smith ME, Ammirati J, Bojantchev D, Abdel-Wahab MA, Zhang M, Tian E, Lu Y-Z, Zhang J-Y, Ma J, Dutta AK, Acharya K, Du T-Y, Xu J, Kim JS, Lim YW, Gerlach A, Zeng N-K, Han Y-X, Razaghi P, Raza M, Cai L, Calabon MS, Jones EBG, Saha R, Kumar TKA, Krishnapriya K, Thomas A, Kaliyaperumal M, Kezo K, Gunaseelan S, Singh SK, Singh PN, Lagashetti AC, Pawar KS, Jiang S, Zhang C, Zhang H, Qing Y, Bau T, Peng X-C, Wen T-C, Ramirez NA, Niveiro N, Li M-X, Yang ZL, Wu G, Tarafder E, Tennakoon DS, Kuo C-H, da Silva TM, Souza-Motta CM, Bezerra JDP, He G, Ji X-H, Suwannarach N, Kumla J, Lumyong S, Wannathes N, Rana S, Hyde KD, Zhou L-W (2024a) Fungal diversity notes 1717–1817: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 124(1): 1–216. <https://doi.org/10.1007/s13225-023-00529-0>
- Liu SL, Zhao P, Cai L, Shen S, Wei HW, Na Q, Han M, Wei R, Ge Y, Ma H, Karunaratna SC, Tibprommab S, Zhang B, Dai D, Lin L, Fan X-L, Luo Z-L, Shen H-W, Lu L, Lu W, Xu R-F, Tohtirjap A, Wu F, Zhou L-W (2024b) Catalogue of fungi in China 1. New taxa of plant-inhabiting fungi. *Mycology*, 1–59. <https://doi.org/10.1080/21501203.2024.2316066>
- Luo KY, Zhao CL (2022) A molecular systematics and taxonomy research on *Trechispora* (Hydnodontaceae, Trechisporales): Concentrating on three new *Trechispora* species from East Asia. *Journal of Fungi (Basel, Switzerland)* 8(10): 1020. <https://doi.org/10.3390/jof8101020>
- Mardones M, Carranza-Velázquez J, Mata-Hidalgo M, Amador-Fernández X, Urbina H (2023) Taxonomy and phylogeny of the genus *Ganoderma* (Polyporales, Basidiomycota) in Costa Rica. *MycKeys* 100: 5–47. <https://doi.org/10.3897/mycokeys.100.106810>
- Miettinen O, Larsson KH (2006) *Trechispora elongata* species nova from North Europe. *Mycotaxon* 96: 193–198.
- Miller MA, Pfeiffer W, Schwartz T (2012) The CIPRES Science Gateway: Enabling high-impact science for phylogenetics researchers with limited resources. *Association for Computing Machinery* 39: 1–8. <https://doi.org/10.1145/2335755.2335836>
- Nylander JAA (2004) MrModeltest v.2. Program Distributed by the Author; Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.
- Ordynets A, Larsson KH, Langer E (2015) Two new *Trechispora* species from La Réunion Island. *Mycological Progress* 14(11): 113. <https://doi.org/10.1007/s11557-015-1133-0>
- Parmasto E (1968) *Conspectus Systematis Corticiacearum*. Institutum Zoologicum et Botanicum Academiae Scientiarum R.P.S.S, Tartu, 1–97.
- Phookamsak R, Hyde KD, Jeewon R, Bhat DJ, Jones EBG, Maharachchikumbura SSN, Raspe O, Karunaratna SC, Wanasinghe DN, Hongsanan S, Doilom M, Tennakoon DS, Machado AR, Firmino AL, Ghosh A, Karunaratna A, Mešić A, Dutta AK, Thongbai B, Devadatha B, Norphanphoun C, Senwanna C, Wei D, Pem D, Ackah FK, Wang G-N, Jiang H-B, Madrid H, Lee HB, Goonasekara ID, Manawasinghe IS, Kušan I, Cano J, Gené J, Li J, Das K, Acharya K, Raj KNA, Latha KPD, Chethana KWT, He M-Q,

- Dueñas M, Jadan M, Martín MP, Samarakoon MC, Dayarathne MC, Raza M, Park MS, Telleria MT, Chaiwan N, Matočec N, de Silva NI, Pereira OL, Singh PN, Manimohan P, Uniyal P, Shang Q-J, Bhatt RP, Perera RH, Alvarenga RLM, Nogal-Prata S, Singh SK, Vadthananat S, Oh S-Y, Huang S-K, Rana S, Konta S, Paloi S, Jayasiri SC, Jeon SJ, Mehmood T, Gibertoni TB, Nguyen TTT, Singh U, Thiyagaraja V, Sarma VV, Dong W, Yu X-D, Lu Y-Z, Lim YW, Chen Y, Tkalčec Z, Zhang Z-F, Luo Z-L, Daranagama DA, Thambugala KM, Tibpromma S, Camporesi E, Bulgakov TS, Dissanayake AJ, Senanayake IC, Dai DQ, Tang L-Z, Khan S, Zhang H, Promputtha I, Cai L, Chomnunti P, Zhao R-L, Lumyong S, Boonmee S, Wen T-C, Mortimer PE, Xu J (2019) Fungal diversity notes 929–1035, taxonomic and phylogenetic contributions on genera and species of fungi. *Fungal Diversity* 95(1): 1–273. <https://doi.org/10.1007/s13225-019-00421-w>
- Rauschert S (1987) Nomenklatorische Studien bei Höheren Pilzen IV. Nichtblätterpilze (Aphylophorales) mit Ausschluss der Porlinge. *Feddes Repertorium* 98(11–12): 657–664. <https://doi.org/10.1002/fedr.19870981116>
- Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* 98(6): 625–634. [https://doi.org/10.1016/S0953-7562\(09\)80409-7](https://doi.org/10.1016/S0953-7562(09)80409-7)
- Rogers DP (1944) The genera *Trechispora* and *Galzinia*. *Mycologia* 36(1): 70–103. <https://doi.org/10.1080/00275514.1944.12017530>
- Rogers DP, Jackson HS (1943) Notes on the synonymy of some North American Thelephoraceae and other resupinates. *Farlowia* 1(2): 263–328. <https://doi.org/10.5962/p.315982>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Ryvarden L (1975) Studies in the *Aphylophorales* of Africa 3. Three new polypores from Zaire. *Bulletin du Jardin Botanique National de Belgique* 45(1/2): 197–203. <https://doi.org/10.2307/3667599>
- Ryvarden L (2002) A note on the genus *Hydnodon* Banker. Some neotropical wood-inhabiting fungi. *Synop Fungorum* 15: 31–33.
- Ryvarden L, Liberta AE (1978) Contribution to the Aphylophorales of the Canary Islands 4. Two new species of *Trechispora* and *Xenmastella*. *Canadian Journal of Botany* 56(20): 2617–2619. <https://doi.org/10.1139/b78-314>
- Ryvarden L, Stokland L, Larsson KH (2003) A critical checklist of Corticioid and Poroid fungi of Norway. *Synop Fungorum* 17: 1–209.
- Sommai S, Pinruan U, Khamsuntorn P, Lueangjaroenkit P, Somrithipol S, Luangsa-ard J (2023) Three new species of *Trechispora* from Northern and Northeastern Thailand. *Mycological Progress* 22(6): 42. <https://doi.org/10.1007/s11557-023-01886-5>
- Telleria MT, Melo I, Dueñas M, Larsson KH, Paz Martín MP (2013) Molecular analyses confirm *Brevicellicium* in Trechisporales. *IMA Fungus* 4(1): 21–28. <https://doi.org/10.5598/imafungus.2013.04.01.03>
- Trichies G, Schultheis B (2002) *Trechispora antipus* sp. nov., une seconde espèce bisporique du genre *Trechispora* (Basidiomycota, Stereales). *Mycotaxon* 82: 453–458.
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Vries BWL de (1987) Some new corticioid fungi. *Mycotaxon* 28(1): 77–90.

- Vu D, Groenewald M, de Vries M, Gehrman T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92(1): 135–154. <https://doi.org/10.1016/j.simyco.2018.05.001>
- Wang CG, Vlasák J, Dai YC (2021) Phylogeny and diversity of *Bjerkandera* (Polyporales, Basidiomycota), including four new species from South America and Asia. *MycKeys* 79: 149–172. <https://doi.org/10.3897/mycokeys.79.63908>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) *PCR protocols: A Guide to Methods And Applications*. Academic Press, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Phillips AJL, Pereira DS, Dai DQ, Aptroot A, Monteiro JS, Druzhinina IS, Cai F, Fan X, Selbmann L, Coleine C, Castañeda-Ruiz RF, Kukwa M, Flakus A, Fiuza PO, Kirk PM, Kumar KCR, Ieperuma Arachchi IS, Suwannarach N, Tang L-Z, Boekhout T, Tan CS, Jayasinghe RPPK, Thines M (2022) Forecasting the number of species of asexually reproducing fungi (Ascomycota and Basidiomycota). *Fungal Diversity* 114(1): 463–490. <https://doi.org/10.1007/s13225-022-00500-5>
- Wu SH, Chen YP, Wei CL, Floudas D, Dai YC (2017) Two new species of *Phanerochaete* (Basidiomycota) and redescription of *P. robusta*. *Mycological Progress* 17(4): 425–435. <https://doi.org/10.1007/s11557-017-1368-z>
- Wu F, Yuan Y, Chen JJ, Cui BK, Zhou M, Dai YC (2020) *Terrestriporiaceae* fam. nov., a new family of Russulales (Basidiomycota). *Mycosphere* 11(1): 2755–2766. <https://doi.org/10.5943/mycosphere/11/1/21>
- Wu F, Tohtirjap A, Fan LF, Zhou LW, Alvarenga RLM, Gibertoni TB, Dai Y-C (2021) Global diversity and updated phylogeny of *Auricularia* (Auriculariales, Basidiomycota). *Journal of Fungi* (Basel, Switzerland) 7(11): 933. <https://doi.org/10.3390/jof7110933>
- Wu F, Zhou LW, Vlasák J, Dai YC (2022) Global diversity and systematics of Hymenochaetaceae with poroid hymenophore. *Fungal Diversity* 113(1): 1–192. <https://doi.org/10.1007/s13225-021-00496-4>
- Xu TM, Chen YH, Zhao CL (2019) *Trechispora yunnanensis* sp. nov. from China. *Phytotaxa* 424(4): 253–261. <https://doi.org/10.11646/phytotaxa.424.4.5>
- Yuan HS, Dai YC (2012) Wood-inhabiting fungi in southern China. 6. Polypores from Guangxi autonomous region. *Annales Botanici Fennici* 49(5–6): 341–351. <https://doi.org/10.5735/085.049.0605>
- Yuan Y, Bian LS, Wu YD, Chen JJ, Wu F, Liu HG, Zeng GY, Dai YC (2023) Species diversity of pathogenic wood-rotting fungi (Agaricomycetes, Basidiomycota) in China. *Mycology* 14(3): 204–226. <https://doi.org/10.1080/21501203.2023.2238779>
- Yurchenko E, Wu SH (2014) Three new species of *Hyphodontia* with peg-like hyphal aggregations. *Mycological Progress* 13(3): 533–545. <https://doi.org/10.1007/s11557-013-0935-1>
- Zhang QY, Liu HG, Papp V, Zhou M, Dai YC, Yuan Y (2023) New insights into the classification and evolution of *Favolaschia* (Agaricales, Basidiomycota) and its potential distribution, with descriptions of eight new species. *Mycosphere* 14(1): 777–814. <https://doi.org/10.5943/mycosphere/14/1/10>
- Zhao CL, Wu ZQ (2017) *Ceriporiopsis kunmingensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. *Mycological Progress* 16(1): 93–100. <https://doi.org/10.1007/s11557-016-1259-8>

- Zhao W, Zhao CL (2021) The phylogenetic relationship revealed three new wood-inhabiting fungal species from genus *Trechispora*. *Frontiers in Microbiology* 12: 650195. <https://doi.org/10.3389/fmicb.2021.650195>
- Zhao CL, Qu MH, Huang RX, Karunarathna SC (2023a) Multi-gene phylogeny and taxonomy of the wood-rotting fungal genus *Phlebia* sensu lato (*Polyporales*, *Basidiomycota*). *Journal of Fungi* (Basel, Switzerland) 9(3): 1–41. <https://doi.org/10.3390/jof9030320>
- Zhao H, Dai YC, Wu F, Liu XY, Maurice S, Krutovsky KV, Pavlov IN, Lindner DL, Martin FM, Yuan Y (2023b) Insights into the ecological diversification of the Hymenochaetales based on comparative genomics and phylogenomics. *Genome Biology and Evolution* 15(8): 1–15. <https://doi.org/10.1093/gbe/evad136>
- Zhou M, Dai YC, Vlasák J, Liu HG, Yuan Y (2023) Updated systematics of *Trichaptum* s.l. (Hymenochaetales, Basidiomycota). *Mycosphere* 14(1): 815–917. <https://doi.org/10.5943/mycosphere/14/1/11>