

SHORT REPORT

Open Access

Polymorphic microsatellite markers in the brown seaweed *Fucus vesiculosus*

Rui Candeias, Pilar Casado-Amezúa, Gareth A Pearson, Ester A Serrão and Sara Teixeira*

Abstract

Background: *Fucus vesiculosus* is a brown seaweed dominant on temperate rocky shores of the northern hemisphere and, is typically distributed in the mid-upper intertidal zone. It is an external fertilizer that reproduces sexually, providing an excellent model to address conflicting theories related to mating systems and sexual selection. Microsatellite markers have been reported for several *Fucus* species, however the genomic libraries from where these markers have been isolated, have originated from two or more species pooled together (*F. vesiculosus* and *F. serratus* in one library; *F. vesiculosus*, *F. serratus* and *Ascophyllum nodosum* in a second library), or when the genomic DNA originated from only one species it was from *Fucus spiralis*. Although these markers cross-amplify *F. vesiculosus* individuals, the level of polymorphism has been low for relatedness studies.

Findings: The microsatellite markers described here were obtained from an enriched genomic library, followed by 454 pyrosequencing. A total of 9 microsatellite markers were tested across 44 individuals from the North of Portugal. The mean number of alleles across loci was 8.7 and the gene diversity 0.67.

Conclusions: The high variability displayed by these microsatellite loci should be useful for paternity analysis, assessing variance of reproductive success and in estimations of genetic variation within and between populations.

Keywords: Intertidal, Genetic diversity, *Fucus vesiculosus*, Brown macroalgae, Microsatellites

Findings

The seaweed genus *Fucus* (Fucaceae, Phaeophyta) dominates the intertidal biomass of the northern hemisphere's shores, where several species co-occur and in some cases hybridize [1]. *Fucus vesiculosus* is a dominant producer on temperate rocky shores of the northern hemisphere, typically distributed in the mid-upper intertidal zone. It is dioecious, reproduces sexually, and has external fertilization, thus providing an excellent model to address conflicting theories related to mating systems and sexual selection. Variance in reproductive success can lead to sexual selection on reproductive traits and even to sexual conflict between the sexes [2]. Currently, nothing is known about the consequences of external fertilization and the role it plays in the evolution of reproductive traits in this species. Appropriate genetic markers, such as variable microsatellite loci, would allow for more detailed studies related to

this topic as well as how mating systems influence patterns of variation within and between populations.

To date, four studies have reported microsatellite markers for different species of the *Fucus* genera [3-6]. However, none of the previously described markers have been isolated from the species *Fucus vesiculosus* alone, rather previous studies have been mainly focused on hybridization and speciation processes that affect several species of this genus. Therefore, genomic libraries with a mixture of two species or more have been developed to obtain common markers between the target species (*F. vesiculosus* and *F. serratus* in one library [6]; *F. vesiculosus*, *F. serratus* and *Ascophyllum nodosum* in a second library [3]). In the studies where the genomic DNA originated from only one species it has been from *Fucus spiralis* [4] and *Fucus guiryi* ([5], published as *F. spiralis* before this new species was described). Although these markers cross-amplify for *F. vesiculosus* individuals, their level of polymorphism has proven low for relatedness studies (unpublished observations, Teixeira S.).

* Correspondence: steixeira@ualg.pt
Centre of Marine Sciences, CCMAR-CIMAR, University of the Algarve,
Gambelas Campus, 8005-139 Faro, Portugal

Here we report the development and characterization of polymorphic microsatellite loci for *Fucus vesiculosus*. These markers will be useful for relatedness and population genetic studies of this species and hence to assess within and between population genetic diversity.

Whole genomic DNA was isolated from sperm cells of three *F. vesiculosus* males using the CTAB method [7]. We used sperm cells as starting material to avoid cross contamination from the abundant microbial communities commonly found in marine organisms. To isolate the microsatellite sequences, a combination of an SSR-enrichment protocol (standard CT/GT) with 454 pyrosequencing was performed by a commercial company (Ecogenics GmbH, Zürich, Switzerland). The same company designed 48 primers that we tested for polymorphisms across a panel of seven individuals obtained from locations in France (Roscoff [48°42'47.53"N; 4°02'32.95"W]) and Portugal (Viana do Castelo [41°41'32.57"N; 8°50'57.57"W]). An M13-tail (TGAAAACGACGCCAGT) was added at the 5' end of all forward primers to enable fluorescent-dye labelling [8]. We identified nine polymorphic loci, and these results are presented in Table 1.

PCR amplification was performed in 10 µL reaction volumes containing 10 ng of genomic DNA, 1x Qiagen HotStart *Taq* buffer, 200 µM of dNTP's, 0.04 µM of forward primer, 0.16 µM of reverse primer and fluorescently-

labeled M13 primer, and 0.5 U of HotStart *Taq* polymerase (Qiagen). All amplifications were conducted in a Perkin-Elmer GeneAmp7200 (Waltham, MA, USA) with the following program: 15 min at 95°C; 30 cycles composed of 30 s denaturation at 95°C, 45 s at the annealing temperature (Table 1) and 45 s elongation at 72°C, followed by an additional 8 cycles composed of 30 s of denaturation at 95°C, 45 s at 53°C, 45 s elongation at 72°C, and a final extra elongation step for 30 min at 72°C. The forward primer for each set was fluorescently labelled with either FAM, ATT550, or HEX, and PCR products were multiplexed. Fragment analysis was conducted on an ABI 3130XL automated sequencer (Applied Biosystems, Foster City, CA, USA) with Rox350 size standard. Alleles were scored using Peak Scanner 1.0 (Applied Biosystems).

Genetic variation for all markers was tested across 44 individuals sampled in the North of Portugal (Viana do Castelo). The number of alleles per locus (n), expected (H_E), observed heterozygosity (H_O) and heterozygote deficiency (F_{IS} ; Table 1) were calculated using the software GENETIX 4.05 [9]. The polymorphic information content, PIC was calculated using PICcalc [10]. The majority of the optimized markers (9) were highly polymorphic, PIC values ranged from 0.41 (Fves11) to 0.8 (Fves1), the number of alleles found for the 9 loci ranged from 5 (Fves11) to 18 (Fves12) alleles; H_E varied from

Table 1 Characterization of the polymorphic microsatellite loci identified in *Fucus vesiculosus*

Locus	Repeat motif	Primer sequence (5'-3')	T _a (°C)	Allele size range (bp)	n	H _E	H _O	F _{IS}	PIC	f NL
Fves1	(ATC) ₉	F: GCAGGATCGACAACCATACC R: GATCCACCGACATGCTTTG	58	103-181	12	0.83	0.23	0.7	0.8	0.56
Fves2	(AC) ₁₂	F: TGTATCAGCATACGGAAAAGC R: ATCAGGCATGTTGTCGTCGG	56	100-112	7	0.76	0.63	0.16	0.71	0.06
Fves5	(TG) ₁₁	F: CGATGGGGGATGAGTGATTG R: TGACATGCGGAGACAAAACG	56	124-180	9	0.65	0.75	-0.15	0.6	0
Fves7	(AC) ₁₁	F: ACTTGATGTCCAGAATGAATGGG R: AGGTGTGCGTGCGATTATTAC	56	191-213	6	0.68	0.25	0.6	0.62	0.25
Fves8	(CA) ₁₆	F: TGGGGAAGTACACACACACG R: TCGTTGCGGTAACACTTTGC	56	137-169	6	0.50	0.4	0.23	0.45	0.09
Fves10	(TG) ₁₁	F: AGGCGCTTGAAAATGACCTG R: CCCGTGATTACTTTGGTATCGTC	56	103-169	6	0.47	0.2	0.6	0.44	0.4
Fves11	(AC) ₁₅	F: TGTGTGACCTTCTCTGTG R: AGTATTGGCGTGTGCAACTG	56	133-147	5	0.45	0.43	0.04	0.41	0
Fves12	(GT) ₁₇	F: AACTTTACCCGTTTTCCACG R: CTCGCGGTAGAACACCTCTC	58	107-159	18	0.82	0.85	-0.05	0.78	0
Fves14	(AG) ₁₇	F: AATGACGGGGCCGGAATG R: GGCCGTCGATCTTACTCAC	56	112-138	9	0.82	0.7	0.19	0.78	0.1

The following details are reported: name, motif, primer sequence and annealing temperature (Ta°C). Also descriptive statistics are presented, based on one population analysed, number of alleles n , expected heterozygosity, H_E , observed heterozygosity, H_O , F_{IS} according to Hardy-Weinberg equilibrium, polymorphic information content, PIC, and null allele frequencies, f_{NL} . Bold numbers indicate significant values at 5% level after q-value correction.

0.45 (Fves11) to 0.83 (Fves1) and H_O from 0.2 (Fves10) to 0.85 (Fves12). Significant heterozygote deficiency was observed for 5 markers (Fves1, 7, 8, 10, and 14), as shown by the high and significant F_{IS} values (Table 1). Null alleles might occur at these loci, as confirmed by the determination of the frequency of null alleles in the dataset using ML-NULLFREQ [11] (Table 1). We tested for linkage disequilibrium between all pairs of loci using the software GENETIX 4.05 [9]. No linkage disequilibrium was found after the correction for multiple tests using the false discovery rate (FDR) approach [12] in QVALUE [13].

Previous studies found for the population of Viana do Castelo and across loci, gene diversities of 0.59 [5] and 0.58 [14], while the mean number of alleles found were 4.3 [5] and 5.6 [14]. The overall gene diversity found for the same population across loci, with our newly developed set of markers was of 0.67 and the mean number of alleles 8.7. The higher variability displayed by these microsatellite loci may be useful for paternity analysis and population genetic studies of this species.

Availability of supporting data

The microsatellite sequences are available through the National Center for Biotechnology Information (see <http://www.ncbi.nlm.nih.gov/>). The accession numbers on the repository are the following: GenBank accession no. KP765803 through KP765811.

Competing interests

The authors declare they have no competing interests.

Authors' contributions

ST, GAP and EAS were responsible for the design and implementation of the study, supervision of the work and processing interpretation of the results. RC, PCA and ST participated in data analysis and microsatellite marker validation, RC and ST drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgments

This study was supported by the Portuguese Science Foundation (FCT) through project PTCD/MAR/104477/2008 (ST), EXCL/AAG-GLO/0661/2012, and postdoctoral fellowship SFRH/BPD/39097/2007 (ST).

Received: 6 January 2015 Accepted: 24 February 2015

Published online: 08 March 2015

References

- Lüning K. Seaweeds: Their Environment, Biogeography and Ecophysiology. New York: John Wiley & Sons; 1990.
- Arnold SJ. Is there a unifying concept of sexual selection that applies to both plants and animals? *Am Nat*. 1994;144:S1–S12.
- Engel CR, Brawley S, Edwards KJ, Serrão E. Isolation and cross-species amplification of microsatellite loci from the furoid seaweeds *Fucus vesiculosus*, *F. serratus*, and *Ascophyllum nodosum* (Heterokontophyta, Fucales). *Mol Ecol Notes*. 2003;3:180–2.
- Wallace AR, Klein AS, Mathieson AC. Determining the affinities of salt marsh furoids using microsatellite markers: evidence of hybridization and introgression between two species of *Fucus* (Phaeophyceae) in a Maine estuary. *J Phycol*. 2004;40:1013–27.
- Perrin C, Daguin C, Van De Vliet M, Engel CR, Pearson GA, Serrão EA. Implications of mating system for genetic diversity of sister algal species:

- Fucus spiralis* and *Fucus vesiculosus* (Heterokontophyta, Phaeophyceae). *Eur J Phycol*. 2007;42:219–30.
- Coyer JA, Hoarau G, Beszerti B, Pearson G, Olsen JL. Expressed sequence tag derived polymorphic SSR markers for *Fucus serratus* and amplification in other species of *Fucus*. *Mol Ecol Res*. 2009;9:168–70.
- Doyle JJ, Doyle JL. Isolation of plant DNA from fresh tissue. *Focus*. 1990;12:13–5.
- Schuelke M. An economic method for the fluorescent labelling of PCR fragments. *Nat Biotech*. 2000;18:233–4.
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F. GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Montpellier (France): Laboratoire Génome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II; 1996.
- Nagy S, Poczai P, Cernák I, Gorji AM, Hegedűs G, Taller J. PICcalc: An online program to calculate polymorphic information content for molecular genetic studies. *Biochem Genet*. 2012;50:670–2.
- Kalinowski ST, Taper ML. Maximum likelihood estimation of the frequency of null alleles at microsatellite loci. *Conserv Genet*. 2006;7:991–5.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc*. 1995;57:289–300.
- Storey JD. A direct approach to false discovery rates. *J R Stat Soc*. 2002;64:479–98.
- Nicastro K, Zardi G, Teixeira S, Neiva J, Serrão E, Pearson GA. Shift happens: trailing edge contraction associated with recent warming trends threatens a distinct genetic lineage in the marine macroalga *Fucus vesiculosus*. *BMC Biol*. 2013;11:6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

