



Therapy of clinical stage IIA and IIB seminoma: a systematic review

Julia Heinzlbecker¹ · Stefanie Schmidt² · Julia Lackner² · Jonas Busch³ · Carsten Bokemeyer⁴ · Johannes Classen⁵ · Annette Dieing⁶ · Oliver Hakenberg⁷ · Susanne Krege⁸ · Alexandros Papachristofilou⁹ · David Pfister¹⁰ · Christian Ruf¹¹ · Hans Schmelz¹² · Heinz Schmidberger¹³ · Rainer Souchon¹⁴ · Christian Winter¹⁵ · Friedemann Zengerling¹⁶ · Sabine Kliesch¹⁷ · Peter Albers¹⁸ · Christoph Oing^{4,19}

Received: 19 July 2021 / Accepted: 25 October 2021 / Published online: 15 November 2021
© The Author(s) 2021

Abstract

Purpose The optimal treatment for clinical stage (CS) IIA/IIB seminomas is still controversial. We evaluated current treatment options.

Methods A systematic review was performed. Only randomized clinical trials and comparative studies published from January 2010 until February 2021 were included. Search items included: seminoma, CS IIA, CS IIB and therapy. Outcome parameters were relapse rate (RR), relapse-free (RFS), overall and cancer-specific survival (OS, CSS). Additionally, acute and long-term side effects including secondary malignancies (SMs) were analyzed.

Results Seven comparative studies (one prospective and six retrospective) were identified with a total of 5049 patients (CS IIA: 2840, CS IIB: 2209). The applied treatment modalities were radiotherapy (RT) ($n = 3049$; CS IIA: 1888, CS IIB: 1006, unknown: 155) and chemotherapy (CT) or no RT ($n = 2000$; CS IIA: 797, CS IIB: 1074, unknown: 129). In CS IIA, RRs ranged from 0% to 4.8% for RT and 0% for CT. Concerning CS IIB RRs of 9.5%–21.1% for RT and of 0%–14.2% for CT have been reported. 5-year OS ranged from 90 to 100%. Only two studies reported on treatment-related toxicities.

Conclusions RT and CT are the most commonly applied treatments in CS IIA/B seminoma. In CS IIA seminomas, RRs after RT and CT are similar. However, in CS IIB, CT seems to be more effective. Survival rates of CS IIA/B seminomas are excellent. Consequently, long-term toxicities and SMs are important survivorship issues. Alternative treatment approaches, e.g., retroperitoneal lymph node dissection (RPLND) or dose-reduced sequential CT/RT are currently under prospective investigation.

Keywords Testicular cancer · Seminoma · CS IIA/B · Systematic review · Treatment · Toxicity

Introduction

Testicular cancer (TC) is the most common solid malignancy in young men [1, 2]. Accounting for only 7% of all TC patients, clinical stage (CS) IIA/B seminoma, defined as disease spread to the retroperitoneal lymph nodes of up to 2 cm (CS IIA) or of more than 2 cm to up to 5 cm (CS IIB) in maximum diameter, is a rare disease [3, 4]. Established national and international TC guidelines recommend radiotherapy (RT) or chemotherapy (CT) for the treatment of CS IIA/B seminoma patients [5, 6]. However, high-level evidence to define the optimal treatment remains elusive

[5, 6]. As survival rates of CS IIA/B patients are excellent and approach 100%, treatment-related acute and long-term toxicities are of considerable interest for shared treatment decision-making [7].

Here, we summarize the available evidence regarding the different treatment modalities of CS IIA/B seminoma including associated acute and long-term toxicities.

Methods

This work is based on a systematic literature search that was conducted for the development of the first German clinical practice guideline [4, 5, 8].

We performed a systematic literature review in accordance with the preferred reporting items for systematic

✉ Julia Heinzlbecker
julia.heinzlbecker@uks.eu

Extended author information available on the last page of the article

reviews and meta-analyses (PRISMA) guidelines [9]. The search was conducted in Medline (via Ovid) and the Cochrane Central Register of Controlled Trials (search period January 2010–February 2021). The detailed methods, including the search strategy, can be found in Suppl. 1.

Randomized controlled trials and comparative studies were considered if they included patients with IIA or IIB seminoma, who received additional treatment after surgical treatment of the primary tumor. For studies to be included, outcome data had to be displayed separately for these patient groups. Non-seminomatous disease as well as patients at other disease stages were excluded. Considered treatments were radiotherapy (RT), chemotherapy (CT), retroperitoneal lymph node dissection (RPLND) or combinations of these treatments. Our endpoints of interest were relapse rate (RR), overall (OS), cancer-specific survival (CSS), as well as adverse events [acute and late toxicities including treatment-related secondary malignancies (SM)].

Results

2830 records were identified through database searching or other sources. Seven comparative studies met our inclusion criteria. These were published between 2011 and 2017 [10–16] (see Table 1). There were one prospective and six retrospective trials including a total of 5049 assessable CS IIA/B patients (CS IIA: 2840; CS IIB: 2209). 3049 patients received RT (CS IIA: 1888; CS IIB: 1006; unknown: 155) and 2000 patients received CT or “no RT” (CS IIA: 797; CS IIB: 1074; unknown: 129) (unfortunately “no RT” was not further defined in the respective study [13], see below). Figure 1 displays the study identification process.

Only three studies reported on the specific RT template (see Table 1a) [10–12]. The applied total cumulative radiation dose was reported in five studies ranging from 25.5 to 36 Gy [10–12, 14, 15]. Concerning CT, only three studies reported on CT regimens with EP (etoposide/cisplatin), EP or BEP (bleomycin/etoposide/cisplatin) and EP, HOP (ifosfamide, vincristine, cisplatin), PVB (cisplatin, vinblastine, bleomycin) or VAB (vinblastine, cyclophosphamide, dactinomycin, bleomycin) being used [10–12]. Of note, two studies used “no RT” as comparator to patients receiving RT, however did not further define “no RT” [13, 16]. Across studies, the median follow-up ranged between 3.8 and 10 years. Four studies were classified as level of evidence (LoE) 2b, three were downgraded to LoE 4, mainly because of missing information on outcome definition, assessment and statistical variance and no control for possible confounders. Details of the risk of bias assessment are given in Suppl. 2.

Outcome for relapse and survival

In terms of outcomes, only two studies assessed RRs (see Table 1a) [10, 12]. One did not include CS IIA patients for CT, the other did not include CS IIB patients for RT. The overall RRs for RT were 10.2% and 24.2%. For CS IIA patients, a RR following RT of 0%–10.3% and for CS IIB patients of 28.6% was reported. For CT, both studies showed an overall RR of 0%. Tandstad et al. showed a statistically significant difference in the RRs of RT and CT (10.3% vs. 0%, $p=0.01$) [10].

In terms of RFS, only two studies reported data (see Table 1a) [10, 11]. Kollmannsberger et al. studied CS II including CS IIC patients, with a 5-year RFS of 91.7% for RT and 95.5% for CT [11]. Tandstad et al. showed a 5-year RFS for CS IIA patients of 88.7% for RT and of 100% for CT (CS IIA/B) [10].

All studies reported on OS and three studies additionally on CSS (see Table 1a). Domont et al. showed for CS IIA/B patients a 5-year OS of 82% following RT versus 88% following CT [12]. Kollmannsberger et al. described a 5-year OS rate of 92.3% following RT and 90.7% following CT [11]. Patel et al. reported 5–15-year OS rates for CS IIA patients of 95.8%–94.1% and for CS IIB patients of 95.9%–87.2% [16]. The other four studies described treatment specific OS data for CS IIA and CS IIB patients [10, 13–15]. In CS IIA patients, the 5-year OS rates following RT were 96%–100% and for CT or “no RT” 88%–100%. Ahmed et al. and Paly et al. reported a significantly reduced OS after CT or “no RT” compared to RT in CS IIA seminoma patients ($p=0.008$, $p<0.01$) [13, 15]. In CS IIB patients, 5-year OS rates following RT were 95.2%–100% and following CT or “no RT” 90%–100%. Ahmed et al. reported reduced OS rates for CS IIB seminoma patients for “no RT” compared to RT-treated patients ($p=0.03$) [13].

Regarding 5-year CSS, Tandstad et al. reported a 5-year CSS of 100% for CS IIA and CS IIB patients treated with RT as CT, respectively [10]. Patel et al. reported a 5–15-year CSS for CS IIA patients of 98.1%, respectively, and for CS IIB patients of 98%–95.9% [16]. Ahmed et al. reported a 5-year CSS of CS IIA patients of 97% after RT versus 96% for “no RT” and for CS IIB patients of 98% versus 98%, respectively [13].

Additionally, Glaser et al., Paly et al. and Patel et al. reported significantly reduced hazards ratios (HR) for death for RT in CS IIA patients, with HRs of 0.22–0.34 ($p=0.005$, $p=0.014$) with regards to RT [14, 16] and a significantly increased HR for death of 13.3 ($p<0.01$) with regards to CT [15] (see Table 1a).

Table 1 Overview of the comparative studies on CS IIA/B seminoma patients included in the systematic review (a) patient and treatment characteristics, outcome parameters for relapse/survival and quality parameters (b) toxicities

Author	Year (study period)	Type of study	n, overall (CS IIA/B)	Treatment modality	RT characteristics		CT regimen [courses]	Median FU in years (range; IQR)	Outcome		Time of reference	LoE	
					n, CS IIA	n, CS IIB			Field	Dose (Gy)			RR (%)
Tandstad	2011 (2000–2006)	Prospective cohort study SWENOTECA Swedish, Norwegian	1384 ^b (102)	RT (29; IIA); CT (73; IIA: 6; IIB: 67)	Paraaortic+iliac	27	EP [4]/BEP in large tumours [NA]	^b 5.2 (0.1–10.2)	RT: 10.3 (IIA); CT: 0 (IIA; IIB)	IIA: RT: 88.7; CT: 100	IIA, IIB: RT, CT: 100	5	2b
				RT (19); CT (24)	Paraaortic+iliac	35	EP [4]/BEP (3)	3.8 (0.2–9.2)	IIA-C: NA	IIA-C: RT: 92.3; CT: 91.7; CT: 95.5 ^b	IIA-C: RT: NA	IIA-C: 97	5
Domont ^c	(1980–2001)	Retrospective cohort study Patients at Institut Gustave Roussy, Villejuif, France	67 ^b (36)	RT (33; IIA: 5; IIB: 28); CT (3; IIB)	Paraaortic+iliac; M/SC	36	EP [4]/HOP [NA]/PVB [NA]/VAB [NA]	^b 9.4 (1–21)	RT: 24.2; IIA: 0; IIB: 28.6; CT: 0; IIA: n.a., IIB: 0	IIA-C: 97	IIA-C: OS after relapse: RT: 82; CT: 88	5	2b
				RT (19); CT (24)	Paraaortic+iliac	35	EP [4]/BEP (3)	3.8 (0.2–9.2)	IIA-C: NA	IIA-C: RT: 92.3; CT: 91.7; CT: 95.5 ^b	IIA-C: RT: NA	IIA-C: 97	5

Table 1 (continued)

Author	Year (study period)	Type of study	n, overall (CS IIA/IB)		Treatment modality	RT characteristics		CT regimens [courses]	Median FU in years (range; IQR)	Outcome		Time of reference	LoE
			n, CS IIA	n, CS IIB		Field	Dose (Gy)			RR (%)	RFS (%)		
Ahmed	2015 (1988–2003)	Retrospective cohort study US SEER database	241		RT (136); no RT (105)	NA	NA	NA	10 (NA)	NA	NA	5, 10, 15	4 ^a
			145	96						IIA: RT: 96, 96, 96; no RT: 88, 77, 77 IIB: RT: 98, 98, 88; no RT: 90, 86, 86 p=0.008	IIA: RT: 97, 97, 97; no RT: 96, RT: 96, 92, 92 IIB: RT: 98, 98, 98; no RT: 98, 96, 88; no RT: 90, 96, 96 p=0.003		
Glaser	2016 (1998–2012)	Retrospective cohort study US National Cancer Data Base	2,437 ^b (1772)	812	RT (1,192; IIA: 750, IIB: 442); CT (580; IIA: 210, IIB: 370)	NA	IIA: 30.9; IIB: 35.5	NA	^b 5.4 (2.8–8.8)	NA	NA	5	2b
			960							IIA: RT: 99; CT: 93 IIB: RT: 95.2; CT: 92.4 HR: IIA: 0.22; IIB: 0.22 p=0.005			
Paly	2016 (1998–2012)	Retrospective cohort study US National Cancer Data Base	1,885	805	RT (1,160; IIA: 780, IIB: 380); CT (725; IIA: 300, IIB: 425)	NA	IIA: 25.5; IIB: 25.5; MBD: 10	NA	4.2 (IQR: 5.6)	NA	NA	5	4 ^a
			1,080							IIA: RT: 99.4; CT: 91.2, p<0.01 IIB: RT: 96.1; CT: 92.8 HR: IIA: 13.3 IIB: 13.3 p<0.01			

Table 1 (continued)

Author	Year (study period)	Type of study	n, overall (CS IIA/B)	n, CS IIA	n, CS IIB	Treatment modality		RT characteristics		CT regimens [courses]	Median FU in years (range; IQR)	Outcome			Time of reference	LoE
						RT (480; IIA: 324, IIB: 156); no RT (490; IIA: 281, IIB: 209)	Field	Dose (Gy)	RR (%)			RFS (%)	OS (%)	CSS (%)		
Patel ^e	2017 (1988–2013)	Retrospective cohort study US SEER database	16,463 ^b (970)	605	365	RT (480; IIA: 324, IIB: 156); no RT (490; IIA: 281, IIB: 209)	NA	NA	NA	NA	7.4 (IQR: 3.3–12.1)	NA	NA	IIA: 95.8, 94.1, 94.1 IIB: 95.9, 98.1 91.7, 87.2 IIB: 98, 97.2, 95.9 vs. no RT: IIA: OS: 0.34 (0.14–0.8), p=0.014	5, 10, 15	4 ^a
(b)																
Author	Side effects															
	Acute toxicity: n (%)															
	Late toxicity: n (%)															
	Secondary malignancy: n (%)															
Domont ^f	^b RT: Grade 1/2: Nausea: 34 (92%) Diarrhea: 19 (51%) Grade 3/4: Nausea: 3 (8%) CT: Grade 3/4: Neutropenia: 7 (22%) Febrile neutropenia: 5 (15%) ^b RT in field: 3 (8%): Colorectal cancer: 1 (3%) Duodenal cancer: 1 (3%) Medullary thyroid cancer: 1 (3%) CT: 2 (7%): Colorectal cancer: 1 (3%) Esophageal cancer: 1 (3%) Otolotoxicity: 2 (5%) Neuropathy: 2 (5%) Fertility Disorders: NA Pulmonary fibrosis: NA Nephrotoxicity: 0 (0%) CT: 8 (27%): Otolotoxicity: 2 (7%) Neuropathy: 1 (3%) Fertility disorders: 2 (7%) Pulmonary fibrosis: 1 (3%) Grade 2 nephrotoxicity: 2 (7%)															
	Risk of SM															
	NA															

Table 1 (continued)

Author	Side effects	Risk of SM
Acute toxicity: n (%)		Secondary malignancy: n (%)
Late toxicity: n (%)		
Patel ^c	NA	<p>^bRT/no RT: overall nontesticular: 82 (0.9%)/30 (0.4%)</p> <p>Oral/pharynx: 0 (0%)/1 (0.01%)</p> <p>Gastrointestinal: 21 (0.2%)/10 (0.1%) Respiratory: 25 (0.3%)/8 (0.1%)</p> <p>Urinary: 7 (0.1%)/0 (0%)</p> <p>Gastrointestinal: 21 (0.2%)/10 (0.1%) Respiratory: 25 (0.3%)/8 (0.1%)</p> <p>Lymphoma/leukemia: 15 (0.2%)/8 (0.1%)</p> <p>Soft tissue/heart: 4 (0.04%)/2 (0.03%)</p> <p>Skin: 4 (0.04%)/0 (0%)</p> <p>Eye/orbit: 1 (0.01%)/0 (0%)</p> <p>Nervous system: 2 (0.02%)/1 (0.01%)</p> <p>Endocrine: 1 (0.01%)/0 (0%)</p> <p>Myeloma: 1 (0.01%)/0</p> <p>Miscellaneous: 1 (0.01%)/0 (0%)</p>


^bEP bleomycin/etoposide/cisplatin, CI confidence interval, CS clinical stage, CSS cancer-specific survival, CT chemotherapy, EP etoposide/cisplatin, FU follow-up, Gy Gray, HOP ifosfamide, vincristine, cisplatin, HR hazard ratio, IQR interquartile range, LoE level of evidence, M mediastinal, MDB median boost dose, n number of patients, NA not announced, n.a. not applicable, OS overall survival, PVB cisplatin, vinblastine, bleomycin, REF reference group, RFS relapse-free survival, RoB risk of bias, RR relapse rate, RT radiotherapy, SC supraclavicular, SEER Surveillance, Epidemiology, and End Results Program, SM secondary malignancy, SWENOTECA Swedish and Norwegian Testicular Cancer Group, US United States, VAB vinblastine, cyclophosphamide, dactinomycin, bleomycin

^aLevel of evidence was downgraded due to study limitations from the risk of bias assessment

^bOther stages as CS IIA/B are included

^cStudies that also reported toxicities

Fig. 1 Flow chart of the study inclusion process for the systematic review according to PRISMA [9]



Outcome for acute, late toxicities and secondary malignancies

Only two studies assessed toxicities (see Table 1b) [12, 16]. As reported by Domont et al., 92% of the RT patients experienced grade 1/2 nausea and 51% grade 1/2 diarrhea [12]. Grade 3/4 nausea occurred in 8% of RT patients. Grade 3/4 neutropenia occurred in 22% of CT patients and febrile neutropenia in 15%. Late toxicities were reported in 11% of RT patients and in 27% of CT patients with neuropathy, fertility disorders, pulmonary fibrosis and nephrotoxicity described for CT patients. In terms of SM, 8% of RT and 7% of CT patients developed a SM, which was colorectal, duodenal and medullary thyroid cancer in the RT and colorectal and oesophageal cancer in the CT groups.

Only Patel et al. reported on SMs, comparing patients who received RT to those who did not receive RT (median follow-up: 8.3 years) [16]. For the overall population (CS I–CS IIC), they reported a higher rate of SMs in patients who received RT (0.9% vs. 0.4%) with a significantly higher relative risk (RIR) of SMs in the RT group (RIR: 1.84).

When adjusting for stage and age this result remained statistically significant for CS IA patients, only (see Table 1b).

Discussion

RT and CT represent the most frequently applied and guideline-endorsed treatment options for CS IIA/B seminoma achieving excellent long-term outcomes [6, 8].

Concerning efficacy, in CS IIA RRs of 0–4.8% following RT at 30 Gy and of 0% following CT have been reported. For CS IIB patients, RRs after RT ranged from 9.5% to 21.1% at a 36 Gy cumulative dose and from 0 to 14.2% after CT. Thus, RT and CT seem equally effective in CS IIA, whereas CT tends to be more effective in CS IIB patients. Long-term outcomes are excellent with 5-year OS rates for CS IIA/B seminoma patients ranging from 90 to 100%, irrespective of the applied treatment. Nevertheless, survival data of CS IIA patients warrant attention as few studies, however with considerable risk of bias, found a reduced OS when CT was applied. The reasons for this so far remain unknown.

RRs and RFS

Of the studies included for this systematic review, only the prospective study of Tandstad et al. included both treatments, RT and CT for CS IIA patients [10]. They reported a significantly higher RR following RT (11.3% vs. 0% following CT), but the applied cumulative radiation dose was only 27 Gy. Today, a radiation dose of 30 Gy is the guideline-endorsed standard dose for CS IIA patients based on several single-arm prospective trials reporting RRs of 0–4.8% [17–19].

Concerning CS IIB patients, only Domont et al. compared RT and CT outcomes [12]. The RR following RT was 28.6% versus 0% in the CT cohort, but only three patients of the study received CT. The applied cumulative radiation dose was 36 Gy in line with the nowadays recommended standard dose for CS IIB patients [17–19]. Reported RRs of single-arm prospective trials assessing RT efficacy both lower (9.5–11.1%), and higher RRs of up to 21.1% with a radiation dose of 36 Gy [17, 18, 20] and thus, the true RR following RT remains elusive but seems unanimously higher than following CT based on prospective study data.

Various other studies not included in this systematic review also addressed the treatment of CS IIA/B seminomas with different CT regimens and various cumulative RT doses [17–41] (see Suppl. 3). Today's guideline-endorsed CT standard is three cycles of BEP [6, 8]. However, the CT regimens of the studies included in the systematic review were highly heterogeneous and do not represent the current standard of care. There exist several prospective single-arm trials evaluating different CT regimens [31, 34, 35]. Arranz Arija et al. evaluated up to four courses of EP in CS IIA/B seminoma patients and found a 3-year RFS of 91% [31]. Garcia-del-Muro et al. evaluated four courses of EP or three courses of BEP with RRs of 0% for CS IIA and 11.1% for CS IIB patients and a 5-year progression-free survival (PFS) of 100% and 87%, respectively [35]. Consequently, four cycles of EP may be considered in patients where Bleomycin has to be omitted, which is also the established approach for advanced metastatic germ cell tumors with a favorable IGC-CCG risk profile [6, 8]. The optimal number of cycles of BEP remains controversial since retrospective analyses of either four or two cycles of BEP also reported a 0% RR [37, 38]. Three to four cycles of Carboplatin AUC7, is not equally effective compared to cisplatin-based combination CT with higher RRs reported in a prospective study by Krege et al. [34].

Taken together, RT and CT appear to be equally effective in terms of RRs in CS IIA. However, CT seems to have lower RRs in CS IIB patients than RT. This may be due to the given heterogeneity of CS IIB comprising small lesions with a diameter of just 2 cm to bulky nodal metastases with a diameter of up to 5 cm.

OS and CSS

All studies included in this systematic review assessed OS as an endpoint and it is obvious that cure rates are unanimously high together with an extremely low rate of cancer-associated deaths [9–15]. Three studies reported on CSS [10, 13, 16]. However, only Tandstad et al. and Ahmed et al. discriminated for CS IIA or IIB and for RT or CT/"no RT". 5-year CSS ranged from 97 to 100% in CS IIA patients treated with RT versus 96%–100% for CS IIA patients treated with CT/no RT [10, 13]. For CS IIB patients, CSS ranged from 98 to 100% for RT and from 98 to 100% for CT/no RT [10, 13]

Ahmed et al. found a statistically significant difference regarding OS for patients who did not receive RT [13]. Nevertheless, these results must be interpreted cautiously as there are certain limitations to the study, the most important being that "other approaches than RT" are not defined and that data on doses in the SEER database are missing (see Suppl. 2). The OS results of two other US National Cancer Database-based projects on CS IIA seminoma patients merit attention [14, 15]. Glaser et al. reported a significantly lower HR for death (HR: 0.22) for CS IIA patients treated with RT, with a 5-year OS of 99% (RT) compared to 93% (CT) [14]. This advantage even persisted on a propensity-adjusted multivariate analysis. However, neither CT regimens nor numbers of applied courses of CT were documented reflecting a substantial source of bias. In line with these results, Paly et al. reported a significantly higher HR of death (HR: 13.3) for CS IIB patients treated with CT, with a 5-year OS of 91.2% (CT) and 99% (RT) [15]. However, the study of Paly et al. also has major limitations due to the limited availability of only 80% of the data on radiation dose and lacking information about CT details. Nevertheless, both results are corroborated by a study of Patel et al. that assessed the SEER database and found a reduced HR for death (HR: 0.34) in CS IIA patients who received RT [16]. Again, also in this study, outcomes were not defined and there was lacking information on treatment details due to the limited data collection within SEER.

With regards to long-term survival, the risk of SM is of interest. Groot et al. evaluated cause-specific mortality among testicular cancer (TC) patients in a large multicentre cohort study with a 17.6-year median follow-up [42]. They found RT and CT to be associated with increased SM-related mortality. However, only the receipt of CT was additionally associated with increased standard mortality ratios (SMR) for ischemic heart disease (IHD) and respiratory disease. Though, when considering RT doses at > 26–32 Gy and > 32–36 Gy or platinum dosage < 400 mg/m², the current standard doses applied in CS IIA/B seminoma patients, both RT and CT were associated with elevated SM mortality (HR: 1.98; 2.55), but CT was no longer associated with IHD or respiratory disease mortality [42].

Alternative treatment approaches

There exist several studies on the combination of sequential CT and RT and on surgery by RPLND for the treatment of CS IIA/B seminoma [21–25, 36, 41]. However, none of them met the prespecified inclusion criteria of our performed systematic review. Patterson et al. evaluated the value of adding Carboplatin to RT on the basis of the SEER database [22]. They reported a RR for CS IIA patients of 7% (CT + RT) versus 13% (RT) and for CS IIB patients of 5% (CT + RT) versus 26% (RT), respectively. 5-year OS rates for CS IIA patients were 91.7% (CT + RT) versus 95.3% (RT) and for CS IIB patients 100% (CT + RT) versus 93.9% (RT) [22]. Horwich et al. reported on a pilot study of 51 CS IIA/B patients who were treated with the combination of a single cycle of Carboplatin CT followed by RT with reduced dose and extent of the radiation fields [21]. After a median follow-up of 4.6 years they reported no relapses. The prospective single-arm phase II SAKK 01/10 trial evaluated 3-year PFS in CS IIA/B seminoma patients treated with one cycle of Carboplatin AUC7 followed by reduced-field involved-node RT with 30 Gy in CS IIA and 36 Gy in CS IIB patients [43–45]. After a median follow-up of 4.5 years, 3-year PFS was 93.7% (CI 85.5–98.5%) in CS IIA and 95.2% (CI 85.2–96.4%) in CS IIB patients, respectively. There were no cancer-specific deaths [45]. Additionally, the single-arm phase II SAKK 01/18 (NCT03937843) trial evaluates 3-year PFS in seminoma patients who receive one cycle of Carboplatin AUC7 (CS IIA) or EP (CS IIB) followed by reduced-field involved-node RT with 24 Gy (CS IIA) or 30 Gy (CS IIB).

Surgical approaches by RPLND for CS IIA/B seminomas have also been reported on. Warszawski et al. reported a RR for CS IIA patients of 0% (RPLND) versus 10% (RT) and for CS IIB patients of 67% (RPLND) versus 20% (RT) [25]. Subsequently, several prospective trials assessed the efficacy of RPLND in CS IIA/B seminomas in an attempt to avoid CT- and RT-related, potentially life-threatening late toxicities [46, 47]. An interim analysis of the prospective phase II PRIMETEST trial, which included 22 CS IIA/B seminoma patients with a mean tumor size of 2.6 cm, displayed an overall 23% RR at a mean follow-up of 24 months. All patients remained relapse free after salvage treatment so far [41]. Preliminary results of the prospective phase II SEMS trial have also been published recently. Here, 55 CS IIA/B seminoma patients underwent RPLND. After a median follow-up of 24 months an overall RR of 18% was reported. The 2-year RFS rate was 85% and the 2-year OS rate 100% [48]. Both trials concluded that RPLND can be a therapeutic option as first-line treatment in early-stage metastatic seminoma. A caveat may be the so far limited follow-up period, as seminomas may relapse later than just two years from surgery.

Toxicities

Only two studies of our systematic review report on treatment-associated toxicities and thus, evidence on treatment-related sequelae in CS IIA/B seminoma patients is scarce [12, 16]. Nausea and diarrhea are typical immediate side effects of RT [12, 17, 19], while nausea/vomiting, alopecia and transient bone marrow suppression are commonly related to platinum-based CT [31, 34, 35, 37] (see Suppl. 4). Treatment-related adverse events of RPLND comprise complications such as lymphocele, chylous ascites or ileus [23, 25, 48] (see Suppl. 4).

Concerning *long-term side effects* other than SMs, gastroesophageal reflux disease, sexual dysfunction, hypogonadism, diabetes and coronary artery disease have been described for RT [22, 33, 39, 49–51] and ototoxicity, neurotoxicity, nephrotoxicity and fertility disorders for CT [7, 12] (see Suppl. 4). A typical long-term side effect of RPLND is retrograde ejaculation; however, in modified template resections, as recommended for CSIIA/B seminoma patients, this risk can be minimized [48].

The most important, hence potentially life-threatening treatment-related long-term toxicities are *cardiovascular disease (CVD)* and *SMs* [7]. In terms of CT, an association with cardiovascular disease during the first year after CT has been described [51, 52]. Lauritsen et al. found that the risk of CVD thereafter decreased to normal levels. However, after ten years from diagnosis, there was an increasing risk for myocardial infarction and cardiovascular fatalities [51]. Nevertheless, it remains questionable if an association also exists for lower cumulative doses of CT as applied in CS IIA/B seminoma patients as Groot et al. found CT at dosages < 400 mg/m² no longer to be associated with ischemic heart disease [42]. Additionally, Haugnes et al. found an increased odds for having intermediate/high risk cardiovascular morbidity or predicted mortality only in patients treated at platinum dosages > 850 mg/m² [53].

SMs have been described as long-term adverse events for both CT and RT among TC patients [7]. Travis et al. report a cumulative risk of developing a secondary solid cancer for a 20-year-old seminoma patient of almost 50% by the age of 75 [54]. There is growing discussion on the amount of harm by the different therapies applied to TC patients accounting for solid SMs, whereas the risk of leukemia is typically ascribed to etoposide-based combination CT [7].

The impact of either CT or RT on SM risk remains controversial. Both studies of the systematic review reporting on toxicities, reported on SM. While Domont et al. found comparable amounts of SMs in a very limited number of patients, Patel et al. reported a significantly higher risk of SMs following RT. However, after adjusting for stage, this remained significant for CS 1A seminoma patients only [12, 16]. Other reports on SM in CS IIA/B seminomas are mostly

from single-arm RT studies reporting on in-field solid SMs mostly affecting the gastrointestinal (GI) and urinary tract organs (see Suppl. 4) [19, 22, 29, 33, 36, 38–40, 49]. Reports from CT or RPLND studies on the risk of SM specifically in CS IIA/B seminomas are scarce (see Suppl. 4) [22, 34, 36].

It, thus, turns out that both RT and CT are associated with increased standard incidence ratios (SIR) of various cancers, whereas there seems to be no risk of solid SMs after surgery alone, except for soft tissue sarcoma [55]. The receipt of CT was associated with an increased risk for solid SMs (HR: 2.4), colorectal (HR: 3.9) and non-colorectal GI SMs (HR: 5.0) [55]. However, at dosages of < 400 mg/m² platinum, there was no statistically significant difference [55]. On the contrary, infradiaphragmatic RT at a cumulative dose of > 26–32 Gy and > 32–36 Gy was still associated with a higher solid SM risk (HR 2.4; 2.5), especially within the infradiaphragmatic regions (HR 3.4; 3.9) [55]. These results are in line with earlier reports of Travis et al. [54]. Among mainly RT-treated seminoma patients, increased SIR for SMs were restricted to infradiaphragmatic sites, with small intestine (SIR: 8.9), pancreatic (SIR: 4.4) and urinary bladder cancers (SIR: 3.4) [54].

Regarding SM mortality, Groot et al. found RT to be associated with a higher SM-related mortality especially due to colorectal, pancreatic and urologic malignancies, whereas CT was associated with a higher SM-related mortality from lung, colorectal, non-colorectal GI malignancies and leukemia [42]. Even at the currently applied doses in CS IIA/B seminomas, RT and CT both were associated with higher SM mortality [42].

There are several limitations to our systematic review. First, most of the studies were retrospective analyses, including data partly derived from the 1980s onwards. Second, level of evidence was mainly downgraded because missing information on outcome definition and assessment, missing information on statistical variance, and/or no control for possible confounders. Third, case numbers were mostly small and not solely including CS IIA/IIB patients. Fourth, treatment data, which are essential for the evaluation of treatment effects and harms, were rarely reported. As a result, essential treatment details were lacking, which did hamper data interpretation and therefore, we decided to not conduct a meta-analysis. Fifth, treatment regimens and diagnostic techniques have changed over time and do not reflect the various approaches assessed in the included studies anymore. This adds to the heterogeneity of identified data.

Conclusion

Long-term outcomes of CSIIA/B seminoma patients are excellent and RT and CT represent equally effective treatment options especially for CS IIA patients, whereas in CS

IIB seminomas, CT may be beneficial in terms of reduced RRs. Surgical and sequential approaches of CT and RT hold the promise to change the treatment landscape in the future. Shared decision-making with patients should be informed by the so far equal efficacy of RT and CT based on relatively low evidence. Particular attention should be paid on treatment-related side effects.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00345-021-03873-5>.

Author contributions JH: protocol/project development, data collection or management, literature search, data analysis, manuscript writing/editing. SS: protocol/project development, data collection or management, literature search, methodological assessment, data analysis, manuscript writing/editing. JL: protocol/project development, data collection or management, literature search, methodological assessment, data analysis, manuscript writing/editing. JB: protocol/project development, manuscript editing. CB: protocol/project development, data analysis, manuscript editing. JC: protocol/project development, manuscript editing. AD: protocol/project development. OH: protocol/project development, manuscript editing. SK: protocol/project development, manuscript editing. AP: protocol/project development, data analysis, manuscript editing. DP: protocol/project development, manuscript editing. CR: protocol/project development. HS: protocol/project development, manuscript editing. HS: protocol/project development, manuscript editing. RS: protocol/project development, data analysis, manuscript editing. CW: protocol/project development. FZ: protocol/project development, manuscript editing. SK: protocol/project development, data analysis, manuscript editing. PA: protocol/project development, data analysis, manuscript editing. CO: protocol/project development, data analysis, manuscript editing. This research did not involve Human Participants and/or Animals. Thus, written informed consent was not necessary.

Funding Open Access funding enabled and organized by Projekt DEAL. This work is based on a clinical guideline program, which was funded by the German Cancer Aid Foundation (DKH) (Reference No. 70112789).

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.



References

- Gurney JK, Florio AA, Znaor A et al (2019) International trends in the incidence of testicular cancer: lessons from 35 years and 41 countries. *Eur Urol* 76:615–623. <https://doi.org/10.1016/j.eururo.2019.07.002>
- Nigam M, Aschebrook-Kilfoy B, Shikanov S, Eggener S (2015) Increasing incidence of testicular cancer in the United States and Europe between 1992 and 2009. *World J Urol* 33:623–631. <https://doi.org/10.1007/s00345-014-1361-y>
- Rothermundt C, Thurneysen C, Cathomas R et al (2018) Baseline characteristics and patterns of care in testicular cancer patients: first data from the Swiss Austrian German Testicular Cancer Cohort Study (SAG TCCS). *Swiss Med Wkly* 148:w14640. <https://doi.org/10.4414/smw.2018.14640>
- Kliesch S, Schmidt S, Wilborn D et al (2021) Management of germ cell tumours of the testis in adult patients. German clinical practice guideline part i: epidemiology, classification, diagnosis, prognosis, fertility preservation, and treatment recommendations for localized stages. *Urol Int*. <https://doi.org/10.1159/000510407>
- Kliesch S, Schmidt S, Wilborn D et al (2021) Management of germ cell tumours of the testes in adult patients: german clinical practice guideline, part II—recommendations for the treatment of advanced, recurrent, and refractory disease and extragonadal and sex cord/stromal tumours and for the management of follow-up, toxicity, quality of life, palliative care, and supportive therapy. *Urol Int*. <https://doi.org/10.1159/000511245>
- Laguna MP, Albers P, Algaba F et al (2020) Guidelines on testicular cancer. EAU guidelines. EAU Guidelines Office, Arnhem, The Netherlands
- Haugnes HS, Bosl GJ, Boer H et al (2012) Long-term and late effects of germ cell testicular cancer treatment and implications for follow-up. *J Clin Oncol* 30:3752–3763. <https://doi.org/10.1200/JCO.2012.43.4431>
- (2020) S3-Leitlinie Diagnostik, Therapie und Nachsorge der Testikulären Keimzelltumoren, Langversion 1.1. Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF)
- Moher D, Liberati A, Tetzlaff J et al (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 6:e1000097. <https://doi.org/10.1371/journal.pmed.1000097>
- Tandstad T, Smaaland R, Solberg A et al (2011) Management of seminomatous testicular cancer: a binational prospective population-based study from the Swedish norwegian testicular cancer study group. *J Clin Oncol* 29:719–725. <https://doi.org/10.1200/JCO.2010.30.1044>
- Kollmannsberger C, Tyldesley S, Moore C et al (2011) Evolution in management of testicular seminoma: population-based outcomes with selective utilization of active therapies. *Ann Oncol* 22:808–814. <https://doi.org/10.1093/annonc/mdq466>
- Domont J, Massard C, Patrikidou A et al (2013) A risk-adapted strategy of radiotherapy or cisplatin-based chemotherapy in stage II seminoma. *Urol Oncol* 31:697–705. <https://doi.org/10.1016/j.urolonc.2011.04.004>
- Ahmed KA, Wilder RB (2015) Stage IIA and IIB testicular seminoma treated postorchietomy with radiation therapy versus other approaches: a population-based analysis of 241 patients. *Int Braz J Urol* 41:78–85. <https://doi.org/10.1590/S1677-5538.IBJU.2015.01.11>
- Glaser SM, Vargo JA, Balasubramani GK, Beriwal S (2016) Stage II testicular seminoma: patterns of care and survival by treatment strategy. *Clin Oncol (R Coll Radiol)* 28:513–521. <https://doi.org/10.1016/j.clon.2016.02.008>
- Paly JJ, Lin CC, Gray PJ et al (2016) Management and outcomes of clinical stage IIA/B seminoma: results from the national cancer data base 1998–2012. *Pract Radiat Oncol* 6:e249–e258. <https://doi.org/10.1016/j.prro.2016.05.002>
- Patel HD, Srivastava A, Alam R et al (2017) Radiotherapy for stage I and II testicular seminomas: secondary malignancies and survival. *Urol Oncol* 35:606.e1–606.e7. <https://doi.org/10.1016/j.urolonc.2017.06.051>
- Classen J, Schmidberger H, Meisner C et al (2003) Radiotherapy for stages IIA/B testicular seminoma: final report of a prospective multicenter clinical trial. *J Clin Oncol* 21:1101–1106. <https://doi.org/10.1200/JCO.2003.06.065>
- Schmidberger H, Bamberg M, Meisner C et al (1997) Radiotherapy in stage IIA and IIB testicular seminoma with reduced portals: a prospective multicenter study. *Int J Radiat Oncol Biol Phys* 39:321–326. [https://doi.org/10.1016/s0360-3016\(97\)00155-7](https://doi.org/10.1016/s0360-3016(97)00155-7)
- Bamberg M, Schmidberger H, Meisner C et al (1999) Radiotherapy for stages I and IIA/B testicular seminoma. *Int J Cancer* 83:823–827. [https://doi.org/10.1002/\(sici\)1097-0215\(19991210\)83:6%3c823::aid-ijc22%3e3.0.co;2-v](https://doi.org/10.1002/(sici)1097-0215(19991210)83:6%3c823::aid-ijc22%3e3.0.co;2-v)
- Weissbach L, Bussar-Maatz R, Löhrs U et al (1999) Prognostic factors in seminomas with special respect to HCG: results of a prospective multicenter study. *Seminoma Study Group Eur Urol* 36:601–608. <https://doi.org/10.1159/000020055>
- Horwich A, Dearnaley DP, Sohaib A et al (2013) Neoadjuvant carboplatin before radiotherapy in stage IIA and IIB seminoma. *Ann Oncol* 24:2104–2107. <https://doi.org/10.1093/annonc/mdt148>
- Patterson H, Norman AR, Mitra SS et al (2001) Combination carboplatin and radiotherapy in the management of stage II testicular seminoma: comparison with radiotherapy treatment alone. *Radiother Oncol* 59:5–11. [https://doi.org/10.1016/s0167-8140\(00\)00240-1](https://doi.org/10.1016/s0167-8140(00)00240-1)
- Hu B, Shah S, Shojaei S, Daneshmand S (2015) Retroperitoneal lymph node dissection as first-line treatment of node-positive seminoma. *Clin Genitourin Cancer* 13:e265–e269. <https://doi.org/10.1016/j.clgc.2015.01.002>
- Mezvrishvili Z, Managadze L (2006) Retroperitoneal lymph node dissection for high-risk stage I and stage IIA seminoma. *Int Urol Nephrol* 38:615–619. <https://doi.org/10.1007/s11255-005-4793-x>
- Warszawski N, Schmücking M (1997) Relapses in early-stage testicular seminoma: radiation therapy versus retroperitoneal lymphadenectomy. *Scand J Urol Nephrol* 31:355–359. <https://doi.org/10.3109/00365599709030619>
- Chung PWM, Gospodarowicz MK, Panzarella T et al (2004) Stage II testicular seminoma: patterns of recurrence and outcome of treatment. *Eur Urol* 45:754–759. <https://doi.org/10.1016/j.eururo.2004.01.020> (discussion 759–760)
- Sridharan S, Chung PWM, Jewett MA et al (2013) Use of radiotherapy for seminoma patients with low-volume infradiaphragmatic nodal disease. *JCO* 31:335–335. https://doi.org/10.1200/jco.2013.31.6_suppl.335
- Warde P, Gospodarowicz M, Panzarella T et al (1998) Management of stage II seminoma. *J Clin Oncol* 16:290–294. <https://doi.org/10.1200/JCO.1998.16.1.290>
- Bauman GS, Venkatesan VM, Ago CT et al (1998) Postoperative radiotherapy for stage I/II seminoma: results for 212 patients. *Int J Radiat Oncol Biol Phys* 42:313–317. [https://doi.org/10.1016/s0360-3016\(98\)00227-2](https://doi.org/10.1016/s0360-3016(98)00227-2)
- Zagars GK, Pollack A (2001) Radiotherapy for stage II testicular seminoma. *Int J Radiat Oncol Biol Phys* 51:643–649. [https://doi.org/10.1016/s0360-3016\(01\)01701-1](https://doi.org/10.1016/s0360-3016(01)01701-1)
- Arranz Arijia JA, García del Muro X, Gumà J et al (2001) E400P in advanced seminoma of good prognosis according to the international germ cell cancer collaborative group (IGCCCC)

- classification: the Spanish germ cell cancer group experience. *Ann Oncol* 12:487–491. <https://doi.org/10.1023/a:1011127715764>
32. Porcaro AB, Antonioli SZ, Maffei N et al (2002) Management of testicular seminoma advanced disease. Report on 14 cases and review of the literature. *Arch Ital Urol Androl* 74:81–85
 33. Garcia-Serra AM, Zlotecki RA, Morris CG, Amdur RJ (2005) Long-term results of radiotherapy for early-stage testicular seminoma. *Am J Clin Oncol* 28:119–124. <https://doi.org/10.1097/O1.coc.0000143843.08585.ce>
 34. Krege S, Boergermann C, Baschek R et al (2006) Single agent carboplatin for CS IIA/B testicular seminoma. A phase II study of the German testicular cancer study group (GTCSG). *Ann Oncol* 17:276–280. <https://doi.org/10.1093/annonc/mdj039>
 35. Garcia-del-Muro X, Maroto P, Gumà J et al (2008) Chemotherapy as an alternative to radiotherapy in the treatment of stage IIA and IIB testicular seminoma: a Spanish germ cell cancer group study. *J Clin Oncol* 26:5416–5421. <https://doi.org/10.1200/JCO.2007.15.9103>
 36. Detti B, Livi L, Scoccianti S et al (2009) Management of stage II testicular seminoma over a period of 40 years. *Urol Oncol* 27:534–538. <https://doi.org/10.1016/j.urolonc.2008.07.034>
 37. Giannis M, Aristotelis B, Vassiliki K et al (2009) Cisplatin-based chemotherapy for advanced seminoma: report of 52 cases treated in two institutions. *J Cancer Res Clin Oncol* 135:1495–1500. <https://doi.org/10.1007/s00432-009-0596-2>
 38. Pichler R, Leonhartsberger N, Stöhr B et al (2012) Two cycles of cisplatin-based chemotherapy for low-volume stage II seminoma: results of a retrospective, single-center case series. *Chemotherapy* 58:405–410. <https://doi.org/10.1159/000345701>
 39. Hallemeier CL, Pisansky TM, Davis BJ, Choo R (2013) Long-term outcomes of radiotherapy for stage II testicular seminoma—the Mayo Clinic experience. *Urol Oncol* 31:1832–1838. <https://doi.org/10.1016/j.urolonc.2012.03.010>
 40. Stein ME, Zidan J, Charas T, Ben-Yosef R (2014) Radiotherapy for stage IIA seminoma: the Northern Israel oncology center experience, 1971–2010. *Rep Pract Oncol Radiother* 19:281–286. <https://doi.org/10.1016/j.rpor.2014.02.004>
 41. Albers P, Hiester A, Grosse Siemer R, Lusch A (2019) The PRIMETEST trial: Interim analysis of a phase II trial for primary retroperitoneal lymph node dissection (RPLND) in stage II A/B seminoma patients without adjuvant treatment. *J Clin Oncol* 37:507–507. https://doi.org/10.1200/JCO.2019.37.7_suppl.507
 42. Groot HJ, van Leeuwen FE, Lubberts S et al (2020) Platinum exposure and cause-specific mortality among patients with testicular cancer. *Cancer* 126:628–639. <https://doi.org/10.1002/cncr.32538>
 43. Papachristofilou A, Bedke J, Hayoz S et al (2020) Treatment compliance and early toxicity in SAKK 01/10: single-dose carboplatin and involved-node radiotherapy for treatment of stage IIA/B seminoma. *J Clin Oncol* 38:405–405. https://doi.org/10.1200/JCO.2020.38.6_suppl.405
 44. Swiss Group for Clinical Cancer Research (2020) Carboplatin chemotherapy and involved node radiotherapy in stage IIA/B seminoma (NCT: 01593241). clinicaltrials.gov
 45. Papachristofilou A, Bedke J, Hayoz S et al (2021) Single-dose carboplatin followed by involved-node radiotherapy as curative treatment for seminoma stage IIA/B: efficacy results from the international multicenter phase II trial SAKK 01/10. *Ann Oncol* 32:1283–1346. <https://doi.org/10.1016/annonc/annonc741>
 46. Heinrich-Heine University, Duesseldorf (2016) Phase II single-arm trial to evaluate progression free survival with primary retroperitoneal lymph-node dissection (pRPLND) only in patients with seminomatous testicular germ cell tumors with clinical stage IIA/B (PRIMETEST) (NCT: 02797626). clinicaltrials.gov
 47. University of Southern California (2021) Surgery in early metastatic seminoma (SEMS): phase II trial of retroperitoneal lymph node dissection as first-line treatment for testicular seminoma with isolated retroperitoneal disease (1–3cm) (NCT: 02537548). clinicaltrials.gov
 48. Daneshmand S, Cary C, Masterson TA et al (2021) SEMS trial: Result of a prospective, multi-institutional phase II clinical trial of surgery in early metastatic seminoma. *J Clin Oncol* 39:375–375. https://doi.org/10.1200/JCO.2021.39.6_suppl.375
 49. Hallemeier CL, Davis BJ, Pisansky TM, Choo R (2014) Late gastrointestinal morbidity in patients with stage I-II testicular seminoma treated with radiotherapy. *Urol Oncol* 32:496–500. <https://doi.org/10.1016/j.urolonc.2013.10.002>
 50. Groot HJ, Gietema JA, Aleman BMP et al (2018) Risk of diabetes after para-aortic radiation for testicular cancer. *Br J Cancer* 119:901–907. <https://doi.org/10.1038/s41416-018-0248-x>
 51. Lauritsen J, Hansen MK, Bandak M et al (2020) Cardiovascular risk factors and disease after male germ cell cancer. *J Clin Oncol* 38:584–592. <https://doi.org/10.1200/JCO.19.01180>
 52. Fung C, Dinh P, Ardeshir-Rouhani-Fard S et al (2018) Toxicities associated with cisplatin-based chemotherapy and radiotherapy in long-term testicular cancer survivors. *Adv Urol* 2018:8671832. <https://doi.org/10.1155/2018/8671832>
 53. Haugnes HS, Aass N, Fosså SD et al (2008) Predicted cardiovascular mortality and reported cardiovascular morbidity in testicular cancer survivors. *J Cancer Surviv* 2:128–137. <https://doi.org/10.1007/s11764-008-0054-1>
 54. Travis LB, Fosså SD, Schonfeld SJ et al (2005) Second cancers among 40,576 testicular cancer patients: focus on long-term survivors. *J Natl Cancer Inst* 97:1354–1365. <https://doi.org/10.1093/jnci/dji278>
 55. Groot HJ, Lubberts S, de Wit R et al (2018) Risk of solid cancer after treatment of testicular germ cell cancer in the platinum era. *J Clin Oncol* 36:2504–2513. <https://doi.org/10.1200/JCO.2017.77.4174>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Julia Heinzlbecker¹  · Stefanie Schmidt² · Julia Lackner² · Jonas Busch³ · Carsten Bokemeyer⁴ · Johannes Classen⁵ · Annette Dieing⁶ · Oliver Hakenberg⁷ · Susanne Krege⁸ · Alexandros Papachristofilou⁹ · David Pfister¹⁰ · Christian Ruf¹¹ · Hans Schmelz¹² · Heinz Schmidberger¹³ · Rainer Souchon¹⁴ · Christian Winter¹⁵ · Friedemann Zengerling¹⁶ · Sabine Kliesch¹⁷ · Peter Albers¹⁸ · Christoph Oing^{4,19} 

¹ Department of Urology and Paediatric Urology, Saarland University Medical Centre and Saarland University Faculty of Medicine, Homburg, Saar, Germany

² UroEvidence@Deutsche Gesellschaft Für Urologie, Berlin, Germany

³ Department of Urology, Vivantes Clinics am Urban, Berlin, Germany

⁴ II. Medical Clinic and Polyclinic, University Hospital Hamburg-Eppendorf, Hamburg, Germany

⁵ Department of Radiotherapy, Radiological Oncology and Palliative Medicine, St. Vincentius-Kliniken, Karlsruhe, Germany

⁶ Clinic for Internal Medicine-Hematology and Oncology, Vivantes Clinics Am Urban, Berlin, Germany

⁷ Urological Clinic and Polyclinic, University Hospital Rostock, Rostock, Germany

⁸ KEM, Protestant Hospital Essen-Mitte, Clinic for Urology, Pediatric Urology and Urological Oncology, Essen, Germany

⁹ Clinic of Radiotherapy and Radiation Oncology, University Hospital Basel, Basel, Switzerland

¹⁰ Department of Urology, University Hospital Cologne, Cologne, Germany

¹¹ Department of Urology, Bundeswehrkrankenhaus (German Federal Armed Forces Hospital), Ulm, Germany

¹² Department of Urology, Bundeswehrkrankenhaus (German Federal Armed Forces Hospital), Koblenz, Germany

¹³ Clinic and Polyclinic for Radiooncology and Radiotherapy, University Hospital Mainz, Mainz, Germany

¹⁴ Department for Radiooncology, University Hospital Tübingen, Tübingen, Germany

¹⁵ Urologie Neandertal (Regional Joint Practice), Erkrath, Germany

¹⁶ Department of Urology, University Hospital Ulm, Ulm, Germany

¹⁷ Centre of Reproductive Medicine and Andrology, Department of Clinical and Surgical Andrology, University Hospital Münster, Münster, Germany

¹⁸ Department of Urology, University Hospital Düsseldorf, Düsseldorf, Germany

¹⁹ Mildred Scheel Cancer Career Centre HaTriCs4, University Cancer Centre Hamburg, University Medical Centre Hamburg-Eppendorf, Hamburg, Germany