

Electronic Thesis and Dissertation Repository

6-25-2013 12:00 AM

Assessing and Analyzing Bat Activity with Acoustic Monitoring: Challenges and Interpretations

Amanda M. Adams
The University of Western Ontario

Supervisor
Brock Fenton
The University of Western Ontario

Graduate Program in Biology

A thesis submitted in partial fulfillment of the requirements for the degree in Doctor of
Philosophy

© Amanda M. Adams 2013

Follow this and additional works at: <https://ir.lib.uwo.ca/etd>



Part of the [Other Ecology and Evolutionary Biology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Adams, Amanda M., "Assessing and Analyzing Bat Activity with Acoustic Monitoring: Challenges and Interpretations" (2013). *Electronic Thesis and Dissertation Repository*. 1333.
<https://ir.lib.uwo.ca/etd/1333>

This Dissertation/Thesis is brought to you for free and open access by Scholarship@Western. It has been accepted for inclusion in Electronic Thesis and Dissertation Repository by an authorized administrator of Scholarship@Western. For more information, please contact wlsadmin@uwo.ca.

ASSESSING AND ANALYZING BAT ACTIVITY WITH ACOUSTIC
MONITORING: CHALLENGES AND INTERPRETATIONS

(Thesis format: Integrated Article)

by

Amanda Murphy Adams

Graduate Program in Biology,
Collaborative Program in Environment & Sustainability

A thesis submitted in partial fulfillment
of the requirements for the degree of
Doctor of Philosophy

The School of Graduate and Postdoctoral Studies
The University of Western Ontario
London, Ontario, Canada

© Amanda M. Adams 2013

Abstract

Acoustic monitoring is a powerful technique for learning about the ecology of bats, but understanding sources of variation in the data collected is important for unbiased interpretation. The objectives of this dissertation were to investigate sources of variation in acoustic monitoring and make recommendations for acoustic survey design and analysis. I addressed this goal in three ways: i) variation resulting from differences in bat detectors, ii) methods for objective identification of peak activity, and iii) the use of stationary transects to address within-site spatial variation.

First, I compared variation of detection of echolocation calls among commonly available bat detectors and found significant differences in distance and angle of detection. Consequently, this source of variation should be taken into account when comparing datasets obtained with different systems. Furthermore, choice of detector should be taken into account when designing new studies.

Second, I investigated two statistical methods for identifying peaks in activity, percentile thresholds and space-time scan statistic (SaTScan). Acoustic monitoring provides a relative measure of activity levels and is rarely evaluated based on objective criteria, so describing bat activity as “high” or “low” is useful only in context of the studies in question. Percentile thresholds allow for peaks to be identified relative to a larger distribution of activity levels. SaTScan identifies peaks in space and time that are significantly higher than the background expectation of the dataset. Both methods are valuable tools for replicable and objective identification of peak activity that can be applied at various temporal and spatial scales.

Third, I examine how within-site spatial variation can impact estimates of bat activity. I used a stationary transect of bat detectors to i) assess variation in patterns of activity at each detector, ii) test whether spatial or temporal factors were more important for explaining variation in activity, iii) explore what sampling effort in space and time is required for species-specific activity levels. The picture of activity differs significantly within a site depending on detector placement so it is important to use multiple detectors simultaneously to collect accurate estimates of activity.

Keywords

Bats, activity levels, echolocation, spatio-temporal variation, acoustic transect, passive acoustic monitoring, bat detectors, peak activity, percentile thresholds, SaTScan.

Co-Authorship Statement

A version of Chapter 2 was published in the *Methods in Ecology and Evolution* with Meredith Jantzen, Rachel Hamilton, and Brock Fenton as co-authors. Ms. Jantzen and Ms. Hamilton contributed to study design, data collection, and data interpretation. Dr. Fenton provided access to equipment. All authors contributed editorial comments on the manuscript.

A version of Chapter 3 has been submitted for publication to *Acta Chiropterologica* with Liam McGuire, Lauren Hooton, and Brock Fenton as co-authors. Dr. McGuire provided logistical support in Saskatchewan and contributed to interpretation of the data. Ms. Hooton helped create the discriminant function analysis for species identification of acoustic data. Dr. Fenton provided access to the acoustic equipment and funding. All authors contributed editorial comments on the manuscript.

A version of Chapter 4 has been submitted for publication to the *Journal of Applied Ecology* with Brock Fenton. Dr. Fenton provided equipment, funding, and editorial comments on the manuscript.

A version of Chapter 5 will be submitted for publication to *Ecology* with Brock Fenton. Dr. Fenton provided funding, equipment, and contributed to study design and editorial comments on the manuscript.

Acknowledgments

First and foremost, I would like to acknowledge my supervisor, Dr. Brock Fenton, for his guidance and support. His limitless enthusiasm and curiosity has motivated me as an academic. Brock's passion for teaching, science, bats, and lifelong learning goes beyond inspiring. Thank you for the opportunities you have opened and for never giving up on me. Some good.

The Department of Biology has been a wonderful environment to work in as a graduate student and I appreciate all of the faculty who took time to support me. I would like to thank the members of my advisory committee, Drs. Nusha Keyghobadi and Jeremy McNeil, for their recommendations, questions, and encouragement. Thanks to Ben Rubin for always being willing to sit and talk about bats and stats with me. I am grateful for my involvement with the Centre for Environment and Sustainability. I had the opportunity to be part of a community of diverse and intelligent people from a wide range of disciplines. Thank you to Dr. Dan Shrubsole who never doubted me and was always happy to talk about anything. Also, thanks to Holly Sanderson for her support in many of my endeavors.

I am grateful for all of my friends and colleagues at Western. Thank you to my labmates, my family while in Canada: Dr. Signe Brinkløv, Lindsey Clairemont, Dr. Beth Clare, Matthew Emrich, Dr. Erin Fraser, Rachel Hamilton, Colin Hayward, Lauren Hooton, Meredith Jantzen, Louis Lazure, Dr. Liam McGuire, Dara Orbach, Leslie Reynolds, Aline Simões, Dr. Mark Skowronski, Toby Thorne, and Nina Veselka. They have been friends, collaborators, and valuable resources. Without their support I would not have made it

through. Special thanks to Liam for always having the enthusiasm and encouragement to keep me going. You are a remarkable person and never cease to amaze me. Thanks to the Iron Chef participants for good food and something to look forward to each month. I would also like to specially thank Erin, Johnston, and Emma for always opening their home and hearts to me.

There are some people who are so important they have a presence in every part of your life. Steve Duncan, Jess Hurley, Dr. Nikhil Lobo, and Dr. Katie Stammer not only helped me maintain my sanity, but also showed me a world beyond bats. Thanks for all of the support, stimulating conversations, laughs, and advice you have given me over the years. Thank you to all of my friends, especially Joette Ball and Susan Morales, who continue to support me, no matter where I am in the world.

Some of the best experiences of my life have been spent doing field work. I would trade a day of relaxation for a night taking bats out of a net while inhaling insects and spending time with great people. I am indebted to those I have worked with in the field: Dr. Beth Clare, Lauren Hooton, Dr. Liam McGuire, and Juliet Nagel. I would also like to thank the Windle family for opening their doors to me and making time spent alone collecting data not so lonely.

I am grateful for the funding I received from Bat Conservation International, Department of Biology, Centre for Environment and Sustainability, Graduate Teaching Assistants' Union, and Society of Graduate Students.

Thank you to everyone I have interacted with during my time at Western. You have made this a special part of my life. Thanks to everyone at the Grad Club; some of the best

conversations I've had were while working. The encouragement everyone showed for their resident "bat lady" made a difference. Being asked, "How are the bats," several times a week let me know you care and I appreciate everyone who was rooting for me to finish.

Finally, I would like to thank all of my family. Thank you especially to my parents, Emily, Grandma Barbara, and Laura Gomez, for their continued support and encouragement. While everyone else thinks I am crazy for being in school for so long their ongoing reminders of how proud they are of me has served as motivation to persevere. I am forever grateful to my parents for instilling love and respect for nature in me. I would not be where I am today without them, even if my dad still doesn't understand my obsession with bats.

Table of Contents

Co-Authorship Statement.....	iv
Acknowledgments.....	v
Table of Contents	viii
List of Tables	xiii
List of Figures	xv
List of Appendices	xviii
Chapter 1	1
1 Introduction	1
1.1 Conservation and ecological sampling	1
1.2 Acoustic monitoring.....	3
1.2.1 Species identification	4
1.2.2 Environmental quality.....	4
1.2.3 Individuals and population structure.....	5
1.2.4 Technology: advancements and limitations.....	5
1.2.5 High frequency sounds	6
1.3 Monitoring bats.....	7
1.3.1 Acoustic monitoring today.....	9
1.4 Variation in acoustic monitoring	10
1.4.1 Variation from attenuation	11
1.4.2 Variation from equipment.....	11
1.4.3 Variation from bats	12
1.5 Bat conservation.....	13
1.5.1 Bats as bioindicators	14

1.5.2	Current threats to bats	14
1.5.3	Standardized protocols.....	16
1.6	Dissertation structure	17
1.7	References.....	19
Chapter 2	35
2	Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats	35
2.1	Introduction.....	35
2.2	Methods.....	39
2.2.1	Optimizing detector recording settings	40
2.2.2	Synthetic call playback	42
2.2.3	Recording free-flying bats	44
2.3	Results.....	44
2.3.1	Synthetic call playback	44
2.3.2	Recording free-flying bats	49
2.4	Discussion	51
2.5	Acknowledgments.....	55
2.6	References.....	55
Chapter 3	59
3	How high is high? Using percentile thresholds to identify peak bat activity.....	59
3.1	Introduction.....	59
3.2	Methods.....	61
3.2.1	Study areas	61
3.2.2	Acoustic monitoring and analysis	61
3.2.3	Statistical analysis	62
3.3	Results.....	64

3.3.1	Among sites	64
3.3.2	Within nights.....	70
3.4	Discussion.....	72
3.5	Acknowledgements.....	75
3.6	References.....	76
Chapter 4	80
4	Identifying peaks in bat activity: a new application of the space-time scan statistic...	80
4.1	Introduction.....	80
4.2	Methods.....	83
4.2.1	Study areas	83
4.2.2	Acoustic sampling.....	83
4.2.3	Statistical analysis.....	84
4.3	Results.....	87
4.3.1	SaTScan description and comparison to percentile thresholds.....	87
4.3.2	Species-specific activity.....	92
4.4	Discussion	94
4.5	Acknowledgements.....	96
4.6	References.....	97
Chapter 5	104
5	Value in variation? Stationary acoustic transects to account for spatial variation in bat activity	104
5.1	Introduction.....	104
5.2	Methods.....	107
5.2.1	Study areas	107
5.2.2	Acoustic sampling.....	107
5.2.3	Statistical methods	112

5.2.4	Generalized linear mixed models.....	112
5.2.5	Percentile thresholds	113
5.2.6	SaTScan	114
5.2.7	Sub-sampling for estimates of activity	115
5.3	Results.....	115
5.3.1	Differences among detectors.....	115
5.3.2	Generalized linear mixed models: variance partitioning of random effects	118
5.3.3	Percentile thresholds: magnitude of activity.....	120
5.3.4	SaTScan: peaks of activity in space and time	123
5.3.5	Sub-sampling for estimates of activity	126
5.4	Discussion	129
5.5	Acknowledgements.....	132
5.6	References.....	132
6	Conclusion.....	138
6.1	Contribution to acoustic studies of bats	138
6.1.1	Variation from detectors	139
6.1.2	Objective methods for detecting peaks in bat activity	139
6.1.3	Variation within sites	140
6.2	Recommendations for future acoustic surveys	140
6.3	Future research directions	143
6.4	Concluding remarks	145
6.5	References.....	146
	Appendices.....	150
	Appendix A: Discriminant function analysis for species identification	150
	Appendix B: Chapter 5 supplementary material.....	156

Appendix C: Permission to reproduce published material.....	168
Appendix D: Animal use protocol approvals.....	169
Appendix E: Research permits.....	172
Curriculum Vitae	175

List of Tables

Table 2.1. A comparison of the features associated with commercially available ultrasonic bat detector systems.	36
Table 2.2. Detector settings used in this study.....	41
Table 2.3. Automated detection parameter settings used for call analysis in callViewer..	43
Table 3.1. Number of calls per hour at each percentile threshold for each species of bat and all species combined for a distribution of activity from three sites in Canada. Half of the activity falls below the 50 th percentile (median) and 10% of activity is above the 90 th percentile threshold.	65
Table 3.2. Summary of bat activity, for A) all species combined, B) <i>Lasiurus borealis</i> , C) <i>L. cinereus</i> , and D) <i>Myotis lucifugus</i> , at three sites in Canada in 2008 and 2009. Activity over the 50 th percentile threshold was considered high and the relative importance of a site was based on the threshold activity exceeded.	68
Table 5.1. Percent canopy cover at each of the detector locations within A) each of the four sites in Ontario, Canada in 2008 and B) Renfrew mine in 2010.....	109
Table 5.2. Classification accuracy for species identification of echolocation calls for bats in Ontario, Canada with quadratic discriminate function analysis (DFA) using cross-validation. Post-DFA filter settings removed echolocation calls identified to species, but outside of the species-specific ranges of duration and minimum frequencies (Fmin) to further improve classification accuracy.	111
Table 5.3. Number of bat echolocation calls per hour at the 50 th and 90 th percentile thresholds for each species and all species combined for a distribution of activity at eight sites in Canada and New York. Activity above the 50 th percentile (median) is considered high and activity above the 90 th percentile threshold is very high.....	121

Table 5.4. Number of nights when at least one hour of bat activity was above the 50th and 90th percentile thresholds at each detector (A-E) over 15 nights at an abandoned mine near Renfrew, Ontario in 2010..... 122

Table 5.5. Mean \pm SD number of nights required to have at least 80% of random samples with mean nightly bat activity within 10-50% of the grand mean of the entire dataset (total calls per night for one detector over 15 nights) for six detectors at a mine near Renfrew, Ontario, Canada in 2010. Data were randomly sampled 100 times to include nightly activity for one detector for two to ten night sample periods. Empty cells are when more than ten nights of sampling were required for all detectors..... 127

Table 5.6. Number of nights required to have at least 80% of random samples with mean nightly bat activity within 30% of the grand mean of the entire dataset at A) four sites in Ontario, Canada in 2008 (five detectors over 12 nights) and B) one site in 2010 (five detectors over 15 nights). Data were randomly sampled 100 times to include nightly activity for one to five detectors for two to ten night sample periods for each site. Empty cells are when more than ten nights of sampling were required for all sites. 128

List of Figures

- Figure 2.1. Mean number of calls detected by each bat detector system at four frequencies at each distance and angle during the synthetic playback experiment. There were 15 calls played for each frequency/distance/angle combination.46
- Figure 2.2. Distance of 50% probability of detection calculated with a logistic regression for each frequency at 0° by each bat detector system during the synthetic playback experiment. Patterns were similar for all detectors at 45° and 90°, but with lower overall probability of detection.47
- Figure 2.3. Performance varied among detectors with a strong effect of frequency. Call detection (arcsine square root transformed number of calls) ± SE by call frequency evaluated at a distance of 22.5 m. Detectors with the same letter superscript were not significantly different from each other within each frequency.48
- Figure 2.4. Mean number of calls ± SE per pass relative to Avisoft for each bat detector from recordings of free-flying *Lasiurus cinereus* on three nights. Batlogger detected more calls than any of the other systems (detectors with the same letter superscript were not significantly different from each other).50
- Figure 3.1. Frequency of activity levels for all bat species combined at three sites in Canada. Dashed, vertical lines are thresholds of the number of calls per hour based on the 50th, 70th, 90th, 95th, and 99th percentiles of the distribution. Activity over the 50th percentile threshold (median) is considered high activity because half of all detections had fewer than 265 calls per hour.66
- Figure 3.2. Comparison of species-specific, within-night bat activity patterns between seasons at sites in Ontario, Canada, highlighting the periods when activity is above certain thresholds for two species. Mean hourly activity for *Lasiurus borealis* at Long Point during A) spring (n = 6 nights) and B) autumn migration (n = 6) and *Myotis lucifugus* activity at Renfrew during C) spring (n = 5) and D) swarming (n = 5). Solid lines are rolling averages (20 increments across each night) of the mean nightly activity

among all nights at a site \pm SD (dashed red); horizontal dashed lines (grey) are percentile thresholds; vertical dashed lines (light grey) are sunrise and sunset.71

Figure 4.1. Nightly activity of bats for all nights at Renfrew during swarming (Site 3b) in 2008 with calls for all species combined. Highlighted time frames are SaTScan high cluster time frames defined by A) various maximum cluster sizes (MCS) within a single night and B-F) MCS of 30% with multivariate scan.88

Figure 4.2. Proportion of overall bat activity in SaTScan high cluster time frames at various maximum cluster sizes (MCS) similar to A) total minutes over the 50th percentile threshold, and B) total nightly activity for three sites in Ontario, Canada during spring (a) and late summer (b) in 2008 with calls for all species combined.90

Figure 4.3. Comparison of nights of peak bat activity (mean calls per minute \pm SD) identified by SaTScan spatial-temporal analysis (grey bars) and with at least 60 minutes of activity over the five percentile thresholds (*50th percentile, **70th percentile, ***90th percentile, ****95th percentile), for all species combined at three sites in Ontario, Canada during spring (a) and late summer (b) in 2008. Note different y-axes.91

Figure 4.4. Mean species-specific bat activity at three sites in Ontario, Canada during spring (a) and late summer (b) for 2008. Shaded time frames are peaks identified by multivariate SaTScan. Note different y-axes. EPFU – *Eptesicus fuscus*, LABO – *Lasiurus borealis*, LACI – *L. cinereus*, MYLU – *Myotis lucifugus*, PESU – *Perimyotis subflavus*.93

Figure 5.1. Mean calls per hour \pm SD for each species of bat at five detectors at Long Point, Ontario (n = 12 nights). Note different y-axes. Detectors of the same letter were paired at the same location and first detectors in the transect (A1/A2) had intermediate openness at the site (20% canopy cover), B1/B2 were the most enclosed (37% canopy cover), and C1 was the most open (5% canopy cover). Detector number denotes height: 1 was low (1.5 m above ground) and 2 was high (4 m). Means, within each species, with the same lowercase letter are not significantly different ($p > 0.05$) according to a Kruskal-Wallis multiple comparison post hoc test.117

Figure 5.2. Proportion of variance explained by spatial (detector, site) and temporal (night) random effects for seven species of bats at four sites in Ontario, Canada. Within-site variation (detector location) is an important component in explaining variation in activity levels of all species combined. The importance of within-site, among-site, and night-to-night effects differs by species. 119

Figure 5.3. Peak periods of bat activity along linear, stationary transects within two sites: A) during late summer (migration) at Long Point, Ontario in 2008 and B) during swarming (August) at an abandoned mine near Renfrew, Ontario in 2010. Species-specific activity patterns vary among detectors. Detectors at Long Point were paired at two heights, low (1.5 m; A1, B1, and C1) and high (4 m; A2 and B2). All detectors at Renfrew were at 1.5 m. There is no evidence of spatial partitioning, but temporal partitioning (no time overlap of peak activity) is evident between several species, such as *L. noctivagans* and *L. borealis*. EPFU – *Eptesicus fuscus*, LANO – *Lasionycteris noctivagans*, LABO – *Lasiurus borealis*, LACI – *L. cinereus*, MYLU – *Myotis lucifugus*, MYSE – *M. septentrionalis*, PESU – *Perimyotis subflavus*. 125

List of Appendices

Appendix A: Discriminant function analysis for species identification	150
Appendix B: Chapter 5 supplementary material.....	156
Appendix C: Permission to reproduce published material.....	168
Appendix D: Animal use protocol approvals.....	169
Appendix E: Research permits.....	172

Chapter 1

1 Introduction

1.1 Conservation and ecological sampling

Preserving biodiversity is a fundamental goal of conservation, but many challenges remain, particularly because of a lack of information about most species. Insufficient knowledge about distribution and natural history are major hurdles because wildlife management and monitoring are basic requirements for the conservation of species. We have to know what is present to conserve it, but collecting realistic data on organisms can be challenging to researchers and conservation efforts. Rare, elusive, and cryptic species can be difficult to sample and while the presence of a species can be confirmed, its absence can only be inferred with a degree of probability (Kéry 2002). Field studies of organisms may be hampered by our limited ability to observe them and/or access their habitats. However, some of these obstacles have been resolved, at least in part, by technological advances (e.g., radio telemetry; Cagnacci *et al.* 2010), but at the same time these techniques may create new challenges to consider.

Ecological studies use many approaches at different spatial and temporal scales to address questions relating to species distribution and abundance. Sampling can provide an estimate rather than an exact measure of what is in the environment, and this is further limited by the trade-off between sampling effort and accuracy of the data obtained. Effort is limited by factors such as time, money, manpower, and habitat accessibility, while accuracy will depend on the sampling effort actually invested. It is important to recognize that balancing the trade-off between effort and accuracy comes from understanding of the

focal organism and complexity of the study system (Loehle 2004). Spatial and temporal complexity will impact the study scale, but in the end, there is no single scale that is appropriate for all ecological studies (Levin 1992).

The ability to address sources of bias will impact data accuracy and play a role in determining investment in sampling effort. Bias can arise from poor practical techniques or non-representative sampling (Sutherland 2006) related to the type of sampling method employed (e.g., visual, capture, acoustic). Poor technique, such as missed direct observations of individuals, can obviously impact the quality of data. Non-representative sampling can cause bias, for example if it is assumed that a single sampling technique will detect all individuals in a population or species in a community equally. Taking into account that variation in the probability of detection may differ among habitats, seasons, species, age or sex of individuals within a species, as well as the sampling techniques deployed, is essential to minimizing the chances of collecting biased data.

For example, visually conspicuous species are often overrepresented, so one must use sampling methods that account for those that cannot simply be seen when walking a transect (e.g., birds and bats in mist-nets, traps or fogging for insects). However, these specialized capture techniques are invasive, as well as time and labor intensive. Acoustic methods provide an efficient, non-intrusive way to study species that use auditory signals, although these record disproportionately more species with high-intensity calls and provide no data on actual population sizes (Flaquer *et al.* 2007). Consequently, to minimize bias, many researchers stress the importance of using multiple sampling techniques simultaneously (Kalko *et al.* 1996, O'Farrell and Gannon 1999, Duffy *et al.* 2000, Milne *et al.* 2004).

Many organisms produce auditory signals, produced intentionally for communication or orientation or unintentionally as a byproduct of activities such as feeding or movement, which researchers can use to document ecology and behavior. Conspicuous auditory signals are one means of detecting some otherwise cryptic organisms and bioacoustics research is an integral part of conservation plans for many animals (Baptista and Gaunt 1997).

1.2 Acoustic monitoring

Acoustic monitoring is fundamental for the study of many organisms traditionally sampled by visual or capture techniques. Songs or calls can be highly reliable taxonomic features, especially for anurans (Taylor *et al.* 1996), bats (Fenton and Bell 1981), birds (Parker 1991, Somervuo *et al.* 2006), cetaceans (Oswald *et al.* 2003), and insects (Riede 1998, Chesmore and Ohya 2004). Acoustic sampling is an important tool for wildlife management and conservation because it can estimate diversity and relative abundances.

Detection of species that can easily be heard but are not easily seen is the true power of acoustic monitoring. For example, high flying bats, such as *Lasiurus cinereus* and *Eurderma maculata*, are rarely included in capture inventories, but are readily detectable in acoustic surveys (O'Farrell and Gannon 1999). Acoustic surveys can provide more accurate estimates of diversity than capture techniques (Dawson and Efford 2009) and thus provide the opportunity to learn about organisms from the community to the individual level.

1.2.1 Species identification

Identification of organisms to species based on their acoustic signals allows researchers to quickly survey the biodiversity of regions (Riede 1998), leading to more accurate species counts and occupancy estimates (Brandes 2008). Birds are perhaps the best known group where species identification with acoustic signals provides the most effective sampling approach (Parker 1991, Riede 1993). For example, Parker (1991) recorded the vocalizations of 85% of the 287 bird species present in the Bolivian Amazon in seven days, compared with 54 days when using captures. The North American Breeding Bird Survey (BBS) heavily relies on acoustic identification and has provided a wealth of information on bird populations and relative abundances, which is used in several conservation efforts (Sauer *et al.* 2003). Standardized protocols incorporating acoustic monitoring, such as the BBS, have provided a powerful tool for management and conservation at various geographic scales.

1.2.2 Environmental quality

Bioindicators are species or communities sensitive to identified environmental stressors or disturbances that may be used to assess the quality of the environment and/or record changes over time (Jones *et al.* 2009). Indicators of environmental change, such as shifts in community structure and species diversity, should be included in conservation plans (Lim and Engstrom 2001). Species that can be monitored acoustically are invaluable indicators of habitat quality because of less invasive sampling methods. Birds and invertebrates are most commonly used as bioindicators (e.g., Browder *et al.* 2002, Mankin *et al.* 2010). For example, acoustic studies of orthopteran communities have been

useful indicators of eutrophication (Fischer *et al.* 1997) and bats are good indicators of habitat quality (Wickramasinghe *et al.* 2003, Kalcounis-Rueppell *et al.* 2007).

1.2.3 Individuals and population structure

Sounds can be used to evaluate fine scale information about communities, including structure of populations and identification of individuals. The low-frequency vocalizations of elephants can vary with group size, composition, and reproductive status, so recordings could provide valuable information on abundance and population structure, especially when they live in densely forested areas (Payne *et al.* 2003). Similarly, blue whale songs can be divided into regional types which can be used to characterize population structure (McDonald *et al.* 2006). Bird dialects can also play an important role in conservation, from obtaining demographic information (Laiolo and Tella 2008) to impacting translocation efforts (e.g., Bradley 2012). Some animals can be identified to the individual-level based on their acoustic signals (e.g., fallow deer, Reby *et al.* 1998; Stellar sea lions, Campbell *et al.* 2002; African wild dog, Hartwig 2005), which can be used for identification in place of physical marks (i.e., tags and bands; Laiolo 2010), providing important information for population monitoring. A paucity of data on individual variation currently precludes identification to the individual level in animals such as insects, anurans, and bats (Obrist 1995, Chesmore 2001).

1.2.4 Technology: advancements and limitations

No matter the focal organism of an acoustic study, technological achievements strongly influence the feasibility of using acoustic techniques. Advances in technology and software have allowed researchers to overcome difficulties with data collection and storage, making it possible to monitor previously inaccessible habitats and develop a

better “picture” of biodiversity. Acoustic studies are not without drawbacks. Analysis of acoustic data can be extremely slow when done manually, requiring highly trained personnel (Chesmore 2004) and limited by the availability of an accurate reference library (Riede 1998). Manual identification of species relies on expert knowledge, which is inherently subjective. Possibly the greatest drawback is the inability to count individuals using acoustic methods (Brandes 2008), as activity does not equal abundance (Hayes 2000). It is not possible to determine whether sounds are coming from one or multiple individuals moving past a microphone, so acoustic data can only provide a relative index of activity levels (Hayes 2000). Interpretation of acoustic data often relies on descriptive terminology, such as “higher” or “lower,” limiting the accuracy of conclusions. There is no framework or guideline on how to interpret specific activity levels beyond relative differences in activity from one sample to the next (Kunz *et al.* 2007b).

1.2.5 High frequency sounds

While many sounds recorded during acoustic monitoring are audible to humans, others are not. Toothed whales (Odontoceti; e.g., dolphins) and bats (Chiroptera) are two taxa that use echolocation for communication, foraging, and orientation by emitting pulses of high frequency sound and gathering information based on the returning echoes. Signals produced by odontocetes travel through water at a speed about four times faster than if produced in air. Dolphins are able to detect small targets from a few meters (Kastelein *et al.* 1999) to over a hundred meters away (Au 1993). Most dolphins emit a combination of whistles and clicks that do not change in duration or shape. The central frequency depends on the intensity of the signal, ranging from 30-60 kHz (low intensity) to ~100 kHz (high intensity, Au 1993). Bat echolocation has a much slower rate of information

transfer than odontocetes because their signals move through air. Bat calls are much more diverse in structure than odontocete signals, varying in duration, from 0.3 to 300 ms; in frequency, from 12 to >200 kHz; and in shape, ranging from broadband to constant frequency (Neuweiler 1989). Detecting both dolphin and bat echolocation calls requires a researcher to rely on technology capable of detecting high frequency sounds. Bats provide additional challenges as they are small, nocturnal animals that cannot be sampled visually and capture success is limited to low flying species.

1.3 Monitoring bats

Our knowledge of bats and their natural history largely arises from the research of Lazzaro Spallanzani in 1794, who provided the first evidence that bats use sounds for orientation and obstacle avoidance. While Spallanzani's methods of blinding and deafening bats were crude, he provided the basis for what Griffin (1944) later coined as "echolocation."

Griffin (1958) was the first to use a "sonic amplifier" to investigate ultrasonic sounds.

This detector was able to show pulse repetition rates, but the next development in detectors showed that these pulses were actually frequency-modulated calls (Griffin 2004). In 1951, when Griffin first made recordings in the field, bat detectors were still barely portable, very expensive, and extremely fragile. Griffin required the use of a station wagon to transport all the necessary equipment. Visual outputs from an oscilloscope were recorded by a video camera and audio output from a portable AM radio (Griffin 2004). Improvements, which led to the first "true" bat detector, allowed for a portable instrument that produced an audio signal of the ultrasonic pulses (Griffin 2004).

Specialized microphones convert high frequency sounds to electric signals, which are then recorded and transformed for storage, playback, and analysis. Bat detectors can be

heterodyne or broadband. Heterodyne detectors allow users to tune in to a specific, narrow range of frequencies. The heterodyne detector converts high frequency sounds to the audible range in real time. The narrow frequency range is limiting, but it provides a quick method to listen for bat presence. Heterodyne detectors provide no information on call structure and require extensive training for species identification in the field. More complete bat survey efforts require broadband sampling because the calls of species in a single community can cover a wide range of frequencies.

The maximum recorded frequency can be no more than half the sampling rate. Originally storage devices were not fast enough to record echolocation calls at a high enough sampling rate, so signals had to be brought down to a frequency range that could be analyzed. Frequency division and time expansion methods allow broadband recording and transformation of high frequencies for users to listen to at the audible frequency range (Parsons and Obrist 2004). Frequency division decreases an incoming signal frequency (kHz) to bring it into the audible range by dividing the frequency by a preset value. A drawback is that all amplitude information is lost from the recorded call and only a single harmonic is analyzed. Time expansion plays signals back at a slower speed by recording a broadband signal and playing it back with the call duration increased and frequencies decreased (Pettersson 2004). Slowed signals are able to be processed at a lower sampling rate, while preserving all call information. The system records and stores a signal, then plays it back, but during playback, the system cannot also be recording, limiting recording for only part of the available time (Parsons and Obrist 2004). While time expansion systems make high quality recordings, including the amplitude and spectral component information, the drawback is they are not capable of sampling continuously.

Until the 1980s, technology was the bottleneck to addressing many questions in acoustic monitoring. Field research of bats flourished once more portable detectors became available, and it was determined that bat species could be identified by their echolocation calls (Ahlén 1981, Fenton and Bell 1981). Recordings were originally made on analogue tape recorders, but by the mid-1980s they could be stored digitally. Once sound cards were capable of higher sampling rates, high frequency signals could be recorded directly without transformations.

1.3.1 Acoustic monitoring today

Today, devices primarily use one of two methods to analyze digitally recorded calls: zero-crossing and Fourier analyses. Zero-crossing analysis, used to analyze frequency division signals, is advantageous because it is simple and fast with low digital storage requirements. However, it suffers from the same limitations as frequency division: loss of amplitude information and analysis of a single harmonic. Fourier analysis, or spectral analysis, is the more common method of analysis for detectors, used for time expansion and untransformed, real-time signals. It generates two outputs: power spectrum and spectrogram, giving information on amplitude, frequency, and time aspects of echolocation calls. Fourier analysis calculates frequency information by averaging blocks of data across the call. Using more blocks of data increases the accuracy of frequency information, but decreases time resolution because of an inverse relationship between frequency and time (Parsons *et al.* 2000). Very little information is lost during spectral analysis and it is relatively insensitive to background noise, but analysis can be computationally demanding (Parsons *et al.* 2000, Parsons and Szewczak 2009).

An ongoing debate persists regarding the value of the two analysis methods. In practice, the quick, simplistic method of zero-crossing analysis detects fewer echolocation calls than the information-rich spectral analysis. Echolocation call features (i.e., lowest frequency and duration) and sensitivity also differ between the two types of analysis (Fenton 2000, Fenton *et al.* 2001). Spectral analysis requires more data storage and battery life than zero-crossing analysis. Full-spectrum devices allow analysis of an entire echolocation call, including harmonics and amplitude information; because bats probably use the entire call they produce, including harmonics, it is important to understand and include the entire call during analysis (Griffin 2004).

There is now a wide variety of commercially available, inexpensive, portable bat detectors, which allow extensive study of bats through acoustic monitoring; more than 500 studies have been published since 2011. Digital technology and miniaturization of electronic components have made many field studies possible, and new computer software is making analysis of recorded signals even more efficient (Parsons and Obrist 2004). Large quantities of data are easily collected with acoustic methods, but processing and analysis of these immense datasets are still problematic. Analysis is moving towards full automation based on statistical models and computer programming techniques, but has not been commercialized in a robust enough manner to be adopted for standardized protocols.

1.4 Variation in acoustic monitoring

There are three levels of variation that can complicate acoustic data interpretation. First, variation is created by the movement of sounds through air. Second, the degree of variation detected can be affected by the equipment used. Lastly, there is variation with

respect to the animals themselves, from echolocation behavior to community-level activity patterns. No matter what the focus of the study, it is vital to recognize all of the possible sources of variation in a dataset.

1.4.1 Variation from attenuation

The movement of sound through air is affected by several factors. All sounds transmitted through air are subject to spreading loss, where sound waves spread out as they move away from the source and thus lose intensity with distance (independent of frequency, Griffin 1971). Atmospheric attenuation occurs when sounds are absorbed by atmospheric moisture which can be affected by the frequency at which the sounds are emitted, as well as humidity, and temperature. Higher frequency sounds have shorter wavelengths, resulting in greater attenuation (Griffin 1971, Lawrence and Simmons 1982), but yield more details to bats about their targets (Griffin 1958, Simmons and Stein 1980). The frequencies dominated in calls of many species (20-60 kHz) suggest a balance between call range due to attenuation and detection resolution (Fenton *et al.* 1998). Species with higher-frequency components in their calls will have a lower effective range of echolocation due to attenuation and thus may be more difficult to detect with detectors.

1.4.2 Variation from equipment

The variation resulting from the use of different equipment is controllable but often underappreciated, and must be accounted for when developing standardized protocols. This is especially true for acoustic bat detectors, as not all systems have the same sensitivity (Forbes and Newhook 1990, Waters and Walsh 1994, Fenton 2000, Fenton *et al.* 2001) or hear the same signals in the same way. There are differences between brands (Forbes and Newhook 1990, Waters and Walsh 1994) and even between individual

detectors of the same model (Larson and Hayes 2000). Differences between time expansion and zero-crossing analysis detectors can be as high as 19 dB in sensitivity, resulting in zero-crossing systems missing quite a high proportion of bat activity, even for species with relatively high-intensity calls because the detection distance of the detector will be shorter for less sensitive microphones (Fenton *et al.* 2001).

1.4.3 Variation from bats

An individual may vary the structure of their echolocation calls in response to habitat structure (Kalko and Schnitzler 1993, Broders *et al.* 2004), insect noise (Gillam and McCracken 2007), and other bats (Obrist 1995). We are only beginning to appreciate individual variation (Masters *et al.* 1995, Betts 1998, Fenton *et al.* 2004), and it seems unlikely that one could identify individuals from echolocation calls in the field as variation in response to ecological conditions overwhelms the amount of variation at the individual level.

Bat activity and community structure are variable in both space and time. Numerous extrinsic factors affect temporal activity patterns of bats, including insect abundance (Hayes 1997, Lee and McCracken 2002), air temperature (Kunz 1973, Lacki 1984, Nagraeff and Brigham 1995, Hayes 1997), rainfall (Fenton *et al.* 1977, Parsons *et al.* 2003), relative humidity (Lacki 1984, Adam *et al.* 1994), and wind (Adam *et al.* 1994). Activity levels can vary annually (Milne 2006), seasonally (Russ *et al.* 2003, Milne *et al.* 2005, Johnson *et al.* 2011), among nights (Hayes and Adam 1996, Krusic *et al.* 1996, Hayes 1997, Broders 2003), and within nights (Maier 1992, Krusic *et al.* 1996, Hayes 1997, Milne 2006). Activity also varies at both large and small spatial scales. Patchiness in activity can be driven by congregations of bats at special locations at specific times of

the year. Some locations attract high numbers of bats, such as bodies of water, where insects are abundant. Maternity colonies, roosting, hibernation, and migration sites are all locations where bats congregate in greater densities and present higher activity levels. While variation among habitats has received the most attention (e.g., Krusic *et al.* 1996, Vaughan *et al.* 1997, Sherwin *et al.* 2000, Loeb and O’Keefe 2006), spatial variation within a site due to habitat heterogeneity is also an important source of variation that must be considered during surveys (Hayes 2000). Multiple bat detectors recording within a site can be important for reliable estimates of activity levels (Britzke 2003, Fischer *et al.* 2009). Flying bats move through three dimensional space and the vertical stratification of bat activity can vary with species and habitat structural complexity (Kalcounis *et al.* 1999, Hayes and Gruber 2000). Few studies have attempted to sample into the canopy because of the logistical limitations of sampling at greater heights (Kalko and Handley 2001, Lim and Engstrom 2001).

1.5 Bat conservation

Bats are an important group of mammals, serving vital ecological roles as nocturnal insect predators, pollinators, seed dispersers, and have been recognized as significant natural resources in recent years (Gannon *et al.* 2003). Management agencies now recognize the need for practical research of bats (Barclay and Brigham 1996) and that the ecosystem services provided by bats are being quantified (Boyles *et al.* 2011, Kunz *et al.* 2011). Boyles *et al.* (2011) estimate that the United States agricultural industry receives \$22.9 billion in ecosystem services annually from insectivorous bats as predators of many crop and forest pests.

1.5.1 Bats as bioindicators

Changes in community structure and species diversity may serve as indicators of environmental change and should be included in conservation plans (Lim and Engstrom 2001). Bats have been identified as good candidates as bioindicators because of the ability of researchers to monitor trends in populations, occupancy by bats of high trophic levels, and widespread distribution of bat taxa (Jones *et al.* 2009). For example, bats are ecological indicators of ecosystem disturbance (Fenton *et al.* 1992, Medellín *et al.* 2000), habitat quality (Kalcounis-Rueppell *et al.* 2007), and have been included as one of the United Kingdom's Biodiversity Indicators (Bat Conservation Trust 2011). Research conducted with bat detectors informs our understanding of bat ecology and behavior and is frequently used to guide important wildlife management decisions (U.S. Fish and Wildlife Service 2012).

1.5.2 Current threats to bats

Over 1200 species of bats have been described (Simmons 2005) and 172 species are threatened (i.e., vulnerable, endangered, or critically endangered) while five are now extinct (IUCN 2012). Bat populations are being impacted around the world, from habitat destruction to climate change (Jones *et al.* 2009), but two current threats facing bats today are wind-energy developments and white-nose syndrome (WNS). Wind turbines are responsible for the death of countless bats, especially for migratory species (Kunz *et al.* 2007a), due to direct collision with turbine blades (Horn *et al.* 2008, Rollins *et al.* 2012) or pulmonary barotrauma as a result of the rapid drops in air-pressure near moving turbine blades (Baerwald *et al.* 2008). Wind energy development is increasing around the world, as are efforts to find viable mitigation options. There is a growing demand for

surveys at wind energy developments to assess potential risks to bats using both acoustic and capture methods, but these efforts are currently not guided by any standardized survey design (Kunz *et al.* 2007a,b).

White-nose syndrome is causing severe declines of bat populations in eastern North America, with many colonies decreasing by 99% within two years of infection (Frick *et al.* 2010). This fungal infection is caused by a cold-adapted fungus, *Geomyces destructans*, resulting in increased arousals during hibernation, leading to dehydration and depleted fat reserves (Cryan *et al.* 2010, Frick *et al.* 2010). Since being identified in upper New York State in 2006, it has spread rapidly, directly by bats as well as by anthropogenic activity (Frick *et al.* 2010) through the persistence on equipment, clothing, and shoes. Acoustic monitoring is an effective way to document population declines without the risk of transmission of WNS with traditional capture methods (Brooks 2011, Dzal *et al.* 2011, Ford *et al.* 2011).

Given the importance of bats as bioindicators and worldwide decline in populations a global monitoring program with standardized methodologies is needed (Jones *et al.* 2009, Stahlschmidt and Brühl 2012). Lack of standardized protocols hinders the establishment of clear guidelines for surveys and effective regulation of assessment efforts by government, such as for wind energy developments. There are long-term monitoring efforts with standardized protocols in the United Kingdom (Walsh *et al.* 2001), which include acoustic surveys consisting of 1 km walking transects with frequency division detectors. New York State has implemented a standardized driving transect protocol to monitor bat populations post-WNS, which is now being adopted by other regions

although there is little research into the effectiveness of moving transects for sampling bat communities (Russ *et al.* 2003, Stahlschmidt and Brühl 2012).

1.5.3 Standardized protocols

Standardization of survey protocols is an important step to make applied research applicable to management and government policies, as the transition of scientific research to environmental policy decisions is often challenging. Policy based on scientific research is crucial for concerted efforts addressing increased conservation needs. Lack of standardization makes it virtually impossible to compare results, making standardized experimental approaches a necessity (Hayes 1997). Techniques to make monitoring more efficient and accurate will greatly benefit science and conservation efforts. Bats are critical to ecosystem function and the need for large-scale, standardized monitoring efforts is greater than ever before in the wake of threats, such as WNS and wind energy developments. Eventually, standardized protocols based on research will outline methods, from detector model, detector deployment, species identification, to statistical methods for evaluating data.

Some research has investigated variation affecting bat monitoring surveys and made recommendations for future surveys (Hayes 1997, Fischer *et al.* 2009, Skalak *et al.* 2012), but not enough studies have been published to provide a basis for standardized methods. Sources of variation — atmospheric attenuation, bat detectors, and bats — must be considered to collect unbiased acoustic data (Hayes 2000). While atmospheric attenuation cannot be controlled, variation from detectors and bats should be. Any standardized protocol must address and account for these sources of variation. In this dissertation my goal is to investigate these two sources of variation and propose methods for replicable,

objective evaluation of bat activity levels. It was my objective to address these specific questions related to the acoustic monitoring of bats: i) how does detector choice impact acoustic monitoring results?, ii) what is peak bat activity and how is it identified?, iii) how does detector placement impact the design and interpretation of acoustic monitoring studies?

1.6 Dissertation structure

Each chapter in my dissertation was prepared for independent publication. Chapter 2 has been published, Chapters 3 and 4 are in review for publication, and Chapter 5 is pending submission for publication. The four chapters are united by a common theme of investigating sources of variation in acoustic bat surveys in order to make recommendations for standardized sampling protocols. Below is a brief outline of each chapter.

In Chapter 2 (Do you hear what I hear? Implications of detector selection for passive acoustic monitoring of bats), I examined how several brands of bat detectors provide different depictions of the same dataset. The purpose was to quantify variation in detection performance among several commercially available bat detector systems and investigate if this is an important source of variation in acoustic monitoring methods. With an unprecedented variety of detectors available, it is crucial for researchers and management officials to understand differences in the performance of each of the detectors, both while selecting detectors to use and interpreting results from studies using different detector brands. I conducted a two-part study using five passive bat detectors: first, a controlled experiment with synthetic calls recorded at fixed frequencies, distances, and angles; and second, a field experiment recording free-flying bats.

In Chapter 3 (How high is high? Using percentile thresholds to identify peak bat activity), I address the question, “What is peak activity and how is it identified?” and illustrate a method for classifying activity levels and investigate how patterns and peaks of activity vary among sites. Describing bat activity as “high” or “low” is useful only in context and is rarely evaluated based on objective criteria. I recorded acoustic bat activity at three sites, spanning a range of situations. I calculated species-specific thresholds of activity levels at six (25th - 99th) percentiles derived from a larger distribution of activity levels among all sites. I used these percentile thresholds to identify important sites for each species based on where I found high activity, defined by objective criteria. It is important to have clear definitions of “high” activity, especially when making conservation and management decisions.

In Chapter 4 (Identifying peaks in bat activity: a new application of the space-time scan statistic), I address the same question as in Chapter 3, but present an alternate solution to the problem of identifying peak bat activity. I propose a new application for the space-time scan statistic (SaTScan) as an objective statistical technique for identifying peak periods of bat activity and compared it to the use of percentile thresholds, at three scales: within nights, among nights at a site, and among sites. I then experimentally tested SaTScan by analyzing species-specific activity at three sites. SaTScan has the potential to be a valuable tool for quickly identifying activity peaks with an objective, replicable, and statistically-sound method that can be applied at many temporal and spatial scales.

In Chapter 5 (Value in variation? Stationary acoustic transects to account for spatial variation in bat activity), I examine how horizontal and vertical variation within a site can impact estimates of bat activity. I measured bat activity with linear, stationary transects of

bat detectors at four sites in Ontario. I assessed variation in patterns and levels of bat activity at each detector with percentile thresholds and SaTScan. I also tested whether spatial or temporal factors were more important for explaining variation in activity. Lastly, I explored what sampling effort is required for species-specific activity levels at each site.

In Chapter 6, I conclude my dissertation with a summary of how addressing variation in acoustic bat surveys is necessary for accurate estimates of activity and make recommendations for sampling protocols. Finally, I highlight future directions for the study of bat ecology with the intention of better conservation and management.

1.7 References

- Adam, M., M. Lacki, and L. Shoemaker. 1994. Influence of environmental conditions on flight activity of *Plecotus townsendii virginianus* (Chiroptera: Vespertilionidae). *Brimleyana* 21:77–85.
- Ahlén, I. 1981. Identification of Scandinavian bats by their sounds. Rapport - Sveriges Lantbruksuniversitet, Institutionen foer Viltkologi 6:1–56.
- Au, W. 1993. The sonar of dolphins. Springer-Verlag, New York, NY.
- Baerwald, E., G. D'Amours, B. Klug, and R. Barclay. 2008. Barotrauma is a significant cause of bat fatalities at wind turbines. *Current Biology* 18:695–696.

- Baptista, L. F., and S. L. L. Gaunt. 1997. Bioacoustics as a tool in conservation. Pages 212–241 in J. R. Clemmons and R. Buchholz, editors. Behavioral Approaches to Conservation in the Wild. Cambridge University Press, Cambridge.
- Barclay, R., and R. Brigham. 1996. Bats and Forests Symposium.
- Bat Conservation Trust. 2011. The state of the UK's bats: National Bat Monitoring Programme population trends 2011. London.
- Betts, B. 1998. Effects of interindividual variation in echolocation calls on identification of big brown and silver-haired bats. *Journal of Wildlife Management* 62:1003–1010.
- Boyles, J., P. Cryan, G. McCracken, and T. Kunz. 2011. Economic importance of bats in agriculture. *Science* 332:11–12.
- Bradley, D. 2012. Evolutionary significance and conservation implications of vocal dialects in North Island Kōkako (*Callaeas wilsoni*). PhD Thesis. University of Waikato.
- Brandes, S. T. 2008. Automated sound recording and analysis techniques for bird surveys and conservation. *Bird Conservation International* 18:S163–S173.
- Britzke, E. 2003. Use of ultrasonic detectors for acoustic identification and study of bat ecology in the eastern United States. PhD Thesis. Tennessee Technological University.

- Broders, H. 2003. Another quantitative measure of bat species activity and sampling intensity considerations for the design of ultrasonic monitoring studies. *Acta Chiropterologica* 5:235–241.
- Broders, H., C. Findlay, and L. Zheng. 2004. Effects of clutter on echolocation call structure of *Myotis septentrionalis* and *M. lucifugus*. *Journal of Mammalogy* 85:273–281.
- Brooks, R. T. 2011. Declines in summer bat activity in central New England 4 years following the initial detection of white-nose syndrome. *Biodiversity and Conservation* 20:2537–2541.
- Browder, S. F., D. H. Johnson, and I. J. Ball. 2002. Assemblages of breeding birds as indicators of grassland condition. *Ecological Indicators* 2:257–270.
- Cagnacci, F., L. Boitani, R. A. Powell, and M. S. Boyce. 2010. Animal ecology meets GPS-based radiotelemetry: a perfect storm of opportunities and challenges. *Philosophical transactions of the Royal Society B* 365:2157–2162.
- Campbell, G. S., R. C. Gisiner, D. A. Helweg, and L. L. Milette. 2002. Acoustic identification of female Steller sea lions (*Eumetopias jubatus*). *Journal of the Acoustical Society of America* 111:2920–2928.
- Chesmore, E. D. 2001. Application of time domain signal coding and artificial neural networks to passive acoustical identification of animals. *Applied Acoustics* 62:1359–1374.

- Chesmore, E. D. 2004. Automated bioacoustic identification of species. *Anais da Academia Brasileira de Ciências* 76:436–440.
- Chesmore, E. D., and E. Ohya. 2004. Automated identification of field-recorded songs of four British grasshoppers using bioacoustic signal recognition. *Bulletin of Entomological Research* 94:319–330.
- Cryan, P. M., C. U. Meteyer, J. G. Boyles, and D. S. Blehert. 2010. Wing pathology of white-nose syndrome in bats suggests life-threatening disruption of physiology. *BMC Biology* 8:135.
- Dawson, D. K., and M. G. Efford. 2009. Bird population density estimated from acoustic signals. *Journal of Applied Ecology* 46:1201–1209.
- Duffy, A. M., L. F. Lumsden, C. R. Caddle, R. R. Chick, and G. R. Newell. 2000. The efficacy of Anabat ultrasonic detectors and harp traps for surveying microchiropterans in south-eastern Australia. *Acta Chiropterologica* 2:127–144.
- Dzal, Y., L. P. McGuire, N. Veselka, and M. B. Fenton. 2011. Going, going, gone: the impact of white-nose syndrome on the summer activity of the little brown bat (*Myotis lucifugus*). *Biology Letters* 7:392–394.
- Fenton, M. B. 2000. Choosing the “correct” bat detector. *Acta Chiropterologica* 2:215–224.
- Fenton, M. B., N. G. H. Boyle, T. M. Harrison, and D. J. Oxley. 1977. Activity patterns, habitat use, and prey selection by some African insectivorous bats. *Biotropica* 9:73–85.

- Fenton, M. B., and G. Bell. 1981. Recognition of species of insectivorous bats by their echolocation calls. *Journal of Mammalogy* 62:233–243.
- Fenton, M. B., L. Acharya, D. Audet, M. B. C. Hickey, C. Merriman, D. M. Obrist, Martin K. Syme, and B. Adkins. 1992. Phyllostomid bats (Chiroptera: Phyllostomidae) as indicators of habitat disruption in the Neotropics. *Biotropica* 24:440–446.
- Fenton, M. B., C. Portfors, I. Rautenbach, and J. Waterman. 1998. Compromises: sound frequencies used in echolocation by aerial-feeding bats. *Canadian Journal of Zoology* 76:1174–1182.
- Fenton, M. B., S. Bouchard, M. J. Vonhof, and J. Zigouris. 2001. Time-expansion and zero-crossing period meter systems present significantly different views of echolocation calls of bats. *Journal of Mammalogy* 82:721–727.
- Fenton, M. B., D. S. Jacobs, E. J. Richardson, P. J. Taylor, and W. White. 2004. Individual signatures in the frequency-modulated sweep calls of African large-eared, free-tailed bats *Otomops martiensseni* (Chiroptera: Molossidae). *Journal of Zoology* 262:11–19.
- Fischer, F., U. Schulz, and H. Schubert. 1997. Quantitative assessment of grassland quality: acoustic determination of population sizes of orthopteran indicator species. *Ecological Applications* 7:909–920.

- Fischer, J., J. Stott, B. S. Law, M. D. Adams, and R. I. Forrester. 2009. Designing effective habitat studies: quantifying multiple sources of variability in bat activity. *Acta Chiropterologica* 11:127–137.
- Flaquer, C., I. Torre, and A. Arrizabalaga. 2007. Comparison of sampling methods for inventory of bat communities. *Journal of Mammalogy* 88:526–533.
- Forbes, B., and E. Newhook. 1990. A comparison of the performance of three models of bat detectors. *Journal of Mammalogy* 71:108–110.
- Ford, W. M., E. R. Britzke, C. a. Dobony, J. L. Rodrigue, and J. B. Johnson. 2011. Patterns of acoustical activity of bats prior to and following white-nose syndrome occurrence. *Journal of Fish and Wildlife Management* 2:125–134.
- Frick, W. F., J. F. Pollock, A. C. Hicks, K. E. Langwig, D. S. Reynolds, G. G. Turner, C. M. Butchkoski, and T. H. Kunz. 2010. An emerging disease causes regional population collapse of a common North American bat species. *Science* 329:679–82.
- Gannon, W., R. Sherwin, and S. Haymond. 2003. On the importance of articulating assumptions when conducting acoustic studies of habitat use by bats. *Wildlife Society Bulletin* 31:45–61.
- Gillam, E. H., and G. F. McCracken. 2007. Variability in the echolocation of *Tadarida brasiliensis*: effects of geography and local acoustic environment. *Animal Behaviour* 74:277–286.
- Griffin, D. R. 1944. Echolocation by blind men, bats and radar. *Science* 100:589–590.

- Griffin, D. R. 1958. Listening in the dark: the acoustic orientation of bats and men. Yale University Press.
- Griffin, D. R. 1971. The importance of atmospheric attenuation for the echolocation of bats (Chiroptera). *Animal Behaviour* 19:55–61.
- Griffin, D. R. 2004. The past and future history of bat detectors. Pages 6–9 in R. Brigham, E. Kalko, G. Jones, S. Parsons, and H. J. G. A. Limpens, editors. *Bat Echolocation Research: tools, techniques and analysis*. Bat Conservation International. Austin, TX.
- Hartwig, S. 2005. Individual acoustic identification as a non-invasive conservation tool: an approach to the conservation of the African wild dog *Lycaon pictus* (Temminck, 1820). *Bioacoustics* 15:35–50.
- Hayes, J. 1997. Temporal variation in activity of bats and the design of echolocation-monitoring studies. *Journal of Mammalogy* 78:514–524.
- Hayes, J. 2000. Assumptions and practical considerations in the design and interpretation of echolocation-monitoring studies. *Acta Chiropterologica* 2:225–236.
- Hayes, J. and M. Adam. 1996. The influence of logging riparian areas on habitat utilization by bats in western Oregon. Pages 228–237 in R. M. R. Barclay and R. M. Brigham, editors. *Bats and Forests Symposium*.
- Hayes, J., and J. Gruber. 2000. Vertical stratification of bat activity in an old-growth forest in western Washington. *Northwest Science* 74:102–108.

- Horn, J. W., E. B. Arnett, and T. H. Kunz. 2008. Behavioral responses of bats to operating wind turbines. *Journal of Wildlife Management* 72:123–132.
- IUCN. 2012. IUCN Red List of Threatened Species. Version 2012.2.
www.iucnredlist.org.
- Johnson, J. B., J. W. Edwards, and W. M. Ford. 2011. Nocturnal activity patterns of Northern Myotis (*Myotis septentrionalis*) during the maternity season in West Virginia (USA). *Acta Chiropterologica* 13:391–397.
- Jones, G., D. Jacobs, T. Kunz, M. Willig, and P. Racey. 2009. *Carpe noctem*: the importance of bats as bioindicators. *Endangered Species Research* 8:93–115.
- Kalcounis, M., and K. Hobson. 1999. Bat activity in the boreal forest: importance of stand type and vertical strata. *Journal of Mammalogy* 80:673–682.
- Kalcounis-Rueppell, M., V. Payne, S. Huff, and A. Boyko. 2007. Effects of wastewater treatment plant effluent on bat foraging ecology in an urban stream system. *Biological Conservation* 138:120–130.
- Kalko, E. K. V., and H. Schnitzler. 1993. Plasticity in echolocation signals of European pipistrelle bats in search flight: implications for habitat use and prey detection. *Behavioral Ecology and Sociobiology* 33:415–428.
- Kalko, E. K. V., C. Handley, and D. Handley. 1996. Organization, diversity, and long-term dynamics of a Neotropical bat community. Pages 503–553 *in* M. L. Cody and J. A. Smallwood, editors. *Long-term studies of vertebrate communities*. Academic Press, San Diego, CA, USA.

- Kalko, E. K. V., and C. Handley. 2001. Neotropical bats in the canopy: diversity, community structure, and implications for conservation. *Plant Ecology* 153:319–333.
- Kastelein, R. A., W. W. Au, H. T. Rippe, and N. M. Schooneman. 1999. Target detection by an echolocating harbor porpoise (*Phocoena phocoena*). *Journal of the Acoustical Society of America* 105:2493–8.
- Kéry, M. 2002. Inferring the absence of a species: a case study of snakes. *Journal of Wildlife Management* 66:330–338.
- Krusic, R., M. Yamasaki, C. D. Neefus, P. J. Pekins, and P. J. Perkins. 1996. Bat habitat use in White Mountain National Forest. *Journal of Wildlife Management* 60:625–631.
- Kunz, T. 1973. Resource utilization: temporal and spatial components of bat activity in central Iowa. *Journal of Mammalogy* 54:14–32.
- Kunz, T., E. Arnett, W. P. Erickson, A. Hoar, G. Johnson, R. P. Larkin, M. D. Strickland, R. Thresher, and M. Tuttle. 2007a. Ecological impacts of wind energy development on bats: questions, research needs, and hypotheses. *Frontiers in Ecology and Environment* 5:315–324.
- Kunz, T. H., E. B. Arnett, B. M. Cooper, W. P. Erickson, R. P. Larkin, T. Mabee, M. L. Morrison, M. D. Strickland, and J. M. Szewczak. 2007b. Assessing impacts of wind-energy development on nocturnally active birds and bats: a guidance document. *Journal of Wildlife Management* 71:2449–2486.

- Kunz, T. H., E. Braun de Torrez, D. Bauer, T. Lobova, and T. H. Fleming. 2011. Ecosystem services provided by bats. *Annals of the New York Academy of Sciences* 1223:1–38.
- Lacki, M. 1984. Temperature and humidity-induced shifts in the flight activity of little brown bats. *The Ohio Journal of Science* 84:264–266.
- Laiolo, P. 2010. The emerging significance of bioacoustics in animal species conservation. *Biological Conservation* 143:1635–1645.
- Laiolo, P., and J. Tella. 2008. Social determinants of songbird vocal activity and implications for the persistence of small populations. *Animal Conservation* 11:433–441.
- Larson, D. J., and J. P. Hayes. 2000. Variability in sensitivity of Anabat II bat detectors and a method of calibration. *Acta Chiropterologica* 2:209–213.
- Lawrence, B. D., and J. A. Simmons. 1982. Measurements of atmospheric attenuation at ultrasonic frequencies and the significance for echolocation by bats. *Journal of the Acoustical Society of America* 71:585–90.
- Lee, Y. F., and G. McCracken. 2002. Foraging activity and food resource use of Brazilian free-tailed bats, *Tadarida brasiliensis* (Molossidae). *Ecoscience* 9:306–313.
- Levin, S. 1992. The problem of pattern and scale in ecology: the Robert H. MacArthur award lecture. *Ecology* 73:1943–1967.
- Lim, B., and M. Engstrom. 2001. Bat community structure at Iwokrama forest, Guyana. *Journal of Tropical Ecology* 17:647–665.

- Loeb, S., and J. O'Keefe. 2006. Habitat use by forest bats in South Carolina in relation to local, stand, and landscape characteristics. *Journal of Wildlife Management* 70:1210–1218.
- Maier, C. 1992. Activity patterns of pipistrelle bats (*Pipistrellus pipistrellus*) in Oxfordshire. *Journal of Zoology* 228:69–80.
- Mankin, R. W., R. D. Hodges, H. T. Nagle, C. Schal, R. M. Pereira, and P. G. Koehler. 2010. Acoustic indicators for targeted detection of stored product and urban insect pests by inexpensive infrared, acoustic, and vibrational detection of movement. *Journal of Economic Entomology* 103:1636–1646.
- Masters, W., K. Raver, and K. Kazial. 1995. Sonar signals of big brown bats, *Eptesicus fuscus*, contain information about individual identity, age and family affiliation. *Animal Behaviour* 50:1243–1260.
- McDonald, M., J. A. Hildebrand, and S. L. Mesnick. 2006. Biogeographic characterisation of blue whale song worldwide: using song to identify populations. *Journal of Cetacean Research and Management* 8:55.
- Medellín, R. A., M. Equihua, and M. A. Amin. 2000. Bat diversity and abundance as indicators of disturbance in Neotropical rainforests. *Conservation Biology* 14:1666–1675.
- Milne, D. J. 2006. Habitat relationships, activity patterns and feeding ecology of insectivorous bats of the Top End of Australia. PhD Thesis. James Cook University.

- Milne, D. J., M. Armstrong, A. Fisher, T. Flores, and C. Pavey. 2004. A comparison of three survey methods for collecting bat echolocation calls and species-accumulation rates from nightly Anabat recordings. *Wildlife Research* 31:57–63.
- Milne, D. J., A. Fisher, I. Rainey, and C. Pavey. 2005. Temporal patterns of bats in the Top End of the Northern Territory, Australia. *Journal of Mammalogy* 86:909–920.
- Negraeff, O., and R. M. Brigham. 1995. The influence of moonlight on the activity of little brown bats (*Myotis lucifugus*). *Zeitschrift für Säugetierkunde* 60:330–336.
- Neuweiler, G. 1989. Foraging ecology and audition in echolocating bats. *Trends in Ecology and Evolution* 4:160–166.
- O’Farrell, M., and W. Gannon. 1999. A comparison of acoustic versus capture techniques for the inventory of bats. *Journal of Mammalogy* 80:24–30.
- Obrist, M. K. 1995. Flexible bat echolocation: the influence of individual, habitat and conspecifics on sonar signal design. *Behavioral Ecology and Sociobiology* 36:207–219.
- Oswald, J., J. Barlow, and T. Norris. 2003. Acoustic identification of nine delphinid species in the eastern tropical Pacific Ocean. *Marine Mammal Science* 19:20–37.
- Parker, T. A. 1991. On the use of tape recorders in avifaunal surveys. *Auk* 108:443–444.
- Parsons, K. N., G. Jones, and F. Greenaway. 2003. Swarming activity of temperate zone microchiropteran bats: effects of season, time of night and weather conditions. *Journal of Zoology* 261:257–264.

- Parsons, S., A. Boonman, and M. Obrist. 2000. Advantages and disadvantages of techniques for transforming and analyzing chiropteran echolocation calls. *Journal of Mammalogy* 81:927–938.
- Parsons, S., and M. K. Obrist. 2004. Recent methodological advances in the recording and analysis of chiropteran biosonar signals in the field. Pages 458–477 in J. A. Thomas, C. F. Moss, and M. Vater, editors. *Echolocation in bats and dolphins*. University of Chicago Press, Chicago, IL.
- Parsons, S., and J. M. Szewczak. 2009. Detecting, recording, and analyzing the vocalizations of bats. Pages 91–111 in T. H. Kunz and S. Parsons, editors. *Ecological and behavioral methods for the study of bats*, 2nd edition. Johns Hopkins University Press, Baltimore, MD.
- Payne, K. B., M. Thompson, and L. Kramer. 2003. Elephant calling patterns as indicators of group size and composition: the basis for an acoustic monitoring system. *African Journal of Ecology* 41:99–107.
- Pettersson, J. 2004. The properties of sound and bat detectors. in R. Brigham, E. Kalko, G. Jones, S. Parsons, and H. J. G. A. Limpens, editors. *Bat Echolocation Research: tools, techniques and analysis*. Bat Conservation International. Austin, TX.
- Reby, D., J. Joachim, J. Lauga, S. Lek, and S. Aulagnier. 1998. Individuality in the groans of fallow deer (*Dama dama*) bucks. *Journal of Zoology* 245:79–84.
- Riede, K. 1993. Monitoring biodiversity: analysis of Amazonian rainforest sounds. *Ambio* 22:546–548.

Riede, K. 1998. Acoustic monitoring of Orthoptera and its potential for conservation.

Journal of Insect Conservation 2:217–223.

Rollins, K. E., D. K. Meyerholz, G. D. Johnson, A. P. Capparella, and S. S. Loew. 2012.

A forensic investigation into the etiology of bat mortality at a wind farm: barotrauma or traumatic injury? *Veterinary Pathology* 49:362–371.

Russ, J. M., M. Briffa, and W. I. Montgomery. 2003. Seasonal patterns in activity and

habitat use by bats (*Pipistrellus* spp. and *Nyctalus leisleri*) in Northern Ireland, determined using a driven transect. *Journal of Zoology* 259:289–299.

Sauer, J., J. Fallon, and R. Johnson. 2003. Use of North American Breeding Bird Survey

data to estimate population change for bird conservation regions. *Journal of Wildlife Management* 67:372–389.

Sherwin, R., W. Gannon, and S. Haymond. 2000. The efficacy of acoustic techniques to

infer differential use of habitat by bats. *Acta Chiropterologica* 2:145–153.

Simmons, J. A., and R. A. Stein. 1980. Acoustic imaging in bat sonar: echolocation

signals and the evolution of echolocation. *Journal of Comparative Physiology A* 135:61–84.

Simmons, N. B. 2005. Order Chiroptera. Pages 312–529 in D. Wilson and D. Reeder,

editors. *Mammal species of the world*, Vol 1, 3rd edn. Johns Hopkins University Press, Baltimore, MD.

- Skalak, S. L., R. E. Sherwin, and R. M. Brigham. 2012. Sampling period, size and duration influence measures of bat species richness from acoustic surveys. *Methods in Ecology and Evolution* 3:490–502.
- Somervuo, P., A. Harma, and S. Fagerlund. 2006. Parametric representations of bird sounds for automatic species recognition. *IEEE Transactions on Speech, and Language Processing* 14:2252–2263.
- Stahlschmidt, P., and C. A. Brühl. 2012. Bats as bioindicators – the need of a standardized method for acoustic bat activity surveys. *Methods in Ecology and Evolution* 3:503–508.
- Sutherland, W. J. (Ed.). 2006. *Ecological census techniques*. 2nd edition. Cambridge University Press, New York. 1–432.
- Taylor, A., G. Watson, G. Grigg, and H. McCallum. 1996. Monitoring frog communities: an application of machine learning. *Proceedings of Eighth Innovative Applications of Artificial Intelligence Conference, Portland Oregon*:1564–1569.
- U.S. Fish and Wildlife Service. 2012. Land-based wind energy guidelines. URL: http://static.whitenosesyndrome.org/sites/default/files/white-nose_syndrome_national_plan_may_2011.pdf.
- Vaughan, N., G. Jones, and S. Harris. 1997. Habitat use by bats (Chiroptera) assessed by means of a broad-band acoustic method. *Journal of Applied Ecology* 34:716–730.

- Walsh, A. L., C. Catto, T. Hutson, P. Racey, P. Richardson, and S. Lanston. 2001. The UK's National Bat Monitoring Programme. Final Report 2001. London, England.
URL: www.bats.org.uk/publications_download.php/335/NBMP01.pdf.
- Waters, D. A., and A. L. Walsh. 1994. The influence of bat detector brand on the quantitative estimation of bat activity. *Bioacoustics* 5:205–221.
- Wickramasinghe, L. P., S. Harris, G. Jones, and N. Vaughan. 2003. Bat activity and species richness on organic and conventional farms: impact of agricultural intensification. *Journal of Applied Ecology* 40:984–993.

Chapter 2

2 Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats¹

2.1 Introduction

Echolocation provides a window through which the behavior and ecology of bats can be evaluated. Specifically, calls used by echolocating bats can be conspicuous to bat detectors, permitting biologists to distinguish among species by their calls and to identify foraging activity. Bat detectors, instruments sensitive to the acoustic frequencies dominating bat calls, have been extensively used in a range of bat studies, from those investigating echolocation behavior, to others documenting patterns of distribution and activity levels. By 2012, the variety of commercially available bat detectors offered a spectrum of features at a range of prices (e.g., weatherproofing, temperature sensors, storage options; Table 2.1) but key features, such as microphone quality, sampling rate, and recording technology will determine the ability to detect bats. Many published articles have used data from bat detectors to address questions about the echolocation behavior of bats, as well as their patterns of activity and habitat use (e.g., Gillam 2007, Collins and Jones 2009, Müller *et al.* 2012).

¹ A version of this chapter has been published and is presented here with permission from John Wiley and Sons.

Citation: Adams, A. M., M. K. Jantzen, R. M. Hamilton, and M. B. Fenton. 2012. Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats. *Methods in Ecology and Evolution* 3:992–998.

Table 2.1. A comparison of the features associated with commercially available ultrasonic bat detector systems.

System and manufacturer feature	AnaBat SD2 Titley Electronics	Avisoft UltraSoundGate 116 w CM16/CPMA Avisoft Bioacoustics	Batcorder 2.0 ecoObs	Batlogger Elekon AG	Song Meter SM2BAT 192 kHz Wildlife Acoustics
Recording technology	Zero-crossing	16-bit, full-spectrum	16-bit, full-spectrum	16-bit, full-spectrum	16-bit, full-spectrum
Sampling rate	N/A	500 kHz	500 kHz	312.5 kHz	192 kHz (384 kHz available)
Sound file type	AnaBat	.wav	.raw	.wav & .xml	.wav & .wac
Storage type	Compact Flash (CF)	External through computer	SDHC	SDHC	SDHC x 4
Storage capacity	128 GB	unlimited	16 GB	32 GB	128 GB
Battery	4 AA batteries	Runs off computer	NiMH 6V 2700mA, rechargeable	LIB 3.7V 4600mAh, rechargeable	4 D batteries
Microphone type	Condenser	Condenser	Electret	Electret	Electret
Omnidirectional microphone?	No	No	Yes	Yes	Yes
Recording schedule?	Yes, through CF Reader	Yes, through Avisoft-RECORDER software	Yes	Yes	Yes
Post-process tools	AnaLook	Avisoft-SASLab Pro	None	BatExplore	Batch noise scrubber, zero-cross converter, Wac2Wav converter
Channels	1	1	1	1	2
Weatherproof enclosure?	No	No	Yes	with StrongBox	Yes
Weatherproof microphone?	No	No	No	No	Yes
GPS?	Can connect externally	No	No	Yes	optional
Temperature sensor?	Internal temperature	No	No	External temperature	Internal and external
Price (USD)	\$2,200†	\$5999*	\$3,273*‡	\$2,035*‡	\$999*

Pricing from company *website or †manufacturer. ‡Price converted to USD.

Acoustic sampling is a common, powerful technique for monitoring the activity of echolocating bats. Bat detectors are widely used by researchers, including those working for government agencies, environmental consulting firms, and academics. Behavioral, presence/absence, and relative abundance data are commonly collected with these devices. The results of research relying on bat detectors inform our understanding of bat ecology and behavior and are frequently used to guide important wildlife management decisions (U.S. Fish and Wildlife Service 2012). Acoustic monitoring is non-intrusive and capable of recording large quantities of data. However, the specific combination of hardware may affect the quality, precision, and quantity of data collected.

Variation in microphone sensitivity and detection algorithms can produce data sets that differ among detectors. Both Downes (1982) and Fenton (2000) noted significant differences in detection sensitivity among brands of narrowband and broadband acoustic detectors. This variation has the potential to affect acoustic monitoring studies and their conclusions; whether the focus is curiosity-driven research or environmental assessments where low bat activity is assumed to equal low numbers of bats and therefore low risk (U.S. Fish and Wildlife Service 2012).

The acoustic nature of bats is highly variable (e.g., frequency, intensity, etc.) which can influence detectability by even the 'best' detectors. Bats using low-intensity echolocation calls dominated by higher frequencies are less detectable than those using high-intensity calls dominated by lower frequencies. Higher-frequency sounds attenuate more quickly and will be detected less frequently than higher intensity, lower-frequency calls (Lawrence and Simmons 1982), resulting in under-representation of these species in

acoustic surveys (Murray *et al.* 2009). Detection bias will be further compounded by the sensitivity and frequency response of the bat detector. Different systems vary in their performance over the range of biologically relevant frequencies. If the microphone has lower sensitivity to high frequencies, the bias caused by atmospheric attenuation will be further exaggerated. The consequences of detection bias will depend on the community being studied; the frequencies of bat calls range from ~8 kHz to > 200 kHz, and this range varies with a given bat community. Researchers must consider the community in question when choosing the most appropriate bat-detecting system for their research (Limpens and McCracken 2004).

Microphones with lower sensitivity will detect bats at shorter distances relative to more sensitive microphones. Detectors with shorter detection ranges will sample a smaller airspace and thus have a lower probability of detecting any bats present. Also, not all detectors are equal in their directionality and the orientation of the detector in relation to the bat affects detection (Britzke *et al.* 2010). When all other factors are equal, detectors with omnidirectional microphones will have a better chance of detecting a bat, compared to more directional microphones. However, a less directional microphone will be less sensitive, giving it a smaller detection range (Limpens and McCracken 2004). The smaller the microphone, the more omnidirectional it will be.

Three levels of variation can confound data acquired with bat detectors. First is the variation associated with the movement of sound through air. Second is that intrinsic to the instruments. Third is variation in echolocation behavior and call design among bats. Whether the focus of a study is echolocation behavior or documenting patterns of habitat use, it is important to distinguish between factors two and three. We presented synthetic

acoustic signals and echolocation calls of free-flying bats in the wild to compare ultrasonic call detection by five commercially available bat detectors. Our goal was to provide data about relative bat detector performance and bat echolocation behavior.

With an increasing number of commercially available bat detectors, it is important to address variation in the technologies. A fundamental factor of any methodology is addressing the capabilities and limitations of the equipment being used. It is vital to be aware of the differences that may result from the use of different equipment even when the same sampling method is employed. To date, no study has examined the differences in the detection efficacy among direct high-speed bat detector models.

2.2 Methods

We simultaneously deployed five direct high-speed bat detectors for recording both synthetic playback and free-flying bats: AnaBat SD2 (Titley Scientific, Ballina, NSW, Australia), Avisoft UltraSoundGate 116 CM16/CMPA (Avisoft Bioacoustics, Berlin, Germany), Batcorder 2.0 (ecoObs, Nuremberg, Germany), Batlogger (Elekon AG, Luzern, Switzerland), and Song Meter SM2BAT (Wildlife Acoustics, Inc, Concord, MA). There are several other commercially available detectors that we were unable to include in this study, for example D500X and D1000X (Pettersson Elektronik) and AR125 (Binary Acoustic Technology). During all trials, microphones were within 10 cm of each other, on a parallel plane. Microphone order and position were rearranged randomly for each trial to change microphone position, but maintain consistent microphone spacing. We avoided variation by recording with only one detector of each model and recording with all detectors at the same time.

2.2.1 Optimizing detector recording settings

We used playback of synthetic signals to optimize detection settings for each system. Our synthetic signal file was 1478 ms in duration, and consisted of 20, 57 ms long, constant frequency (CF) signals, five signals at each of four frequencies: 25, 55, 85, and 115 kHz. For playback, we used a laptop running Avisoft RECORDER-NiDAQmx software connected to an ultrasonic playback interface with an integrated D/A power amplifier (UltraSoundGate Player 116). The interface was connected to an UltraSoundGate Dynamic Speaker ScanSpeak (hardware and software: Avisoft Bioacoustics, Berlin, Germany), which we did not calibrate. When possible, we recorded with all combinations of setting configurations for each detector. When combinations were prohibitively large (>100) we recorded in intervals spanning the full range of configurations. For each configuration, we played synthetic signals 5 m from each device. We analyzed each recording visually to find the optimum settings for recording conditions. In cases where multiple configurations were equal, we chose the settings closest to the default settings for the detector. These settings were used for the remainder of our experiments (Table 2.2).

Table 2.2. Detector settings used in this study.

AnaBat SD2	Avisoft UltraSoundGate 116	Batcorder 2.0	Batlogger	Song Meter SM2BAT
Gain: 7 Data Div - 16	Gain: 7 Trigger: permanent (continuous) Sampling rate: 500 kHz Format: 16 bit Buffer: 0.050 No. Buffers: 4	Critical frequency: 14 kHz Threshold: -36 dB Post trigger: 800 ms Quality: 40	Crest minCrest: 5 minRMS: 2 minPeak: 5 HighPass: 6	Sampling rate: 192 kHz Compression: WAC0 Gain: 36 dB Dig HPF: fs/16 Dig LPF: Off Trigger Level: 15 SNR Trigger Win Right: 1 s Div Ratio: 16

See each respective detector manual (available online) for the setting description.

2.2.2 Synthetic call playback

We played the synthetic CF signals three times at 5 m intervals (5 – 40 m) and three angles (0°, 45°, 90°) in an open field. This resulted in 15 calls of each frequency played at each distance and angle (24 combinations). We used the automated detection feature (Table 2.3) of callViewer (v. 18, Skowronski and Fenton 2008), to count the number of calls detected by each system and manually inspected each recording to ensure that there were no false positives. CallViewer is a custom echolocation sound analysis program written with MATLAB software (The MathWorks, Natick, Massachusetts). Because AnaBat file formats are not compatible with callViewer software, we visually inspected these recordings in AnaLook (v. 3.8, Titley Electronics, Ballina, Australia). We used general linear models to analyze the number of signals detected (considering each frequency separately) with angle, detector, distance and all two-way interactions. To compare among detectors we generated pair-wise comparisons of the estimated marginal means, controlling for the effect of distance and angle. We used a similar approach to compare the effect among the three angles. We estimated the detection range by modeling the probability of detection of each signal frequency at each angle by all detectors with a logistic regression in PASW18 (SPSS Inc., Chicago, IL). From the fitted logistic regression we determined the distance corresponding to a detection probability of 0.50 as our estimate of detection range (i.e., beyond this distance there is less than a 50% chance that the signal would be detected).

Table 2.3. Automated detection parameter settings used for call analysis in callViewer.

Parameter	Setting
Minimum link length	10
Window length (ms)	0.3
Frame rate (fps)	10000
Chunk size (sec)	1
Minimum energy (dB)	14
Echo filter threshold (dB)	10
UPPER cutoff freq. (kHz)	Inf
LOWER cutoff freq. (kHz)	15
Window type	Blackman
Delta size (+/- frames)	1

2.2.3 Recording free-flying bats

Free-flying bats produce complex, frequency-modulated calls that vary in intensity in contrast to the simple, constant-frequency signals we used for the synthetic playback experiment. To introduce the variability that is present in natural settings we recorded free-flying bats. We deployed the detectors for two hours per night on three separate nights in a suburban area in London, Ontario, Canada. The Avisoft system detected more bat echolocation calls than any of the other detectors so we used the data from it as a baseline. We chose 26 easily identifiable passes (minimum seven consecutive calls), from hoary bats (*Lasiurus cinereus*), and counted the number of calls in each pass. We manually counted all calls recorded regardless of call quality or completeness. We used callViewer to analyze the full spectrum system calls and AnaLook to analyze calls from AnaBat. We calculated the proportion of calls detected per pass relative to Avisoft, arcsine-square root transformed the data, and compared detector performance with ANOVA and Tukey's post hoc test in PASW18 after finding no effect of recording night.

2.3 Results

2.3.1 Synthetic call playback

Overall, Avisoft detected the most signals (1067 signals, 25% of all signals presented), and AnaBat detected the fewest (240 signals, 5% of all presented). Avisoft was the only system that detected the 115 kHz signal and only at 5 m (Fig. 2.1A). AnaBat did not detect CF signals at 85 kHz and 115 kHz (Fig. 2.1E). The other detectors only recorded 85 kHz signals at 5 m, except Avisoft which recorded these signals at 10 m (Fig. 2.1). All

systems detected the 55 kHz signals, but detection range varied from 7 m to 16 m at 0° (Fig. 2.2). Song Meter did not detect 115 kHz signals because the frequency is outside of this model's detection capabilities. A model with a higher sampling frequency is available and would likely have detected higher frequency signals.

The number of signals detected at 25 kHz varied significantly among detectors ($F_{4,348} = 21.32$, $p < 0.001$; Fig. 2.3), except Batcorder and Song Meter. AnaBat recorded the fewest 25 kHz CF signals. There were also differences among detectors in the number of 55 kHz signals detected ($F_{4,346} = 22.74$, $p < 0.001$; Fig. 2.3); Avisoft recorded more than Song Meter and AnaBat, while Batcorder recorded more than AnaBat. Batlogger recorded significantly more signals than any other detector for at 25 kHz and 55 kHz.

There was a significant interaction between detector and distance for both 25 kHz and 55 kHz signals ($F_{4,348} = 9.42$, $p < 0.001$; $F_{4,346} = 13.63$, $p < 0.001$; Fig. 2.1). For 25 kHz, Batcorder and Song Meter detections reflected a greater rate of attenuation with distance than AnaBat, Avisoft, and Batlogger. For 55 kHz, AnaBat had the greatest rate of attenuation with distance and Batlogger had the lowest (Fig. 2.1).

Overall, there was an effect of angle for both 25 kHz and 55 kHz signals ($F_{2,348} = 24.92$, $p < 0.001$; $F_{2,346} = 21.06$, $p < 0.001$; Fig. 2.1); the number of signals detected declined as the angle increased. The effect of angle was the same among all detectors ($p > 0.05$). There was no interaction between angle and distance for 25 kHz signals ($p > 0.05$), but there was an interaction for 55 kHz signals ($F_{2,346} = 12.62$, $p < 0.001$). For 55 kHz signals, there was no difference between 0° and 45°, but these two angles had a greater rate of decline in number of signals over distance than 90°.




Figure 2.1. Mean number of calls detected by each bat detector system at four frequencies at each distance and angle during the synthetic playback experiment. There were 15 calls played for each frequency/distance/angle combination.




Figure 2.2. Distance of 50% probability of detection calculated with a logistic regression for each frequency at 0° by each bat detector system during the synthetic playback experiment. Patterns were similar for all detectors at 45° and 90° , but with lower overall probability of detection.




Figure 2.3. Performance varied among detectors with a strong effect of frequency. Call detection (arcsine square root transformed number of calls) \pm SE by call frequency evaluated at a distance of 22.5 m. Detectors with the same letter superscript were not significantly different from each other within each frequency.

2.3.2 Recording free-flying bats

Batlogger recorded significantly more hoary bat echolocation calls (relative to Avisoft) than any other system ($F_{3, 100} = 45.26$, $p < 0.001$; Fig. 2.4), while AnaBat, Batcorder, and Song Meter did not differ significantly from each other. Only AnaBat and Batcorder failed to detect all 26 passes; both of these systems did not record any calls from two passes. One of the 26 passes included a feeding buzz that was recorded by all of the detectors. Avisoft, Batcorder, Batlogger, and Song Meter recorded more calls (23 – 25 calls) in the feeding buzz than AnaBat (11 calls).




Figure 2.4. Mean number of calls \pm SE per pass relative to Avisoft for each bat detector from recordings of free-flying *Lasiurus cinereus* on three nights. Batlogger detected more calls than any of the other systems (detectors with the same letter superscript were not significantly different from each other).

2.4 Discussion

Our results demonstrate that there is significant variation in detection efficacy among commercially available bat detectors. The differences in the detection abilities of these microphones, particularly in relation to differing frequency sensitivity, illustrate the hazards of comparing data collected by different detecting systems. Our results show that detection of different frequencies varied among detector systems and was affected by the distance and angle of the signal from the detector. Avisoft and Batlogger detected more of the highest frequency signals we tested than the other detectors, but as expected, these signals were detected at much shorter ranges. Detection distance for the 55 kHz synthetic signals (detected by all systems) is particularly relevant because this frequency is in the range of most species of bats that occur in temperate regions. In Hawaii, where only one species of bat occurs (*L. cinereus semotus*), any of the systems we used would suffice, although each would provide quite a different view of bat activity. In Newfoundland, where two species occur (*Myotis lucifugus*, *M. septentrionalis*) any of the systems we tested would suffice for *M. lucifugus* (echolocation call frequency of most energy ~40 kHz, maximum frequency ~81 kHz), but only some would accurately document activity by *M. septentrionalis*, which uses calls dominated by higher frequencies (frequency of most energy ~60 kHz, maximum frequency ~126 kHz; (Faure *et al.* 1993, Ratcliffe and Dawson 2003). In Newfoundland, some systems would be better than others. In other parts of the world, some bat species use echolocation calls dominated by frequencies >85 kHz. For these bat communities, the detection distance of the 85 kHz synthetic signals in our study is important to consider. Monitoring the activity of vespertilionid bats in the

subfamilies Kerivoulinae and Murininae would be difficult with any of the systems we tested because these species produce high frequency (80 – 200 kHz), frequency-modulated sweeps.

Variation in detection distance among detectors has important practical implications. For many studies, it is particularly important to understand the volume of airspace being sampled, such as when interpreting the results of pre-construction acoustic surveys conducted at potential wind energy facility sites where high bat mortality is a concern (Kunz *et al.* 2007). On modern wind turbines, the lower edge of the blade swept area is ~20 m above-ground (Barclay *et al.* 2007). Our data demonstrate detection ranges of 7 – 16 m, and therefore, none of the ground-based microphone systems we tested can detect bats flying in the area swept by the blades of wind turbines. Even a detector placed on the nacelle of a turbine (in the center of the blade swept area) would sample no more than one-third of the area swept by 50 m long blades (Kunz *et al.* 2007).

When we focus on detection of echolocation calls from free-flying bats, bat detectors fell into one of two performance groups. AnaBat, Batcorder, and Song Meter did not differ significantly in the number of hoary bat echolocation calls detected. These bats produce high intensity echolocation calls with a minimum frequency which is typically ~17 kHz (Obriest 1995). The minimum frequency of hoary bat calls is lower than the lowest frequency of our synthetic calls. Consequently, our free-flying bat results represent a best-case scenario; we used only high intensity, low-frequency calls and our sampling method, counting all calls regardless of quality, presented the most optimistic view of activity. In reality, many species are much less detectable and the quality of many recorded calls is too poor to be identified to species or counted as a bat call. Using

automated detection algorithms with recording quality standards will provide more objective call counts when measuring activity. If we had looked at passes from any of the Ontario *Myotis* species (calls with a minimum frequency range of ~34 - 40 kHz; Thomas *et al.* 1987), it is likely that the results from our free-flying passes would have mirrored the results from our synthetic call trials.

Among the detectors we tested, AnaBat is unique in that it is the only detector to use zero-crossing analysis which may (Corben and Fellers 2001) or may not (Fenton 2000) provide an adequate picture of bat activity. Our data contributes to this discussion, demonstrating that AnaBat is capable of performing similarly to a full-spectrum detector (Fig. 2.4), but in most cases it detects fewer calls (Fig. 2.3). Therefore, we emphasize the importance of considering the research questions and local bat fauna. While our results from the synthetic-call trials agree that full-spectrum detectors are more sensitive, our free-flying bat trial showed that there are circumstances where the differences are not substantial. Ultimately, the specific hypotheses and objectives of a study will dictate the suitability of various detectors (Limpens and McCracken 2004). No one recording system is ideal for all situations and thus it is the responsibility of the researcher (and the reader) to consider how the performance of the recording system will impact the results and conclusions of the study.

It is important to note that regardless of recording system, all microphones detect only a subset of the calls present in the environment (e.g. in our playback experiment the best system detected only 25% of the calls we played). However, our findings show that some subsets are significantly larger than others. This discrepancy is essential to remember when attempting to compare datasets collected with different detecting systems. Even

when comparing multiple detectors of the same model, the microphones must be calibrated to ensure comparable performance (Larson and Hayes 2000). With an increasing number of threats to bat populations (e.g., wind turbines, white-nose syndrome) there may be a drive to develop more rigorous monitoring programs with standardized protocols for bat surveys. Our results highlight the importance of considering the specific detector used, and the variation that may arise from different microphones.

As technology continues to evolve, the number of commercially available detectors will increase. As with the current proliferation in detectors on the market, many brands will persist (e.g., AnaBat, Avisoft) and new brands will emerge (e.g., Batlogger). In such a specialized market there will probably be few dramatic changes in the technology; we would expect to see increases in microphone sensitivity, battery life, and storage capacity, along with continued software upgrades to improve detection algorithms. With a high diversity of detectors, each with a wide range of settings and technical capabilities, it is now necessary to report not only the type of detector used, but also the settings chosen (e.g., Table 2.2) and as many hardware details as possible. The extent that detector-specific settings have on performance and accuracy between detectors of the same brand remains to be seen. Finally, it comes to the issue of comparability of results; different detectors will give different results, which must be taken into account.

Whether the bat-detecting system you are using hears the same signals as the one I am using depends upon the echolocation calls. There are numerous factors that contribute to variation in datasets from acoustic monitoring; our results demonstrate that the detector plays a role in this variation. Ultimately, it is crucial that differences in detector

performance be considered when designing studies and comparing results from different detectors, whether among models included in our study, other extant models, or those yet to be invented. No detector is ideal for all research questions and methods, and conversely, not all detectors are appropriate for a given question or methodology.

2.5 Acknowledgments

We thank L. P. McGuire, E. E. Fraser, J. F. Miller, and A. Cameron for their support and input for this study. Thanks to L. Lazure and M. D. Skowronski for creating the synthetic calls. D. Morningstar, M. Gumprich, M. K. Obrist, R. M. R. Barclay, and E. F. Baerwald generously allowed us to use their equipment.

2.6 References

- Barclay, R. M. R., E. F. Baerwald, and J. C. Gruver. 2007. Variation in bat and bird fatalities at wind energy facilities: assessing the effects of rotor size and tower height. *Canadian Journal of Zoology* 85:381–387.
- Britzke, E. R., B. A. Slack, M. P. Armstrong, and S. C. Loeb. 2010. Effects of orientation and weatherproofing on the detection of bat echolocation calls. *Journal of Fish and Wildlife Management* 1:136–141.
- Collins, J., and G. Jones. 2009. Differences in bat activity in relation to bat detector height: implications for bat surveys at proposed windfarm sites. *Acta Chiropterologica* 11:343–350.

- Corben, C., and G. Fellers. 2001. Choosing the “correct” bat detector—a reply. *Acta Chiropterologica* 3:245–256.
- Downes, C. 1982. A comparison of sensitivities of three bat detectors. *Journal of Mammalogy* 63:343–345.
- Faure, P. A., J. H. Fullard, and J. W. Dawson. 1993. The gleaning attacks of the northern long-eared bat, *Myotis septentrionalis*, are relatively inaudible to moths. *Journal of Experimental Biology* 178:173–89.
- Fenton, M. B. 2000. Choosing the “correct” bat detector. *Acta Chiropterologica* 2:215–224.
- Gillam, E. H. 2007. Eavesdropping by bats on the feeding buzzes of conspecifics. *Canadian Journal of Zoology* 85:795–801.
- Kunz, T., E. Arnett, W. P. Erickson, A. Hoar, G. Johnson, R. P. Larkin, M. D. Strickland, R. Thresher, and M. Tuttle. 2007. Ecological impacts of wind energy development on bats: questions, research needs, and hypotheses. *Frontiers in Ecology and Environment* 5:315–324.
- Larson, D. J., and J. P. Hayes. 2000. Variability in sensitivity of Anabat II bat detectors and a method of calibration. *Acta Chiropterologica* 2:209–213.
- Lawrence, B. D., and J. A. Simmons. 1982. Measurements of atmospheric attenuation at ultrasonic frequencies and the significance for echolocation by bats. *Journal of the Acoustical Society of America* 71:585–90.

- Limpens, H. J. G. A., and G. F. McCracken. 2004. Choosing a bat detector: theoretical and practical aspects. Pages 28–37 in R. M. Brigham, E. K. V Kalko, G. Jones, and H. J. G. A. Limpens, editors. *Bat Echolocation Research: tools, techniques, and analysis*. Bat Conservation International, Austin, TX.
- Müller, J., M. Mehr, C. Bässler, M. B. Fenton, T. Hothorn, H. Pretzsch, H.-J. Klemmt, and R. Brandl. 2012. Aggregative response in bats: prey abundance versus habitat. *Oecologia* 169:673–684.
- Murray, K. L., E. Fraser, C. Davy, T. H. Fleming, and M. B. Fenton. 2009. Characterization of the echolocation calls of bats from Exuma, Bahamas. *Acta Chiropterologica* 11:415–424.
- Obrist, M. K. 1995. Flexible bat echolocation: the influence of individual, habitat and conspecifics on sonar signal design. *Behavioral Ecology and Sociobiology* 36:207–219.
- Ratcliffe, J. M., and J. W. Dawson. 2003. Behavioural flexibility: the little brown bat, *Myotis lucifugus*, and the northern long-eared bat, *M.septentrionalis*, both glean and hawk prey. *Animal Behaviour* 66:847–856.
- Skowronski, M. D., and M. B. Fenton. 2008. Model-based automated detection of echolocation calls using the link detector. *Journal of the Acoustical Society of America* 124:328–336.

Thomas, D., G. Bell, and M. Fenton. 1987. Variation in echolocation call frequencies recorded from North American vespertilionid bats: a cautionary note. *Journal of Mammalogy* 68:842–847.

U.S. Fish and Wildlife Service. 2012. Land-based wind energy guidelines. URL: http://www.fws.gov/windenergy/docs/WEG_final.pdf.

Chapter 3

3 How high is high? Using percentile thresholds to identify peak bat activity²

3.1 Introduction

Researchers passively monitor bats by eavesdropping on their echolocation calls (Hooper 1966), providing insight into many aspects of ecology and behavior (e.g., Fenton 2003, Neuweiler 1989). Determining the timing and location of peak activity levels is important for an understanding of ecology and behavior, and has management implications.

Activity levels of bats can vary dramatically temporally with increased activity during certain life history events or at ecologically relevant sites, such as riparian foraging areas (e.g., Rautenbach *et al.* 1996, Hayes 1997), maternity colonies (e.g., Murray and Kurta 2004), migration stopover sites (e.g., Barclay 1984, Dzal *et al.* 2009), or pre-hibernation swarming sites (Parsons *et al.* 2003). Conversely, areas used for commuting or dispersed foraging are likely to have lower activity levels.

There is no reliable way to convert the number of echolocation calls or passes recorded into the number of bats present, so acoustic monitoring only provides relative indications of low or high bat activity. However, these classifications are subjective and there is no framework or guideline with respect to interpreting activity levels, beyond activity in one sample being relatively higher than another (Kunz *et al.* 2007b). Previously, several

² A version of this chapter is in review.

Citation: Adams, A. M., L. P. McGuire, L. A. Hooton, and M. B. Fenton. How high is high? Using percentile thresholds to identify peak bat activity. *Acta Chiropterologica*: in review.

methods have been used to identify peaks of activity, including visual identification with activity plots (Hayes 1997), choosing an arbitrary level of activity as ‘high’ (Broders 2003, Brooks and Ford 2005), or calculating the number of calls that are above a certain threshold (Gorresen *et al.* 2009, Hamilton 2012). Since acoustic monitoring provides a relative indication of bat activity it is important to define objective criteria for making comparisons among sites or time periods.

Our goal was to determine a method for objectively identifying peak bat activity with the purpose of examining the use of percentile thresholds for this task. Percentile thresholds have been used to identify high occupancy sites and make inferences about habitat attributes (Gorresen *et al.* 2009). Percentiles are simply 100 regular intervals in any cumulative distribution with the median at the 50th percentile where half of all observations fall below this threshold. Percentile thresholds are not affected when data differs in dispersion pattern, and are less susceptible to outliers than traditional statistics, such as analysis of variance (Sokal and Rohlf 1981). When applied to acoustic bat activity, the 99th percentile represents the most infrequent activity levels, with the greatest number of calls, where these periods of high activity only occur 1% of the time (e.g., Fig. 1). Ecological data are often overdispersed (O’Hara and Kotze 2010), with many observations with few calls and few with many calls, which violates key assumptions of common statistical methods. To illustrate the use of percentile thresholds, we used acoustic recordings from several sites in Canada. The aims of this paper are: (1) explore the suitability of percentile thresholds for identifying peak bat activity within- and among-nights and (2) compare the use of percentile thresholds to traditional statistical procedures.

3.2 Methods

3.2.1 Study areas

Data were recorded at (i) Long Point, Ontario, a 35 km long peninsula extending from the north shore of Lake Erie, for six nights during both spring (June 2008) and autumn migration (August/September 2008); (ii) an abandoned mine near Renfrew, Ontario for five nights during both spring (May/June 2008) and autumn swarming (August 2008); and (iii) along the Battle Creek in Cypress Hills Interprovincial Park, Saskatchewan for five nights during summer (July 2009).

We considered three species in our analysis, *Lasiurus borealis*, *L. cinereus*, and *Myotis lucifugus*. Given the migratory nature of *L. borealis* and *L. cinereus* we predicted that we would observe high activity in autumn at Long Point as it is an important site for bats during migratory periods (Dzal *et al.* 2009, McGuire *et al.* 2012). Cypress Hills is a forested region where all three species occur (Willis and Brigham 2003, 2005), although *L. borealis* is rare (Willis and Brigham 2003) and we expected this site to be the least important to *L. borealis*. *Myotis lucifugus* is known to swarm and hibernate at Renfrew (Fenton 1969), therefore we expected to observe high activity only during the autumn swarming period.

3.2.2 Acoustic monitoring and analysis

We recorded continuously from dusk until dawn at all locations, on nights with no rain, using externally polarized condenser microphones (Avisoft CM16/CMPA) connected to an Avisoft UltraSoundGate 416-200 or UltraSoundGate 116 (Avisoft Bioacoustics,

Berlin, Germany) at 8 bit with a 250 kHz sampling rate, and gain at seven. The system was operated with Avisoft Recorder USG software.

We identified echolocation calls in all files using the automated detection feature in callViewer (v. 18; Skowronski 2008), a custom sound analysis program written with MATLAB software (The MathWorks, Natick, Massachusetts). We filtered the data to eliminate noise and weak or fragmented calls, only including detections with duration 0.99 – 30 ms and minimum frequency (Fmin) 15 - 60 kHz. The filter parameters were selected based on conservative estimates of the echolocation call structure of the bat species present at our recording sites. We identified calls to species using quadratic discriminant function analysis (DFA, Appendix A), which compared our unidentified data to a training dataset that included seven species: *Eptesicus fuscus*, *Lasionycteris noctivagans*, *L. borealis*, *L. cinereus*, *M. lucifugus*, *M. septentrionalis*, and *Perimyotis subflavus*. All species were weighted equally in the classification analysis. Classification of each call was based on 11 call parameters extracted by the automated detection feature of callViewer (Skowronski and Fenton 2008). Cross-validation indicated the species classification accuracy was greater than 88% for the three species included in our analysis (Table A1). To further improve classification accuracy, and because DFA does not assign calls to an “unknown” category, we applied a post-hoc, species-specific filter to remove any data that were above or below typical durations and minimum frequencies for each species (Table A2).

3.2.3 Statistical analysis

The sampling unit in our analyses was the number of calls of a given species recorded each hour. We also summed the calls per hour, of the seven species post-DFA and

filtering, for total number of calls per hour for all species combined. We calculated percentile thresholds of activity for *L. borealis*, *L. cinereus*, *M. lucifugus*, and all species combined based on the distribution of the number of calls per hour for all nights regardless of site. We considered six thresholds, the 25th, 50th (median), 70th, 90th, 95th, and 99th percentiles, for each species and all species combined. Passive acoustic monitoring often includes many time periods with no echolocation calls recorded, resulting in issues of zero-inflation in statistical analysis (McCullagh and Nelder 1989); to avoid these issues, we excluded time periods when no bats were recorded, thus framing our analysis in terms of ranking activity given the presence of bats. We recognize that zero calls per hour is informative about activity levels at a site and differs from non-data, but since thresholds are used as a measure of how high activity is it is not necessary to have a descriptor when they are not present. The absence of echolocation calls in acoustic recordings is unambiguous and therefore, does not need to be included in the definition of relative activity level thresholds. By creating a large dataset of recordings from a wide range of ecological situations we created a null distribution to generate percentile thresholds. We then compared results from a particular site to the percentile thresholds of the larger distribution. We counted the number of nights with at least one hour above each percentile threshold. We also calculated differences in number of calls per hour, including time periods with zero activity, among sites with Kruskal-Wallis ($\alpha = 0.05$) using *kruskal.test* in R (v. 2.13.1; R Development Core Team 2011) and a pair-wise post-hoc test (*kruskalmc* in *pgirmess* package in R; Giraudoux 2011) because data were positively skewed, where many hours contained few echolocation calls.

We used the previously calculated percentile thresholds to evaluate patterns of within-night activity, when peaks occurred and the degree of variation within a night, specifically for *L. borealis* at Long Point and *M. lucifugus* at Renfrew. We identified when activity was over the median to see if peaks occurred at dusk (first hour of the night), in the middle of the night, or at dawn (within one hour of sunrise), looking for patterns of unimodal, bimodal, constant, or irregular activity. To measure the degree of variation of activity levels within-nights, we calculated the proportion of the night with activity above or below the median, where constant activity would always be above or below the median.

3.3 Results

3.3.1 Among sites

We recorded for a total of 258 hours and for all seven species combined, after excluding samples when no calls were detected, the resulting samples size was 219 hours. Half of the time which contained fewer than 265 calls per hour (50th percentile), while the top 10% of activity ranged from 4504 to 28358 calls per hour (above the 90th percentile, Table 3.1, Fig. 3.1). All sites had activity levels above the 50th percentile, but Renfrew was the only site to have activity above the 90th percentile threshold and even exceeded the 99th percentile threshold (Table 3.2A) during the swarming period, when activity was significantly higher than all other sites ($H_4 = 32.01$, $p < 0.001$; Table. 3.2A).

Table 3.1. Number of calls per hour at each percentile threshold for each species of bat and all species combined for a distribution of activity from three sites in Canada. Half of the activity falls below the 50th percentile (median) and 10% of activity is above the 90th percentile threshold.

	Number of calls at percentile threshold					
	25 th	50 th	70 th	90 th	95 th	99 th
Species combined	100	265	493	4504	10616	19510
<i>Lasiurus borealis</i>	7	15	31	66	81	158
<i>L. cinereus</i>	20	63	115	266	396	534
<i>Myotis lucifugus</i>	25	88	314	5818	11865	19404




Figure 3.1. Frequency of activity levels for all bat species combined at three sites in Canada. Dashed, vertical lines are thresholds of the number of calls per hour based on the 50th, 70th, 90th, 95th, and 99th percentiles of the distribution. Activity over the 50th percentile threshold (median) is considered high activity because half of all detections had fewer than 265 calls per hour.

Lasiurus borealis was detected in 109 hours of recording, and half of the time these were brief passes with less than seven calls per hour (Table 3.1). Activity exceeded the 90th percentile at Long Point and Renfrew (swarming), while Long Point (migration) was the only site with activity levels that exceeded the 99th percentile threshold (Table 3.2B). Nightly activity was not significantly different among sites ($H_4 = 45.85$, $p = <0.001$, Table 3.2B).

Lasiurus cinereus was detected during 147 hours of recording. Long Point was the only site with activity above the 70th percentile threshold (Table 3.2C). Nightly activity varied among sites ($H_4 = 149.78$, $p < 0.001$) and was significantly higher at Long Point (migration) than Renfrew and Cypress Hills (Table 3.2C).

Myotis lucifugus was detected during 177 hours of recording. Activity was above the median at all sites, but only above the 90th percentile threshold at Renfrew (swarming). Activity was significantly higher at Renfrew (swarming; $H_4 = 75.02$, $p < 0.001$; Table 3.2D).

Table 3.2. Summary of bat activity, for A) all species combined, B) *Lasiurus borealis*, C) *L. cinereus*, and D) *Myotis lucifugus*, at three sites in Canada in 2008 and 2009. Activity over the 50th percentile threshold was considered high and the relative importance of a site was based on the threshold activity exceeded.

A) All species combined				Number of nights with at least one hour above percentile threshold					
Site	No. nights	Mean calls per hour \pm SD	Peak time of night (hour after sunset) \pm SD	25th	50th	70th	90th	95th	99th
Cypress Hills	5	231.2 \pm 309.9 b	1.6 \pm 0.5	3	3	3	0	0	0
Long Point spring	6	277.2 \pm 224.7 b	3.6 \pm 1.8	6	6	4	0	0	0
Long Point migration	6	246.1 \pm 237.5 b	2.8 \pm 1.9	6	4	3	0	0	0
Renfrew spring	5	484.3 \pm 1019.3 b	3.6 \pm 0.5	5	4	2	1	0	0
Renfrew swarming	5	5503.5 \pm 6910.9 a	4.0 \pm 0	5	5	5	5	4	2

B) <i>Lasiurus borealis</i>				Number of nights with at least one hour above percentile threshold					
Site	No. nights	Mean calls per hour \pm SD	Peak time of night (hour after sunset) \pm SD	25th	50th	70th	90th	95th	99th
Cypress Hills	5	4.6 \pm 12.6 b	2.8 \pm 1.3	4	3	3	0	0	0
Long Point spring	6	9.7 \pm 17.0 b	4.5 \pm 1.9	6	6	4	1	0	0
Long Point migration	6	25.4 \pm 39.9 a	4.0 \pm 3.0	6	6	5	3	3	1
Renfrew spring	5	6.3 \pm 12.8 b	3.2 \pm 0.8	6	4	2	0	0	0
Renfrew swarming	5	5.2 \pm 18.7 b	9.5 \pm 1.0	4	4	1	1	1	0

C) <i>L. cinereus</i>				Number of nights with at least one hour above percentile threshold					
Site	No. nights	Mean calls per hour \pm SD	Peak time of night (hour after sunset) \pm SD	25th	50th	70th	90th	95th	99th
Cypress Hills	5	45.5 \pm 59.5 b	2.2 \pm 1.1	5	3	2	0	0	0
Long Point spring	6	149.6 \pm 148.3 a	3.3 \pm 1.9	6	6	6	5	4	0
Long Point migration	6	78.0 \pm 111.3 ab	3.5 \pm 2.3	6	5	5	2	1	1
Renfrew spring	5	0.4 \pm 1.5 c	--	0	0	0	0	0	0
Renfrew swarming	5	0.2 \pm 0.2 c	--	0	0	0	0	0	0

D) <i>Myotis lucifugus</i>				Number of nights with at least one hour above percentile threshold					
Site	No. nights	Mean calls per hour \pm SD	Peak time of night (hour after sunset) \pm SD	25th	50th	70th	90th	95th	99th
Cypress Hills	5	139.5 \pm 239.3 b	1.6 \pm 0.5	5	3	3	0	0	0
Long Point spring	6	7.5 \pm 19.9 c	4.7 \pm 1.9	3	1	0	0	0	0
Long Point migration	6	87.8 \pm 109.5 b	5.7 \pm 3.4	6	4	2	0	0	0
Renfrew spring	5	424.4 \pm 1007.7 b	3.6 \pm 0.5	5	5	3	0	0	0
Renfrew swarming	5	5409.4 \pm 6826.4 a	4.0 \pm 0	5	5	5	5	4	1

Note: Means in second column followed by the same letter are not significantly different ($p < 0.05$) according to Kruskal-Wallis pairwise post-hoc test.

3.3.2 Within nights

Lasiurus borealis within-night activity at Long Point was less variable during spring than during autumn migration, with the majority of activity below the median ($75 \pm 11\%$ below median, Fig. 3.2A) and bimodal peaks above the median during spring. During migration, activity fluctuated more above and below the median ($53 \pm 31\%$ below median) and peaks of activity, above the median, were consistently in the middle of the night and at dawn (Fig. 3.2B).

Myotis lucifugus had activity above the 50th percentile four hours after sunset on every night during swarming at Renfrew (Table 3.2D) and the majority of activity was above the median ($74 \pm 18\%$ above median), all in the middle of the night lasting until dawn (Fig. 3.2D). During spring, activity fluctuated more ($38 \pm 33\%$ above median), with unimodal peaks of activity in the middle of the night (Fig. 3.2C).




Figure 3.2. Comparison of species-specific, within-night bat activity patterns between seasons at sites in Ontario, Canada, highlighting the periods when activity is above certain thresholds for two species. Mean hourly activity for *Lasiurus borealis* at Long Point during A) spring (n = 6 nights) and B) autumn migration (n = 6) and *Myotis lucifugus* activity at Renfrew during C) spring (n = 5) and D) swarming (n = 5). Solid lines are rolling averages (20 increments across each night) of the mean nightly activity among all nights at a site \pm SD (dashed red); horizontal dashed lines (grey) are percentile thresholds; vertical dashed lines (light grey) are sunrise and sunset.

3.4 Discussion

Percentile thresholds allowed for objective identification of peak activity levels at sites, while taking into account among-site and within-night variation of activity. Any activity above the median could be considered ‘high’ activity since it is higher than the majority of activity levels among all sampling units (Sokal and Rohlf 1981). All thresholds above the median are simply establishing ‘how high’ activity is. One way to interpret this scale is to identify sites based on the highest percentile threshold activity levels exceeded. For example, during swarming at Renfrew *M. lucifugus* had activity levels regularly exceeding the 95th percentile, while no other site exceeded the 70th percentile (Table 3.2D). It could be interpreted that sites with activity levels exceeding the highest percentile thresholds are important sites to bats. An international program identifies important bird areas based on criteria: significant numbers of threatened, range-restricted, migratory, or congregatory species. By the same logic, sites with activity over the highest thresholds could be defined as important areas for bats, with the assumption that high acoustic activity links to high abundance.

Examining overall bat activity, data for all species combined, Long Point does not appear to be a particularly important site. However when evaluating at the species-level it is apparent that Long Point is very important to migratory species, *L. borealis* and *L. cinereus*, especially during migration (Table 3.2B). Activity for all species combined (Tables 3.1,3.2A) was most influenced by *M. lucifugus* because of the nature of its echolocation, having shorter call duration and pulse interval (Miller 2001), resulting in more calls per individual. Because species differ in detectability and frequency of their

echolocation calls it is important to conduct species-specific analyses or use activity indices for more accurate comparisons among species (Miller 2001).

Traditional statistical methods, such as Kruskal-Wallis, do not go to a level of detail necessary for further identification of degrees of activity levels. Kruskal-Wallis results showed that sites with activity exceeding the 95th percentile had significantly higher activity, but did not provide as much information about differences of importance among sites, such as the magnitude of species-specific activity at each site. Percentile thresholds allowed us to evaluate skewed distributions and draw conclusions objectively, while looking at a finer scale.

Using percentile thresholds to define high activity is a replicable method of describing within-night activity patterns, including important times of night and degree of variation, which allows for interpretation of how a site is used by bats and its potential significance to them. From an applied perspective, it is crucial for mitigation and management decisions that methods be clearly defined when identifying times and locations with high or low activity. While a bimodal distribution of activity is typical of many insectivorous bat species, with a peak at dusk during initial foraging and a smaller peak at dawn (Kunz 1973, Hayes 1997), it is important to specify how these peaks in activity are identified. Percentile thresholds are a method to not only identify these peaks, but also describe the magnitude of activity levels relative to a larger dataset.

Environmental assessment surveys represent one area where using acoustic recordings to monitor bat activity is of particular importance. With increasing development of wind energy facilities and associated bat mortality (Kunz *et al.* 2007a), there is a growing

demand for environmental consultants to conduct pre- and/or post-construction acoustic surveys of local bat communities. There is an impetus for increasing knowledge and data about variation in activity levels, which could lead to viable mitigation options. Most pre- and post-construction surveys are not guided by a standardized survey design (Kunz *et al.* 2007a). Consequently, it is difficult to compare among acoustic monitoring surveys conducted by different groups at different times in different locations. The principle behind conducting pre-construction surveys is to assess potential risks of wind turbines to bats in an area (Kunz *et al.* 2007b). Essentially, environmental consulting agencies make recommendations based on the activity levels at a site, aiming to not erect turbines in areas of high bat activity. The lack of an agreed upon definition of 'high' bat activity and a lack of standardization in survey methodology make this a futile expectation.

Assessment recommendations are typically made by comparing site activity levels to sites in a region. Unless these comparisons are made based on clearly defined thresholds they run the risk of being subjective, leading to unsubstantiated conclusions.

The practice of establishing a definition of 'high' activity for a given site (e.g., Broders 2003) is worthwhile, but one must clearly define which criteria are being used to measure peak activity relative to a baseline. It is a difficult practice to remove subjectivity when making decisions based on relative data, such as activity levels from acoustic recordings. Communicating the relative activity level based on percentile thresholds is an objective method and allows us to move away from subjective practices.

We have demonstrated the concept of applying percentile thresholds for identifying sites important for bats at a relatively small scale. Our analysis is effective because our dataset included a wide range of activity levels. However, percentile thresholds depend entirely

on the dataset, and underlying distribution of activity levels they are based on. If generating percentile thresholds for a species is based on a limited range of activity levels, such as *M. lucifugus* from Renfrew during swarming, no other site would be considered to have ‘high’ activity. Increasing the number of sampling points contributing to the overall distribution of activity levels increases the power of this analysis. The next step is to move towards a null distribution to generate percentile thresholds. Our ability to place a given survey in the broader context will continue to improve as the database increases to include more natural variation (i.e., seasonal, annual, geographic, meteorological). Such an endeavor is far too extensive for any one group to undertake, but through collaboration and technological advancements, it is possible that such a database could be realized. Creating a public repository of acoustic datasets in order to evaluate activity of a species in the context of its entire range would allow us to standardize terms such as ‘high’ activity in an objective manner.

3.5 Acknowledgements

We thank A. Costello for her feedback on the manuscript. Thanks to the Long Point Bird Observatory, M. Brigham of the University of Regina, and the Windle family for housing and access to their properties. Funding was provided by a student research scholarship from Bat Conservation International to A.M.A., Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery grant to M.B.F. and NSERC post-doctoral fellowship to L.P.M.

3.6 References

- Barclay, R. 1984. Observations on the migration, ecology and behavior of bats at Delta Marsh, Manitoba. *Canadian Field-Naturalist* 98:331–336.
- Broders, H. 2003. Another quantitative measure of bat species activity and sampling intensity considerations for the design of ultrasonic monitoring studies. *Acta Chiropterologica* 5:235–241.
- Brooks, R., and W. Ford. 2005. Bat activity in a forest landscape of central Massachusetts. *Northeastern Naturalist* 12:447–462.
- Dzal, Y., L. A. Hooton, E. L. Clare, and M. B. Fenton. 2009. Bat activity and genetic diversity at Long Point, Ontario, an important bird stopover site. *Acta Chiropterologica* 11:307–315.
- Fenton, M. B. 1969. Summer activity of *Myotis lucifugus* (Chiroptera: Vespertilionidae) at hibernacula in Ontario and Quebec. *Canadian Journal of Zoology* 47:597–602.
- Fenton, M. B. 2003. Eavesdropping on the echolocation and social calls of bats. *Mammal Review* 33:193–204.
- Giraudoux, P. 2011. pgirmess: Data analysis in ecology. R package version 1.5.2. <http://CRAN.R-project.org/package=pgirmess>.
- Gorresen, P. M., F. J. Bonaccorso, and C. A. Pinzari. 2009. Habitat occupancy and detection of the Pacific sheath-tailed bat (*Emballonura semicaudata*) on Aguiguan,

- Commonwealth of the Northern Mariana Islands. *Acta Chiropterologica* 11:331–342.
- Hamilton, R. 2012. Spatial and temporal activity of migratory bats at landscape features. MSc Thesis. Western University.
- Hayes, J. P. 1997. Temporal variation in activity of bats and the design of echolocation-monitoring studies. *Journal of Mammalogy* 78:514–524.
- Hooper, J. 1966. The ultrasonic “voices” of bats. *New Scientist*:496–497.
- Kunz, T. H. 1973. Resource utilization: temporal and spatial components of bat activity in central Iowa. *Journal of Mammalogy* 54:14–32.
- Kunz, T. H., E. Arnett, W. P. Erickson, A. Hoar, G. Johnson, R. P. Larkin, M. D. Strickland, R. Thresher, and M. Tuttle. 2007a. Ecological impacts of wind energy development on bats: questions, research needs, and hypotheses. *Frontiers in Ecology and Environment* 5:315–324.
- Kunz, T. H., E. B. Arnett, B. M. Cooper, W. P. Erickson, R. P. Larkin, T. Mabee, M. L. Morrison, M. D. Strickland, and J. M. Szewczak. 2007b. Assessing impacts of wind-energy development on nocturnally active birds and bats: a guidance document. *Journal of Wildlife Management* 71:2449–2486.
- McCullagh, P., and J. Nelder. 1989. *Generalized linear models*. Chapman and Hall, New York, NY.

- McGuire, L. P., C. G. Guglielmo, S. A. Mackenzie, and P. D. Taylor. 2012. Migratory stopover in the long-distance migrant silver-haired bat, *Lasionycteris noctivagans*. *Journal of Animal Ecology* 81:377–85.
- Miller, B. W. 2001. A method for determining relative activity of free flying bats using a new activity index for acoustic monitoring. *Acta Chiropterologica* 3:93–105.
- Murray, S. W., and A. Kurta. 2004. Nocturnal activity of the endangered Indiana bat (*Myotis sodalis*). *Journal of Zoology* 262:197–206.
- Neuweiler, G. 1989. Foraging ecology and audition in echolocating bats. *Trends in Ecology and Evolution* 4:160–166.
- O'Hara, R. B., and D. J. Kotze. 2010. Do not log-transform count data. *Methods in Ecology and Evolution* 1:118–122.
- Parsons, K. N., G. Jones, and F. Greenaway. 2003. Swarming activity of temperate zone microchiropteran bats: effects of season, time of night and weather conditions. *Journal of Zoology* 261:257–264.
- R Development Core Team. 2011. R: a language and environment for statistical computing. R Foundation for Statistical Computing. v. 2.13.1. R Foundation for Statistical Computing, Vienna, Austria.
- Rautenbach, I. L., M. B. Fenton, and M. J. Whiting. 1996. Bats in riverine forests and woodlands: a latitudinal transect in southern Africa. *Canadian Journal of Zoology* 74:312–322.

Skowronski, M. D. 2008. callViewer. v. 18. London, Ontario, Canada.

Skowronski, M. D., and M. B. Fenton. 2008. Model-based automated detection of echolocation calls using the link detector. *Journal of the Acoustical Society of America* 124:328–336.

Sokal, R., and F. Rohlf. 1981. *Biometry*. WH Freeman and Company, 2nd edition. WH Freeman and Company, New York, NY.

Willis, C., and R. Brigham. 2003. New records of the eastern red bat, *Lasiurus borealis*, from Cypress Hills Provincial Park, Saskatchewan: a response to climate change? *Canadian Field-Naturalist* 117:651–654.

Willis, C., and R. Brigham. 2005. Physiological and ecological aspects of roost selection by reproductive female hoary bats (*Lasiurus cinereus*). *Journal of Mammalogy* 86:85–94.

Chapter 4

4 Identifying peaks in bat activity: a new application of the space-time scan statistic³

4.1 Introduction

An important first step in identifying peak bat activity is establishing the range of local variation in activity as this may vary temporally within or among nights, as well as seasonally or annually. Such scales of variability are the result of numerous factors, including insect abundance (Taylor and O'Neill 1988, Hayes 1997, Lee and McCracken 2002), air temperature (Kunz 1973, Lacki 1984, Negraeff and Brigham 1995, Hayes 1997), rainfall (Fenton *et al.* 1977, Parsons *et al.* 2003), relative humidity (Lacki 1984, Adam *et al.* 1994), wind (Adam *et al.* 1994), and species-specific life history factors such as reproductive timing (e.g., Maier 1992, Johnson *et al.* 2011) or seasonal movements (e.g., Barclay 1984, Parsons *et al.* 2003). By understanding patterns and variation in activity we can better understand the behavior of bats.

Understanding bat activity patterns and identifying periods of peak activity at various time scales, within-night to annual patterns can be important for basic research, monitoring or management. Acoustic monitoring of echolocation calls is a commonly used method of measuring bat activity. Although there is no demonstrated quantitative relationship between the numbers of calls and number of bats, the data indicate relative

³ A version of this chapter is in review.

Citation: Adams. A. M. and M. B. Fenton. Identifying peaks in bat activity: a new application of the space-time scan statistic. *Journal of Applied Ecology*: in review.

levels of bat activity. However, there is no currently no definition of high activity or objective means of identifying periods of high activity.

Activity patterns may permit inference on how bats use a site. For example, activity at a roost may produce two clear peaks of activity corresponding to emergence and return at dusk and dawn (Kunz 1973, Hayes 1997), while monitoring the same site when young are nursing may show a unimodal peak of activity while females are pregnant (Maier 1992). At a swarming site, when bats congregate in August and September for pre-hibernation mating, activity may peak several hours after sunset reflecting the times of arrival of bats that have travelled long distances to the site (Fenton 1969). Migration stopover sites could have nightly unimodal peaks of activity at dawn during migratory periods as species arrive at sites along migration routes (McGuire *et al.* 2012).

Previously, peaks in activity have been identified visually with activity plots (Hayes 1997), with selection of an arbitrary level of activity as ‘high’ (Broders 2003, Brooks and Ford 2005), or by the number of calls above percentile thresholds (Adams *et al.* submitted, Gorresen *et al.* 2009). Percentile thresholds identify peak acoustic bat activity (Adams *et al.* submitted) by comparing when activity exceeds thresholds based on a larger distribution of activity from a range of sites, placing activity levels in a larger context.

We propose the use of the space-time scan statistic (SaTScan) as an analytical tool for identifying spatial and temporal peaks of bat activity. SaTScan was originally developed for monitoring the spread of disease by detecting localized clusters of infection in space and time where the number of cases differed significantly from the background expectation (Kulldorff 1997). However, it has also been applied to other fields (see

Kulldorff 1994 for a partial bibliography), including ecology (e.g., Coulston and Riitters 2003, Rubin and MacFarlane 2008, Vadrevu 2008, Tuia *et al.* 2008).

Autocorrelation is a general statistical property of variables observed along geographic space and time-series (Koenig 1999). Spatial and temporal autocorrelation present a statistical problem because the data violate the assumption of independence inherent in most standard statistical procedures (e.g., ANOVA; Legendre 1993). As space or time intervals decrease, the dependence between successive observations usually increases (Legendre 1993, Koenig 1999). Autocorrelation can be problematic, for studies using radio telemetry (Rooney *et al.* 1998, Dray *et al.* 2010) or those examining macro-scale patterns of species diversity (Legendre 1993, Diniz-Filho *et al.* 2003), because it inflates Type I errors and could bias environmental factors with higher spatial autocorrelation (Lennon 2000). Only a few studies have accounted for autocorrelation in spatial or temporal datasets of bat activity (Audet and Fenton 1988, Gorresen and Willig 2004, Loeb and O'Keefe 2006, Stevens *et al.* 2007, Hein *et al.* 2009). Autocorrelation may be less of a concern because bats are highly mobile and able to fly substantial distances in short periods of time (Henry *et al.* 2002), but the potential for violation of independence in temporal and spatial datasets remains because much of their behavior keeps them in a restricted area for an extended period of time. Even if autocorrelation is less likely with bat data, a technique that removes this concern would make one more confident in conclusions drawn. SaTScan controls for spatial and temporal trends, whether autocorrelated or not, by adjusting for purely spatial and purely temporal variation (Kulldorff 2004).

Our goal was to examine the validity of SaTScan as a method for identifying peaks in bat activity, by comparing it with the percentile threshold approach. We also applied SaTScan to assess temporal patterns of activity of five species of bats at three sites.

4.2 Methods

4.2.1 Study areas

We recorded bat echolocation calls in 2008 at three sites in Ontario, Canada that were at least 131 km apart, on clear nights in both spring and fall: i) Long Point, an important location for bats during migratory periods (Adams *et al.* submitted, Dzal *et al.* 2009, McGuire *et al.* 2012), on six nights during both spring (June, Site 1a) and fall migration (August/September, Site 1b); ii) along the shore of Lake Opinicon at the Queen's University Biological Station (QUBS), where bats forage and roost (Barclay 1982, Arh 2009), for five nights during both spring (May/June, Site 2a) and summer (August, Site 2b); iii) an abandoned mine near Renfrew, an important swarming site and hibernaculum (Fenton 1969, 1970) housing up to 30,000 bats during the winter, for five nights during both spring (May/June, Site 3a) and fall swarming (August, Site 3b).

4.2.2 Acoustic sampling

Each night we recorded continuously from dusk until dawn using an Avisoft UltraSoundGate System (Avisoft Bioacoustics, Berlin, Germany) at 8 bit with a sampling rate of 250 kHz, trigger set to continuous recording, and gain at seven. Externally polarized condenser microphones (Avisoft CM16/CMPA) connected to an Avisoft UltraSoundGate 416-200 were operated with Avisoft Recorder USG software.

All acoustic data were analyzed using callViewer (v. 18, Skowronski 2008), a custom echolocation sound analysis program written with MATLAB software (The MathWorks, Natick, Massachusetts), using the automated detection feature to identify echolocation calls in acoustic recordings. We filtered and identified the calls to species using discriminant function analysis (DFA, following Adams *et al.* submitted, Appendix A) for seven species in Ontario, *Eptesicus fuscus*, *Lasionycteris noctivagans*, *Lasiurus borealis*, *L. cinereus*, *Myotis lucifugus*, *M. septentrionalis*, and *Perimyotis subflavus*.

4.2.3 Statistical analysis

We summed the calls per minute, post-DFA and filtering, for all species combined (overall activity for all seven species). We also calculated nightly activity as number of calls per minute for each of five species (*E. fuscus*, *L. borealis*, *L. cinereus*, *M. lucifugus*, and *P. subflavus*). We used two methods to identify peak bat activity, SaTScan and percentile thresholds.

We used SaTScan software (v. 9.1.1, Kulldorff *et al.* 2005, freeware available online), to detect peaks in activity at two levels at each site, within- and among-nights. SaTScan can identify high or low clusters, where event occurrence is significantly more likely within the cluster than outside the cluster ($\alpha = 0.01$). In the case of acoustic bat activity, SaTScan would identify a cluster of high activity in space and/or time where the null hypothesis is that activity levels are always constant everywhere. Significance of a potential high cluster is determined based on a likelihood ratio $\lambda = L/L_0$, where L is the max likelihood and L_0 is the max likelihood constrained to a true null hypothesis. Higher λ means greater support for the alternate hypothesis that activity is greater inside the

cluster than outside. Monte Carlo hypothesis testing assigns the degree of significance of each cluster by determining the probability of obtaining a value that is at least as high as the observed value from randomized data (Kulldorff *et al.* 2005). Purely temporal scan statistic analysis moves a scanning window over each possible time point across an entire time frame. The window size ranges from the size of a single time point (e.g., one minute in this study) up to a user-defined maximum cluster size (MCS), which limits the maximum size of the cluster to a percentage of the total sampling period, most commonly the window size includes no more than half the total time period (MCS of 50%, Kulldorff 1997). We ran SaTScan five times for each analysis, specifying the temporal MCS at 1, 5, 10, 30, and 50%. A high MCS gives more power but low specificity in describing the boundaries of clusters (Rubin and MacFarlane 2008). The ability to search for peaks with various MCS helps give a better measure of the strength of a peak because a high MCS considers both large and small clusters, while a low MCS only considers small clusters. To apply this method to both space and time simultaneously SaTScan uses cylinders rather than one-dimensional windows. The base of the scanning window represents space and the height represents time, with both the width and the height varying up to the limits of the MCS.

The scan statistic can use different probability models depending on the nature of the data. We applied both purely temporal and spatial-temporal analysis to analyze bat activity. Purely temporal analysis, for within-site analyses, used the discrete Poisson model (Kulldorff 1997), while spatial-temporal analysis, for among-site analysis, used the space-time permutation model (Kulldorff *et al.* 2005). We used the retrospective analysis option because we had data with a fixed geographic region and fixed temporal

study period. For both models, we input case files including the number of calls per minute (count data), minute after sunset (MAS), and site location. For the space-time permutation model we also included a coordinates file with the latitude and longitude of each site location. We specified a one-tailed analysis to look for clusters with unexpectedly high numbers of calls and used 999 Monte Carlo replications. The analysis output included the “most likely cluster,” which described the locations and time frame(s) that were significantly higher than the background expectation with the observed and expected number of cases and p-value. We also used multivariate SaTScan, which allows for multiple datasets to be searched simultaneously for clusters (Kulldorff *et al.* 2007), with each night within a site/season as a separate dataset to detect the peak time period among all nights for a single site/season.

We calculated threshold values at five percentile thresholds (50th, 70th, 90th, 95th, and 99th percentiles, following Adams *et al.* submitted) based on the distribution of number of calls per minute for all nights, at all sites, for all species combined, excluding all minutes with no activity from the distribution. We compared peaks identified by percentile thresholds and SaTScan by calculating the proportion of minutes above the 50th percentile threshold (median) that was included in each SaTScan high cluster time period. We also compared among-night activity between the two methods. With SaTScan, we identified peak nights with spatial-temporal analysis of mean calls per minute for each night and compared it to which nights had at least 60 minutes of activity above each percentile threshold.

To demonstrate application of SaTScan we used multivariate SaTScan to identify peaks in species-specific activity. Peak time frames for each species at each site/season were

defined by the lowest MCS that included the most nights. When all nights were not included in the peak time frame we reran multivariate analysis, only including the nights that were not included in the result of the previous analysis. We compared the peak time frames among sites and seasons for each species and among species within each site/season. We excluded species at sites/seasons with fewer than two nights of activity. Comparisons of peak time frames among sites and seasons were considered different when there was no overlap in MAS.

4.3 Results

4.3.1 SaTScan description and comparison to percentile thresholds

SaTScan defined time frames that differed in length depending on the MCS (e.g. Fig. 4.1A). The high cluster time frame for a MCS of 50% ranged from 2 – 282 minutes, while a MCS of 1% ranged from 1 – 6 minutes. When there was a strong, singular peak of activity the time frame could be the same, regardless of MCS. Sometimes, the peak time frame did not include the maximum minute of activity, but the maximum five minutes of activity was always included in the peak time frame. Multivariate SaTScan identified peaks for multiple nights within a site and season (e.g. Fig. 4.1B – F). A MCS of 50% and 30% identified peak time frames that included all nights at Site 1 and Site 3, but when the MCS was reduced some nights were no longer included. Temporal variation in activity was high at Site 2, so multivariate SaTScan was not able to identify a time frame that worked for all nights at the site/season.




Figure 4.1. Nightly activity of bats for all nights at Renfrew during swarming (Site 3b) in 2008 with calls for all species combined. Highlighted time frames are SaTScan high cluster time frames defined by A) various maximum cluster sizes (MCS) within a single night and B-F) MCS of 30% with multivariate scan.

The number of calls per minute, for all species combined, at each percentile threshold levels were: 29 at the 50th, 70 at the 70th, 256 at the 90th, 366 at the 95th, and 592 at the 99th percentile, respectively. SaTScan and percentile thresholds differed in the number of minutes of peak activity identified (Fig. 4.2A). Percentile thresholds identified any minutes with activity over the threshold levels, while SaTScan identified a single time period that included activity that was higher than expected. SaTScan peaks were most similar to percentile thresholds at Site 3, where there was a clear, unimodal peak of activity. SaTScan was less similar to percentile thresholds at Sites 1 and 2 because activity was more evenly spread throughout the night. SaTScan with a MCS of 50% and multivariate SaTScan with a MCS of 30% included the majority of minutes above the 50th percentile at Sites 1 and 3 (Fig. 4.2A) and the majority of the total nightly activity at Sites 2 and 3 (Fig. 4.2B). The maximum minute of activity for a night was included in the SaTScan time frame on 67% of the nights at Site 1, all nights at Site 2, and 90% of the nights at Site 3.

Peak nights differed between percentile thresholds and SaTScan (Fig. 4.3). Sites 2a and 3b were the most important to bats based on percentile thresholds because they both had nights that exceeded the 90th and 95th percentile thresholds. SaTScan would identify Site 1a as an important site even though only one night had 60 minutes of activity over the 50th percentile threshold. SaTScan peak nights more closely reflect within-site peaks, rather than comparisons among sites. Peaks based on percentile thresholds do not consider space, by adding in the spatial component with SaTScan, sites that are farther apart are weighted differently.




Figure 4.2. Proportion of overall bat activity in SaTScan high cluster time frames at various maximum cluster sizes (MCS) similar to A) total minutes over the 50th percentile threshold, and B) total nightly activity for three sites in Ontario, Canada during spring (a) and late summer (b) in 2008 with calls for all species combined.




Figure 4.3. Comparison of nights of peak bat activity (mean calls per minute \pm SD) identified by SaTScan spatial-temporal analysis (grey bars) and with at least 60 minutes of activity over the five percentile thresholds (*50th percentile, **70th percentile, ***90th percentile, ****95th percentile), for all species combined at three sites in Ontario, Canada during spring (a) and late summer (b) in 2008. Note different y-axes.

4.3.2 Species-specific activity

We found no indication of temporal partitioning among species that was consistent among all sites and seasons (Fig.4.4). *Lasiurus cinereus* always had the earliest peaks of activity in the night at sites where it was present. All species had overlapping peaks of activity at Sites 2 and 3a. The majority of peak activity occurred within the first five hours of the night for all species at all sites, except *L. borealis* at Site 3b (Fig. 4.4). Peaks of activity at Site 1a were very inconsistent, with different timing of peaks among nights for most species (Fig. 4.4G,S,Y). *Lasiurus borealis* activity patterns differed between Site 1a and 1b with timing of peak activity being more consistent at Site 1b (Fig.4.4H). The majority of *M. lucifugus* peaks of activity were two to four hours after sunset and was most consistent at Site 3b, being the only case when all nights were included at all MCS (50 – 1%).




Figure 4.4. Mean species-specific bat activity at three sites in Ontario, Canada during spring (a) and late summer (b) for 2008. Shaded time frames are peaks identified by multivariate SaTScan. Note different y-axes. EPFU – *Eptesicus fuscus*, LABO – *Lasiurus borealis*, LACI – *L. cinereus*, MYLU – *Myotis lucifugus*, PESU – *Perimyotis subflavus*.

4.4 Discussion

SaTScan precisely identified peaks of activity, but cannot identify every individual minute of high activity within a night. There are two main advantages of SaTScan. First, multivariate SaTScan identifies peaks that occur among all datasets. Identifying common peaks within each site/season can be extremely valuable for recognizing important time periods for bat activity, whether to focus sampling efforts or comparing species-specific temporal patterns. Second, the true strength of SaTScan is the ability to simultaneously account for both spatial and temporal patterns and it works with data at any spatial and temporal scales. We analyzed our data by minute because it allowed SaTScan to define more specific peak time periods, but any time scale is possible. Our analysis was at a fairly large spatial scale with coarse resolution, but this method has the potential to be more useful at a much smaller spatial scale without concerns of autocorrelation during analysis.

SaTScan and percentile thresholds both identify peak activity objectively, but do so in different ways making it difficult to compare results. SaTScan identifies a single cluster of peak activity, while percentile thresholds identify any time frame with activity above a defined threshold. Using multiple percentile thresholds provides a measure of the magnitude of activity at a site and can be useful for identifying important sites (Adams *et al.* submitted), while SaTScan is best at identifying peak time periods, especially among multiple nights. We recommend using the two methods in combination to have a more complete picture of activity levels and activity patterns at a site. Percentile thresholds will be better for identifying important sites, while SaTScan will identify peaks in activity

among nights. For example, Site 3b is the most important to bats because the majority of nights had overall activity that exceeded the 90th percentile threshold (Fig. 4.3), while it also stood apart from the rest of the sites with peak of activity, identified by SaTScan, three to four hours after sunset. In contrast, Sites 1, 2, and 3a had few to no nights with activity above the 90th percentile and peak activity starting one to two hours after sunset.

We identified peaks in activity with SaTScan that were comparable to findings in other studies. For example, peaks in foraging activity were consistent with foraging patterns reported by temperate species at other sites (Kunz 1973), however, at our sites in Ontario *M. lucifugus* did not have peaks of activity immediately after sunset as observed in Nova Scotia (Broders *et al.* 2003) potentially due to timing of insect prey (Rautenbach *et al.* 1996) or proximity of roosting habitat. The consistency of peaks among nights with SaTScan can potentially indicate bats' use of a site, for example peak activity was most consistent during migration (Site 1b) and swarming (Site 3b) and more variable among nights during foraging (Sites 1a and 2).

A limitation is that SaTScan will not identify peaks of activity when activity is consistently high or low throughout a night, but using percentile thresholds in combination with SaTScan would identify when this is the case. SaTScan is also not able to detect multiple temporal peaks, if there are bimodal peaks only one will be identified. Foraging sites are most commonly observed with bimodal activity, with a peak at dusk when bats forage on crepuscular insects and another peak at dawn when returning to their day roost (Kunz 1974, Rydell 1993, Kunz *et al.* 1995, Hayes 1997). It is possible to recognize bimodal patterns of activity with SaTScan by running multiple analyses, each time excluding the peak time period and searching for a secondary peak. For example, it

was possible to identify a secondary peak of *L. borealis* activity just before dawn at Site 1b after excluding the primary peak at dusk (Fig. 4.4H). We did not detect secondary peaks of activity for all species combined at Sites 1 or 3, rather just extensions of the primary peak time period.

SaTScan is a valuable tool to quickly identify peaks with an objective, replicable, and statistically sound method that can be applied at various temporal and spatial scales. As bat detector technologies improve, allowing all night recordings over long periods of time at many locations, it is more difficult to analyze the vast amounts of data. SaTScan identifies when and where bats are most active, which has applications for basic and applied research, such as comparing peaks in activity among habitat types, commercial developments (e.g., wind energy), or years (e.g., meta-analysis of annual fluctuations pre- and post- white-nose syndrome).

4.5 Acknowledgements

We thank Drs. E. Clare, E. Fraser, J. McNeil, B. Rubin, and M. Skowronski for comments on earlier versions of the manuscript. Thanks to the Long Point Bird Observatory, Queen's University Biological Station, and the Windle family for housing and access to their property. We thank Bat Conservation International for funding A.M.A. and the Natural Sciences and Engineering Research Council (Canada) for supporting M.B.F.

4.6 References

- Adam, M., M. Lacki, and L. Shoemaker. 1994. Influence of environmental conditions on flight activity of *Plecotus townsendii virginianus* (Chiroptera: Vespertilionidae). *Brimleyana* 21:77–85.
- Adams, A. M., L. P. McGuire, L. A. Hooton, and M. B. Fenton. How high is high? Using percentile thresholds to identify peak bat activity. submitted.
- Arh, M. R. 2009. Foraging ecology of a bat assemblage. MSc Thesis. University of Toronto.
- Audet, D., and M. Fenton. 1988. Heterothermy and the use of torpor by the bat *Eptesicus fuscus* (Chiroptera: Vespertilionidae): a field study. *Physiological Zoology* 61:197–204.
- Barclay, R. M. R. 1982. Night roosting behavior of the little brown bat, *Myotis lucifugus*. *Journal of Mammalogy* 63:464–474.
- Barclay, R. M. R. 1984. Observations on the migration, ecology and behavior of bats at Delta Marsh, Manitoba. *Canadian Field-Naturalist* 98:331–336.
- Broders, H. 2003. Another quantitative measure of bat species activity and sampling intensity considerations for the design of ultrasonic monitoring studies. *Acta Chiropterologica* 5:235–241.

- Broders, H., G. Quinn, and G. Forbes. 2003. Species status, and the spatial and temporal patterns of activity of bats in southwest Nova Scotia, Canada. *Northeastern Naturalist* 10:383–398.
- Brooks, R., and W. Ford. 2005. Bat activity in a forest landscape of central Massachusetts. *Northeastern Naturalist* 12:447–462.
- Coulston, J. W., and K. H. Riitters. 2003. Geographic analysis of forest health indicators using spatial scan statistics. *Environmental Management* 31:764–773.
- Diniz-Filho, J. A., L. M. Bini, and B. A. Hawkins. 2003. Spatial autocorrelation and red herrings in geographical ecology. *Global Ecology and Biogeography* 12:52–64.
- Dray, S., M. Royer-Carenzi, and C. Calenge. 2010. The exploratory analysis of autocorrelation in animal-movement studies. *Ecological Research* 25:673–681.
- Dzal, Y., L. A. Hooton, E. L. Clare, and M. B. Fenton. 2009. Bat activity and genetic diversity at Long Point, Ontario, an important bird stopover site. *Acta Chiropterologica* 11:307–315.
- Fenton, M. B. 1969. Summer activity of *Myotis lucifugus* (Chiroptera: Vespertilionidae) at hibernacula in Ontario and Quebec. *Canadian Journal of Zoology* 47:597–602.
- Fenton, M. B. 1970. Population studies of *Myotis lucifugus* (Chiroptera: Vespertilionidae) in Ontario. *Life Sciences Contributions Royal Ontario Museum* 77:1–34.

- Fenton, M. B., N. G. H. Boyle, T. M. Harrison, and D. J. Oxley. 1977. Activity patterns, habitat use, and prey selection by some African insectivorous bats. *Biotropica* 9:73–85.
- Gorresen, P. M., and M. Willig. 2004. Landscape responses of bats to habitat fragmentation in Atlantic Forest of Paraguay. *Journal of Mammalogy* 85:688–697.
- Gorresen, P. M., F. J. Bonaccorso, and C. A. Pinzari. 2009. Habitat occupancy and detection of the Pacific sheath-tailed bat (*Emballonura semicaudata*) on Aguiguan, Commonwealth of the Northern Mariana Islands. *Acta Chiropterologica* 11:331–342.
- Hayes, J. P. 1997. Temporal variation in activity of bats and the design of echolocation-monitoring studies. *Journal of Mammalogy* 78:514–524.
- Hein, C. D., S. B. Castleberry, and K. V. Miller. 2009. Site-occupancy of bats in relation to forested corridors. *Forest Ecology and Management* 257:1200–1207.
- Henry, M., D. Thomas, R. Vaudry, and M. Carrier. 2002. Foraging distances and home range of pregnant and lactating little brown bats (*Myotis lucifugus*). *Journal of Mammalogy* 83:767–774.
- Johnson, J. B., J. W. Edwards, and W. M. Ford. 2011. Nocturnal activity patterns of Northern Myotis (*Myotis septentrionalis*) during the maternity season in West Virginia (USA). *Acta Chiropterologica* 13:391–397.

- Koenig, W. 1999. Spatial autocorrelation of ecological phenomena. *Trends in Ecology & Evolution* 14:22–26.
- Kulldorff, M. 1997. A spatial scan statistic. *Communications in Statistics - Theory and Methods* 26:1481–1496.
- Kulldorff, M. 2004. SaTScan user's guide. v. 9.1.1. Boston, MA.
- Kulldorff, M., R. Heffernan, J. Hartman, R. Assunção, and F. Mostashari. 2005. A space-time permutation scan statistic for disease outbreak detection. *PLoS Medicine* 2:e59.
- Kulldorff, M., F. Mostashari, L. Duczmal, W. K. Yih, K. Kleinman, and R. Platt. 2007. Multivariate scan statistics for disease surveillance. *Statistics in Medicine* 26:1824–1833.
- Kunz, T. 1973. Resource utilization: temporal and spatial components of bat activity in central Iowa. *Journal of Mammalogy* 54:14–32.
- Kunz, T. 1974. Feeding ecology of a temperate insectivorous bat (*Myotis velifer*). *Ecology* 55:693–711.
- Kunz, T., J. Whitaker, and M. Wadanoli. 1995. Dietary energetics of the insectivorous Mexican free-tailed bat (*Tadarida brasiliensis*) during pregnancy and lactation. *Oecologia* 101:407–415.
- Lacki, M. 1984. Temperature and humidity-induced shifts in the flight activity of little brown bats. *Ohio Journal of Science* 84:264–266.

- Lee, Y. F., and G. McCracken. 2002. Foraging activity and food resource use of Brazilian free-tailed bats, *Tadarida brasiliensis* (Molossidae). *Ecoscience* 9:306–313.
- Legendre, P. 1993. Spatial autocorrelation: trouble or new paradigm? *Ecology* 74:1659–1673.
- Lennon, J. J. 2000. Red-shifts and red herrings in geographical ecology. *Ecography* 23:101–113.
- Loeb, S., and J. O’Keefe. 2006. Habitat use by forest bats in South Carolina in relation to local, stand, and landscape characteristics. *Journal of Wildlife Management* 70:1210–1218.
- Johnson, J. B., J. W. Edwards, and W. M. Ford. 2011. Nocturnal activity patterns of Northern Myotis (*Myotis septentrionalis*) during the maternity season in West Virginia (USA). *Acta Chiropterologica* 13:391–397.
- McGuire, L. P., C. G. Guglielmo, S. A. Mackenzie, and P. D. Taylor. 2012. Migratory stopover in the long-distance migrant silver-haired bat, *Lasionycteris noctivagans*. *Journal of Animal Ecology* 81:377–85.
- Negraeff, O., and R. M. Brigham. 1995. The influence of moonlight on the activity of little brown bats (*Myotis lucifugus*). *Zeitschrift für Säugetierkunde* 60:330–336.
- Parsons, K. N., G. Jones, and F. Greenaway. 2003. Swarming activity of temperate zone microchiropteran bats: effects of season, time of night and weather conditions. *Journal of Zoology* 261:257–264.

- Rautenbach, I. L., M. B. Fenton, and M. J. Whiting. 1996. Bats in riverine forests and woodlands: a latitudinal transect in southern Africa. *Canadian Journal of Zoology* 74:312–322.
- Rooney, S., A. Wolfe, and T. Hayden. 1998. Autocorrelated data in telemetry studies: time to independence and the problem of behavioural effects. *Mammal Review* 28:89–98.
- Rubin, B., and D. MacFarlane. 2008. Using the space-time permutation scan statistic to map anomalous diameter distributions drawn from landscape-scale forest inventories. *Forest Science* 54:523–533.
- Rydell, J. 1993. Variation in foraging activity of an aerial insectivorous bat during reproduction. *Journal of Mammalogy* 74:503–509.
- Skowronski, M. D. 2008. *callViewer*. v. 18. London, Ontario, Canada.
- Stevens, R. D., C. López-González, and S. J. Presley. 2007. Geographical ecology of Paraguayan bats: spatial integration and metacommunity structure of interacting assemblages. *Journal of Animal Ecology* 76:1086–1093.
- Taylor, R., and M. O’Neill. 1988. Summer activity patterns of insectivorous bats and their prey in Tasmania. *Wildlife Research* 15:533–539.
- Tuia, D., F. Ratle, R. Lasaponara, L. Telesca, and M. Kanevski. 2008. Scan statistics analysis of forest fire clusters. *Communications in Nonlinear Science and Numerical Simulation* 13:1689–1694.

Vadrevu, K. P. 2008. Analysis of fire events and controlling factors in eastern India using spatial scan and multivariate statistics. *Geografiska Annaler: Series A, Physical Geography* 90:315–328.

Chapter 5

5 Value in variation? Stationary acoustic transects to account for spatial variation in bat activity

5.1 Introduction

Variation is a challenge to anyone studying ecology. The ability to identify and account for different sources of variation at a particular location can impact the conclusions drawn from the data. This is true whether asking questions about habitat associations or making recommendations for environmental policy. Frequently, high variation in bat activity has been reported (e.g., Hayes and Adam 1996, Milne *et al.* 2009). It is important to account for variation to maximize the chances of obtaining unbiased measures of bat activity (Hayes 1997, 2000) and to detect specific species (Broders 2003, Skalak *et al.* 2012). Acoustic surveys are common practice for studying bat ecology (Britzke *et al.* 2013) and a key assumption is that activity levels recorded from a single bat detector reflect a broader set of locations and times (Hayes 2000). To capture acoustic activity that accurately describes local activity levels, it is essential to account for variation from detectors used (Adams *et al.* 2012), temporal variability within and among nights (Milne *et al.* 2005, Skalak *et al.* 2012), and spatial variation within sites (Hayes 2000, Fischer *et al.* 2009). Both field work and analysis of acoustic data are labor intensive, resulting in a trade-off between sampling effort and collecting sufficient data so as to accurately describe bat activity at a site. A compromise can be made by measuring and understanding activity variability present at a site (Fischer *et al.* 2009). Where should we draw the line when making the trade-off between effort and accuracy?

Decisions on sampling effort are typically based on limitations in funding and personnel, but are ultimately impacted by the research question being asked. The number of sampling nights necessary to accurately estimate levels of bat activity at a site differs among locations. Hayes (1997) recommended six to eight nights of sampling to obtain accurate estimates of overall activity at sites in Oregon, USA, while Broders (2003) found that 14+ nights of sampling were required for accurate estimates of *Myotis lucifugus* activity at sites in New Brunswick, Canada. Identification of species-specific activity may play a role in these different recommendations of sampling effort (Broders 2003).

The validity of the assumption that extrapolation of recording activity from a single detection point reflects activity for an entire site/habitat has received little attention (Hayes 2000, Britzke *et al.* 2013), considering that a single detector has a limited range of 5 m to 40 m for high frequencies (Adams *et al.* 2012). To account for horizontal spatial variation, multiple detectors can be deployed within a site (Krusic *et al.* 1996b, Gannon *et al.* 2003, Duchamp *et al.* 2006, Fischer *et al.* 2009), or a single detector can be moved to a new location within the site each night (Fischer *et al.* 2009). Vertical spatial variation can be addressed with multiple detectors deployed at various heights (Hayes and Gruver 2000, Reynolds 2006, Fischer *et al.* 2009, Staton and Poulton 2012a). The use of multiple detectors simultaneously can increase the probability of detecting different species of bats at large (Skalak *et al.* 2012) and small (Duchamp *et al.* 2006) spatial scales, but there is little evidence of how multiple detectors within a site impact estimates of activity levels (Fischer *et al.* 2009).

It is also important to accurately detect patterns of activity to increase understanding of bat ecology and behavior. Nightly activity patterns vary by species (Kunz 1973, Broders

et al. 2003, Milne *et al.* 2005, Skalak *et al.* 2012) and with extrinsic factors, such as temperature (e.g., Hayes 1997) and insect abundance (e.g., Ciechanowski *et al.* 2007). Temporal and spatial partitioning may underlie resource partitioning among species of bats with similar ecology and morphology (Kunz 1973, Nicholls and Racey 2006, Adams and Thibault 2006) and could be another reason for variation in spatio-temporal activity patterns. If patterns of activity differ spatially within a site, then the placement of detectors could impact depictions of activity patterns.

Our goal was to quantify small-scale spatio-temporal variation in bat activity as determined by monitoring echolocation calls and to demonstrate the effect on acoustic sampling design and interpretation. We did this by addressing three objectives: i) testing whether space, including detection height, or time, explained more of the variation in bat activity to prioritize sampling efforts; ii) assessing within-site spatio-temporal variation in activity among species and how detector location impacted depictions of activity at each site; and iii) investigating how many nights and detectors were necessary to have accurate estimates of mean nightly activity and how this varied by species. We expected that estimates of activity would be more accurate with more detectors within a site. We also expected that activity levels and patterns would vary along a linear, stationary acoustic transect and would require multiple sampling points within a site to accurately assess bat activity. We predicted that species would partition resources in space and time and that species-specific peaks of activity would exhibit little or no overlap in time and space.

5.2 Methods

5.2.1 Study areas

We recorded bat echolocation calls at four sites in Ontario, Canada: first, at Long Point for a total of 12 nights (June, August, September 2008), which is an important site for bats during migratory periods (Dzal *et al.* 2009, McGuire *et al.* 2012); second, at the Queen's University Biological Station (QUBS), for 11 nights (May, June, August 2008), a site where bats forage and roost nearby (Barclay 1982, Arh 2009); third, at an abandoned mine near Renfrew for 12 nights in 2008 (May, June, August) and 15 consecutive nights in 2010 (August), which is an important swarming site and hibernaculum (Fenton 1969, 1970) housing up to 30,000 bats during the winter; lastly, along the shore of a lake 500 m from the abandoned mine at Renfrew for 12 nights (May, June, August 2008). Throughout the summer we rotated from site to site, recording for three clear, consecutive nights before moving to the next site.

5.2.2 Acoustic sampling

We recorded continuously from dusk until dawn with batcorders 1.0 (ecoObs, Nuremberg, Germany). Batcorders have a sampling rate of 500 kHz and 16 bit sampling resolution and were set at a critical frequency of 14 kHz (i.e., a trigger event prompted recordings of sound in the frequency range of 14 - 250 kHz). Recordings were activated at a low detector sensitivity threshold of -36 dB (1.6% of the microphone's maximum amplitude) to increase the recording range of the detectors, similar to increasing gain on other detectors. Batcorders have a pre-trigger of 50 ms and we adjusted the post-trigger to 800 ms, which is the interval between successive detected sounds written into the same file, the higher value maximizes the number of below-threshold calls recorded. We set

quality to 40, an intermediate value where higher values of the quality detection algorithm are less conservative, to allow for recognition of sounds that are less like echolocation calls. Recordings were saved to an HCSD card in .RAW format and then converted to .WAV with a custom conversion program created with MATLAB software (The MathWorks, Natick, Massachusetts) for future analysis.

We placed detectors along a linear transect 40 m apart, corresponding to the minimum sampling distance without overlap (Stahlschmidt and Brühl 2012). All four sites were forested, but ranged in levels of canopy cover throughout the site (Table 5.1). Transect lines within each site were chosen so detector locations ranged in amount of canopy cover. Detectors were oriented upward at a 45° angle, facing the area with the greatest opening. In 2008, we set out three sets of paired detectors with one low detector at 1.5 m above ground level and one high detector at 4 m in an 80 m transect. One high detector malfunctioned for half of the field season and was not included in the majority of the analysis. In 2010, five detectors were 1.5 m high in a 160 m transect. We measured canopy cover with a densitometer at each detector location. Each night we recorded temperatures at every detector with iButton Thermochron temperature data loggers (Maximum Integrated, San Jose, California) and wind speed and relative humidity with a Kestrel 4000 (Nielsen-Kellerman, Boothwyn, Pennsylvania) at a single sampling location in the site.

Table 5.1. Percent canopy cover at each of the detector locations within A) each of the four sites in Ontario, Canada in 2008 and B) Renfrew mine in 2010.

A)	Detector locations		
Sites	A	B	C
Long Point	20%	37%	5%
QUBS	36%	57%	43%
Renfrew lake	38%	30%	68%
Renfrew mine	75%	69%	39%

B)	Detector locations				
Site	A	B	C	D	E
Renfrew mine	30%	75%	67%	39%	70%

We analyzed acoustic data using callViewer (v. 18, Skowronski 2008), a custom echolocation sound analysis program written with MATLAB. We used the automated detection feature of callViewer to identify echolocation calls in acoustic recordings. We then filtered and identified our data to species using quadratic discriminant function analysis (DFA) in R (v. 2.13.1, R Development Core Team 2011, following Adams *et al.* submitted, Appendix A) for seven species in Ontario: *Eptesicus fuscus*, *Lasionycteris noctivagans*, *Lasiurus borealis*, *L. cinereus*, *Myotis lucifugus*, *M. septentrionalis*, and *Perimyotis subflavus*. Classification accuracy from cross-validation was high (Table 5.2), but to further improve classification accuracy and because DFA is not capable of assigning calls to an “unknown” category, we applied a post hoc, species-specific filter to remove any data that were above or below specific durations and minimum frequencies for each species (Table 5.2).

Table 5.2. Classification accuracy for species identification of echolocation calls for bats in Ontario, Canada with quadratic discriminate function analysis (DFA) using cross-validation. Post-DFA filter settings removed echolocation calls identified to species, but outside of the species-specific ranges of duration and minimum frequencies (Fmin) to further improve classification accuracy.

Species	DFA classification accuracy	Post-DFA filter	
		Duration	Fmin
<i>Eptesicus fuscus</i>	78 %	3 – 12 ms	20 – 30 kHz
<i>Lasionycteris noctivagans</i>	94 %	4 – 12 ms	21 – 30 kHz
<i>Lasiurus borealis</i>	88 %	5 – 17 ms	29 – 43 kHz
<i>L. cinereus</i>	90 %	8 – 30 ms	15 – 29 kHz
<i>Myotis lucifugus</i>	90 %	3 – 8 ms	30 – 43 kHz
<i>M. septentrionalis</i>	82 %	1 – 3.5 ms	25 – 60 kHz
<i>Perimyotis subflavus</i>	93 %	5 – 15 ms	36 – 46 kHz

5.2.3 Statistical methods

We totaled the number of calls per minute after sunset (MAS), hour after sunset (HAS), and per night for each detector per species. We also summed the calls per MAS, HAS, and night for all seven species, post-DFA and filtering, for activity of all species combined. There were five parts to our analysis: (i) Kruskal-Wallis non-parametric test for overall differences in activity among detectors, (ii) generalized linear mixed models (GLMM) to partition variance components among random effects, (iii) percentile thresholds for magnitude of activity, (iv) space-time scan statistic (SaTScan) for identifying peaks of activity, and (v) sub-sampling to test the effect of sampling effort on estimates of activity. To test the overall significant differences in activity levels (mean calls per hour) among detectors within a site, we used Kruskal-Wallis rank sum test and a pair-wise post-hoc test (*kruskalmc* in *pgirmess* package in R; Giraudoux 2011).

5.2.4 Generalized linear mixed models

To test the relative contribution of spatial and temporal variation with non-normal data including random effects, we used a GLMM fit by the Laplace approximation and a Poisson distribution (*glmer* in *lme4* package in R; Bolker *et al.* 2009, Bates *et al.* 2011). Acoustic data were not transformed to normality because transformations of count data can increase bias that is negligible when using an appropriate model, such as the Poisson distribution (O'Hara and Kotze 2010). We analyzed total calls per night (nightly activity) for each species at each site and detector. Random effects were site (2008 only), detector nested in site, and night nested in site. Fixed effects were canopy cover and detector height (2008 only), each detector-level effects. Like many ecological studies with count data, our dataset was overdispersed (Richards 2008), with greater variability than

expected (variance/mean > 1). We used additive overdispersion to account for overdispersion in our model by including an individual-level random effect, adding a random intercept with one level per observation that captures overdispersion (Elston *et al.* 2001, Browne *et al.* 2005). Residual variation was not included because Poisson GLMM only has one parameter, with a known mean-variance relationship, so there is no estimate for residual variation (Sokal and Rohlf 1981), but the individual-level effect is the variation not explained by other random effects. First we compared models with and without fixed effects; the difference in variance between the models provided an estimate of detector-level variance for the fixed effect (Elston *et al.* 2001). We then continued model selection by dropping random effects with non-significant variance estimates and model comparison with likelihood ratio tests and Akaike's Information Criterion (AIC). We calculated the proportion of variation explained by each factor in the full model in order to test if space or time explains the most variation in activity levels.

5.2.5 Percentile thresholds

To detect peaks and patterns in activity among detectors within nights, we used two methods: percentile thresholds (Adams *et al.* submitted) and SaTScan (Adams and Fenton submitted). We calculated percentile thresholds (50th and 90th percentiles) of activity for each species with a larger distribution of activity from 14,898 detector hours, after excluding hours without activity. We created this larger distribution using data from our four Ontario sites in 2008 and 2010, four sites in Saskatchewan, Canada in 2009, and one site near the Hudson River in New York, USA in 2009. When the number of calls per hour is greater than the 50th percentile threshold, we consider activity to be "high" since it is above the majority of the hourly activity for a species from the larger distribution.

Activity over the 90th percentile threshold was considered “very high” activity. We then identified the number of hours above each threshold for each species on every night at all detectors and sites in Ontario.

5.2.6 SaTScan

We identified peaks in activity with SaTScan software package (v. 9.1.1, Kulldorff *et al.* 2005, following Adams and Fenton submitted), which can identify clusters of activity in space and time that are significantly different from the background expectation. We specified a one-tailed analysis to look for clusters with unexpectedly high numbers of calls with the space-time permutation model (Kulldorff *et al.* 2005). We used retrospective analysis, specified spatial and temporal maximum cluster sizes (MCS) at 50%, and used 999 Monte Carlo replications. We input case files consisting of the number of calls per minute, MAS, and detector location within the site. The coordinates file included the Cartesian coordinates (x, y, z) of each detector location. The analysis output included the “most likely cluster” and “secondary clusters” which described the locations (detectors) and time frame(s) when activity was significantly higher than the background expectation with the observed and expected number of cases and p-value. We analyzed activity for each species individually. We used multivariate SaTScan (Kulldorff *et al.* 2007) with case files for each night at a site as separate datasets. For 2008, we ran multivariate SaTScan twice for each site, for the first six nights (spring) and the last six nights (summer). Since multivariate SaTScan is limited to 12 datasets, we ran the 15 consecutive nights from 2010 in two batches, the first 12 nights and again with nights four to 15. We then compared locations and timing of peak activity that occurred on the majority of nights among all species with Gantt charts.

5.2.7 Sub-sampling for estimates of activity

To determine the effect of number of nights and amount of space sampled on estimates of activity levels, we randomly sampled subsets of nights to compare the mean activity level of the subsets to a grand mean of the total sample (following Hayes 1997). We sampled two to ten night subsets 100 times for each species and all species combined for each detector individually. We created a loop to run *sample* without replacement 100 times in R for each subset of nights. We calculated the mean nightly activity for each subset and compared it to the grand mean of the full dataset of all 15 nights at each detector in 2010. We determined the proportion of subsamples within 10-50% of the mean of the grand mean. We then expanded the analysis for the full model to include mean activity of five detectors over all nights (2008 = 12 nights, 2010 = 15 nights). We sampled one to five detector subsets for each two to ten night subset 100 times and determined the proportion of subsamples within 30% of the grand mean. We based recommendations on how many nights had at least 80% of the subsamples within 30% of the grand mean (following Hayes 1997).

5.3 Results

5.3.1 Differences among detectors

Every species was present at all sites, but species-specific activity levels varied dramatically among and within sites. Overall activity, for all species combined, was significantly different among detectors within each site, except Renfrew mine in 2008 (Long Point $H_4 = 182.49$, $p < 0.001$; QUBS $H_4 = 45.46$, $p < 0.001$; Renfrew lake $H_4 = 96.66$, $p < 0.001$; Renfrew mine 2010 $H_4 = 45.08$, $p < 0.001$). The majority of species-specific activity was significantly different among detectors within each site, except *L.*

noctivagans, and *M. septentrionalis* at QUBS and Renfrew lake (Table B1). Overall, estimates of activity levels at a site differed substantially depending on detector placement (Fig. 5.1, B1).




Figure 5.1. Mean calls per hour \pm SD for each species of bat at five detectors at Long Point, Ontario ($n = 12$ nights). Note different y-axes. Detectors of the same letter were paired at the same location and first detectors in the transect (A1/A2) had intermediate openness at the site (20% canopy cover), B1/B2 were the most enclosed (37% canopy cover), and C1 was the most open (5% canopy cover). Detector number denotes height: 1 was low (1.5 m above ground) and 2 was high (4 m). Means, within each species, with the same lowercase letter are not significantly different ($p > 0.05$) according to a Kruskal-Wallis multiple comparison post hoc test.

5.3.2 Generalized linear mixed models: variance partitioning of random effects

The proportion of variance explained by each random effect differed among species in 2008 (Fig. 5.2). Detector position and night-to-night differences significantly explained of activity levels for every species ($p < 0.05$), but site was not significant for all species (Fig. 5.2, Table B2). Spatial components had the greatest association with variation in activity levels for *L. noctivagans*, *L. borealis*, and *M. lucifugus*. Detector height was only significant for explaining variation in *L. noctivagans* activity ($p = 0.0251$, Table B2) with activity levels higher at higher detectors. Canopy cover explained a significant amount of variation for all species ($p < 0.05$, Table B2) with activity increasing as canopy cover decreased, but *M. septentrionalis* was more active with increased canopy cover. Canopy cover had the greatest impact on *L. borealis*, with 27% of the variation attributable to detector location due to the effects of canopy cover.

During 2010, the majority of within-site variation was associated with the individual-level effect, except for *M. septentrionalis* where the majority of variation was explained by differences among detector locations. The two species with the least activity at the site in 2010, *E. fuscus* and *L. noctivagans*, had the most variation associated with the individual-level effect. Between detector location and night-to-night differences, spatial effects contributed more to variation in activity for *L. borealis*, *L. cinereus*, and *M. septentrionalis*. Spatial and temporal effects equally contributed to variation in activity for *M. lucifugus* and *P. subflavus*. Canopy cover was a significant factor for *L. borealis* and *L. cinereus*, explaining 37% and 21% of the detector-level effects, with activity increasing as canopy cover decreased.




Figure 5.2. Proportion of variance explained by spatial (detector, site) and temporal (night) random effects for seven species of bats at four sites in Ontario, Canada. Within-site variation (detector location) is an important component in explaining variation in activity levels of all species combined. The importance of within-site, among-site, and night-to-night effects differs by species.

5.3.3 Percentile thresholds: magnitude of activity

The number of nights that each species had high activity, above the 50th and 90th percentile thresholds (Table 5.3), differed among all detectors at all sites (Table 5.4,B3). The “picture” of activity differed depending on detector placement. For example, at Renfrew mine in 2010 both *Myotis* species had high activity on the majority of the nights at all detectors. However, if only detector C were present, it would not be evident that *M. lucifugus* and *L. borealis* had very high activity at the site (Table 5.4). If not all detectors were present at the site in 2010 then the magnitude of activity for each species would not be evident. It was not necessary to have detectors at both heights in 2008 because the number of nights with high activity between paired detectors (1.5 and 4 m) was always the same or within one to two nights, except at QUBS (*L. noctivagans*, *L. cinereus*, and *M. septentrionalis*) and Long Point (*L. noctivagans*) where there was higher activity on more nights at one of the high detectors (Table B3).

Table 5.3. Number of bat echolocation calls per hour at the 50th and 90th percentile thresholds for each species and all species combined for a distribution of activity at eight sites in Canada and New York. Activity above the 50th percentile (median) is considered high and activity above the 90th percentile threshold is very high.


Species	Number of calls per hour at percentile thresholds	
	50 th	90 th
Species combined	218	2793
<i>Eptesicus fuscus</i>	35	308
<i>Lasionycteris noctivagans</i>	6	32
<i>Lasiurus borealis</i>	20	166
<i>L. cinereus</i>	31	327
<i>Myotis lucifugus</i>	87	1426
<i>M. septentrionalis</i>	42	960
<i>Perimyotis subflavus</i>	21	140

Table 5.4. Number of nights when at least one hour of bat activity was above the 50th and 90th percentile thresholds at each detector (A-E) over 15 nights at an abandoned mine near Renfrew, Ontario in 2010.

Species	50th percentile Detectors					90th percentile Detectors				
	A	B	C	D	E	A	B	C	D	E
Species combined	10	15	15	15	13	7	14	5	5	2
<i>Eptesicus fuscus</i>	1	1		2		1				
<i>Lasionycteris noctivagans</i>	4	1		1		1	1		1	
<i>Lasiurus borealis</i>	9	5		15	8		1		13	
<i>L. cinereus</i>	2	1		7	1				1	
<i>Myotis lucifugus</i>	10	15	15	15	14	6	7		7	2
<i>M. septentrionalis</i>	10	15	15	13	15	4	14	10		1
<i>Perimyotis subflavus</i>	3	4		10	3	1	1		1	

5.3.4 SaTScan: peaks of activity in space and time

Peaks of activity, detected by SaTScan, differed among detector locations, even between paired detectors at two heights, at all sites (Fig. 5.3,B1). Although activity patterns varied, we were unable to detect evidence of spatial partitioning among species, every species had overlapping activity on at least one detector at each site. All species had peaks of activity at every detector during at least one season, except *E. fuscus* that never had peaks of activity at detectors B1/B2 Long Point (Fig. 5.3A) or detector C1 at QUBS (Fig. B1.2). There was some degree of temporal partitioning among species within sites, for example *Myotis lucifugus* and *M. septentrionalis* did not have overlapping activity peaks at any detector at Long Point (Fig. 5.3A); neither did *L. noctivagans*, *L. borealis*, and *P. subflavus* (Fig. 5.3A). At Renfrew 2010 only three species had significant peaks of activity on the majority of nights and timing of peaks overlapped among all three species (Fig. 5.3B). *Lasionycteris noctivagans* and *L. borealis* peaks of activity rarely overlapped temporally in 2008 (Fig. B1).






Figure 5.3. Peak periods of bat activity along linear, stationary transects within two sites: A) during late summer (migration) at Long Point, Ontario in 2008 and B) during swarming (August) at an abandoned mine near Renfrew, Ontario in 2010. Species-specific activity patterns vary among detectors. Detectors at Long Point were paired at two heights, low (1.5 m; A1, B1, and C1) and high (4 m; A2 and B2). All detectors at Renfrew were at 1.5 m. There is no evidence of spatial partitioning, but temporal partitioning (no time overlap of peak activity) is evident between several species, such as *L. noctivagans* and *L. borealis*. EPFU – *Eptesicus fuscus*, LANO – *Lasionycteris noctivagans*, LABO – *Lasiurus borealis*, LACI – *L. cinereus*, MYLU – *Myotis lucifugus*, MYSE – *M. septentrionalis*, PESU – *Perimyotis subflavus*.

5.3.5 Sub-sampling for estimates of activity

As the number of nights increased so did the probability of obtaining mean estimates of activity more similar to the grand mean (Table 5.5,5.6). When sampling with one detector and species identification was not taken into account, it required three to nine nights to have at least 80% of the subsamples within 30% of the grand mean in 2010 (Table 5.5). When we accounted for species, the number of nights increased; the most active species, *M. lucifugus* and *M. septentrionalis*, required fewer nights of sampling than any of the other species at the site (Table 5.5). When the grand mean included multiple detectors within a site, we required at least four detectors recording for a minimum of four nights to have accurate estimates of overall activity across a site (Table 5.6A,B). We required sampling for at least five nights with five detectors for the most active species within a season (Table 5.6B). Even within a season it was not possible for a single detector moving locations for fifteen nights to be within 30% of the grand mean. Fewer nights were required when all nights in the grand mean were during a single season (Table 5.6B), as opposed to during multiple periods in the year (Table 5.6A).

Table 5.5. Mean \pm SD number of nights required to have at least 80% of random samples with mean nightly bat activity within 10-50% of the grand mean of the entire dataset (total calls per night for one detector over 15 nights) for six detectors at a mine near Renfrew, Ontario, Canada in 2010. Data were randomly sampled 100 times to include nightly activity for one detector for two to ten night sample periods. Empty cells are when more than ten nights of sampling were required for all detectors.

Species	Percent deviation from grand mean				
	$\leq 10\%$	$\leq 20\%$	$\leq 30\%$	$\leq 40\%$	$\leq 50\%$
Species combined		8.4 \pm 3.4	5.6 \pm 3.4	4.3 \pm 1.7	3.0 \pm 1.4
<i>Eptesicus fuscus</i>					9.2 \pm 1.8
<i>Lasionycteris noctivagans</i>					8.8 \pm 4.7
<i>Lasiurus borealis</i>			8.4 \pm 4.2	6.8 \pm 4.1	5.6 \pm 3.8
<i>L. cinereus</i>				10.6 \pm 1.7	8.0 \pm 2.5
<i>Myotis lucifugus</i>			6.8 \pm 3.7	5.8 \pm 4.0	4.4 \pm 2.9
<i>M. septentrionalis</i>			7.6 \pm 3.6	5.6 \pm 3.7	4.4 \pm 3.8
<i>Perimyotis subflavus</i>			10.5 \pm 2.1	8.5 \pm 1.5	7.4 \pm 1.8

Table 5.6. Number of nights required to have at least 80% of random samples with mean nightly bat activity within 30% of the grand mean of the entire dataset at A) four sites in Ontario, Canada in 2008 (five detectors over 12 nights) and B) one site in 2010 (five detectors over 15 nights). Data were randomly sampled 100 times to include nightly activity for one to five detectors for two to ten night sample periods for each site. Empty cells are when more than ten nights of sampling were required for all sites.

A)					
Species	Number of detectors in subsample				
	1	2	3	4	5
Species combined		10.5 ± 3	10.3 ± 3.5	9.8 ± 4.5	7.3 ± 4.0
<i>Eptesicus fuscus</i>				11.0 ± 2.0	9.5 ± 3.3
<i>Lasionycteris noctivagans</i>			11.3 ± 1.5	9.8 ± 2.6	8.5 ± 4.1
<i>Lasiurus borealis</i>				10.5 ± 3.0	8.8 ± 3.0
<i>L. cinereus</i>			11.2 ± 2.0	8.8 ± 3.8	9.8 ± 4.5
<i>Myotis lucifugus</i>					10.3 ± 3.5
<i>M. septentrionalis</i>			10.3 ± 3.5	8.5 ± 4.1	7.8 ± 5.1
<i>Perimyotis subflavus</i>					11.0 ± 1.7

B)					
Species	Number of detectors in subsample				
	1	2	3	4	5
Species combined				4	3
<i>E. fuscus</i>					
<i>L. noctivagans</i>					
<i>L. borealis</i>					4
<i>L. cinereus</i>					
<i>M. lucifugus</i>				9	5
<i>M. septentrionalis</i>					5
<i>P. subflavus</i>					9

5.4 Discussion

Patterns of bat activity vary within a habitat in response to a variety of biotic (e.g., prey abundance) and abiotic factors (e.g., habitat structure). Our results demonstrate significantly different depictions of bat activity depending on where a detector was placed within a site, even when relatively close together (40 m), including differences in number of calls, magnitude of activity, and temporal patterns of activity. While there was some effect of detector height and canopy cover on activity levels, these factors did not explain the majority of the variation from detector location. Differences in where peaks of activity occurred among detectors indicate that it is important to place multiple detectors throughout a site in order to capture spatial variation in activity.

Our results differed from those in other regions, suggesting that the relative importance of sources of variation can vary considerably regionally. Variation among sites was less important in our study than in Australia (e.g., Fischer *et al.* 2009), accounting for only 13% of variation on average. Within-site heterogeneity was most important at our sites, accounting for two-thirds of variation in activity levels, while it was not as significant in other studies (Krusic *et al.* 1996b, Fischer *et al.* 2009). It is clear that experimental design will impact what we see in snap-shots of bat activity at a site. The goal is to have the most accurate estimate of activity, while the extent of variation determines where effort needs to be focused to increase accuracy. When interested in overall activity levels, it is most important to sample for fewer nights and more locations within a site, but the number of nights sampled must increase when estimating species-specific activity.

Detectors placed at greater heights did not help to account for variation in activity levels, except for one species. It is possible that the small difference in heights (2.5 m) and low

vertical vegetation structure was not biologically meaningful or high enough to reveal a difference. However, detectors with a small difference in height have shown disparity of activity levels (Fischer *et al.* 2009), although these differences could be attributed to slightly different detector locations within the site. There is value in sampling at heights greater than 1.5 m (e.g., Krusic *et al.* 1996, Reynolds 2006), but our study suggests this is only worthwhile if detectors are at heights greater than 4 m. While we saw differences between detector heights in timing of activity peaks (e.g., Fig. 5.3) and the number of nights with activity above various percentile thresholds, we suggest a better use of multiple detectors would be to sample at more points throughout the site to account for horizontal heterogeneity in similar regions.

The only evidence of temporal partitioning was between *L. noctivagans* and *L. borealis*, which corresponds to findings by Kunz (1973) in Iowa, most likely driven by preferences for different insect prey. It is possible that this finding is an artifact of relatively low activity levels of *L. noctivagans* at the majority of our sites, but warrants further investigation into these species' activity patterns. Sampling for the entire night is a requirement for any study interested in activity levels (Hayes 1997, Skalak *et al.* 2012). While many studies detect a peak of activity in the first two hours after sunset (e.g., Kunz 1973, Hayes 1997, Broders *et al.* 2003), only *L. cinereus* regularly had peaks of activity in this period during our study. Where and when peaks in species-specific activity occurred differed enough among detectors to suggest that sampling at a single point or for a portion of the night would give only a partial picture of bat activity at a given site.

Greater sampling effort, in both space and time, will always lead to more accurate estimates of activity levels. When sampling with a single detector, regardless of species,

our findings agreed with those of Hayes (1997), which recommended sampling for no less than six nights. It appears we had greater variation among detectors within a site (Table 5.5) than Hayes (1997) did among sites and years. Once species-specific estimates of activity for a site include multiple sampling points within a site, it is not possible to have accurate estimates of activity with a single detector. Species with relatively high activity levels require fewer nights of sampling; the number of nights sampled at a site would need to be either based on a particular species of interest or on the least active species at the site. Acoustically rare species, with lower detection probabilities, have higher variation in activity levels, more time with no activity, requiring more nights to simply detect these species (Skalak *et al.* 2012). Using one detector and moving it to a different location each night, while an option at a large spatial scale (Skalak *et al.* 2012), is not as desirable at a small scale. We required fewer nights when recording with multiple detectors simultaneously within a site. Blocked sampling designs are more efficient than completely randomized designs for comparisons among sites, requiring fewer nights of sampling (Hayes 1997), which was also the case for sampling within sites.

There is growing concern of inadequate accounts of variation in acoustic studies (Hayes 2000, Sherwin *et al.* 2000, Gannon *et al.* 2003, Skalak *et al.* 2012). Considering our results, the use of multiple detectors within a site should be standard practice for all acoustic surveys (Duchamp *et al.* 2006, Fischer *et al.* 2009, Stahlschmidt and Brühl 2012). Our findings caution extrapolation of results from a single detector to estimate activity of a site. The importance of among- and within-site variation differs among regions and it may be possible to use fewer detectors with less structural complexity at a site (Britzke 2003). If using fewer detectors it would be worthwhile conducting a

preliminary field season to see where important sources of variation are in order to focus sampling efforts more efficiently in future seasons. If experimental designs do not adequately account for within-site variation, activity estimates will be biased.

5.5 Acknowledgements

We thank Dr. S. Brinkløv, Dr. E. Fraser, R. Hamilton, M. Jantzen, and Dr. K. Stammer for feedback on this manuscript. Thanks to the Long Point Bird Observatory, Queen's University Biological Station (QUBS), and the Windle family for housing and access to their properties. Funding was provided by a student research scholarship from Bat Conservation International to A.M.A. and Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery grant to M.B.F.

5.6 References

- Adams, A. M., M. K. Jantzen, R. M. Hamilton, and M. B. Fenton. 2012. Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats. *Methods in Ecology and Evolution* 3:992–998.
- Adams, A. M., and M. B. Fenton. Identifying peaks in bat activity: a new application of the space-time scan statistic. submitted.
- Adams, A. M., L. P. McGuire, L. A. Hooton, and M. B. Fenton. How high is high? Using percentile thresholds to identify peak bat activity. submitted.
- Adams, R. A., and K. M. Thibault. 2006. Temporal resource partitioning by bats at water holes. *Journal of Zoology* 270:466–472.

- Arh, M. R. 2009. Foraging ecology of a bat assemblage. MSc Thesis. University of Toronto.
- Barclay, R. M. R. 1982. Night roosting behavior of the little brown bat, *Myotis lucifugus*. *Journal of Mammalogy* 63:464–474.
- Bates, D., M. Maechler, and B. Bolker. 2011. lme4: Linear mixed-effects models using S4 classes. R package version 0.999375-42. <http://CRAN.R-project.org/package=lme4>.
- Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution* 24:127–35.
- Britzke, E. 2003. Use of ultrasonic detectors for acoustic identification and study of bat ecology in the eastern United States. PhD Thesis. Tennessee Technological University.
- Britzke, E. R., E. H. Gillam, and K. L. Murray. 2013. Current state of understanding of ultrasonic detectors for the study of bat ecology. *Acta Theriologica* 58:109–117.
- Broders, H. 2003. Another quantitative measure of bat species activity and sampling intensity considerations for the design of ultrasonic monitoring studies. *Acta Chiropterologica* 5:235–241.
- Broders, H., G. Quinn, and G. Forbes. 2003. Species status, and the spatial and temporal patterns of activity of bats in southwest Nova Scotia, Canada. *Northeastern Naturalist* 10:383–398.

- Browne, W., S. Subramanian, K. Jones, and H. Goldstein. 2005. Variance partitioning in multilevel logistic models that exhibit overdispersion. *Journal of the Royal Statistical Society Series A* 168:599–613.
- Ciechanowski, M., T. Zając, a. Biłas, and R. Dunajski. 2007. Spatiotemporal variation in activity of bat species differing in hunting tactics: effects of weather, moonlight, food abundance, and structural clutter. *Canadian Journal of Zoology* 85:1249–1263.
- Duchamp, J., M. Yates, R. Muzika, and R. Swihart. 2006. Estimating probabilities of detection for bat echolocation calls: an application of the double-observer method. *Wildlife Society Bulletin* 34:408–412.
- Dzal, Y., L. A. Hooton, E. L. Clare, and M. B. Fenton. 2009. Bat activity and genetic diversity at Long Point, Ontario, an important bird stopover site. *Acta Chiropterologica* 11:307–315.
- Elston, D., R. Moss, T. Boulinier, C. Arrowsmith, and X. Lambin. 2001. Analysis of aggregation, a worked example: numbers of ticks on red grouse chicks. *Parasitology* 122:563–569.
- Fenton, M. B. 1969. Summer activity of *Myotis lucifugus* (Chiroptera: Vespertilionidae) at hibernacula in Ontario and Quebec. *Canadian Journal of Zoology* 47:597–602.
- Fenton, M. B. 1970. Population studies of *Myotis lucifugus* (Chiroptera: Vespertilionidae) in Ontario. *Life Sciences Contributions Royal Ontario Museum* 77:1–34.

- Fischer, J., J. Stott, B. S. Law, M. D. Adams, and R. I. Forrester. 2009. Designing effective habitat studies: quantifying multiple sources of variability in bat activity. *Acta Chiropterologica* 11:127–137.
- Gannon, W., R. Sherwin, and S. Haymond. 2003. On the importance of articulating assumptions when conducting acoustic studies of habitat use by bats. *Wildlife Society Bulletin* 31:45–61.
- Giraudoux, P. 2011. *pgirmess: Data analysis in ecology*. R package version 1.5.2. <http://CRAN.R-project.org/package=pgirmess>.
- Hayes, J. 1997. Temporal variation in activity of bats and the design of echolocation-monitoring studies. *Journal of Mammalogy* 78:514–524.
- Hayes, J. 2000. Assumptions and practical considerations in the design and interpretation of echolocation-monitoring studies. *Acta Chiropterologica* 2:225–236.
- Hayes, J. and M. Adam. 1996. The influence of logging riparian areas on habitat utilization by bats in western Oregon. Pages 228–237 *in* R. M. R. Barclay and R. M. Brigham, editors. *Bats and Forests Symposium*.
- Hayes, J., and J. Gruver. 2000. Vertical stratification of bat activity in an old-growth forest in western Washington. *Northwest Science* 74:102–108.
- Krusic, R., M. Yamasaki, C. D. Neefus, and P. J. Perkins. 1996. Bat habitat use in White Mountain National Forest. *Journal of Wildlife Management* 60:625–631.

- Kulldorff, M., R. Heffernan, J. Hartman, R. Assunção, and F. Mostashari. 2005. A space-time permutation scan statistic for disease outbreak detection. *PLoS Medicine* 2:e59.
- Kulldorff, M., F. Mostashari, L. Duczmal, W. Katherine Yih, K. Kleinman, and R. Platt. 2007. Multivariate scan statistics for disease surveillance. *Statistics in Medicine* 26:1824–33.
- Kunz, T. 1973. Resource utilization: temporal and spatial components of bat activity in central Iowa. *Journal of Mammalogy* 54:14–32.
- McGuire, L. P., C. G. Guglielmo, S. A. Mackenzie, and P. D. Taylor. 2012. Migratory stopover in the long-distance migrant silver-haired bat, *Lasionycteris noctivagans*. *Journal of Animal Ecology* 81:377–85.
- Milne, D., A. Fisher, I. Rainey, and C. Pavey. 2005. Temporal patterns of bats in the Top End of the Northern Territory, Australia. *Journal of Mammalogy* 86:909–920.
- Nicholls, B., and P. Racey. 2006. Habitat selection as a mechanism of resource partitioning in two cryptic bat species *Pipistrellus pipistrellus* and *Pipistrellus pygmaeus*. *Ecography* 5:697–708.
- O’Hara, R. B., and D. J. Kotze. 2010. Do not log-transform count data. *Methods in Ecology and Evolution* 1:118–122.
- R Development Core Team. 2011. R: a language and environment for statistical computing. R Foundation for Statistical Computing. v. 2.13.1. R Foundation for Statistical Computing, Vienna, Austria.

- Reynolds, D. S. 2006. Monitoring the potential impact of a wind development site on bats in the northeast. *Journal of Wildlife Management* 70:1219–1227.
- Sherwin, R., W. Gannon, and S. Haymond. 2000. The efficacy of acoustic techniques to infer differential use of habitat by bats. *Acta Chiropterologica* 2:145–153.
- Skalak, S. L., R. E. Sherwin, and R. M. Brigham. 2012. Sampling period, size and duration influence measures of bat species richness from acoustic surveys. *Methods in Ecology and Evolution* 3:490–502.
- Sokal, R., and F. Rohlf. 1981. *Biometry*. WH Freeman and Company, 2nd edition. WH Freeman and Company, New York, NY.
- Stahlschmidt, P., and C. A. Brühl. 2012. Bats as bioindicators – the need of a standardized method for acoustic bat activity surveys. *Methods in Ecology and Evolution* 3:503–508.
- Staton, T., and S. Poulton. 2012. Seasonal variation in bat activity in relation to detector height: a case study. *Acta Chiropterologica* 14:401–408.

6 Conclusion

The main goal of this dissertation was to investigate sources of variation in acoustic monitoring of bat activity and how this impacts acoustic survey design. I addressed this goal in three ways: i) variation from bat detectors, ii) methods for objective identification of peaks of activity, and iii) the use of stationary, linear transects to address within-site spatial and temporal variation. While several studies have examined variation in acoustic monitoring of bats and made recommendations for future surveys (Hayes 1997, 2000, Fischer *et al.* 2009, Skalak *et al.* 2012), more are needed to understand sources of variation and improve methods to develop standardized sampling protocols. Combining the results of this research, I have provided insight into how sources of variation and methods of data analysis can impact interpretation of acoustic data of bat activity. Detailed discussions of specific experimental results are presented in the pertinent chapters. Here, I summarize the cumulative findings of my dissertation and make recommendations for future acoustic bat surveys. Finally, I conclude with a description of challenges and gaps in our knowledge challenges to our greater understanding of bat activity, and suggestions for future research directions.

6.1 Contribution to acoustic studies of bats

An integral component of conducting effective acoustic surveys is accounting for sources of variation, an essential step to collecting unbiased data (Hayes 2000), while using objective methods to analyze that data. This information is necessary for effective sampling design, data analysis, and interpretation of results. I have contributed to this field in three ways:

6.1.1 Variation from detectors

Many factors can increase variation in an acoustic dataset, one of which is the detector used (Adams *et al.* 2012/Chapter 2). I found that detector performance varied among systems, differing in sensitivity and directionality, and is an important source of variation in acoustic monitoring methods that must be accounted for when creating sampling protocols and comparing datasets. The value of these results is not simply as a comparison of different detectors, but as a demonstration of how different technologies can give different results; an issue that needs to be considered when designing and evaluating any studies using these technologies.

6.1.2 Objective methods for detecting peaks in bat activity

A limitation of acoustic monitoring is the relative nature of data interpretation. Previously, conclusions have been based on subjective assessments about the relative importance of sites or species-specific activity patterns. I proposed two methods for objectively identifying peaks in bat activity at various scales: percentile thresholds (Chapter 3) and SaTScan (Chapter 4). Using percentile thresholds to assess acoustic data permits an unbiased measure of the importance of a site and is a replicable method of describing within-night activity patterns. The strength of this method is evaluating activity levels at several thresholds based on a larger distribution of activity among sites. SaTScan is a valuable tool for quickly identifying peaks with an objective, replicable, and statistically-sound method that can be applied at various temporal and spatial scales. Using these two methods in combination permits a thorough investigation of activity

levels and patterns at a site, from the magnitude of species-specific activity to comparison of timing of peaks among species or sites.

6.1.3 Variation within sites

Bat activity can vary temporally (e.g, Hayes 1997, Milne *et al.* 2005), but within-site spatial variation has been too often overlooked (Britzke 2003, Fischer *et al.* 2009). I found that within site factors are very important for understanding variation in bat activity, being as or more important than differences among sites (Chapter 5). The high degree of variation within sites can affect sampling design, including necessary sampling effort, and requires the use of multiple detectors recording simultaneously within a site. Detector placement within a site dramatically impacts the depictions of activity, in turn impacting estimates of levels and patterns of activity. An *a priori* understanding of the survey effort necessary should ensure statistically powerful sampling designs, clearer data interpretation, and more successful management and conservation actions.

6.2 Recommendations for future acoustic surveys

To use acoustic monitoring to address ecological questions, it is important to know how sources of variation affect data collection and thus the data itself. While there is no simple formula for what constitutes an ideal survey effort, it is clear that additional effort will result in more precise estimates of activity. Accuracy increases with the number of nights sampled and detectors deployed. It is important to first clearly define the research question and decide on the best study design to test the predictions. If the aim of a study is to determine overall activity levels at a site then a site in Ontario would require

sampling for at least four nights with four detectors within a season, but would require an increased sampling effort when evaluating species-specific activity. It is difficult to extrapolate from my results because the degree of habitat heterogeneity differs among sites. I recommend using preliminary studies to determine the number of detectors and nights necessary to obtain an accurate estimate of activity before establishing a long-term monitoring program. I echo the recommendations of other authors (Hayes 1997, Skalak *et al.* 2012) that monitoring should be done continuously through the night. Ideally, sampling should occur for as long as possible; this is relatively easy with passive methods, but long-term datasets can be inhibiting in terms of analysis.

It is important to use a single brand of detector for a monitoring program and to report detector settings in publications to ensure comparable results among locations and years (Adams *et al.* 2012/Chapter 2). Detectors should be calibrated to reduce variation among detectors of the same brand and among sampling periods (Larson and Hayes 2000).

Passive detection systems with an automatic trigger are best for developing standardized sampling protocols because they remove biased sampling methods and require little effort for deployment (Stahlschmidt and Brühl 2012). Choice of bat detector will depend on the research question being asked and potentially be influenced by budgetary constraints.

Study location and focal species will determine which detectors are appropriate based on their frequency response. Wildlife Acoustics' SongMeter SM2BAT has two different models that differ in sampling rate and the lower sampling rate model would not be adequate to record all species present in the Neotropics. Full-spectrum detectors are a better choice for the majority of research questions since they are more sensitive, with greater detection ranges (Adams *et al.* 2012/Chapter 2), and collect more information than

frequency division systems, leading to more accurate species identification (Fenton 2000). If asking questions about echolocation behavior then a more sensitive and calibrated microphone will be important. Research questions about activity levels at a particular site will require decisions on a trade-off between the more expensive detector (i.e., Avisoft, Batlogger) that detects calls in a larger volume of airspace at a given location or a less expensive option (i.e., SongMeter). Also involved in the decision is the importance of simultaneously monitoring multiple locations within a site. Sampling area heterogeneity and access to multiple detectors will impact this decision.

Successful application of acoustic monitoring to detect within-site variation requires the use of multiple detectors simultaneously (Chapter 5). Understanding structural heterogeneity at a site can determine the number of detectors necessary to capture vertical and horizontal variation in bat activity. A reasonable survey effort will depend on the objectives of a particular study. While my recommendations are for surveys sampling patterns and levels of activity, they are relatively in line with surveys for species richness. Recording continuously for the entire night together with increased sampling effort for more nights at more locations will increase chances of detecting rare species (Skalak *et al.* 2012).

Finally, it is necessary to use objective analytic methods for acoustic data because of the already inherent relative nature of the data. Use of programs, such as SaTScan, makes analysis consistent and replicable. It is necessary to measure activity levels relative to a large distribution, which is closer to the ground truth of what is present in nature. The next step is to create a public repository of acoustic datasets to evaluate activity of a species in the context of its entire range, allowing standardization of terms such as “high

activity.” Standardization makes it possible to review methods used for environmental assessments and creation of protocols for unified monitoring programs among regions.

6.3 Future research directions

It is clear from my results (Chapter 5) and those of Fischer *et al.* (2009) that activity from a single location does not reflect all locations within a site. My specific findings about within-site variation are unlikely to be directly applicable to other regions because of varying habitat heterogeneity. Vertical spatial partitioning is evident in many habitats (Hecker and Brigham 1999, Kalcounis *et al.* 1999, Hayes and Gruber 2000), but it has not been established what acoustic sampling effort is necessary to detect these patterns.

Sampling vertical distributions is limited by the logistics of raising detectors to greater heights; most successful is opportunistic placement on manmade structures, such as towers and poles (Kalcounis *et al.* 1999, Hamilton 2012) and wind turbines (Reynolds 2006), or attaching them to trees (Staton and Poulton 2012b). Further research into patterns of vertical and horizontal spatial variation among habitats is necessary, especially with respect to how habitat structure and differences in insect distributions play roles in determining bat activity. Ideally, future research will provide insights into how transferable findings regarding sampling effort are to other regions with the goal of establishing what measures are necessary to determine sampling efforts in new regions without extensive preliminary study.

Most bat surveys in temperate areas primarily use acoustic methods because of the detectability of the echolocation calls of most insectivorous species, allowing development of standardized protocols based on acoustic monitoring in these regions.

However, acoustic monitoring is not a “silver-bullet” for sampling all bat communities; capture methods are required to sample whispering bats with low intensity echolocation calls and those that do not echolocate at all (Griffin 1958, Fenton 2003). Creating standardized sampling protocols for regions with greater species diversity, such as the tropics, will require an understanding of factors influencing capture success (Kalko and Handley 2001) and how recommendations for acoustic sampling effort would differ.

A major limitation to conservation and management efforts is knowledge of where important sites are for bats. Research into the detectability of special sites, such as roosts, hibernacula, swarming sites, and migration stopover sites, would be extremely valuable. Understanding horizontal spatial variation will play a role, with how patterns of activity drop off as bats move away from these special sites and the proximity to a special site required to detect levels of activity high enough to be notable.

Walking and driving transects are methods used increasingly for standardized bat surveys. The UK’s National Bat Monitoring Program uses volunteers with detectors walking 1 km transects to sample bats and has been successful at detecting population trends over time (Walsh *et al.* 2001). At least 17 states (Herzog and Britzke 2009) and one province use driving transects to sample bat activity levels post-white-nose syndrome (WNS, Britzke and Herzog). Efforts to collect long-term datasets in a standardized and comparable fashion are laudable, but there is little scientific literature to support the use of this method. Russ *et al.* (2003) describe the use of a driving transect and discuss its validity, but make no effort to compare the method, and this is likely what most driving surveys protocols are based on. Stahlschmidt and Brühl (2012) have been the only researchers to compare moving transects to stationary detectors, finding that walking transects fail to

represent the heterogeneous bat activity patterns and stationary detectors have the greatest potential for standardized surveys. There is an urgent need for research into the feasibility of moving transect surveys.

While much research is focused on how to survey at wind energy developments to determine which sites will be high-risk for bats (Reynolds 2006, Arnett *et al.* 2011, Korner-Nievergelt *et al.* 2011), there is still little information available to guide policy and permit critical evaluation of wind energy development proposals and environmental assessment reports. Percentile thresholds (Chapter 3) are the first proposed method for objectively comparing the importance of a site to a species of bat, but we need continued development and research into how environmental recommendations, with the potential to impact survival of numerous bats species, are determined.

6.4 Concluding remarks

Acoustic monitoring studies have a number of inherent limitations and assumptions, but the use of bat detectors can be a very powerful tool for insights into the ecology and behavior of bats. Design of any study includes trade-offs between research objectives and logistics. As acoustic monitoring is increasingly used for large-scale management and conservation efforts in response to growing threats to bat populations, the need for research to support standardized protocols is also rising. The results of my dissertation have provided an increased understanding of how variation plays a role in sampling bat activity. I hope that the information presented here will provide a platform for continued research into objective methods and standardized protocols of studying bats in order to streamline bat conservation efforts of this ecologically important group of mammals.

6.5 References

- Adams, A. M., M. K. Jantzen, R. M. Hamilton, and M. B. Fenton. 2012. Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats. *Methods in Ecology and Evolution* 3:992–998.
- Arnett, E. B., M. M. Huso, M. R. Schirmacher, and J. P. Hayes. 2011. Altering turbine speed reduces bat mortality at wind-energy facilities. *Frontiers in Ecology and the Environment* 9:209–214.
- Britzke, E. R. 2003. Use of ultrasonic detectors for acoustic identification and study of bat ecology in the eastern United States. PhD Thesis. Tennessee Technological University.
- Britzke, E. R., and C. Herzog. Using acoustic surveys to monitor population trends in bats. <http://fw.ky.gov/pdf/acoustictransectguidance.pdf>. Date Accessed: 24 March 2013.
- Fenton, M. B. 2000. Choosing the “correct” bat detector. *Acta Chiropterologica* 2:215–224.
- Fenton, M. B. 2003. Science and the conservation of bats: where to next? *Wildlife Society Bulletin* 31:6–15.
- Fischer, J., J. Stott, B. S. Law, M. D. Adams, and R. I. Forrester. 2009. Designing effective habitat studies: quantifying multiple sources of variability in bat activity. *Acta Chiropterologica* 11:127–137.

- Griffin, D. R. 1958. *Listening in the dark: the acoustic orientation of bats and men*. Yale University Press.
- Hamilton, R. 2012. *Spatial and temporal activity of migratory bats at landscape features*. MSc Thesis. Western University.
- Hayes, J. 1997. Temporal variation in activity of bats and the design of echolocation-monitoring studies. *Journal of Mammalogy* 78:514–524.
- Hayes, J. 2000. Assumptions and practical considerations in the design and interpretation of echolocation-monitoring studies. *Acta Chiropterologica* 2:225–236.
- Hayes, J., and J. Gruber. 2000. Vertical stratification of bat activity in an old-growth forest in western Washington. *Northwest Science* 74:102–108.
- Hecker, K., and R. Brigham. 1999. Does moonlight change vertical stratification of activity by forest-dwelling insectivorous bats? *Journal of Mammalogy* 80:1196–1201.
- Herzog, C., and E. Britzke. 2009. Progress update for acoustic monitoring of summer bats.
[http://whitenosesyndrome.org/sites/default/files/resource/herzognysdecbritzkeusacep
rogessupdateacousticmonitoringsummerbats.pdf](http://whitenosesyndrome.org/sites/default/files/resource/herzognysdecbritzkeusacep
rogessupdateacousticmonitoringsummerbats.pdf). Date accessed: 24 March 2013.
- Kalcounis, M., and K. Hobson. 1999. Bat activity in the boreal forest: importance of stand type and vertical strata. *Journal of Mammalogy* 80:673–682.

- Kalko, E., and C. Handley. 2001. Neotropical bats in the canopy: diversity, community structure, and implications for conservation. *Plant Ecology* 153:319–333.
- Korner-Nievergelt, F., P. Korner-Nievergelt, O. Behr, I. Niermann, R. Brinkmann, and B. Hellriegel. 2011. A new method to determine bird and bat fatality at wind energy turbines from carcass searches. *Wildlife Biology* 17:350–363.
- Larson, D. J., and J. P. Hayes. 2000. Variability in sensitivity of Anabat II bat detectors and a method of calibration. *Acta Chiropterologica* 2:209–213.
- Milne, D., A. Fisher, I. Rainey, and C. Pavey. 2005. Temporal patterns of bats in the Top End of the Northern Territory, Australia. *Journal of Mammalogy* 86:909–920.
- Reynolds, D. S. 2006. Monitoring the potential impact of a wind development site on bats in the northeast. *Journal of Wildlife Management* 70:1219–1227.
- Russ, J. M., M. Briffa, and W. I. Montgomery. 2003. Seasonal patterns in activity and habitat use by bats (*Pipistrellus* spp. and *Nyctalus leisleri*) in Northern Ireland, determined using a driven transect. *Journal of Zoology* 259:289–299.
- Skalak, S. L., R. E. Sherwin, and R. M. Brigham. 2012. Sampling period, size and duration influence measures of bat species richness from acoustic surveys. *Methods in Ecology and Evolution* 3:490–502.
- Stahlschmidt, P., and C. A. Brühl. 2012. Bats as bioindicators – the need of a standardized method for acoustic bat activity surveys. *Methods in Ecology and Evolution* 3:503–508.

Staton, T., and S. Poulton. 2012. Seasonal variation in bat activity in relation to detector height: a case study. *Acta Chiropterologica* 14:401–408.

Walsh, A. L., C. Catto, T. Hutson, P. Racey, P. Richardson, and S. Lanston. 2001. The UK's National Bat Monitoring Programme. Final Report 2001. London, England.
URL: www.bats.org.uk/publications_download.php/335/NBMP01.pdf.

Appendices

Appendix A: Discriminant function analysis for species identification

Passive acoustic monitoring makes it easy to collect large amounts of data; the challenge is analyzing all of it. Identification of echolocation calls to species is a daunting task when done manually, but automated methods make processing many recordings quick and efficient. Automated methods are more consistent and predictable, and can be more accurate than manual analysis (Jennings *et al.* 2008). A common technique for automated identification of bat calls is discriminant function analysis (DFA), it is not only a classic statistical technique, available in all statistical software packages, but is also very effective for classifying echolocation calls to species. DFA has outperformed other classic statistical methods (cluster analysis) and nonconventional methods (classification and regression trees, and neural networks) with higher classification accuracy of identifying calls to species (Preatoni *et al.* 2005).

I developed a DFA for the seven species of bats in Ontario, Canada to analyze my data (Chapters 3, 4, 5). The reference dataset of search-phase calls from recordings of free-flying bats recorded outside of known roosts or at foraging sites where species and individual bats were identified unambiguously (assembled by Lauren Hooton, MSc). No reference recordings came from hand-released individuals. The reference dataset included calls for seven species of bats in Ontario (Table A1). An eighth species (*Myotis leibii*) is found in Ontario, but was not included in the DFA due to the relative rarity of the species,

lack of verified recording for the reference dataset, and difficulty differentiating calls from the much more common *M. lucifugus*. The training dataset for the DFA included one, randomly selected call per individual in order to avoid pseudoreplication (Mundry and Sommer 2007). I had unequal sample sizes among species (Table A1), but this is not a problem for DFA. Because of unequal covariance I used quadratic DFA (Vaughan *et al.* 1997b, Parsons and Jones 2000).

The DFA included 11 predictor variables (call parameters), minimum frequency (Fmin), maximum frequency (Fmax), duration, frequency of most energy (FME), 10th percentile of energy (F10), 60th percentile of energy (F60), 90th percentile of energy (F90), median frequency slope (dFmedian), median energy slope (dEmedian), median frequency smoothness (sFmedian), and median energy smoothness (sEmedian). All call parameters were automatically extracted by the automated detection feature in callViewer (v. 18; Skowronski 2008). CallViewer is a custom echolocation sound analysis program written with MATLAB software (The MathWorks, Natick, Massachusetts). The automated detection parameters were the default settings, except minimum link length was set at 10, minimum energy was set at 14 dB, echo filter threshold was set at 10 dB, and lower frequency cutoff was set to 14 kHz. I chose call parameters to maximize classification accuracy with backwards, stepwise selection, starting with all 21 variables extracted by callViewer and removing them one at a time. The DFA should have fewer predictor variables than the smallest sample size; following the 1/3 rule I reduced down to 11 variables. Classification accuracy with leave-one-out cross-validation was high (Table A1), even between *Eptesicus fuscus* and *Lasionycteris noctivagans*, which cannot be discriminated with manual analysis (Betts 1998).

Table A1. Species-specific details for discriminant function analysis (DFA) to identify unclassified bat echolocation calls to species. The sample size per species was the number of echolocation calls per species included in the reference database (training data) for DFA. Each call was from a different individual. Classification accuracy of the DFA with leave-one-out cross-validation is the percent correct identifications of the training data.

Species	Sample size	Classification accuracy (%)
<i>Eptesicus fuscus</i>	50	78
<i>Lasionycteris noctivagans</i>	50	94
<i>Lasiurus borealis</i>	58	88
<i>L. cinereus</i>	52	90
<i>Myotis lucifugus</i>	50	90
<i>M. septentrionalis</i>	33	94
<i>Perimyotis subflavus</i>	42	93

Before identification of unidentified recordings with DFA, I filtered all acoustic data to eliminate noise and weak or fragmented calls, only including detections with duration 0.99 – 30 ms and Fmin 15 - 60 kHz. The filter parameters were selected based on conservative estimates of the echolocation call structure of the species of bats present in Ontario. The DFA compared our unidentified data to the training dataset and identified each call to species. To further improve classification accuracy, and because DFA does not assign calls to an “unknown” category, I applied a post-hoc, species-specific filter to remove any data that were above or below typical durations and minimum frequencies for each species (Table A2).

DFA and filters were performed in R (v. 2.13.1, R Development Core Team 2011).

Table A2. Post-discriminant function analysis filter to increase classification accuracy of automated identification of bat echolocation calls. The filter removed identified calls that were outside the species-specific ranges of call duration and minimum frequency.


Species	Duration (ms)	Minimum frequency (kHz)
<i>Eptesicus fuscus</i>	3 – 12	20 – 30
<i>Lasionycteris noctivagans</i>	4 - 12	21 - 30
<i>Lasiurus borealis</i>	5 - 17	29 – 43
<i>L. cinereus</i>	8 - 30	15 – 29
<i>Myotis lucifugus</i>	3 - 8	30 – 43
<i>M. septentrionalis</i>	1 – 3.3	25 - 60
<i>Perimyotis subflavus</i>	5 - 15	36 - 46

References


- Betts, B. 1998. Effects of interindividual variation in echolocation calls on identification of big brown and silver-haired bats. *Journal of Wildlife Management* 62:1003–1010.
- Jennings, N., S. Parsons, and M. J. O. Pocock. 2008. Human vs. machine: identification of bat species from their echolocation calls by humans and by artificial neural networks. *Canadian Journal of Zoology* 86:371–377.
- Mundry, R., and C. Sommer. 2007. Discriminant function analysis with nonindependent data: consequences and an alternative. *Animal Behaviour* 74:965–976.
- Parsons, S., and G. Jones. 2000. Acoustic identification of twelve species of echolocating bat by discriminant function analysis and artificial neural networks. *Journal of Experimental Biology* 203:2641–2656.
- Preatoni, D. G., M. Nodari, R. Chirichella, G. Tosi, L. A. Wauters, and A. Martinoli. 2005. Identifying bats from time-expanded recordings of search calls: comparing classification methods. *Journal of Wildlife Management* 69:1601–1614.
- R Development Core Team. 2011. R: a language and environment for statistical computing. R Foundation for Statistical Computing. v. 2.13.1. R Foundation for Statistical Computing, Vienna, Austria.
- Skowronski, M. D. 2008. callViewer. v. 18. London, Ontario, Canada.
- Vaughan, N., G. Jones, and S. Harris. 1997. Identification of British bat species by multivariate analysis of echolocation call parameters. *Bioacoustics* 7:189–207.

Appendix B: Chapter 5 supplementary material


1)




2a)




2b)




3a)



3b)



4a)






Figure B1. Peak periods of bat activity along linear, stationary transects within four sites in Ontario, Canada in 2008: 1) spring at Long Point, 2a) spring at QUBS, 2b) late summer at QUBS, 3a) spring at Renfrew lake, 3b) late summer at Renfrew lake, 4a) spring at Renfrew mine, and 4b) late summer (swarming) at Renfrew mine. Detectors were paired at two heights, low (1.5 m; A1, B1, and C1) and high (4 m; A2 and B2).

Table B1. Comparison of mean hourly bat activity among detectors for each species within sites in Ontario, Canada in 2008: A) Long Point, B) QUBS, C) Renfrew lake, D) Renfrew mine, and E) Renfrew mine in 2010 (df = 4). Significant Kruskal-Wallis tests indicate that activity was significantly different among detectors within a site.

A) Long Point		
Species	H statistic	p-value
<i>E. fuscus</i>	130.94	< 0.001
<i>L. noctivagans</i>	104.21	< 0.001
<i>L. borealis</i>	252.66	< 0.001
<i>L. cinereus</i>	182.01	< 0.001
<i>M. lucifugus</i>	133.94	< 0.001
<i>M. septentrionalis</i>	146.94	< 0.001
<i>P. subflavus</i>	202.91	< 0.001

B) QUBS		
Species	H statistic	p-value
<i>E. fuscus</i>	58.45	< 0.001
<i>L. noctivagans</i>	8.11	0.09
<i>L. borealis</i>	64.53	< 0.001
<i>L. cinereus</i>	36.80	< 0.001
<i>M. lucifugus</i>	34.41	< 0.001
<i>M. septentrionalis</i>	3.34	0.50
<i>P. subflavus</i>	42.23	< 0.001

C) Renfrew lake		
Species	H statistic	p-value
<i>E. fuscus</i>	48.40	< 0.001
<i>L. noctivagans</i>	7.51	0.11
<i>L. borealis</i>	172.92	< 0.001
<i>L. cinereus</i>	19.55	0.001
<i>M. lucifugus</i>	118.41	< 0.001
<i>M. septentrionalis</i>	3.98	0.41
<i>P. subflavus</i>	54.54	< 0.001

D) Renfrew mine		
Species	H statistic	p-value
<i>E. fuscus</i>	29.77	< 0.001
<i>L. noctivagans</i>	37.11	< 0.001
<i>L. borealis</i>	80.88	< 0.001
<i>L. cinereus</i>	33.76	< 0.001
<i>M. lucifugus</i>	18.34	0.001
<i>M. septentrionalis</i>	34.56	< 0.001
<i>P. subflavus</i>	11.64	0.02

E) Renfrew mine 2010

Species	H statistic	p-value
<i>E. fuscus</i>	14.93	0.005
<i>L. noctivagans</i>	24.16	< 0.001
<i>L. borealis</i>	239.68	< 0.001
<i>L. cinereus</i>	50.69	< 0.001
<i>M. lucifugus</i>	47.14	< 0.001
<i>M. septentrionalis</i>	157.46	< 0.001
<i>P. subflavus</i>	40.95	< 0.001

Table B2. Final models of species-specific hourly bat activity (activity) among detectors and four sites in Ontario, Canada in 2008. Generalized linear mixed models (GLMM) included random and fixed effects. Random effects were detector location nested in site (detector), night nested in site (night), and sites (site). Fixed effects were percent canopy cover (canopy) and detector height (height), both at the detector-level. Models were selected by likelihood ratio tests and AIC. Parameter estimates, standard error, and p-values are included for significant fixed effects in the models.

Species	GLMM model	Fixed effect explanatory variable	Parameter Estimate	Standard error	p-value
Species combined	activity ~ (1 detector) + (1 night)	Intercept	8.183	0.379	< 0.001
<i>E. fuscus</i>	activity ~ canopy + (1 detector) + (1 night)	Intercept	7.291	0.831	< 0.001
		canopy	-7.534	1.728	< 0.001
<i>L. noctivagans</i>	activity ~ canopy + height + (1 detector) + (1 night) + (1 site)	Intercept	2.152	1.128	0.056
		canopy	-6.950	2.047	< 0.001
		height	1.385	0.591	0.019
<i>L. borealis</i>	activity ~ canopy + (1 detector) + (1 night) + (1 site)	Intercept	7.880	1.312	< 0.001
		canopy	-11.628	2.276	< 0.001
<i>L. cinereus</i>	activity ~ canopy + (1 detector) + (1 night)	Intercept	7.915	0.620	< 0.001
		canopy	-8.441	1.299	< 0.001
<i>M. lucifugus</i>	activity ~ canopy + (1 detector) + (1 night) + (1 site)	Intercept	7.662	1.13	< 0.001
		canopy	-4.921	1.764	0.005
<i>M. septentrionalis</i>	activity ~ canopy + (1 detector) + (1 night) + (1 site)	Intercept	0.819	2.449	0.014
		canopy	6.417	1.490	< 0.001
<i>P. subflavus</i>	activity ~ canopy + (1 detector) + (1 night)	Intercept	4.503	1.051	< 0.001
		canopy	-7.703	2.248	< 0.001

Table B3. Number of nights when bat activity was over the 50th and 90th percentile thresholds at each detector at four sites in Ontario, Canada in 2010: A) Long Point, B) QUBS, C) Renfrew lake, D) Renfrew mine. Detectors were paired at two heights, low (1.5 m; A1, B1, C1) and high (4 m; A2, B2).

A) Long Point	50 th percentile Detectors					90 th percentile Detectors				
	Species	A1	A2	B1	B2	C1	A1	A2	B1	B2
Species combined	12	12	11	9	12	7	7			1
<i>E. fuscus</i>	12	12	10	11	12	11	10	2	3	6
<i>L. noctivagans</i>	8	9	5	4	5	2	2			
<i>L. borealis</i>	12	12	8	10	12	11	11	1	1	10
<i>L. cinereus</i>	10	11			10	3	3			
<i>M. lucifugus</i>	12	12	1	1	7	4	6			1
<i>M. septentrionalis</i>	12	12	6	7	7	2	10			3
<i>P. subflavus</i>	1		10	12						

B) QUBS	50 th percentile Detectors					90 th percentile Detectors				
	Species	A1	A2	B1	B2	C1	A1	A2	B1	B2
Species combined	3	7	4	4	8		2			
<i>E. fuscus</i>	1	1	2	3	9		1			4
<i>L. noctivagans</i>	5	7	4	5	3		2			
<i>L. borealis</i>	5	9		1	10					5
<i>L. cinereus</i>	6	8			5	3	5			1
<i>M. lucifugus</i>	2	7	1	3	6					1
<i>M. septentrionalis</i>		2		1						
<i>P. subflavus</i>	3	6	6	7	1					

C) Renfrew lake	50 th percentile Detectors					90 th percentile Detectors				
	Species	A1	A2	B1	B2	C1	A1	A2	B1	B2
Species combined	11	11	8	8		2	3			
<i>E. fuscus</i>	11	11	6	7		7	5	1	1	
<i>L. noctivagans</i>	11	11	5	7		3	3			
<i>L. borealis</i>	8	9	7	8	2		1			
<i>L. cinereus</i>	11	11		3		4	3			
<i>M. lucifugus</i>	7	8				3	2			
<i>M. septentrionalis</i>	4	6	3	4	1					
<i>P. subflavus</i>	7	7	9	9	5					

D) Renfrew mine	50th percentile					90th percentile				
	Detectors					Detectors				
Species	A1	A2	B1	B2	C1	A1	A2	B1	B2	C1
Species combined	4	6	10	11	10	1	2	6	5	6
<i>E. fuscus</i>	3	4	3	1	7		3			
<i>L. noctivagans</i>	4	6	9	10	10		1	5	4	6
<i>L. borealis</i>	1	3	1	1	7		1			1
<i>L. cinereus</i>		2		1	11		1			7
<i>M. lucifugus</i>		2	1		5		2			
<i>M. septentrionalis</i>		1		1						
<i>P. subflavus</i>	5	6	11	12	9	4	3	6	6	

Appendix C: Permission to reproduce published material

JOHN WILEY AND SONS LICENSE TERMS AND CONDITIONS

Mar 08, 2013

This is a License Agreement between Amanda Adams ("You") and John Wiley and Sons ("John Wiley and Sons") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by John Wiley and Sons, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

License Number	3104470967139
License date	Mar 08, 2013
Licensed content publisher	John Wiley and Sons
Licensed content publication	Methods in Ecology & Evolution
Licensed content title	Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats
Licensed copyright line	© 2012 The Authors. Methods in Ecology and Evolution © 2012 British Ecological Society
Licensed content author	Amanda M. Adams, Meredith K. Jantzen, Rachel M. Hamilton, Melville Brockett Fenton
Licensed content date	Sep 18, 2012
Start page	992
End page	998
Type of use	Dissertation/Thesis
Requestor type	Author of this Wiley article
Format	Print and electronic
Portion	Full article
Will you be translating?	No
Order reference number	
Total	0.00 USD
Terms and Conditions	

TERMS AND CONDITIONS

This copyrighted material is owned by or exclusively licensed to John Wiley & Sons, Inc. or one of its group companies (each a "Wiley Company") or a society for whom a Wiley Company has exclusive publishing rights in relation to a particular journal (collectively WILEY). By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the billing and payment terms and conditions established by the Copyright Clearance Center Inc., ("CCC's Billing and Payment terms and conditions"), at the time that you opened your Rightslink account (these are

Appendix D: Animal use protocol approvals



April 2, 2008

This is the Original Approval for this protocol
 A Full Protocol submission will be required in 2012

Dear Dr. Fenton:

Your Animal Use Protocol form entitled:
 Behavioural Ecology of Bats
 Funding Agency NSERC - Grant #R3516A03

has been approved by the University Council on Animal Care. This approval is valid from **April 2, 2008 to April 30, 2009**. The protocol number for this project is **#2008-003-04 and replaces #2004-027-03**.

1. This number must be indicated when ordering animals for this project.
2. Animals for other projects may not be ordered under this number.
3. If no number appears please contact this office when grant approval is received.
 If the application for funding is not successful and you wish to proceed with the project, request that an internal scientific peer review be performed by the Animal Use Subcommittee office.
4. Purchases of animals other than through this system must be cleared through the ACVS office. Health certificates will be required.

ANIMALS APPROVED FOR 1 YR.

Species	Strain	Other Detail	Pain Level	Animal # Total for 1 Year
Other, add to detail	Bats	various species MF	C	~ 1500

REQUIREMENTS/COMMENTS

Please ensure that individual(s) performing procedures on live animals, as described in this protocol, are familiar with the contents of this document.

1. Please ensure that the approximate number of bats used under this protocol is submitted to the AUS office by December each year for the annual CCAC report.

c.c. Approved Protocol - B. Fenton, L. McGuire, J. Weber, D. Cheshuk
 Approval Letter - L. McGuire, J. Weber, D. Cheshuk





05.01.09

*This is the 1st Renewal of this protocol
 *A Full Protocol submission will be required in 2012

Dear Dr. **Fenton**

Your Animal Use Protocol form entitled:

Behavioural Ecology of Bats

has had its yearly renewal approved by the Animal Use Subcommittee.

This approval is valid from **05.01.09 to 04.30.10**

The protocol number for this project remains as **2008-003**

1. This number must be indicated when ordering animals for this project.
2. Animals for other projects may not be ordered under this number.
3. If no number appears please contact this office when grant approval is received.
 If the application for funding is not successful and you wish to proceed with the project, request that an internal scientific peer review be performed by the Animal Use Subcommittee office.
4. Purchases of animals other than through this system must be cleared through the ACVS office. Health certificates will be required.

REQUIREMENTS/COMMENTS

Please ensure that individual(s) performing procedures on live animals, as described in this protocol, are familiar with the contents of this document.

The holder of this *Animal Use Protocol* is responsible to ensure that all associated safety components (biosafety, radiation safety, general laboratory safety) comply with institutional safety standards and have received all necessary approvals. Please consult directly with your institutional safety officers.



c.c. L McGuire, J Wasylenko

The University of Western Ontario
 Animal Use Subcommittee / University Council on Animal Care
 Health Sciences Centre, • London, Ontario • CANADA – N6A 5C1
 PH: 519-661-2111 ext. 86770 • FL 519-661-2028 • www.uwo.ca / animal



05.01.2010

*This is the 2nd Renewal of this protocol

*A Full Protocol submission will be required in 04.30.2012

Dear Dr. Fenton

Your Animal Use Protocol form entitled:

Behavioural ecology of bats

has had its yearly renewal approved by the Animal Use Subcommittee.

This approval is valid from **05.01.2010 to 05.01.2011**The protocol number for this project remains as **2008-003**

1. This number must be indicated when ordering animals for this project.
2. Animals for other projects may not be ordered under this number.
3. If no number appears please contact this office when grant approval is received.
If the application for funding is not successful and you wish to proceed with the project, request that an internal scientific peer review be performed by the Animal Use Subcommittee office.
4. Purchases of animals other than through this system must be cleared through the ACVS office. Health certificates will be required.

REQUIREMENTS/COMMENTS

Please ensure that individual(s) performing procedures on live animals, as described in this protocol, are familiar with the contents of this document.

The holder of this *Animal Use Protocol* is responsible to ensure that all associated safety components (biosafety, radiation safety, general laboratory safety) comply with institutional safety standards and have received all necessary approvals. Please consult directly with your institutional safety officers.

c.c. J. Wasylenko

The University of Western Ontario
Animal Use Subcommittee / University Council on Animal Care
Health Sciences Centre, • London, Ontario • CANADA – N6A 5C1

Appendix E: Research permits

MAY-12-2008 11:31

MNR PLANNING PTBO

705 755 3292 P.002

**Ministry of
Natural Resources**
Planning Unit Manager
Southern Region
P.O. Box 7000, 300 Water Street
Peterborough, Ontario K9J 8M5

**Ministère des
Richesses naturelles**
Telephone: (705) 755-5164
Facsimile: (705) 755-3292



May 9, 2008

Mr. Brock Fenton
University of Western Ontario
Department of Biology

Dear Mr. Fenton:

**Subject: Wildlife Scientific Collector's Authorization
Authorization Number: 1045694**

Enclosed please find Wildlife Scientific Collector's Authorization form number 1045694. This authorization is valid only for the person(s), species, numbers, areas and calendar year indicated.

Please review, sign and return the WSC Authorization and attachments to the following address:

*Manager
Southern Region Planning Unit
Ministry of Natural Resources*

Once returned, the authorization will be given final approval and the original mailed back to you. Please note that you are required to carry a copy of the authorization at all times while collecting specimens.

A written report, as outlined in the Authorization Conditions, covering the operation of the preceding year must be submitted within 30 days of the termination date to the Manager of the Southern Region Planning Unit.

The report shall contain a statement outlining the objectives of the operations, the methods used, the number and species of wildlife caught and their fate as well as a map indicating where the collections took place. The submission of a satisfactory report is a prerequisite to any subsequent Wildlife Scientific Collector's Authorization renewals.

If you have any questions please do not hesitate to contact Michael Gatt, Senior Regional Wildlife Biologist, at [REDACTED]

Yours truly,

[REDACTED]
Monique Rolf von den Baumen-Clark
Manager
Southern Region Planning Unit

Attachments

APR-30-2009 12:01

MNR PLANNING PTBU

705 755 3292 P.002

Ministry of Natural
Resources

Southern Region

P.O. Box 7000
300 Water Street
Peterborough, ON
K9J 8M5
Tel: 705-755-3243
Fax: 705-755-3292

Ministère des Richesses
naturelles

Région du Sud

P.O. Box 7000
300, rue Water
Peterborough (Ontario)
K9J 8M5
Tél.: 705-755-3243
Télé.: 705-755-3292



April 17, 2009

Brock Fenton
University of Western Ontario
Department of Biology

Dear Mr. Fenton

Subject: Wildlife Scientific Collector's Authorization
Authorization Number: 1050823

Enclosed please find Wildlife Scientific Collector's (WSC) Authorization form number 1050823. This authorization is valid only for the person(s), species, numbers, areas and calendar year indicated.

Please review, sign and return the WSC Authorization and attachments to the following address:

*Manager
Southern Region Planning Unit
Ministry of Natural Resources*

Once returned, the authorization will be given final approval and the original mailed back to you. Please note that you are required to carry a copy of the authorization at all times while collecting specimens.

A written report, as outlined in the Authorization Conditions, covering the operation of the preceding year must be submitted within 30 days of the termination date to the Manager of the Southern Region Planning Unit.

The report shall contain a statement outlining the objectives of the operations, the methods used, the number and species of wildlife caught and their fate as well as a map indicating where the collections took place. The submission of a satisfactory report is a prerequisite to any subsequent Wildlife Scientific Collector's Authorization renewals.

If you have any questions please do not hesitate to contact Michael Gatt, Senior Regional Wildlife Biologist,

Yours truly,

Monique Rolf von den Baumen-Clark
Manager
Southern Region Planning Unit

Attachments

Ministry of Natural
Resources

Southern Region

P.O. Box 7000
300 Water Street
Peterborough, ON
K9J 8M5
Tel: 705-755-3243
Fax: 705-755-3292

Ministère des Richesses
naturelles

Région du Sud

P.O. Box 7000
300, rue Water
Peterborough (Ontario)
K9J 8M5
Tél.: 705-755-3243
Télééc.: 705-755-3292



April 30, 2010

Brock Fenton
University of Western Ontario
Dept. of Biology & Geological Sciences Bldg.
London, ON N6A 5B7

Dear Mr. Fenton

Subject: Wildlife Scientific Collector's Authorization
Authorization Number: 1055825

Enclosed please find Wildlife Scientific Collector's (WSC) Authorization form number 1055825. This authorization is valid only for the person(s), species, numbers, areas and calendar year indicated.

Please review, sign and return the WSC Authorization and attachments to the following address:

*Manager
Southern Region Planning Unit
Ministry of Natural Resources*



Once returned, the authorization will be given final approval and the original mailed back to you. Please note that you are required to carry a copy of the authorization at all times while collecting specimens.

A written report, as outlined in the Authorization Conditions, covering the operation of the preceding year must be submitted within 30 days of the termination date to the Manager of the Southern Region Planning Unit.

The report shall contain a statement outlining the objectives of the operations, the methods used, the number and species of wildlife caught and their fate as well as a map indicating where the collections took place. The submission of a satisfactory report is a prerequisite to any subsequent Wildlife Scientific Collector's Authorization renewals.

If you have any questions please do not hesitate to contact Michael Gatt, Senior Regional Wildlife Biologist, ([redacted])

Yours truly,



Bob McColm
Manager
Regional Operations

Attachments

Curriculum Vitae

Name: Amanda Adams

Post-secondary Education and Degrees:

University of Western Ontario
London, Ontario, Canada
2007-2013 Ph.D.

University of California, San Diego
La Jolla, California, USA
2004-2006 M.Sc.

University of California, San Diego
La Jolla, California, USA
2001-2005 B.Sc.

Selected Honors and Awards:

Centre for Environment & Sustainability Excellence Award
2010-2011, 2011-2012

Bat Conservation International Student Scholarship
2010

Graduate Teaching Assistant Union Community Service Award
2009-2010

Related Work Experience:

Teaching Assistant
University of Western Ontario
2007-2013

Course Instructor
Continuing Education, University of Western Ontario
2012

Course Development
Distance Studies, University of Western Ontario
2010

Research Assistant
University of Michigan
2007

Intern/Lab Technician
Nautilus Environmental, LLC
2004-2006

Publications:

Published

Adams, A. M., M. K. Jantzen, R. M. Hamilton, and M. B. Fenton. 2012. A comparison of bat detectors: performance with synthetic playback and free-flying bats. *Methods in Ecology and Evolution* 3: 992–998.

Clare, E. L., **A. M. Adams***, A. Maya-Simoes, J. Eger, P. D. N. Hebert, and M. B. Fenton. 2013. Diversification and reproductive isolation: cryptic species in the only New World high-duty cycle bat, *Pteronotus parnellii*. *BMC Evolutionary Biology*, 13: doi:10.1186/1471-2148-13-26.**

Mills, S.C., **A.M. Adams**, R.D. Phoenix. 2013. Bat Species Diversity in the Boreal Forest of Northeastern Ontario, Canada. *Northeastern Naturalist*, 20(2): 309–324.

* Co-first author

** rated as highly accessed by Bio-Medical-Central

Accepted

Clare, E.L., H.R. Goerlitz, V.A. Drapeau, M.W. Holderied, **A.M. Adams**, J. Nagel, E.R. Dumont, P.D.N. Hebert, M.B. Fenton. Trophic niche flexibility in *Glossophaga soricina*: how a nectar seeker sneaks an insect snack. *Functional Ecology*: accepted.

Submitted

Adams, A. M., L. P. McGuire, and M. B. Fenton. How high is high? Using percentile thresholds to identify peak bat activity. *Acta Chiropterologica*: in review.

Adams, A. M. and M. B. Fenton. Identifying peaks in bat activity: a new application of the space-time scan statistic. *Journal of Applied Ecology*: in review.