

A comparative study of molecular mutations in 381 patients with myelodysplastic syndrome and in 4130 patients with acute myeloid leukemia

Ulrike Bacher, Torsten Haferlach, Wolfgang Kern, Claudia Haferlach, Susanne Schnittger

From the Bone Marrow Transplant Unit, University Hospital of Hamburg-Eppendorf, Hamburg, Germany (UB); MLL, Munich Leukemia Laboratory, Munich, Germany (TH, WK, CH, SS).

Acknowledgments: we would like to thank participating centers of the AMLCG study group and other centers for sending bone marrow or blood samples to our laboratory for diagnosis, and for submitting clinical data. This study was performed in part in the Laboratory for Leukemia Diagnostics, III. Medical Department, Ludwig-Maximilians-University of Munich (Head: Prof. Dr. med. W. Hiddemann).

Manuscript received October 3, 2006.

Manuscript accepted March 14, 2007.

Correspondence:
Susanne Schnittger, MLL, Munich Leukemia Laboratory, Max-Lebsche-Platz 31 81377 Munich, Germany.
E-mail: susanne.schnittger@mllo-online.com

ABSTRACT

Background and Objectives

The precise relationship between myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML) is unclear and the role of molecular mutations in leukemic transformation in MDS is controversial. The aim of this study was to clarify the relationship between AML and MDS by comparing the frequency of molecular mutations in the two conditions.

Design and Methods

We compared the frequency of *FLT3*-length mutations (*FLT3*-LM), *FLT3*-TKD, *MLL*-partial tandem duplications (*MLL*-PTD), *NRAS*, and *KITD816* in 381 patients with MDS refractory anemia with excess blasts [RAEB] $n=49$; with ringed sideroblasts [RARS] $n=310$; chronic monomyelocytic leukemia [CMML] $n=22$ and in 4130 patients with AML (*de novo*: $n=3139$; secondary AML [s-AML] following MDS: $n=397$; therapy-related [t-AML]: $n=233$; relapsed: $n=361$).

Results

All mutations were more frequent in s-AML than in MDS and all but the *FLT3*-TKD were more frequent in RAEB than in RA/RARS. The higher incidences in s-AML were significant for *FLT3*-TKD ($p=0.032$), *MLL*-PTD ($p=0.034$), and *FLT3*-LM (RA/RARS: 0/45; RAEB: 8/293; 2.7%; s-AML: 45/389; 11.6%; $p<0.0001$). The incidence of *NRAS*-mutations increased from 17/272 (6.3%) in MDS to 41/343 in s-AML (12.0%) and that of *KITD816*-mutations from 2/290 (0.7%) to 5/341 (1.5%) ($p=n.s.$). *FLT3*-LM-acquisition occurred in 3/22 cases (13.6%) during MDS transformation; *NRAS*-acquisition occurred in 1/24 (4.2%). *FLT3*-LM and *MLL*-PTD were more frequent in AML relapse than in *de novo* AML or s-AML ($p<0.0001$).

Interpretation and Conclusions

The increase of molecular mutations from low- to high-risk MDS, to s-AML, and to relapsed AML emphasizes the value of these mutations as markers of progressing disease. Finally, we found a low rate of 5q- in the molecularly mutated cases in MDS which might explain the stability of this subtype.

Key words: myelodysplastic syndromes (MDS), acute myeloid leukemias (AML), molecular mutations, progression, leukemogenesis.

Haematologica 2007; 92:744-752

©2007 Ferrata Storti Foundation

From clinical, cytomorphologic, cytogenetic, and molecular aspects the myelodysplastic syndromes (MDS) are heterogeneous diseases. If left untreated, the survival of patients with MDS varies between a few months and 20 years. The World Health Organization (WHO) classification (based on the French-American British [FAB] classification of 1982) subdivides MDS into eight subtypes according to the percentages of bone marrow blasts, ringed sideroblasts, and the number of dysplastic cell lineages.¹⁻⁴ Karyotype represents a strong prognostic parameter:⁵ 5q-, -Y, and 20q- as sole abnormalities are associated with a favorable prognosis. Complex aberrant karyotype (defined by ≥ 3 chromosomal abnormalities) and chromosome 7 abnormalities predict a poor prognosis, whereas all other chromosome abnormalities are associated with an intermediate prognosis. The International Prognostic Scoring System (IPSS) categorizes MDS patients into four risk groups based on blast percentage, karyotype, and the number of cell lines showing cytopenia.^{3,5} However, cytogenetic abnormalities are observed in only 30-50% of *de novo* MDS cases.^{6,7} Thus, molecular mutations may serve as potential markers to extend the spectrum of diagnostic and prognostic parameters in MDS. As MDS and AML are conceived as end points of a stepwise process of leukemogenesis in some patients, research in MDS should concentrate as well on the analysis of molecular mutations which occur frequently in AML.^{8,9}

Mutations of the *FLT3*-gene, a member of the class-III-receptor tyrosine kinase family, play a central role in AML.¹⁰⁻¹⁴ The *FLT3* length mutations (LM) (internal tandem duplications or insertions) (20%-27%) are, together with *NPM1*,¹⁵ the most frequent mutations in AML. While *FLT3*-LM are prognostically unfavorable,¹⁶⁻¹⁸ *NPM1* mutations are associated with a favorable outcome.^{15,19} Furthermore, small mutations in the tyrosine kinase domain (TKD) of *FLT3* (*FLT3*-TKD) have been found in 5%-8% of all AML cases.^{11,12,20-22} Their prognostic impact in AML is not yet clarified.^{16,22} In MDS the *FLT3*-mutations are less frequent. LM were found in 2%-3% (Horiike *et al.* n=58; Shih *et al.* n=198),^{23,24} and TKD mutations in 3%, as shown by Yamamoto *et al.* (to our knowledge the only analysis of this marker in MDS; n=29).¹²

Mutations of the *KIT*-proto-oncogene are a further example of class-III-receptor tyrosine kinase mutations in AML. *KITD816*-mutations occur with a frequency of 2% in unselected AML, are localized in the intracellular protein tyrosine kinase domain,^{25,26} and have an unfavorable prognostic impact in the subgroup of AML with t(8;21)/*AML1-ETO*.²⁷⁻²⁹ In MDS, the single study focused on this mutation reported a frequency of 3/39 (6%) when combining all cytomorphologic subtypes in MDS.³⁰ Mutations of the *NRAS*-proto-oncogene are identified in 10-15% of cases of

AML. These mutations increase the activity of the RAS-pathway and lead to cell proliferation and reduction of apoptosis.³¹⁻³⁵ Their influence on prognosis in AML seems dependent on cytogenetics and additional molecular mutations.³⁶ They show a favorable trend in CBF-leukemias and normal karyotype AML lacking *FLT3*-LM and partial tandem duplications of the *MLL* gene (*MLL*-PTD). In MDS frequencies of *NRAS* mutations were reported to be between 7% and 48% in previous studies including series of up to 220 patients.³⁷⁻⁴² *MLL*-PTD occur in 10% of AML with normal karyotype and are associated with a poor prognosis.⁴³⁻⁴⁶

Here we performed a study on the incidence of *FLT3*-LM, *FLT3*-TKD, *MLL*-PTD, *NRAS*- and *KITD816*-mutations in different cytomorphologic subtypes of MDS (n=381) and compared these data to those for AML (n=4130) in order to gain a better understanding of the leukemic transformation of MDS. In addition, we analyzed the correlation of these mutations with cytogenetics in MDS.

Design and Methods

Bone marrow samples – in many cases accompanied by peripheral blood samples – from 381 consecutive patients with MDS at diagnosis and from 4130 patients with AML (*de novo* AML at diagnosis: n=3139; secondary AML [s-AML] at diagnosis: n=397; therapy-related [t-AML] at diagnosis: n=233; and relapsed AML: n=361) were included in the study. The patients' clinical data and biological characteristics are shown in Table 1. The cytomorphological classification was made according to the FAB-classification, as the cohort was analyzed in part before WHO-criteria were defined in these patients.^{2,47} The patients with refractory anemia (RA) and refractory anemia with ringed sideroblasts (RARS) were combined to form the RA/RARS cohort. The cohort with refractory anemia with excess blasts (RAEB) included patients with RAEB-1 ($\leq 10\%$ of blasts) and with RAEB-2 ($< 20\%$ of blasts). The third MDS subgroup was represented by the dysplastic subtype of chronic myelomonocytic leukemia (CMML). Only patients with MDS at diagnosis were included. AML patients were subdivided according to the history of disease: *de novo* AML, secondary AML (s-AML) following MDS, and therapy-related AML (t-AML) in association with previous chemotherapy or radiotherapy of a malignant disease. Screening for *FLT3*-LM, *FLT3*-TKD, *MLL*-PTD, *NRAS*- and *KITD816*-mutations was performed as described before. All methods for mutation analysis have been reported in detail. Briefly, screening for *FLT3*-LM was performed by gel electrophoresis¹⁷ and fragment analysis²⁰ in parallel; *MLL*-PTD were analyzed by quantitative real-time polymerase chain reaction (PCR),⁴⁸ and

analysis for *FLT3*-TKD, *NRAS*- and *KITD816*-mutations was performed using melting curve based Light Cycler analysis and subsequent sequencing of the positive samples.²⁹ In addition, chromosome banding analyses and fluorescence *in situ* hybridization were performed as previously described.⁴⁹

The cytogenetic subgroups were categorized as follows: normal karyotype, reciprocal translocations, complex aberrant karyotype (≥ 3 chromosomal anomalies), deletion of 5q (5q-), chromosome 7 abnormalities, numerical gain of 8 (+8), deletion of 20q (20q-), loss of Y (-Y), inv(3)/t(3;3)(q21;q26), and other aberrations.

Results

Distribution of molecular mutations in low grade MDS, RAEB, and s-AML

First, the distribution of the mutations in the total cohorts of MDS and AML was analyzed. The complete results of the molecular analyses within the different subgroups are shown in Table 2. In MDS *NRAS*-mutations were detected in 17/272 patients (6.3%) and, from among the analyzed mutations, was the one with the highest frequency in MDS, followed by the *MLL*-PTD (10/368; 2.7%) and by the *FLT3*-LM (8/367; 2.2%). In contrast, in the total AML cohort *FLT3*-LM was the most frequent mutation of all those analyzed

Table 1. Clinical data and cytomorphologic subtypes of 381 patients with MDS at diagnosis and of 4130 patients with AML at diagnosis or relapse.

MDS	
Sex	240 males, 141 females
Age	18.1-97.8 years (median 68.1)
RA/RARS	49
RAEB	310
CMML	22
MDS total	381
AML	
Sex	2260 males, 1870 females
Age	16.2-96.8 years (median 61.2)
<i>de novo</i> AML	3139
s-AML	397
t-AML	233
Relapsed AML	361
AML total	4130
Total MDS + AML	4511

(783/3718; 21.1%), followed by the *NRAS* mutation (290/2856; 10.2%), and *FLT3*-TKD (144/3052; 4.7%).

Considering the different cytomorphologic subtypes of MDS, in RA/RARS a *FLT3*-TKD-mutation was observed in 1 of 28 cases whereas *FLT3*-LM, *NRAS*, and *KITD816* were not observed at all. In RAEB *NRAS* mutations represented the most frequent molecular marker (15/223; 6.7%), followed by the *FLT3*-LM

Table 2. Mutated cases in the different cytomorphologic subtypes of MDS and in the cohorts with AML.

Mutation	RA/RARS	RAEB	CMML	Total MDS	<i>de novo</i> AML	s-AML	t-AML	Relapsed AML	Total AML
<i>FLT3</i>-LM									
N. mutated	0	8	0	8	629	45	26	83	783
Total number	45	293	29	367	2813	389	216	300	3718
%	0.0	2.7	0.0	2.2	22.4	11.6	12.0	27.7	21.1
	(<0.0001)	(<0.0001)	(0.034)		(<0.0001)	(<0.0001)	(0.004)	(<0.0001)	
<i>FLT3</i>-TKD									
N. mutated	1	0	0	1	130	6	3	5	144
Total number	28	209	19	256	2357	322	167	206	3052
%	3.6	0.0	0.0	0.4	5.5	1.9	1.8	2.4	4.7
	(1.000)	(<0.0001)	(1.000)		(0.030)	(0.032)	(0.064)	(0.096)	
<i>MLL</i>-PTD									
N. mutated	1	8	1	10	176	28	7	25	85
Total number	46	292	30	368	2735	378	213	303	1197
%	2.2	2.7	3.3	2.7	6.0	7.4	3.3	8.3	7.1
	(0.517)	(0.032)	(0.627)		(0.017)	(0.034)	(0.166)	(0.036)	
<i>NRAS</i>									
N. mutated	0	15	2	17	209	41	21	19	290
Total number	29	223	20	272	2128	343	174	211	2856
%	0.0	6.7	10.0	6.3	9.8	12.0	12.1	9.0	10.2
	(0.128)	(0.108)	(1.000)		(1.000)	(0.177)	(0.294)	(0.810)	
<i>KITD816</i>									
N. mutated	0	2	0	2	39	5	2	3	49
Total number	32	237	21	290	2136	341	172	213	2862
%	0.0	0.8	0.0	0.7	1.8	1.5	1.2	1.4	1.7
	(1.000)	(0.431)	(1.000)		(0.227)	(1.000)	(1.000)	(1.000)	

The *p*-values (in brackets) indicate whether the frequencies of the molecular mutations differ significantly from those in the other MDS or AML stages.

(8/293; 2.7%) and *MLL*-PTD (8/292; 2.7%).

Subsequently the distribution of the respective markers in the different hematologic subgroups was analyzed. The *FLT3*-TKD were heterogeneously distributed, most probably because of the very low frequency of this mutation overall. Statistically significant differences in distribution were found for *FLT3*-LM ($p < 0.0001$) and *MLL*-PTD ($p = 0.004$), whereas the distribution of the *NRAS*- and *KITD816*-mutations did not vary significantly. The incidence of the markers in early MDS categories (RA/RARS) was compared with that in the advanced stages (RAEB) and in s-AML. The incidence of *FLT3*-LM, *MLL*-PTD, *NRAS*-, and *KITD816*-mutations increased from RA/RARS to RAEB and to s-AML. The differences were statistically significant for *FLT3*-LM ($p < 0.0001$), *FLT3*-TKD mutations ($p < 0.0001$), and *MLL*-PTD ($p = 0.032$). The sharpest increase was observed for *FLT3*-LM which were found in no case with RA/RARS (0/45; 0.0%), in 8/293 (2.7%) cases of RAEB, and in 45/389 (11.6%) cases of s-AML ($p < 0.0001$) (Figure 1).

Molecular mutations in CMML

In our small series of CMML with dysplastic subtype *NRAS*-mutations were found in 2/20 (10%) rates, which was similar to the frequency in AML. The other mutations analyzed were rarely (*FLT3*-TKD) or never (*FLT3*-LM and *KITD816*) observed in CMML (Table 2).

Incidence of molecular mutations with respect to history of AML

FLT3-LM were significantly more frequent in *de novo* AML and in AML at relapse than in s-AML or t-AML ($p < 0.0001$). *FLT3*-TKD were significantly more frequent in *de novo* AML ($p = 0.030$) and in s-AML ($p = 0.032$) than in t-AML or in relapsed AML. The frequencies of *MLL*-PTD, *NRAS*-, and *KITD816*-mutations did not differ significantly between the different AML cohorts. These results confirmed those of two previous studies (Table 2).^{17,36}

Concomitant molecular mutations in MDS

The concomitant presence of different molecular mutations was observed in only 2/381 MDS patients (0.5%): one case with RAEB showed *FLT3*-LM and *MLL*-PTD, whereas another RAEB patient had *FLT3*-LM, *MLL*-PTD, and *NRAS*-mutations.

Acquisition of molecular mutations during progression of MDS

Finally, we analyzed whether leukemic transformation of MDS was accompanied by acquisition of the molecular markers. For this analysis, 25 paired MDS/AML cases were available. We found that the mutation status was stable in all cases screened for *FLT3*-TKD (n=24), *MLL*-PTD (n=22), and *KITD816* (n=24). However, acquisition of *FLT3*-LM was

observed in 3/22 cases (13.6%) during leukemic transformation and acquisition of the *NRAS* mutation was observed in 1/24 cases (4.2%) during leukemic transformation.

Distribution of chromosomal abnormalities in molecularly mutated MDS cases


We analyzed the frequency of chromosomal abnormalities in the molecularly mutated MDS cases. The cases with *FLT3*-LM, *NRAS* mutation, and *MLL*-PTD showed a high incidence of normal karyotype (*FLT3*-LM: 4/8 [50%]; *NRAS* mutation: 11/14 [79%]; *MLL*-PTD: 7/9 [78%]). We analyzed whether deletions of 5q or monosomy 7 showed any association with *FLT3*-LM or *NRAS* mutations: del(5q) was rare in *FLT3*-LM positive MDS (1/8; 12%) and was not found in *NRAS*- or *MLL*-PTD mutated cases. Chromosome 7 abnormalities were not detected in any of the *FLT3*-LM-, *NRAS*-, or *MLL*-PTD-positive MDS cases (Table 4).

Discussion


Given the new therapeutic options for MDS, such as allogeneic stem cell transplantation with reduced intensity conditioning for elderly patients,⁵⁰⁻⁵² intensive chemotherapy regimens for high-risk MDS,⁵³ and new compounds including azacitidine⁵⁴ and lenalidomide,⁵⁵ risk assessment and prognostic stratification in MDS have become increasingly important. As cytogenetics included in the IPSS provide the basis for prognostic predictions only in some patients,^{6,7} additional parameters are needed for a more detailed characterization of the biology and prognosis of this heterogeneous disorder. In AML it has been established that 80-85% of all cases with normal karyotype can be further characterized by molecular markers, which are found alone or in combination with others (*NPM1*: 50%, *MLL*-PTD: 10%, *CEBPA*: 15%, *FLT3*-LM: 35%, *FLT3*-TKD: 6%, *NRAS*: 10%). In contrast, in MDS screening for molecular mutations is not currently included in routine practice, as the frequency and prognostic impact of these mutations are less well determined. However, there are many questions also with respect to the role of molecular mutations in the leukemic transformation process of MDS. Thus, in this study, we focused not only on the incidence of different molecular mutations in MDS, but also compared the distribution of these markers within the different stages of MDS and in AML.

In consideration of the central role of *FLT3*-mutations in AML ($\geq 30\%$ of all AML patients show either internal tandem duplications or mutations of the tyrosine kinase domain), *FLT3*-mutations have been hypothesized to be important in MDS transformation.^{11,12,20-22} Indeed, our study gives additional support to an association of *FLT3*-LM with progression. These


A




B



C



D



E




Figure 1A-E. Frequency of the different mutations within the different cytomorphic subtypes of MDS (RA/RARS; RAEB; CMML) and of AML (s-AML; AML at relapse). The x-axis shows the different MDS and AML cohorts. The numbers of the analyzed patients are in parentheses. The percentage of mutated cases is given on the y-axis.

Table 3. Frequency of molecular mutations in previous studies.

<i>FLT3-LM</i>	<i>This study</i>	<i>Shih et al., 2004²⁴</i>	<i>Horiike et al., 1997²³</i>	<i>Total</i>
RA	0% (0/45)	0% (0/27)	0% (0/13)	0.0% (0/98)
RARS			0%	0%
RAEB	2.7% (8/293)	2% (2/99)	0% (0/20)	2.4% (10/412)
RAEB-t	–	–	8% (1/12)	8% (1/12)
CMML	0% (0/29)	6% (3/51)	11% (1/9)	4.5% (4/89)
MDS total	2.2% (8/367)	2.5% (5/198)	3% (2/58)	0.2% (15/623)

<i>FLT3-TKD</i>	<i>This study</i>	<i>Yamamoto et al., 2001¹²</i>	<i>Total</i>
RA	3.6% (1/28)	–	3.6% (1/28)
RARS		–	
RAEB	0% (0/209)	0% (0/6)	0% (0/215)
RAEB-t	–	8% (1/13)	8% (1/13)
CMML	0% (0/19)	0% (0/10)	0% (0/29)
MDS total	0.4% (1/256)	3% (1/29)	0.7% (2/285)

<i>NRAS</i>	<i>This study</i>	<i>Nakagawa et al., 1992⁴¹</i>	<i>Paquette et al., 1993⁵⁷</i>	<i>Mitani et al., 1997⁵⁸</i>	<i>Padua et al., 1998⁴²</i>	<i>Total</i>
RA	0% (0/29)	0% (0/10)	8% (6/72)	0% (0/9)	35% (6/17)	10.0% (20/201)
RARS		0% (0/1)	9% (4/46)	0% (0/1)	25% (4/16)	
RAEB	6.7% (15/223)	67% (2/3)	8% (5/63)	0% (0/8)	50% (5/10)	8.8% (27/307)
RAEB-t	–	100% (1/1)	13% (3/23)	9% (2/23)	not specified (6/47)	12.8% (6/47)
CMML	10.0% (2/20)	20% (1/5)	12% (2/16)	–	66% (21/32)	35.6% (26/73)
MDS total	6.3% (17/272)	20% (4/20)	9% (20/220)	2% (2/44)	48% (36/75)	12.5% (79/631)

<i>KITD816</i>	<i>This study</i>	<i>Lorenzo et al., 2006³⁰</i>	<i>Total</i>
RA	0.0% (0/32)	0% (0/10)	0.0% (0/42)
RARS		–	
RAEB	0.8% (2/237)	0% (0/10)	0.8% (2/247)
RAEB-t	–	13% (2/15)	13% (2/15)
CMML	0.0% (0/21)	0% (0/15)	0.0% (0/36)
MDS total	0.7% (2/290)	4% (2/49)	1.2% (4/339)

<i>MLL-PTD</i>	<i>This study</i>
RA	1/46 (2.2%)
RARS	
RAEB	8/292 (2.7%)
RAEB-t	–
CMML	1/30 (3.3%)
MDS total	2.7% (10/368)

Table 4. Distribution of chromosomal aberrations in the molecularly mutated MDS cases.

	<i>FLT3-LM</i>	<i>FLT3-TKD</i>	<i>cKITD816</i>	<i>NRAS</i>	<i>MLL-PTD</i>
Normal karyotype	50% (4/8)	100% (1/1)	0% (0/2)	79% (11/14)	78% (7/9)
Reciprocal translocations	0% (0/8)	0% (0/1)	0% (0/2)	0% (0/14)	0% (0/9)
Complex aberrant	12% (1/8)	0% (0/1)	0% (0/2)	7% (1/14)	0% (0/9)
inv(3)/t(3;3)	0% (0/8)	0% (0/1)	50% (1/2)	0% (0/14)	0% (0/9)
del(5q)	12% (1/8)	0% (0/1)	0% (0/2)	0% (0/14)	0% (0/9)
Chromosome 7 anomalies	0% (0/8)	0% (0/1)	0% (0/2)	7% (1/14)	0% (0/9)
+8	25% (2/8)	0% (0/1)	0% (0/2)	0% (0/14)	22% (0/9)
del(20q)	0% (0/8)	0% (0/1)	0% (0/2)	0% (0/14)	0% (0/9)
-Y	0% (0/8)	0% (0/1)	0% (0/2)	0% (0/14)	0% (0/9)
Others	0% (0/8)	0% (0/1)	0% (0/2)	7% (1/14)	0% (0/9)

mutations were not found in low-risk MDS, but their incidence increased over the proceeding stages (RAEB; RAEB-t) to s-AML, following MDS. In $\geq 10\%$ of all cases the progression of MDS to AML is accompanied by the acquisition of *FLT3-LM* (this study; Shih *et al.*).^{23,56} The occurrence of *FLT3-LM* at diagnosis of MDS is associated with leukemic transformation and shorter survival.⁵⁶ Furthermore, the incidence of *FLT3-LM* was significantly higher in relapsed AML than in *de novo* and s-AML at first diagnosis, underlining the importance of the *FLT3-LM* also in AML progression (Table 3). Thus, rather than being considered as initial events in the development of MDS, *FLT3-LM* should be considered as secondary events involved in MDS progression. The inclusion of the respective mutation status into MDS risk assessment at diagnosis and during follow-up might improve the identification of patients who may benefit from therapy intensification and might be considered also in routine diagnostics.

The role of *FLT3-TKD* mutations in MDS progression is less clear. *FLT3-TKD* are also significantly more frequent in s-AML than in MDS as shown by this study and by Yamamoto *et al.*,¹² and slightly more frequent in AML relapse than in s-AML at diagnosis (*this study*). This points to a role for *FLT3-TKD* in the transformation of MDS and possibly also in relapse of AML.¹² However, definite conclusions cannot yet be drawn as the numbers of cases and studies are too low – so far only two cases of *FLT3-TKD* mutations in

MDS have been reported: one patient with RA in this study and one patient with RAEB in transformation in the study by Yamamoto *et al.*¹²

Due to the rather high incidence of *NRAS*-mutations in AML, interest was focused on the role of this marker in MDS.³¹⁻³⁶ In this study, as in most previous analyses, *NRAS*-mutations were among the most frequent mutations in MDS ($\geq 6.5\%$ of all cases),^{38,41,42,57} and more frequent than *FLT3*-LM ($\leq 3\%$).^{23,24} Although the reported incidences of *NRAS* mutations in MDS range widely (probably due to different proportions of MDS subtypes in the various analyses), this study and all mentioned previous analyses found higher frequencies of *NRAS* mutations in the advanced stages of MDS than in the initial stages.^{41,42,57,58} This demonstrates an association between *NRAS* mutations and MDS transformation. *NRAS* mutations were further shown to be associated with karyotype evolution, e.g. with the acquisition of monosomy 7, during MDS transformation,^{59,60} and with inferior survival in MDS.⁵⁷ Based on these results the inclusion of *NRAS*-screening at diagnosis and during follow-up in MDS might be discussed. With respect to the *MLL*-PTD, our data showed a significantly higher incidence in AML than in MDS, whereas the frequency of *MLL*-PTD did not vary significantly within the diverse cytomorphic MDS subtypes. To our knowledge there are no further studies on this molecular marker in MDS, so the role of *MLL*-PTD in MDS and in leukemogenesis needs further clarification.

KITD816 mutations play a minor role in AML. In MDS, these mutations seem to be restricted to the advanced stages of MDS, as found both in this study and a study by Lorenzo *et al.*,³⁰ suggesting involvement in the transformation towards AML. We found a slightly higher frequency of *KITD816* in AML than in MDS, but the numbers are too small to comment on this fact. We found no influence of AML progression to relapse on the incidence of *KITD816* mutations.

Another aim of our study was an analysis of the cytogenetic characteristics in the molecularly mutated MDS cases. The high rates of normal karyotype in *NRAS*-mutated cases (79% of all *NRAS*-mutated MDS patients in this study, 57% in the study by De Souza *et al.*)⁶¹ support the hypothesis that *NRAS* mutations might represent the initial event in a proportion of MDS cases while additional aberrations induce leukemic transformation.⁶¹ Some authors have suggested a co-operation of chromosome 7 abnormalities with *RAS* and *FLT3*-LM mutations in leukemogenesis. Side *et al.* found monosomy 7 in two patients who progressed from t-MDS to AML with a positive *NRAS* or *FLT3*-LM mutation status.⁶² In a report by Stephenson *et al.*, *RAS*

mutations occurred in three out of seven patients with RAEB in transformation and monosomy 7.⁶⁰ A single case of *NRAS*-positive RAEB with -7 was reported by de Souza *et al.*⁶¹ In contrast to these reports, we found no association between *NRAS* mutations or *FLT3*-LM and chromosome 7 abnormalities in MDS in our study. Therefore, a co-operation of -7 with these molecular markers can be discussed in single cases, but general conclusions should not be drawn at this time due to the low number of reported cases.

We found a 5q- syndrome in 12% of cases with *FLT3*-LM but in no case of MDS with *NRAS* or *MLL*-PTD mutations. This observation corresponds to that of Fidler *et al.* who found no case of *FLT3*-LM, *NRAS*, or *p53* mutations in four patients with the 5q- syndrome and thus suggested that the stability of this MDS syndrome might be a consequence of the absence of other molecular mutations.⁶³

In conclusion, the progression from the initial stages of MDS to secondary AML can be accompanied by the acquisition of molecular mutations which are known to play an important role in AML, such as the *FLT3*-LM or *NRAS*-mutations. This allows the interpretation of these mutations as markers of progression in MDS and supports the two-hit theory, according which at least two different types of mutations are needed for the development of AML: the class I mutations (which are frequently represented by mutations of receptor tyrosine kinases) mediate myeloproliferation, while the class II mutations lead to an arrest in differentiation in hematopoiesis.⁶⁴⁻⁶⁶ It can, therefore, be hypothesized that a single molecular event leads to the early stages of MDS but additional mutations are needed to cause leukemic transformation.

Finally, further evaluation of molecular markers in MDS, especially *FLT3*-LM, *NRAS*, *FLT3*-TKD, *MLL*-PTD, and *KITD816* mutations can be recommended. Such an evaluation should, of course, be completed by analysis of other mutations which are frequent in MDS, such as point mutations in the *AML1/RUNX* gene^{67,68} or mutations of *TP53*.⁶⁹ These studies may lead to new approaches to the subclassification of MDS and to early detection of progression to AML. Thus, complete understanding of the picture of molecular markers in MDS may also be of therapeutic value.

Authors' Contributions

UB: principal investigator. TH, WK, CH: contribution to the design of the study, conducting the work, interpretation of results, and revision of the manuscript. Primary responsibility for the publication, for the tables and figures: UB. Supervision of study: SS.

Conflict of Interest

The authors reported no potential conflicts of interest.

References

1. Aul C, Giagounidis A, Heinsch M, Germing U, Ganser A. Prognostic indicators and scoring systems for predicting outcome in patients with myelodysplastic syndromes. *Rev Clin Exp Hematol* 2004; 8:E1.
2. Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DA, Gralnick HR, et al. Proposals for the classification of the myelodysplastic syndromes. *Br J Haematol* 1982;51: 189-99.
3. Bennett JM. A comparative review of classification systems in myelodysplastic syndromes (MDS). *Semin Oncol* 2005; 32:S3-10.
4. Jaffe ES, Harris NL, Stein H, Vardiman JW. World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues Lyon: IARC Press; 2001.
5. Greenberg P, Cox C, LeBeau MM, Fenaux P, Morel P, Sanz G, et al. International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood* 1997; 89: 2079-88.
6. Fenaux P. Chromosome and molecular abnormalities in myelodysplastic syndromes. *Int J Hematol* 2001; 73:429-37.
7. Fenaux P, Morel P, Lai JL. Cytogenetics of myelodysplastic syndromes. *Semin Hematol* 1996; 33: 127-38.
8. Raskind WH, Steinmann L, Najfeld V. Clonal development of myeloproliferative disorders: clues to hematopoietic differentiation and multistep pathogenesis of cancer. *Leukemia* 1998; 12:108-16.
9. Hellstrom-Lindberg E, Willman C, Barrett AJ, Sauntharajah Y. Achievements in understanding and treatment of myelodysplastic syndromes. *Hematology (Am Soc Hematol Educ Program)* 2000:110-32.
10. Lyman SD, Jacobsen SE. c-kit ligand and Flt3 ligand: stem/progenitor cell factors with overlapping yet distinct activities. *Blood* 1998;91:1101-34.
11. Abu-Duhier FM, Goodeve AC, Wilson GA, Gari MA, Peake IR, Rees DC, et al. FLT3 internal tandem duplication mutations in adult acute myeloid leukaemia define a high-risk group. *Br J Haematol* 2000; 111: 190-5.
12. Yamamoto Y, Kiyoi H, Nakano Y, Suzuki R, Kodaera Y, Miyawaki S, et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood* 2001;97:2434-9.
13. Frohling S, Schlenk RF, Breitnick J, Benner A, Kreitmeier S, Tobis K, et al. Prognostic significance of activating FLT3 mutations in younger adults (16 to 60 years) with acute myeloid leukemia and normal cytogenetics: a study of the AML Study Group Ulm. *Blood* 2002;100:4372-80.
14. Levis M, Small D. Novel FLT3 tyrosine kinase inhibitors. *Expert Opin Investig Drugs* 2003;12:1951-62.
15. Falini B, Mecucci C, Tiacci E, Alcalay M, Rosati R, Pasqualucci L, et al. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med* 2005;352:254-66.
16. Kiyoi H, Naoe T, Nakano Y, Yokota S, Minami S, Miyawaki S, et al. Prognostic implication of FLT3 and N-RAS gene mutations in acute myeloid leukemia. *Blood* 1999; 93: 3074-80.
17. Schnittger S, Schoch C, Dugas M, Kern W, Staib P, Wuchter C, et al. Analysis of FLT3 length mutations in 1003 patients with acute myeloid leukemia: correlation to cytogenetics, FAB subtype, and prognosis in the AMLCG study and usefulness as a marker for the detection of minimal residual disease. *Blood* 2002; 100:59-66.
18. Kottaridis PD, Gale RE, Langabeer SE, Frew ME, Bowen DT, Linch DC. Studies of FLT3 mutations in paired presentation and relapse samples from patients with acute myeloid leukemia: implications for the role of FLT3 mutations in leukemogenesis, minimal residual disease detection, and possible therapy with FLT3 inhibitors. *Blood* 2002;100:2393-8.
19. Schnittger S, Schoch C, Kern W, Mecucci C, Tschulik C, Martelli MF, et al. Nucleophosmin gene mutations are predictors of favorable prognosis in acute myelogenous leukemia with a normal karyotype. *Blood* 2005;106:3733-9.
20. Thiede C, Steudel C, Mohr B, Schaich M, Schakel U, Platzbecker U, et al. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* 2002;99:4326-35.
21. Abu-Duhier FM, Goodeve AC, Wilson GA, Care RS, Peake IR, Reilly JT. Identification of novel FLT-3 Asp835 mutations in adult acute myeloid leukaemia. *Br J Haematol* 2001;113:983-8.
22. Bacher U, Schoch C, Kern W, Haferlach T, Schnittger S. Prognostic relevance of FLT3-TKD mutations in AML: the combination matters- an analysis of 3082 patients (*in preparation*).
23. Horike S, Yokota S, Nakao M, Iwai T, Sasai Y, Kaneko H, et al. Tandem duplications of the FLT3 receptor gene are associated with leukemic transformation of myelodysplasia. *Leukemia* 1997;11:1442-6.
24. Shih LY, Huang CF, Wang PN, Wu JH, Lin TL, Dunn P, et al. Acquisition of FLT3 or N-ras mutations is frequently associated with progression of myelodysplastic syndrome to acute myeloid leukemia. *Leukemia* 2004;18:466-75.
25. Blume-Jensen P, Hunter T. Oncogenic kinase signalling. *Nature* 2001; 411:355-65.
26. Reilly JT. Receptor tyrosine kinases in normal and malignant haematopoiesis. *Blood Rev* 2003; 17:241-8.
27. Care RS, Valk PJ, Goodeve AC, Abu-Duhier FM, Geertsma-Kleinekoort WM, Wilson GA, et al. Incidence and prognosis of c-KIT and FLT3 mutations in core binding factor (CBF) acute myeloid leukaemias. *Br J Haematol* 2003; 121:775-7.
28. Cairoli R, Grillo R, Beghini A, Tedeschi A, Ripamonti CB, Larizza L, et al. C-Kit point mutations in core binding factor leukemias: correlation with white blood cell count and the white blood cell index. *Leukemia* 2003;17:471-2.
29. Schnittger S, Kohl TM, Haferlach T, Kern W, Hiddemann W, Spiekermann K, et al. KIT-D816 mutations in AML1-ETO-positive AML are associated with impaired event-free and overall survival. *Blood* 2006;107: 1791-9.
30. Lorenzo F, Nishii K, Monma F, Kuwagata S, Usui E, Shiku H. Mutational analysis of the KIT gene in myelodysplastic syndrome (MDS) and MDS-derived leukemia. *Leuk Res* 2006;30:1235-9.
31. Radich JP, Kopecky KJ, Willman CL, Weick J, Head D, Appelbaum F, et al. N-ras mutations in adult de novo acute myelogenous leukemia: prevalence and clinical significance. *Blood* 1990;76:801-7.
32. Neubauer A, Dodge RK, George SL, Davey FR, Silver RT, Schiffer CA, et al. Prognostic importance of mutations in the ras proto-oncogenes in de novo acute myeloid leukemia. *Blood* 1994;83:1603-11.
33. Coghlan DW, Morley AA, Matthews JP, Bishop JF. The incidence and prognostic significance of mutations in codon 13 of the N-ras gene in acute myeloid leukemia. *Leukemia* 1994;8: 1682-7.
34. Stirewalt DL, Kopecky KJ, Meshinchi S, Appelbaum FR, Slovak ML, Willman CL, et al. FLT3, RAS, and TP53 mutations in elderly patients with acute myeloid leukemia. *Blood* 2001;97:3589-95.
35. Illmer T, Thiede C, Fredersdorf A, Stadler S, Neubauer A, Ehninger G, et al. Activation of the RAS pathway is predictive for a chemosensitive phenotype of acute myelogenous leukemia blasts. *Clin Cancer Res* 2005;11:3217-24.
36. Bacher U, Haferlach T, Schoch C, Kern W, Schnittger S. NRAS mutations in AML: biology, cytogenetics, and prognosis - a study on 2502 patients. *Blood* 2006;107:3847-53.
37. Hirai H, Kobayashi Y, Mano H, Hagiwara K, Maru Y, Omine M, et al. A point mutation at codon 13 of the N-ras oncogene in myelodysplastic syndrome. *Nature* 1987; 327: 430-2.
38. Bar-Eli M, Ahuja H, Gonzalez-Cadavid N, Foti A, Cline MJ. Analysis of N-RAS exon-1 mutations in myelodysplastic syndromes by polymerase chain reaction and direct sequencing. *Blood* 1989; 73:281-3.
39. Yunis JJ, Boot AJ, Mayer MG, Bos JL. Mechanisms of ras mutation in myelodysplastic syndrome. *Oncogene* 1989;4:609-14.
40. van Kamp H, de Pijper C, Verlaan-de Vries M, Bos JL, Leeksa CH, Kerkhofs H, et al. Longitudinal analysis of point mutations of the N-ras proto-oncogene in patients with

- myelodysplasia using archived blood smears. *Blood* 1992;79:1266-70.
41. Nakagawa T, Saitoh S, Imoto S, Itoh M, Tsutsumi M, Hikiji K, et al. Multiple point mutation of N-ras and K-ras oncogenes in myelodysplastic syndrome and acute myelogenous leukemia. *Oncology* 1992; 49:114-22.
 42. Padua RA, Guinn BA, Al-Sabah AI, Smith M, Taylor C, Pettersson T, et al. RAS, FMS and p53 mutations and poor clinical outcome in myelodysplasias: a 10-year follow-up. *Leukemia* 1998;12:887-92.
 43. Caligiuri MA, Strout MP, Lawrence D, Arthur DC, Baer MR, Yu F, et al. Rearrangement of ALL1 (MLL) in acute myeloid leukemia with normal cytogenetics. *Cancer Res* 1998; 58: 55-9.
 44. Schnittger S, Kinkel U, Schoch C, Heinecke A, Haase D, Haferlach T, et al. Screening for MLL tandem duplication in 387 unselected patients with AML identify a prognostically unfavorable subset of AML. *Leukemia* 2000;14:796-804.
 45. Dohner K, Tobis K, Ulrich R, Frohling S, Benner A, Schlenk RF, et al. Prognostic significance of partial tandem duplications of the MLL gene in adult patients 16 to 60 years old with acute myeloid leukemia and normal cytogenetics: a study of the Acute Myeloid Leukemia Study Group Ulm. *J Clin Oncol* 2002; 20:3254-61.
 46. Strout MP, Marcucci G, Bloomfield CD, Caligiuri MA. The partial tandem duplication of ALL1 (MLL) is consistently generated by Alu-mediated homologous recombination in acute myeloid leukemia. *Proc Natl Acad Sci USA* 1998; 95:2390-5.
 47. Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DA, Gralnick HR, et al. Proposed revised criteria for the classification of acute myeloid leukemia. A report of the French-American-British Cooperative Group. *Ann Intern Med* 1985; 103:620-5.
 48. Weisser M, Kern W, Schoch C, Hiddemann W, Haferlach T, Schnittger S. Risk assessment by monitoring expression levels of partial tandem duplications in the MLL gene in acute myeloid leukemia during therapy. *Haematologica* 2005; 90:881-9.
 49. Schoch C, Kern W, Schnittger S, Hiddemann W, Haferlach T. Karyotype is an independent prognostic parameter in therapy-related acute myeloid leukemia (t-AML): an analysis of 93 patients with t-AML in comparison to 1091 patients with de novo AML. *Leukemia* 2004;18: 120-5.
 50. Larson RA. Myelodysplasia: when to treat and how. *Best Pract Res Clin Haematol* 2006; 19:293-300.
 51. Shimoni A, Hardan I, Shem-Tov N, Yeshurun M, Yerushalmi R, Avigdor A, et al. Allogeneic hematopoietic stem-cell transplantation in AML and MDS using myeloablative versus reduced-intensity conditioning: the role of dose intensity. *Leukemia* 2006;20:322-8.
 52. Deeg HJ, Shulman HM, Anderson JE, Bryant EM, Gooley TA, Slattery JT, et al. Allogeneic and syngeneic marrow transplantation for myelodysplastic syndrome in patients 55 to 66 years of age. *Blood* 2000;95: 1188-94.
 53. Kantarjian H, Beran M, Cortes J, O'Brien S, Giles F, Pierce S, et al. Long-term follow-up results of the combination of topotecan and cytarabine and other intensive chemotherapy regimens in myelodysplastic syndrome. *Cancer* 2006;106:1099-109.
 54. Silverman LR, Demakos EP, Peterson BL, Kornblith AB, Holland JC, Odchimar-Reissig R, et al. Randomized controlled trial of azacitidine in patients with the myelodysplastic syndrome: a study of the cancer and leukemia group B. *J Clin Oncol* 2002;20:2429-40.
 55. List A, Kurtin S, Roe DJ, Buresh A, Mahadevan D, Fuchs D, et al. Efficacy of lenalidomide in myelodysplastic syndromes. *N Engl J Med* 2005; 352:549-57.
 56. Shih LY, Lin TL, Wang PN, Wu JH, Dunn P, Kuo MC, et al. Internal tandem duplication of fms-like tyrosine kinase 3 is associated with poor outcome in patients with myelodysplastic syndrome. *Cancer* 2004;101: 989-98.
 57. Paquette RL, Landaw EM, Pierre RV, Kahan J, Lubbert M, Lazcano O, et al. N-ras mutations are associated with poor prognosis and increased risk of leukemia in myelodysplastic syndrome. *Blood* 1993;82:590-9.
 58. Mitani K, Hangaishi A, Imamura N, Miyagawa K, Ogawa S, Kanda Y, et al. No concomitant occurrence of the N-ras and p53 gene mutations in myelodysplastic syndromes. *Leukemia* 1997;11:863-5.
 59. Horiike S, Misawa S, Nakai H, Kaneko H, Yokota S, Taniwaki M, et al. N-ras mutation and karyotypic evolution are closely associated with leukemic transformation in myelodysplastic syndrome. *Leukemia* 1994;8:1331-6.
 60. Stephenson J, Lizhen H, Mufti GJ. Possible co-existence of RAS activation and monosomy 7 in the leukaemic transformation of myelodysplastic syndromes. *Leuk Res* 1995; 19:741-8.
 61. de Souza Fernandez T, Menezes de Souza J, Macedo Silva ML, Tabak D, Abdelhay E. Correlation of N-ras point mutations with specific chromosomal abnormalities in primary myelodysplastic syndrome. *Leuk Res* 1998; 22:125-34.
 62. Side LE, Curtiss NP, Teel K, Kratz C, Wang PW, Larson RA, et al. RAS, FLT3, and TP53 mutations in therapy-related myeloid malignancies with abnormalities of chromosomes 5 and 7. *Genes Chromosomes Cancer* 2004; 39:217-23.
 63. Fidler C, Watkins F, Bowen DT, Littlewood TJ, Wainscoat JS, Boulwood J. NRAS, FLT3 and TP53 mutations in patients with myelodysplastic syndrome and a del(5q). *Haematologica* 2004; 89: 865-6.
 64. Knudson AG. Two genetic hits (more or less) to cancer. *Nat Rev Cancer* 2001;1:157-62.
 65. Gilliland DG. Hematologic malignancies. *Curr Opin Hematol* 2001; 8:189-91.
 66. Gilliland DG, Griffin JD. The roles of FLT3 in hematopoiesis and leukemia. *Blood* 2002;100:1532-42.
 67. Christiansen DA, Pedersen-Bjergaard J. Mutations of AML1 are common in therapy-related myelodysplasia following therapy with alkylating agents and are significantly associated with deletion or loss of chromosome 7q and with subsequent leukemic transformation. *Blood* 2004;104:1474-81.
 68. Harada H, Harada Y, Niimi H, Kyo T, Kimura A, Inaba T. High incidence of somatic mutations in the AML1/RUNX gene in myelodysplastic syndrome and low blast percentage myeloid leukemia with myelodysplasia. *Blood* 2004; 103: 2316-24.
 69. Kita-Sasai Y, Horiike S, Misawa S, Kaneko H, Kobayashi M. International prognostic scoring system and TP53 mutations are independent prognostic indicators for patients with myelodysplastic syndrome. *Br J Haematol* 2001; 115:309-12.