

Managing the leaching of water-soluble herbicides in soils using eco-compatible nanocarriers

Original

Managing the leaching of water-soluble herbicides in soils using eco-compatible nanocarriers / Tosco, Tiziana; Granetto, Monica; Fogliatto, Silvia; Vidotto, Francesco. - ELETTRONICO. - (2024). (Intervento presentato al convegno EGU General Assembly 2024 tenutosi a Vienna nel 14-19 april 2024) [10.5194/egusphere-egu24-11500].

Availability:

This version is available at: 11583/2989302 since: 2024-06-04T09:01:33Z

Publisher:

Copernicus

Published

DOI:10.5194/egusphere-egu24-11500

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Managing the leaching of water-soluble herbicides in soils using eco-compatible nanocarriers

Tiziana Tosco¹, Monica Granetto¹, Silvia Fogliatto², and Francesco Vidotto²

¹Politecnico di Torino, Department of Environment, Land and Infrastructure Engineering - DIATI, Torino, Italy
(tiziana.tosco@polito.it)

²University of Torino, Department of Agricultural, Forest and Food Sciences - DISAFA, Grugliasco, Italy

Pesticide use plays a crucial role in achieving high crop yields in the context of a global population growth scenario. However, the extensive application of pesticides has progressively led to the contamination of environmental matrices, particularly soils and groundwater, posing potential risks to human health, flora, and fauna. Nanopesticides can be instrumental in mitigating pesticide pollution, especially for highly soluble and volatile active ingredients. They are formed by nanoparticles (nanocarriers) containing an active ingredient, sometimes shielded by a coating, and dispersed in a colloidal suspension. The nanoformulations proposed in this study utilize low-cost mineral materials (such as montmorillonite, zeolite, kaolin) and food-grade biopolymers to incorporate two distinct herbicides, namely dicamba and S-metolachlor, characterized by high solubility (and thus high migration potential in the subsoil) and, for dicamba only, moderate volatility.

The efficacy of the newly developed nanoherbicides in terms of reduced mobility in porous media, reduced persistency, and efficacy toward target weeds was assessed in the laboratory against the pure herbicide and commercial formulations containing the same active ingredients. Specifically, the mobility in porous media was tested through column transport experiments under both saturated and unsaturated conditions, using sand and standard soils (representative, respectively, of top soil and aquifers). These tests were conducted at various scales, ranging from small columns (1.6 cm diameter, 10 cm length) to a laboratory lysimeter (30 cm diameter, 70 cm length). Batch degradation tests in soils indicated comparable DT50 values for the nanoformulation and the commercially available product. The efficacy of the nanopesticides was also examined against conventional products in greenhouse settings through dose-response tests on selected sensitive weeds. The greenhouse tests revealed that clay-based nanoformulations do not impede the effectiveness of dicamba against target weeds, showing efficacy comparable to the commercial competitor for both dicamba and S-Metolachlor, although variations were observed depending on the treated species.

Despite the small scale of the tests conducted in the laboratory and greenhouse, these initial results suggest the promising efficacy of the proposed nanoformulation approach in controlling the environmental spread of soluble herbicides without compromising efficacy against target

species.

This research was conducted within the Nanograss project, co-funded by the Compagnia di San Paolo Foundation.

Reference

Granetto M., Serpella L., Fogliatto S., Re L., Bianco C., Vidotto F., Tosco T. (2022). Natural clay and biopolymer-based nanopesticides to control the environmental spread of a soluble herbicide. *Science of The Total Environment* 806(3),151199, <https://doi.org/10.1016/j.scitotenv.2021.151199>