

Boise State University
ScholarWorks

College of Arts and Sciences Presentations

2015 Undergraduate Research and Scholarship
Conference

4-20-2015

Zircon Chemistry in a Gabbro Pluton at House Mountain, Idaho

Danielle Terhaar

Mark Schmitz

ZIRCON CHEMISTRY IN A GABBRO PLUTON AT HOUSE MOUNTAIN, ID

Danielle Terhaar and Dr. Mark Schmitz
Department of Geosciences, Boise State University




WHAT WE LEARNED

How can we reconstruct the evolution of a magma as it cools and crystallizes minerals?

Zircon crystals form over a range of magma conditions and their chemical signatures record changes in melt composition through cooling and crystallization.

Zircon geochemistry records comprehensive details of magma composition and evolution.

BACKGROUND




An igneous pluton at the base of House Mountain, Idaho is composed of a gabbro body with cross-cutting granophyre veins (Alexander, 2007). From field relationships and mineralogy, I hypothesize that the granophyre evolved from the gabbro, and have tested the ability of zircon geochemistry to record this relationship.

Key Concepts


- House Mountain HM14-04 granophyre is the residual melt that segregated from HM14-03 gabbro and represents a late stage of magma evolution.
- Magma composition changes through time due to cooling and crystallization (White, 2013).
- Zircon compositions are related to the composition and temperature of the host magma, as well as the degree of crystallization of the magma.

RESULTS




HM14-03 is a gabbro (a mafic igneous intrusive rock) with quartz & feldspar segregations around prismatic hornblende phenocrysts.

HM14-04 is a granophyre (a felsic igneous intrusive rock) with elongate skeletal hornblende phenocrysts surrounded by a quartz + feldspar-rich matrix.



- Ti-in-zircon temperatures are relatively low in both samples, near the solidus.**
Although the gabbro zircons are slightly higher T on average, both rocks record a similar range of temperatures, suggesting that zircon is a very late crystallizing phase in felsic segregations.
- The granophyre records larger Eu anomalies (lower Eu/Eu^*) related to increased feldspar crystallization relative to the gabbro.**
 Eu/Eu^* is a measure of degree of feldspar crystallization; the observed anomalies are consistent with the petrography of the samples.
- The granophyre zircons are enriched in trace elements (ΣREE , Ta, Nb, Hf) compared to the gabbro zircons, consistent with greater degree of crystallization.**
As minerals crystallize and are removed from the bulk melt, trace elements concentrate in melt fraction; zircon incorporates these trace elements.



Methods

- Zircons were separated from the rocks by density and magnetic methods, mounted in epoxy, polished to their centers and imaged using cathodoluminescence.
- Chemical compositions were measured on 25 μ m diameter spot analyses with laser ablation inductively coupled mass spectrometry (LA-ICPMS).
- Trace elements including U, Th, Nb, Ta, Hf, and rare earth elements (REE) were compared on bivariate diagrams with Ti-in-zircon thermometry (Ferry and Watson, 2007).

References

- Alexander, J., 2007, Geochronology of the House Mountain Gneiss Complex in the Atlanta Lobe of the Idaho Batholith. M.S. Geology, Boise State University.
- Bachmann, O., & Bergantz, G. W., 2004, On the origin of crystal-poor rhyolites: Extracted from batholithic crystal mushes. *Journal of Petrology*, v.45, i.8, p.1565-1582.
- Ferry, J.M., and Watson, E.B., 2007, New thermodynamic models and revised calibrations for the Ti-in-zircon and Zr-in-rutile thermometers. *Contributions to Mineralogy and Petrology*, v. 154, no. 4, p. 429-437.
- Gaschnig, R. M., Vervoort, J. D., Lewis, R. S., & Tikoff, B., 2011, Isotopic Evolution of the Idaho Batholith and Challis Intrusive Province, Northern US Cordillera. *Journal of Petrology*, v.52, i.12, p. 2397-2429.
- White, William M., 2013, *Geochemistry*, Wiley-Blackwell: Chichester, UK, p. 268-320.