

1 **REMOVAL OF INORGANIC MERCURY FROM AQUEOUS SOLUTIONS BY**  
2 **BIOMASS OF THE MARINE MACROALGA *CYSTOSEIRA BACCATA***

3 Roberto Herrero<sup>\*</sup>, Pablo Lodeiro, Carlos Rey-Castro, Teresa Vilariño, Manuel E. Sastre  
4 de Vicente.

5 Departamento de Química Física e Enxeñería Química I, University of A Coruña, C/  
6 Alejandro de la Sota, 1, 15008 A Coruña, Spain.

7 <sup>\*</sup>Corresponding author e-mail address: [erob@udc.es](mailto:erob@udc.es); fax: +34 981 167065.

8

9 **Abstract**

10 The ability of *Cystoseira baccata* algal biomass to remove Hg(II) from aqueous  
11 solutions is investigated. The mercury biosorption process is studied through batch  
12 experiments at 25°C with regard to the influence of contact time, initial mercury  
13 concentration, solution pH, salinity and presence of several divalent cations. The acid-  
14 base properties of the alga are also studied, since they are related to the affinity for  
15 heavy metals. The studies of the pH effect on the metal uptake evidence a sharp  
16 increasing sorption up to a pH value around 7.0, which can be ascribed to changes both  
17 in the inorganic Hg(II) speciation and in the dissociation state of the acid algal sites.  
18 The sorption isotherms at constant pH show uptake values as high as 178 mg.g<sup>-1</sup> (at pH  
19 4.5) and 329 mg.g<sup>-1</sup> (at pH 6.0). The studies of the salinity influence on the Hg(II)  
20 sorption capacity of the alga exhibit two opposite effects depending on the electrolyte  
21 added; an increase in concentration of nitrate salts (NaNO<sub>3</sub>, KNO<sub>3</sub>) slightly enhances  
22 the metal uptake, on the contrary, the addition of NaCl salt leads to a drop in the  
23 sorption. The addition of different divalent cations to the mercury solution, namely  
24 Ca<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, Cd<sup>2+</sup>, Pb<sup>2+</sup> and Cu<sup>2+</sup>, reveals that their effect on the uptake process is  
25 negligible. Finally, the equilibrium sorption results are compared with predictions

26 obtained from the application of a simple competitive chemical model, which involves a  
27 discrete proton binding constant and three additional constants for the binding of the  
28 main neutral inorganic Hg(II) complexes, Hg(Cl)<sub>2</sub>, HgOHCl and Hg(OH)<sub>2</sub>, to the algal  
29 surface sites.

30

31 **Keywords:** *Cystoseira baccata*, mercury, adsorption, marine macroalga.

32

### 33 **Introduction**

34 Water pollution by toxic metals in industrial wastewaters has become a major  
35 issue throughout the world. In particular, mercury compounds, whose effects on human  
36 health and aquatic life are regarded as harmful, must be removed from sewage down to  
37 extremely low concentrations. Even though the flux of mercury into the aquatic system  
38 has declined in recent years, there is still a lack of effective and cheap resources for  
39 such wastewater treatment.

40 The effluents from chlorine and chlor-alkali manufacturing processes, via  
41 electrolysis in mercury cells, represent one of the most important sources of mercury  
42 pollution. Other contributions come from pulp, oil refining, plastic and battery  
43 manufacturing industries.

44 Adsorption as a wastewater treatment process has been found to be an  
45 economically feasible alternative for metal removal. Activated carbon is one of the most  
46 well-known adsorbents [Bello et al. (1999); Budinova et al. (2003); Gómez-Serrano et  
47 al. (1998)] but the high costs of the process has limited its use. A search for a low-cost  
48 and easily available adsorbent has led to the investigation of materials of biological  
49 origin as potential metal biosorbents. The variety of materials tested includes bark,  
50 chitin, lignin, modified wool and seaweeds [Bailey et al. (1999)]. They can be used for

51 the effective removal and recovery of several species from wastewater streams. Among  
52 these materials, a few brown marine macroalgae species exhibit higher uptake values  
53 than those of activated carbon and natural zeolite, and are comparable to those of some  
54 synthetic ion exchange resins.

55 The algal cell wall plays an important role in metal binding [Crist et al. (1988)],  
56 due to its high content in polysaccharides with acid functional groups. In brown algae,  
57 the cell wall is mainly comprised of alginates, which usually constitute about 20-40% of  
58 the total dry weight, in addition to fucoidans [Percival and McDowell (1967)]. The  
59 carboxyl groups of alginates are likely to be the main functionalities involved in metal  
60 binding reactions [Schiewer and Wong (2000)] because of their higher abundance with  
61 regard to both carboxyl and amine groups of the proteins.

62 The present work reports a study of the mercury adsorption by non-living  
63 biomass of the brown marine macroalga *Cystoseira baccata* originated from the  
64 Galician coast (NW, Spain). The biosorption process has been analyzed through batch  
65 experiments at 25°C with regard to the influence of the initial metal concentration, the  
66 pH and salinity of the solution, and the presence of different divalent cations. The acid-  
67 base properties of the alga have been studied as they are related to its capacity of  
68 binding heavy metals. Finally, a complexation model for mercury sorption on  
69 *Cystoseira baccata* has been developed in order to explain the adsorption behavior and  
70 the different effects studied on the metal uptake.

71

## 72 **Materials and methods**

73 The brown alga *Cystoseira baccata* was collected from the coasts of A Coruña  
74 (Galicia, NW Spain). The alga was washed twice with running water and once with  
75 deionized water. The washed biomass was oven-dried at 60°C for 24 hours, crushed

76 with an analytical mill, sieved (size fraction of 0.5-1 mm) and stored in polyethylene  
77 bottles until use. The samples for the potentiometric titrations were acid-treated with  
78 diluted HCl following the procedure described elsewhere [Rey-Castro et al. (2003)], in  
79 order to transform the biomass into its fully protonated form.

80 All chemicals used in this work were purchased from Merck. Cellulose nitrate  
81 membrane filters were from Whatman and Albet. Throughout this work, all the  
82 experiments were conducted at least in duplicate.

83

#### 84 *Kinetic studies*

85 Kinetic studies of Hg adsorption by the alga *Cystoseira baccata* were  
86 accomplished to estimate the time necessary to reach the sorption equilibrium. The  
87 experiments were carried out adding 0.25g of dried biomass over 100 ml of a solution  
88 that contained 500 mg·L<sup>-1</sup> of Hg(II) (from HgCl<sub>2</sub>) and sufficient NaNO<sub>3</sub> salt to keep the  
89 ionic strength constant at 0.05M. The mercury content of the solution was checked by  
90 taking aliquots at certain time intervals. The concentration of dissolved Hg in each  
91 aliquot was then analysed by inductively coupled plasma mass spectrometry (ICP-MS).

92

#### 93 *Potentiometric titrations*

94 For each titration, ca. 0.5 g of protonated *C. baccata* biomass were placed in a  
95 thermostated glass cell at a temperature of 25.0±0.1 °C, and 100 mL of 0.05 M NaNO<sub>3</sub>  
96 solution were added to keep ionic strength constant. HCl was also added to yield an  
97 initial pH value around 2.0. The suspension, magnetically stirred, was allowed to  
98 equilibrate and the titration was started solely once the electromotive force  
99 measurement was stable. A NaOH solution, prepared with boiled deionized water, was  
100 employed as titrant, which was added from a Crison microBu 2031 automatic burette.

101 Emf measurements were done by a Crison micropH 2000 meter equipped with a  
102 GK2401C Radiometer combined glass electrode (saturated Ag/AgCl as reference).  
103 After each addition of base, the system was allowed to equilibrate before a stable  
104 reading was obtained. A whole titration typically took 6-7 h.

105 The glass electrode was calibrated in solutions of known proton concentration at  
106 a constant ionic strength following the procedure described elsewhere [Brandariz et al.  
107 (1998); Fiol et al. (1992)]. All titration experiments were done under a nitrogen stream,  
108 intended to remove dissolved O<sub>2</sub> and CO<sub>2</sub>.

109

#### 110 *Adsorption isotherms*

111 The isotherm experiments were carried out at three different values of pH (4.5,  
112 6.0 and 8.0), adjusted by the addition of NaOH or HNO<sub>3</sub> solutions. Different amounts of  
113 HgCl<sub>2</sub> were dissolved in deionized water to prepare ten mercury solutions in a  
114 concentration range from 20 to 2000 mg·L<sup>-1</sup>. A volume of 40 mL of the metal solution  
115 was then added to a 100 mL Erlenmeyer flask containing 0.1 g of alga. The mixtures  
116 were stirred in a rotary shaker at 175 rpm for 4 hours until equilibrium was reached.  
117 Afterwards the suspension was filtered through a 0.45 µm pore size cellulose nitrate  
118 membrane filter. The filtrates were analysed for mercury and sodium contents by ICP-  
119 MS and for chloride ion (which constitutes a key factor in the inorganic mercury  
120 speciation) by capillary electrophoresis.

121 The amount of mercury sorbed at equilibrium,  $q_{eq}$  (mg·g<sup>-1</sup>), which represents the  
122 metal uptake, was calculated from the difference in metal concentration in the aqueous  
123 phase before and after adsorption, according to the following equation:

$$q_{eq} = \frac{V \cdot (C_i - C_{eq})}{1000 \cdot m_s} \quad (1)$$

124 where  $V$  is the volume of mercury solution (mL),  $C_i$  and  $C_{eq}$  are the initial and  
125 equilibrium concentration of mercury in solution ( $\text{mg}\cdot\text{L}^{-1}$ ), respectively, and  $m_s$  is the  
126 mass of dry alga (g).

127

### 128 *Influence on the metal adsorption of pH, salinity and addition of divalent cations*

129 The dependence of *Cystoseira baccata* metal uptake on pH was studied through  
130 batch sorption experiments in the pH range from 0.7 to 9.0, with an initial mercury  
131 concentration of  $500 \text{ mg}\cdot\text{L}^{-1}$ . The pH was adjusted by addition of NaOH and  $\text{HNO}_3$   
132 solutions.

133 The effect of salinity on the adsorption of mercury was tested by addition of  
134 different salts (NaCl,  $\text{NaNO}_3$  and  $\text{KNO}_3$ ) to the solution. The concentration of each salt  
135 ranged from 0.001 to 1 M. The initial mercury concentration and pH were  $500 \text{ mg/L}$   
136 and 6.0, respectively, for all the experiments.

137 The competition effect of several divalent cations, namely Cd(II), Mg(II), Zn(II),  
138 Ca(II), Cu(II) and Pb(II), was tested through batch sorption experiments conducted with  
139 0.1 g of *Cystoseira baccata* in contact with binary mixtures composed of  $50 \text{ mg}\cdot\text{L}^{-1}$   
140 Hg(II) and the competitor metal ion at two different concentrations (500 or  $1000 \text{ mg}\cdot\text{L}^{-1}$   
141 <sup>1</sup>) prepared from the respective metal nitrate salt. Throughout the experiments, the pH  
142 was kept constant at 4.5 to avoid hydrolysis of the cations, specially Cu(II) and Pb(II).

143

## 144 **Results and discussion**

### 145 *Kinetics of the adsorption process: Effect of Contact Time*

146 The batch experiments carried out to study the relationship between contact time  
147 and mercury uptake by *Cystoseira baccata* show that the equilibrium time is always  
148 reached in less than 100 minutes (see Figure 1 as an example). It can be noticed that the

149 contact time significantly affects the Hg uptake; the metal adsorption increases sharply  
150 in the first 50 minutes and tapers off there after, as equilibrium is approached. This  
151 relatively rapid mercury uptake indicates that the sorption process occurs mainly on the  
152 surface of the adsorbent.

153 According to these results, it was set a contact time of 4 hours in order to ensure  
154 that equilibrium conditions are attained. This equilibrium time is clearly shorter than  
155 those usually employed for the adsorption of Hg by other adsorbent materials. Times of  
156 24 hours are proposed for the adsorption by chemically modified chitosan [Jeon and  
157 Höll (2003)], pinus pinaster bark [Vázquez et al. (2002)] or ion exchange resins [Chiarle  
158 et al. (2000)]. Even longer times, from 80 to 120 hours, are necessary with some  
159 carbonaceous materials [Cox et al. (2000)]. However, similar equilibrium times were  
160 found for other heavy metal adsorptions by different macroalgae [Cordero et al. (2004);  
161 Cruz et al. (2004); Lodeiro et al. (2004a)].

162 The rapid kinetics has a significant practical importance, as it will facilitate  
163 smaller reactor volumes ensuring efficiency and economy.

164

#### 165 *Acid-base properties of Cystoseira baccata*

166 The total amount of active sites in protonated *C. baccata* biomass was estimated  
167 by potentiometric titration with a standard solution of NaOH. The number of acid  
168 groups per gram of alga,  $[A]_T$  ( $\text{mmol}\cdot\text{g}^{-1}$ ), was calculated from the maximum of the first  
169 derivative of the titration curves (Figure 2).

170 Sulfate groups are known to be present in the algal cell wall [Percival and  
171 McDowell (1967)]; however, no evidence of their presence was found in the titration  
172 curves. The total number of weak acid groups determined in 0.05M  $\text{NaNO}_3$  was  $2.2 \pm$   
173  $0.1 \text{ mmol g}^{-1}$ . From titrations in different saline media, it was concluded that ionic

174 strength does not influence the number of acidic groups titrated, but strongly affects  
175 their apparent pK values. A physicochemical model based on the Donnan formalism has  
176 recently been proposed by Rey-Castro *et al.* [Rey-Castro et al. (2003)], [Rey-Castro et  
177 al. (2004b)] in order to account for the effects of pH and ionic strength on the proton  
178 binding equilibria of seaweed biomass.

179

#### 180 *Effect of pH on Hg(II) biosorption*

181 Solution pH values have a significant influence on mercury uptake by *Cystoseira*  
182 *baccata*. Several other researchers have already reported a strong dependency of heavy  
183 metal biosorption on pH [Volesky (2003); Wase and Forster (1997)]. As seen from  
184 Figure 3 (a), the mercury uptake is small at low pH. Between pH values 2.0 and 7.0 the  
185 metal adsorption increases sharply, attaining values that remain almost constant for  
186 higher pH values.

187 As a general rule, the pH influence on metal uptake by seaweeds is closely  
188 related to the ionic states of the cell wall functional groups as well as to the metal  
189 speciation in solution. In the case of Hg biosorption, the pH dependence is slightly  
190 different to that observed for other metals. As an example, Cd(II) also presents an S-  
191 shape curve but the maximum uptake is reached at pH 4.5 [Lodeiro et al. (2004b)]. It  
192 can well be assumed that cadmium is present in its free ionic form,  $Cd^{+2}$ , all along the  
193 pH range studied. Therefore, the cadmium biosorption depends on the protonation or  
194 deprotonation of the cell wall functional groups, mainly carboxylic groups, whose  $pK_a$   
195 values are about 2 to 4 [Rey-Castro et al. (2004a)]. On the contrary, the mercury  
196 biosorption process is not only affected by the acid-base properties of the cell wall but  
197 also by the metal chemical speciation, which is rather more complex than that of Cd(II),  
198 and hence, it may play an important role (see Figure 3 b).



199

200 *Adsorption isotherms*

201           The metal distribution between the alga and the aqueous solution at equilibrium  
202 is of importance in determining the maximum adsorption capacity of the alga for  
203 mercury. Figure 4 illustrates the adsorption of mercury on biomass at different pH  
204 values. It can be noted that the uptake rises dramatically with increasing pH. The plots  
205 of uptake vs. aqueous Hg(II) obtained at pH 4.5 and 6.0 are smooth, continuous curves  
206 that eventually reach a saturated value, which suggests monolayer coverage of mercury  
207 on the adsorbent surface, a typical behaviour of most metal ions. At pH 8.0, however,  
208 together with much greater saturation values, an anomalous S-shaped increasing curve  
209 is obtained, which may be explained by a combination of phenomena. On the one hand,  
210 similar S-shaped isotherms have already been described in several studies concerning  
211 Hg(II) sorption to soils with high contents of organic matter (see Drexel et al. [Drexel et  
212 al. (2002)] and references therein), which was attributed to the binding of metal to  
213 soluble organic matter released from the sorbent. In fact, there is an important organic  
214 leaching from seaweed biomass above pH 6, due mainly to alginate solubilization. On  
215 the other hand, the large maximum uptake values determined at pH 8.0 could be  
216 assigned to the presence of surface precipitation processes, which may occur at metal  
217 concentrations close to the limit of bulk precipitation, leading to an increasing slope of  
218 the adsorbed vs. free metal plots (departing from the classical Langmuir isotherm shape)  
219 [Schneider et al. (2001)].

220           From the experiments at constant pH, it can be noticed that an increase in the  
221 initial mercury concentration leads to a larger sorption capacity of the biomass at  
222 equilibrium, while the percentage of mercury removed from solution shows the opposite  
223 trend. Nevertheless, the alga is still able to remove percentages greater than 80% of

224 mercury in solution for relatively high initial mercury concentrations using small  
225 amounts of biosorbent.

226         It can be then concluded that the biomass of the marine alga *Cystoseira baccata*  
227 can efficiently remove high concentrations of mercury in solution over a broad range of  
228 pH, highlighting its potential for effluent treatment processes. The high sorption  
229 capacity of this seaweed is comparable to or even larger than other natural and synthetic  
230 materials, such as fungal biomass [Saglam et al. (1999)] ( $61 \text{ mg}\cdot\text{g}^{-1}$ ), the green alga  
231 *Ulva lactuca* [Zeroual et al. (2003)] ( $149 \text{ mg}\cdot\text{g}^{-1}$ ), the aquaphyte *Potamogeton natans*  
232 [Lacher and Smith (2002)] ( $180 \text{ mg}\cdot\text{g}^{-1}$ ), active carbons from different sources  
233 [Budinova et al. (2003); Yardin et al. (2003)] ( $132\text{-}174 \text{ mg}\cdot\text{g}^{-1}$ ), the synthetic resin  
234 Duolite GT-73 [Chiarle et al. (2000)] ( $362 \text{ mg}\cdot\text{g}^{-1}$ ) or chitosan [Jeon and Höll (2003);  
235 Masri et al. (1974); McKay et al. (1989)] ( $460, 1123$  and  $815 \text{ mg}\cdot\text{g}^{-1}$ , respectively).

236

#### 237 *Effect of salinity on mercury uptake*

238         The studies of the salinity influence on the Hg(II) sorption capacity of the alga  
239 exhibit two opposite effects depending on the electrolyte added, see Figure 5 (a-b). An  
240 increase in concentration of nitrate salts ( $\text{NaNO}_3$ ,  $\text{KNO}_3$ ) causes greater mercury uptake,  
241 from 5% up to 12-15% as the salt concentration varies from 0.01M to 1M, respectively.  
242 On the contrary, a larger presence of NaCl salt leads to a drop in the sorption, which is  
243 decreased by more than 80% at salt concentrations around 1 M, whereas low values of  
244 salinity have small influence on the metal uptake (declined by 8% at NaCl concentration  
245 0.01M).

246         It is often stated in literature that light metal ions, such as sodium or potassium,  
247 compete with divalent cations for the electrostatic binding to the biomass [Schiewer and  
248 Wong (2000)]. Therefore, if a similar mechanism was involved in the present study, the

249 mercury sorption should decrease as the concentration of light metal ions increases.

250 In view of the results of this work (see Figure 5), it can be concluded that the  
251 major effect on the mercury biosorption capacity of the alga is caused by the salt anions,  
252 not by the light metal cations. This turns out to be clear from the comparison of the  
253 mercury sorption that occurs in two saline solutions containing the same concentration  
254 of sodium ion but different type of counterion. As an example, the mercury uptake  
255 decreases by 83% in presence of chloride ions whereas it increases by 15% in presence  
256 of nitrate, although sodium salt concentration is 1M in both cases. Such behavior can be  
257 explained taking into account that the mercury speciation is highly affected by the  
258 presence of chloride ions, which induce the formation of  $\text{HgCl}_3^-$  and  $\text{HgCl}_4^{2-}$  complexes  
259 with low sorption capacity on the algal cell wall. This imposes a limitation in the  
260 application of this technology to waters containing high chloride concentrations, such as  
261 seawater or wastewaters from brine industries. Nevertheless, the Hg(II) adsorption  
262 capacity of algal biomass continues being very high in saline media whose chlorine  
263 concentration does not exceed 0.01M.

264

#### 265 *Effect of the competition of divalent cations on Hg uptake*

266 Figure 6 shows the results of the mercury adsorption experiments carried out in  
267 the presence of 500 or 1000  $\text{mg}\cdot\text{L}^{-1}$  of divalent cation – Cd(II), Mg(II), Zn(II), Ca(II),  
268 Cu(II) and Pb(II) –. It can be observed that the uptake remains practically unaffected  
269 due to the presence of these divalent metal ions at a much greater concentration than  
270 mercury. Only Cu(II) ions seem to decrease slightly the mercury sorption, whereas  
271 Pb(II) ions increase it. Such results could be explained in terms of the different chemical  
272 speciation of metals at the pH studied, which have a significant effect in the adsorption  
273 mechanism. Whereas mercury is largely present as a neutral species,  $\text{HgCl}_2$ , the other

274 metals appear as divalent ions,  $M^{+2}$ , which will interact with the algal cell walls mainly  
275 through an electrostatic mechanism, thus avoiding any competition with the sorption  
276 process of the neutral species of mercury. In fact, different authors have assumed that  
277  $Hg(OH)_2$  behaves as a ligand in aqueous solution [Daughney et al. (2002); Sarkar et al.  
278 (1999)], and hence a similar behavior may be proposed also for  $HgCl_2$ . In this way, its  
279 interaction with the active groups of the algal wall (represented by  $A^-$ ) can be illustrated  
280 with a reaction of the type:



282 Such mechanism does not lead to any change in the charge of the algal surface,  
283 which, hence, could still bind any other metal cation present in solution. Such specific  
284 adsorption of the neutral species  $HgCl_2$  is also supported by the small changes found in  
285 mercury adsorption when the effect of ionic strength was studied.

286 Of all the divalent cations studied, only copper appears to compete slightly with  
287 mercury for sorption sites. Due to its relatively “soft” character,  $Cu^{2+}$  would tend to  
288 form covalent bonds more readily than “hard” cations such as  $Ca^{2+}$  or  $Mg^{2+}$ , which are  
289 mainly electrostatically bound [Schiewer and Wong (1999)]. In addition, copper ions  
290 start undergoing hydrolysis at pH values as low as 4.5, yielding some amount of neutral  
291 species (of higher softness). Since  $Hg(II)$  has a strong “soft” character (especially in the  
292 form of neutral species like  $HgCl_2$ ), it may therefore be expected that  $Cu(II)$  species will  
293 compete to some extent with mercury in the complexation process to the algal wall. On  
294 the contrary,  $Pb(II)$  seems to reinforce the mercury uptake, but authors have not found  
295 any consistent explanation.

296 It is of interest to underline that these studies have been carried out at  
297 concentration values of divalent cations significantly higher than those found in natural  
298 media. The results obtained make clear the high capacity of *Cystoseira baccata* for the

299 disposal of mercury from polluted natural waters, even in presence of high contents of  
300 other divalent ions, which is likely to occur, especially in hard water.

301

### 302 **Modelling of Hg adsorption equilibria**

303 The purpose of this section is to explore how well a simple thermodynamically  
304 consistent chemical model is able to describe approximately most of the experimental  
305 results that have been discussed in previous sections, and to account for the observed  
306 influence of the environmental variables (pH, ionic strength, amount of chloride, etc.).  
307 Such a model would be useful for the prediction of the sorbent performance under  
308 practical conditions.

309 The basic assumptions of this model are:

310 (a) The sorption of the inorganic Hg(II) species involves at least the binding to an  
311 ionized acid group of the seaweed. Therefore, protons and mercury are assumed to  
312 compete with each other for the same acid sites on the algal surface. This is supported  
313 by the fact that the observed variation of the Hg(II) biosorption with pH cannot be  
314 entirely ascribed to the change in mercury speciation.

315 (b) Inorganic Hg(II) essentially appears as neutral species, mainly  $\text{HgCl}_2$ ,  $\text{Hg}(\text{OH})\text{Cl}$   
316 and  $\text{Hg}(\text{OH})_2$ , which behave as ligands in the sorption process, as discussed above. In  
317 fact, calculations of the inorganic Hg(II) speciation using the MINTEQA2  
318 database[Allison et al. (1991)] showed that more than 98% of the total dissolved  
319 mercury appears as neutral species between pH 1 and 9, for the range of Hg  
320 concentrations of this study.

321 (c) For the sake of simplicity, a discrete constant for the binding of each mercury  
322 species is considered, through the corresponding ideal Langmuir isotherms. It must be  
323 admitted that this assumption is quite rough, as shown from potentiometric studies of

324 the acid-base properties [Rey-Castro et al. (2004b); Rey-Castro et al. (2003)]. The algal  
 325 biomass was proved to behave as a heterogenous proton sorbent with acid sites  
 326 exhibiting a continuous distribution of affinities for protons, this being a result of  
 327 different factors (polyelectrolytic effect, intrinsic chemical heterogeneity of the sorbent,  
 328 etc.). Therefore, a model reflecting a distribution of site affinities would surely be more  
 329 accurate. However, this option implies the use of a relatively large number of empirical  
 330 parameters with regard to the amount of available data. On the contrary, the  
 331 consideration of a discrete binding constant for each neutral Hg(II) species is able to  
 332 describe roughly the experimental trends observed, with the smallest number of  
 333 empirical parameters.

334 Finally, it must be pointed out that the simple model proposed in this study does  
 335 not take into account other possible mechanisms for the mercury uptake by the biomass  
 336 postulated in bibliography, such as the sequestering of neutral mercuric species in the  
 337 lipid environment of the algal cellular membranes [Mason et al. (1995)]. As will be  
 338 explained below, this mechanism alone would not explain the observed effect of pH on  
 339 mercury uptake.


340

#### 341 *Analytical isotherms*

342 Assumptions (a)-(c) lead us to the consideration of the following chemical  
 343 model, in which each of the neutral mercury species interacts with at least one acid  
 344 functional group of the algae ( $A^-$ ):



345 where  $M = \text{Hg}(\text{Cl})_2, \text{HgOHCl}$  or  $\text{Hg}(\text{OH})_2$ ,  $K_i$  ( $i=1,2,3$ ) represents the binding constant  
 346 of each species, respectively, and  $n$  is the average number of binding sites per metal  
 347 bound. For instance, the complexation of  $\text{Hg}(\text{OH})_2$  (assuming a 1:1 stoichiometry)  
 348 would be represented by:



349 In this work, only the ideal stoichiometries 1:1 and 1:2 have been considered, for  
 350 simplicity.

351 From these equilibria and the mass balance for the ligand, an analytical  
 352 expression for the sorption isotherm may be easily derived for 1:1 or 1:2  
 353 stoichiometries. The 1:1 model leads to

$$\begin{aligned} [\text{A}]_T &= [\text{AH}] + [\text{A}^-] + [\text{AHg}(\text{Cl})_2^-] + [\text{AHgOHCl}^-] + [\text{AHg}(\text{OH})_2^-] = \\ &= [\text{A}^-] \left( 1 + K_H [\text{H}^+] \right) + \sum [\text{AM}^-] \end{aligned} \quad (5)$$

354 By solving the equilibrium mercury complexation equations ( 3 ) the  
 355 concentration of free ligand can be expressed as

$$[\text{A}^-] = \frac{[\text{AHg}(\text{Cl})_2^-]}{K_1 [\text{Hg}(\text{Cl})_2]} = \frac{[\text{AHgOHCl}^-]}{K_2 [\text{HgOHCl}]} = \frac{[\text{AHg}(\text{OH})_2^-]}{K_3 [\text{Hg}(\text{OH})_2]} \quad (6)$$

356 An expression for the overall concentration of bound mercury is readily obtained  
 357 from eqs.( 5 ) and ( 6 ):

$$\sum [\text{AM}^-] = q_{\text{eq}} = [\text{A}]_T \frac{\sum K_i [\text{M}]}{1 + K_H [\text{H}^+] + \sum K_i [\text{M}]} \quad (7)$$

358 where  $q_{\text{eq}}$  and  $[\text{A}]_T$  must be expressed in the same unities ( $\text{mmol} \cdot \text{g}_{\text{alga}}^{-1}$  or  $\text{mg}_{\text{Hg}} \cdot \text{g}_{\text{alga}}^{-1}$ ),  
 359 and the rest of the concentrations are expressed in  $\text{mol} \cdot \text{L}^{-1}$ .

360 Analogously, when a 1:2 stoichiometry, i.e.  $n=2$  in eq. ( 3 ), is considered, one  
 361 gets:

$$q_{\text{eq}} = 0.5[A]_{\text{T}} \frac{\sum (K_i [M])^{0.5}}{1 + K_{\text{H}} [H^+] + \sum (K_i [M])^{0.5}} \quad (8)$$

362 Eqs. ( 7 ) and ( 8 ) represent very simple competitive ideal Langmuir isotherms  
 363 that involve the assumption of the algal binding sites behaving as homogenous ligands,  
 364 with regard to the binding of protons and mercury. These analytical expressions must be  
 365 combined with an accurate prediction of the distribution of mercury species in solution.

366

### 367 *Inorganic Hg(II) speciation*

368 The chemistry of mercury in aqueous solution is quite complex. Both the  
 369 distribution of the Hg(II) species and the mercury oxide precipitation are very sensitive  
 370 to the solution variables, such as pH or chloride concentration. Furthermore, some  
 371 disagreements about the formation constants of the different inorganic complexes of  
 372 Hg(II) are still found in bibliography. In this work, the inorganic speciation of Hg(II)  
 373 was estimated through MINEQL+ [Schecher and McAvoy (1992)] using the  
 374 thermodynamic database from MINTEQA2 [Allison et al. (1991)]. The activity  
 375 coefficients were calculated through the Davies equation [Sastre de Vicente (1997)]  
 376 applied to the bulk ionic strength values estimated either from the acid and/or base  
 377 additions made for pH adjustment, or from the background salt addition. The  
 378 experimental measurements of mercury, chloride and sodium in the filtrate solutions  
 379 were taken as the total concentrations in the MINEQL+ input.

380

### 381 *Choice of model parameters and comparison with experimental results*

382 The total number of functional groups,  $[A]_{\text{T}}$ , was set to  $2.2 \text{ mmol}\cdot\text{g}^{-1}$ , the value  
 383 determined from the acid-base potentiometric titrations in the absence of Hg species.  
 384 The fit of the proton binding data to an isotherm model allowed the estimation of an



385 average acid constant  $K_H$ , referred to  $\text{NaNO}_3$  0.05 M, with a value of  $10^{3.6}$ . The  
386 variation of this constant with ionic strength was estimated by means of an empirical  
387 Donnan expression derived elsewhere [Rey-Castro et al. (2004b)] using potentiometric  
388 titration data in different electrolytes [Rey-Castro et al. (2003)].

389 The values of the binding constants for the mercury species were chosen to  
390 provide the best simultaneous description of the experiments at pH 4.5 and 6.0, as well  
391 as the plots of Hg(II) uptake vs. pH and the influence of salinity. The values of  $K_1$ - $K_3$   
392 were first optimized by least-squares fit for each data set, and then average values (see  
393 Table 1) were used to plot the model calculations shown in Figure 3 (a), Figure 4 and  
394 Figure 5 (a,b). The isotherm obtained at pH 8.0 was excluded from the model discussion,  
395 since, on the one hand, it does not display the typical Langmuir shape (probably due to  
396 the reasons explained above) and, on the other hand, the maximum uptake values  
397 obtained exceed the total number of acid groups, which can not be explained by the  
398 present complexation model.

399 It is clear from Figure 4 that the 1:1 model is able to reproduce the shift in the  
400 isotherms from pH 4.5 to 6.0, although at low mercury concentrations (and high algal  
401 sites to mercury ratio) the uptakes are underestimated. On the contrary, the best fit to the  
402 data at pH 4.5 is provided by the 1:2 model, although the latter can not explain the  
403 uptake values above  $1.1 \text{ mmol}\cdot\text{g}^{-1}$  obtained at pH 6.0 (1:2 model plot for pH 6.0 not  
404 shown).

405 Figure 3 (b) shows the inorganic speciation of mercury predicted by MINEQL+,  
406 together with the estimated fraction of ionized acid sites on the alga, determined from a  
407 simple Langmuir equation and a more accurate Langmuir-Freundlich model, both  
408 involving an average proton binding constant of  $\log K_H = 5$  (corresponding to the  
409 estimated average ionic strength of the experiments). It is noticed that (i)  $\text{HgCl}_2$  is the

410 predominant species below pH 6.0; (ii) the fraction of chloride complex remains almost  
411 constant throughout this range, and yet the Hg(II) uptake increases in a remarkable way,  
412 thus confirming the postulate that metal speciation is not the only factor in the pH  
413 dependence of the Hg(II) uptake; (iii) the observed amount of mercury bound (Figure  
414 3a) varies in parallel with the fraction of dissociated sites (Figure 3b), in agreement with  
415 the assumption that these sites are involved in the metal binding; (iv) the hydroxyl  
416 mercury complexes are only relevant at the highest pH values, although the Hg(OH)Cl  
417 seems to be less important.

418         The 1:1 and 1:2 model descriptions of the pH dependence are shown in the  
419 Figure 3 (a). Note that both models reflect the increasing trend of the experimental data,  
420 which is explained as a combination of two factors, namely the growing fraction of  
421 dissociated algal sites and the formation of hydroxyl complexes of mercury, with higher  
422 binding affinity. It can be observed that both models underestimate the results below pH  
423 4.0, where the fraction of dissociated sites predicted by a discrete acid-base constant is  
424 very small. This can be a consequence of neglecting the heterogenous nature of the  
425 sorbent with regard to the proton binding. In fact, if the proton binding is represented by  
426 the Langmuir-Freundlich isotherm (which assumes a continuous distribution of  
427 affinities) then it turns out that a significant fraction of acid sites are already dissociated  
428 below pH 4 (see dotted line in Figure 3 b). Therefore, this effect could explain the  
429 observed uptakes at low pH. However, the more simple discrete constant description  
430 was preferred in this work, with the aim of using the lowest number of empirical  
431 parameters.

432         The discrepancy between the calculated uptakes and the experimental values at  
433 low pH may also be attributed to the contribution of a constant (although small)  
434 “background” accumulation of Hg due to the solubilization of neutral species in the

435 phospholipid environment of the algal cells. Mason *et al.* [Mason et al. (1995)] showed  
436 that neutral inorganic complexes of Hg(II) are fairly hydrophobic compounds, with  
437 octanol/water partition coefficients following the trend  $\text{HgCl}_2 > \text{Hg}(\text{OH})\text{Cl} > \text{Hg}(\text{OH})_2$ .  
438 However, this dependence of the lipid solubility on the inorganic speciation of Hg does  
439 not agree with the experimental variation in mercury uptake with pH, which is just the  
440 opposite, i.e., the amount of metal sorbed becomes larger as the fraction of the less  
441 lipophilic species increases. Therefore, the mechanism of lipid solubilization alone is  
442 not enough to explain the experimental results.

443 Since the model assumes the binding of a neutral species to an acid site of the  
444 biomass, it is expected that a change in ionic strength would only influence the  
445 inorganic metal speciation in solution and the dissociation state of the algal sites, but not  
446 the binding reaction, eq. ( 3 ). Therefore, the observed decrease in Hg(II) sorption with  
447 the concentration of added NaCl is ascribed to the formation of  $\text{HgCl}_3^-$  and  $\text{HgCl}_4^{2-}$   
448 complexes, which are favored by high chloride concentrations. In fact, the uptake values  
449 correlate well with the fraction of  $\text{Hg}(\text{Cl})_2$  (the main neutral species at pH 6.0) estimated  
450 through MINEQL+ in the filtrate solutions (Figure 5 c). The model predictions for the  
451 uptake vs. added NaCl data are shown in Figure 5 (a). Note that the 1:1 model agrees  
452 well with the observed trend, whereas the 1:2 model overestimates the experimental  
453 uptakes, although it is still able to reproduce the correct trend.

454 In contrast with these results, the experiments performed in  $\text{KNO}_3$  and  $\text{NaNO}_3$   
455 (Figure 5 b) show a slight increase in the uptake with background salt concentration. In  
456 this case, the background salt anions do not tend to form strong mercury complexes of  
457 any kind. Recall that if the sorption mechanism implicated the binding of  $\text{M}^{2+}$  ions and  
458 ionized surface sites, then the electrostatic effect of the supporting electrolyte would  
459 cause a sharp reduction in the amount of metal bound with ionic strength. On the other

460 hand, in a model involving complexation of neutral species to ionized acid sites, the  
461 factors that may explain the influence of nitrate concentration are the minor changes in  
462 mercury speciation (which are already taken into account in MINEQL+ calculations)  
463 and the decrease in  $K_H$  with ionic strength (incorporated in the model through the  
464 Donnan empirical expression). The latter means that an increasing fraction of  
465 dissociated sites is formed. These effects seem to account for the trend observed in the  
466 experimental data, regardless the stoichiometry considered (Figure 5 b). In addition, the  
467 possible contribution of a “salting out” mechanism for the transfer of neutral Hg  
468 complexes from saline solution to the algal surface cannot be discarded either. In fact, a  
469 similar effect was proposed by Turner *et al.* [Turner et al. (2001)] in order to explain the  
470 increase in sediment/water partitioning coefficients with salinity observed for mercury  
471 in estuaries.

472

### 473 **Conclusions**

474 The results obtained in this study demonstrated that the macroalgae *Cystoseira*  
475 *baccata* could compete with commercial biosorbents for the removal of Hg(II) from  
476 wastewaters because of its low cost, among the several reasons studied in this paper  
477 summarized below.

478 The Hg(II) sorption kinetic is relatively fast, reaching equilibrium in 100  
479 minutes. The total number of weak acid groups determined by potentiometric titration  
480 was  $2.2 \pm 0.1 \text{ mmol g}^{-1}$ .

481 *C. baccata* shows a very high Hg(II) uptake capacity. It is able to remove  
482 percentages greater than 80% of mercury in solution for relatively high initial mercury  
483 concentrations ( $120 \text{ mg}\cdot\text{L}^{-1}$  at pH 4.5), using small amounts of biosorbent.

484           The studies of the salinity influence showed that an increase in concentration of  
485 nitrate salts (NaNO<sub>3</sub>, KNO<sub>3</sub>) causes greater mercury uptake, while a larger presence of  
486 NaCl salt leads to a drop in the sorption, which is decreased by more than 80% at salt  
487 concentrations around 1 M, whereas low values of salinity have small influence on the  
488 metal uptake. Moreover, the addition of different divalent cations to the mercury  
489 solution, namely Ca<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, Cd<sup>2+</sup>, Pb<sup>2+</sup> and Cu<sup>2+</sup>, reveals that their effect on the  
490 uptake process is negligible.

491           The simple competitive model proposed in the present study is able to reflect the  
492 major characteristics of the experimental Hg(II) sorption data, with regard to  
493 equilibrium mercury concentration, pH and presence of background salts in solution.  
494 However, the stoichiometry of the mercury complexes cannot be clearly assessed. On  
495 the one hand, the 1:2 isotherm is able to fit very well the experimental data at low ratios  
496 of metal to algal acid sites, on the other hand, the maximum mercury loadings at a  
497 relatively high pH can only be explained by a 1:1 relationship. The actual binding  
498 mechanism would very probably involve a combination of both mono and bidentate  
499 sites, or even some degree of interaction between mercury and non-ionized sites of the  
500 algal surface.

501

## 502 **Acknowledgements**

503           This work was funded by the projects BQU2002-02133 (from the Ministerio de  
504 Ciencia y Tecnología of Spain) and PGDIT02TAM10302PR (from the Xunta de  
505 Galicia). The authors would like to thank Dr. I. Bárbara and Dr. J. Cremades (U. of A  
506 Coruña) for the collection and classification of the species.

507

508

508 FIGURE CAPTIONS

509 **Figure 1**

510 Sorption of Hg(II) on *C. baccata* vs. time. Biomass concentration:  $2.5 \text{ g}\cdot\text{L}^{-1}$ ;  
511 initial  $\text{HgCl}_2$  concentration:  $500 \text{ mg}\cdot\text{L}^{-1}$ ; saline medium:  $0.05\text{M NaNO}_3$ . The final pH at  
512 equilibrium was around 4.

513 **Figure 2**

514 Acid-base potentiometric titration of an acid-treated biomass sample of *C.*  
515 *baccata* ( $5 \text{ g}\cdot\text{L}^{-1}$ ) in  $0.05 \text{ M NaNO}_3$  at  $25^\circ\text{C}$ . Symbols: pH vs. titrant volume; solid line:  
516 first derivative (in arbitrary units).

517 **Figure 3**

518 Influence of pH on Hg(II) uptake. (a): experimental mercury uptake values  
519 (symbols) and 1:1 (solid line) or 1:2 (dashed line) model estimates, calculated through  
520 eqs. ( 7 ) and ( 8 ), respectively, using the parameters listed in Table 1 ( $r^2= 0.92$  in both  
521 cases). Biomass concentration:  $2.5 \text{ g}\cdot\text{L}^{-1}$ . (b): inorganic speciation of mercury in the  
522 aqueous phase at equilibrium with the biomass, calculated by MINEQL+ (solid lines),  
523 and fraction of dissociated acid sites on the alga, calculated from Langmuir (dashed  
524 line) and Langmuir-Freundlich (dotted line) isotherms.

525 **Figure 4**

526 Hg(II) uptake by *C. baccata* as a function of the aqueous Hg(II) equilibrium  
527 concentration at pH 4.5 (circles), 6.0 (triangles) and 8.0 (diamonds). Biomass  
528 concentration:  $2.5 \text{ g}\cdot\text{L}^{-1}$ . Each symbol represents the average of two replicate batch  
529 sorption experiments performed with the same initial mercury concentration. Lines  
530 represent the uptake values calculated through the 1:1 (solid line) and 1:2 (dashed line)  
531 complexation models, using the parameter values listed in Table 1.

532


533 **Figure 5**

534 Sorption of mercury (initial metal concentration:  $500 \text{ mg}\cdot\text{L}^{-1}$ ; biomass  
535 concentration:  $2.5 \text{ g}\cdot\text{L}^{-1}$ ) as a function of the concentration of added (a) NaCl and (b)  
536  $\text{KNO}_3$  and  $\text{NaNO}_3$  at pH 6.0. Lines represent the expected uptake values estimated from  
537 the 1:1 (solid line) and 1:2 (dashed line) competitive complexation models using the  
538 parameters listed in Table 1. The reference value 100% indicates the mercury sorption  
539 in absence of added salt. (c): correlation between the metal uptake of the experiments  
540 shown in Figure (a) and the fraction of the total Hg(II) present in aqueous phase as the  
541 neutral complex  $\text{HgCl}_2$ . The numbers over each experimental point correspond to the  
542 amounts of added NaCl, in  $\text{mol}\cdot\text{L}^{-1}$ . The error bars represent the difference between two  
543 replicates of the same experiment.

544 **Figure 6**

545 Effect of the presence of several divalent cations (concentrations shown in the  
546 legend) on the Hg(II) uptake by *C. baccata* at pH 4.5. Initial mercury concentration:  $50$   
547  $\text{mg}\cdot\text{L}^{-1}$ ; biomass concentration:  $2.5 \text{ g}\cdot\text{L}^{-1}$ . The reference value 100% indicates the  
548 mercury sorption in absence of competing cations.


549



550

551

Figure 1




552

553

Figure 2




(a)



554

(b)



555

556

Figure 3




Figure 4

557

558

559

560

561

562

563

564

565

566

567

568

569


570

571

572


573

(a)




574

(b)



575

(c)




576

577

Figure 5

578



579

580

**Figure 6**

581

582

583

584

585

586

587

588

589

590

591

592

593

594

## 595 TABLES

596 **Table 1**

597 Parameters of the 1:1 and 1:2 competitive Langmuir models for the binding of  
 598 neutral Hg(II) species, eqs. ( 7 ) and ( 8 ). <sup>a</sup> Determined from the equivalence point of  
 599 the potentiometric titrations; <sup>b</sup> maximum and minimum values in the range 0.004 to 1.0  
 600 M ionic strength, estimated through an empirical Donnan expression [Rey-Castro et al.  
 601 (2004b)] using data from this and previous works [Rey-Castro et al. (2003)]; <sup>c</sup> Average  
 602 values of the metal binding constants optimized by least squares fit for each set of  
 603 experiments. These parameters are assumed to remain constant with ionic strength.  
 604 Errors are shown in brackets.

605

Model	Parameters				
	[A] <sub>T</sub> (mmol·g <sup>-1</sup> ) <sup>a</sup>	logK <sub>H</sub> <sup>b</sup>	logK <sub>1</sub> <sup>c</sup>	logK <sub>2</sub> <sup>c</sup>	logK <sub>3</sub> <sup>c</sup>
<b>1:1</b>	2.2 (0.1)	5.0 - 3.2	2.8 (0.1)	4.0 (0.1)	4.0 (0.1)
<b>1:2</b>	2.2 (0.1)	5.0 - 3.2	4.0 (0.1)	4.5 (0.5)	5.5 (0.2)

606

607

608

609

610

611

612

613

614 **References**

- 615 Allison J. D., Brown D. S. and Novo-Gradac K. J. (1991) *MINTEQA2/PRODEFA2, A*  
616 *geochemical assessment model for environmental systems: version 3.0 user's manual*;  
617 U.S. Environmental Protection Agency: Washington, DC.
- 618 Bailey S. E., Olin T. J., Bricka R. M. and Adrian D. D. (1999) A review of potentially  
619 low-cost sorbents for heavy metals. *Water Research* **33** (11). 2469-2479.
- 620 Bello G., Cid R., García R. and Arriagada R. (1999) Retention of Cr(VI) and Hg(II) in  
621 Eucalyptus globulus- and peach stone-activated carbons. *Journal of Chemical*  
622 *Technology and Biotechnology* **74** 904-910.
- 623 Brandariz I., Vilariño T., Alonso P., Herrero R., Fiol S. and Sastre de Vicente M. E.  
624 (1998) Effect of ionic strength on the formal potential of the glass electrode in various  
625 saline media. *Talanta* **46** 1469-1477.
- 626 Budinova T., Savova D., Petrov N., Razvigorova M., Minkova V., Ciliz N., Apak E.  
627 and Ekinçi E. (2003) Mercury adsorption by different modifications of furfural  
628 adsorbent. *Industrial & Engineering Chemistry Research* **42** (10). 2223-2229.
- 629 Chiarle S., Ratto M. and Rovatti M. (2000) Mercury removal from water by ion  
630 exchange resins adsorption. *Water Research* **34** (11). 2971-2978.
- 631 Cordero B., Lodeiro P., Herrero R. and Sastre de Vicente M. E. (2004) Biosorption of  
632 cadmium by *Fucus spiralis*. *Environmental Chemistry* **1** 180-187.
- 633 Cox M., El-Shafey E. I., Pichugin A. A. and Appleton Q. (2000) Removal of  
634 mercury(II) from aqueous solution on a carbonaceous sorbent prepared from flax shive.  
635 *Journal of Chemical Technology and Biotechnology* **75** 427-435.
- 636 Crist R. H., Oberholser K., Schwartz D., Marzoff J., Ryder D. and Crist D. R. (1988)  
637 Interactions of metals and protons with algae. *Environmental Science & Technology* **22**  
638 (7). 755-760.

639 Cruz C. C. V., Costa A. C. A., Henriques C. A. and Luna A. S. (2004) Kinetic modeling  
640 and equilibrium studies during cadmium biosorption by dead *Sargassum* sp biomass.  
641 *Bioresource Technology* **91** 249-257.

642 Daughney C. J., Siciliano S. D., Rencz A. N., Lean D. and Fortin D. (2002) Hg(II)  
643 adsorption by bacteria: a surface complexation model and its application to shallow  
644 acidic lakes and wetlands in Kejimikujik national park, Nova Scotia, Canada.  
645 *Environmental Science & Technology* **36** (7). 1546-1553.

646 Drexel R. T., Haitzer M., Ryan J. N., Aiken G. R. and Nagy K. L. (2002) Mercury(II)  
647 sorption to two Florida Everglades peats: evidence for strong and weak binding and  
648 competition by dissolved organic matter released from the peat. *Environmental Science*  
649 *& Technology* **36** 4058-4064.

650 Fiol S., Arce F., Armesto X. L., Penedo F. and Sastre de Vicente M. E. (1992) Analysis  
651 of systematic errors in calibrating glass electrodes with H<sup>+</sup> as a concentration probe.  
652 *Fresenius Journal of Analytical Chemistry* **343** 469-472.

653 Gómez-Serrano V., Macías-García A., Espinosa-Mansilla A. and Valenzuela-Calahorro  
654 C. (1998) Adsorption of mercury, cadmium and lead from aqueous solution on heat-  
655 treated and sulphurized activated carbon. *Water Research* **32** (1). 1-4.

656 Jeon C. and Höll W. H. (2003) Chemical modification of chitosan an equilibrium study  
657 for mercury ion removal. *Water Research* **37** 4770-4780.

658 Lacher C. and Smith R. W. (2002) Sorption of Hg(II) by *Potamogeton natans* dead  
659 biomass. *Minerals Engineering* **15** 187-191.

660 Lodeiro P., Cordero B., Barriada J. L., Herrero R. and Sastre de Vicente M. E. (2004a)  
661 Biosorption of cadmium by biomass of brown marine macroalgae. *Bioresource*  
662 *Technology* **in press**.

663 Lodeiro P., Cordero B., Grille Z., Herrero R. and Sastre de Vicente M. E. (2004b)  
664 Physicochemical studies of Cadmium (II) biosorption by the invasive alga in Europe,  
665 *Sargassum muticum*. *Biotechnology and Bioengineering* **88** (2). 237-247.

666 Mason R. P., Reinfelder J. R. and Morel F. M. M. (1995) Bioaccumulation of mercury  
667 and methylmercury. *Water, Air and Soil Pollution* **80** (1-4). 915-921.

668 Masri M. S., Reuter F. W. and Friedman M. (1974) Binding of metal cations by natural  
669 substances. *Journal of Applied Polymer Science* **18** (3). 675-681.

670 McKay G., Blair H. S. and Findon A. (1989) Equilibrium studies for the sorption of  
671 metal ions onto chitosan. *Indian Journal of Chemistry* **28A** (5). 356-360.

672 Percival E. and McDowell R. H. (1967) *Chemistry and enzymology of marine algal*  
673 *polysaccharides*; Academic Press: London New York; pp. xii, 219.

674 Rey-Castro C., Herrero R. and Sastre de Vicente M. E. (2004a) Gibbs-Donnan and  
675 specific ion interaction theory descriptions of the effect of ionic strength on proton  
676 dissociation of alginic acid. *Journal of Electroanalytical Chemistry* **564** 223-230.

677 Rey-Castro C., Herrero R. and Sastre de Vicente M. E. (2004b) Surface charge and  
678 permeable gel descriptions of the ionic strength influence on proton binding to seaweed  
679 biomass. *Chemical Speciation and Bioavailability* **16** (1-2). 61-69.

680 Rey-Castro C., Lodeiro P., Herrero R. and Sastre de Vicente M. E. (2003) Acid-base  
681 properties of brown seaweed biomass considered as a Donnan Gel. A model reflecting  
682 electrostatic effects and chemical heterogeneity. *Environmental Science & Technology*  
683 **37** (22). 5159-5167.

684 Saglam N., Say R., Denizli A., Patir S. and Arica M. Y. (1999) Biosorption of inorganic  
685 mercury and alkylmercury species on to *Phanerochaete chrysosporium* mycelium.  
686 *Process Biochemistry* **34** 725-730.



687 Sarkar D., Essington M. E. and Misra K. C. (1999) Adsorption of mercury (II) by  
688 variable charge surfaces of quartz and gibbsite. *Soil Science Society of America Journal*  
689 **63** 1626-1636.

690 Sastre de Vicente M. E. (1997) Ionic strength effects on acid-base equilibria. A review.  
691 *Current Topics in Solution Chemistry* **2** 157-181.

692 Schecher W. D. and McAvoy D. C. (1992) MINEQL+: A software environment for  
693 chemical equilibrium modeling. *Computers, Environment and Urban Systems* **16** (1).  
694 65-76.

695 Schiewer S. and Wong M. H. (1999) Metal binding stoichiometry and isotherm choice  
696 in biosorption. *Environmental Science & Technology* **33** 3821-3828.

697 Schiewer S. and Wong M. H. (2000) Ionic strength effects in biosorption of metals by  
698 marine algae. *Chemosphere* **41** 271-282.

699 Schneider I. A. H., Rubio J. and Smith R. W. (2001) Biosorption of metals onto plant  
700 biomass: exchange adsorption or surface precipitation? *International Journal of Mineral*  
701 *Processes* **62** 111-120.

702 Turner A., Millward G. E. and Le Roux S. M. (2001) Sediment-water partitioning of  
703 inorganic mercury in estuaries. *Environmental Science & Technology* **35** 4648-4654.

704 Vázquez G., González-Álvarez J., Freire S., López-Lorenzo M. and Antorrena G.  
705 (2002) Removal of cadmium and mercury ions from aqueous solution by sorption on  
706 treated *Pinus pinaster* bark: kinetics and isotherms. *Bioresource Technology* **82** 247-  
707 251.

708 Volesky B. (2003) *Sorption and biosorption*; BV Sorbex: St. Lambert, Quebec; pp. xii,  
709 316.

710 Wase J. and Forster C. F. (1997) *Biosorbents for metal ions*; Taylor & Francis: London;  
711 pp. x, 238.

712 Yardin F., Budinova T., Ekinici E., Petrov N., Razvigorova M. and Minkova V. (2003)  
713 Removal of mercury (II) from aqueous solution by activated carbon obtained from  
714 furfural. *Chemosphere* **52** 835-841.

715 Zeroual Y., Moutaouakkil A., Dzairi F. Z., Talbi M., Chung P. U., Lee K. and Blaghen  
716 M. (2003) Biosorption of mercury from aqueous solution by *Ulva lactuca* biomass.  
717 *Bioresource Technology* **90** 349-351.

718