

Cadernos Lab. Xeolóxico de Laxe  
Coruña. 2000. Vol. 25, pp. 147-149

# **A Systematic typology of granitoid rocks from major element composition. I: The Upper Silica Range**

## **Tipología sistemática de granitoides a partir de su composición en elementos mayores. I: El rango más alto de Si**

F. BEA (1); L.G. CORRETGÉ (2); G.B. FERSHTATER (3).

(1) Department of Mineralogy and Petrology. University of Granada. Campus Fuentenueva.  
18002 Granada, Spain.

(2) Department of Geology. University of Oviedo. 33005 Oviedo, Spain.

(3) Institute of Geology and Geochemistry of the Acad. Sciences of Russia; Pochtovi per. 7, Ekaterinburg,  
620151, Russia.

The composition of granitic rocks is a complex function of many parameters, among which the nature of source materials, P-T-Xvolatiles conditions of partial melting, and the efficiency of melt segregation are of remarkable importance. All these factors, in turn, are closely related to the geotectonic environment where granites originated. The recognition of such relationship gave a great impulse to the understanding of granitic rocks variety (Pitcher, 1982), and is the basis of granite typology as it is currently understood (Pitcher, 1979; Pitcher, 1982), which establishes the following fundamental granite typologies: I- and S- types (Chappell and White, 1974, A-type (Loiselle and Wones, 1979) and M-type (Pitcher, 1983). This S-I-A-M scheme reflects in a simple way broad genetical concepts concerning the source rocks in spite of the, sometimes troublesome, indetermination in their characteristic and definition (Miller, 1985; Clarke, 1992). The classification focuses as well attention on granite types, tectonic setting, source mineralogy and sometimes, metallogenic specialization regardless of whether or not the ore specialization of granites depends upon other criteria (Hannah and Stein, 1990).

This paper is the first result of a worldwide survey on granite chemistry undertaken in order to recognize the different compositional types of granites of the S- I- A- M- types on a statistical basis. It focuses on the silicic margin of the granitoids compositional spectrum. A data set of 4,200 samples with  $\text{SiO}_2 > 70$  wt. % from all over the world has been worked out. Only major elements from this data set

were used in the classification. We tested three methods of distinguishing between granites: The raw major elements, their normative components and various petrochemical parameters. The best results have been obtained by using four parameters: (a) Aluminium Saturation Index,  $\text{ASI} = \text{mol.} (\text{Al}_2\text{O}_3 / [\text{CaO} + \text{Na}_2\text{O} + \text{K}_2\text{O}])$ ; (b) Calc-Alkaline Index,  $\text{CAI} = \text{mol.} (\text{CaO} / [\text{CaO} + \text{Na}_2\text{O} + \text{K}_2\text{O}])$ ; (c) Potassium-Sodium Index,  $\text{KNaI} = \text{mol.} (\text{K}_2\text{O} / [\text{K}_2\text{O} + \text{Na}_2\text{O}])$ ; and (d) Iron-Magnesium Index,  $\text{FeMgI} = \text{mol.} (\text{FeO} / [\text{FeO} + \text{MgO}])$ . In order to reduce the dimensionality of the problem, we performed a principal components analysis on these four parameters. Scores of samples on the Principal Components are given by:

$$\text{FD1} = 10 * (0.1601 * \text{ASI} - 0.0354 * \text{CAI} + 0.048 * \text{KNaI} - 0.026 * \text{FeMgI}); [1]$$

$$\text{FD2} = 10 * (0.0061 * \text{ASI} - 0.0434 * \text{CAI} + 0.0145 * \text{KNaI} + 0.1236 * \text{FeMgI}); [2]$$

The first principal component is dominated by the Aluminium Saturation Index and accounts for almost 50 % of the total variance of the system. The second principal component is dominated by the Iron-Magnesium Index and accounts for nearly 30% of the total variance. The plane defined by both principal components is used as the main tool for classification. The plot of a subset of samples whose typology is well known allows one to split this area into five regions, in which to plot different granite types: (a) Primitive-types; (b) A-types; (c) I-types; (d) SC-types (low  $\text{Fe}/[\text{Fe} + \text{Mg}]$  S-types); (e) SG-types (high  $\text{Fe}/[\text{Fe} + \text{Mg}]$  S-types). Discrimination between primitive-types, is also possible by means of a  $\text{K}_2\text{O}$  vs  $\text{Na}_2\text{O}$  plot, in which fields for M-types, I-types, Magnitogorsk-type granites and ophiolitic plagiogranites are clearly identified.




Figura 1.

## BIBLIOGRAFÍA

- CLARKE, D. B. (1992). *Granitoid Rocks*. London, Chapman & Hall.
- CHAPPELL, B. W. AND WHITE, A. J. R. (1974). Two contrasting granite types. *Pac. Geology*, 8: 173-174.
- HANNAH, J. L. AND STEIN, H. J. (1990). Magmatic and hydrothermal processes in ore-bearing systems. Ore-bearing granite systems, petrogenesis and mineralizing processes. H. J. Stein and J. L. Stein. *Geological Society of America*, 246: 1-10.
- LOISELLE, M. C. AND WONES, D. R. (1979). Characteristics and origin of anorogenic granites. *Geol. Soc. Am. Abst. Prog.* 11: 468.
- MILLER, C. F. (1985). Are strongly peraluminous magmas derived from pelitic sources? *Journal of Geology*, 93: 673-689.
- PITCHER, W. S. (1979). The nature, ascent and emplacement of granite magmas. *J. Geol. Soc. Lond.* 136: 627-662.
- PITCHER, W. S. (1982). Granite types and tectonic environment. *Mountain Building Processes* (ed. K. Hsu). Academic Press, London: 19-40.
- PITCHER, W. S. (1983). Granite: typology, geological environment and melting relationships. *Migmatites, Melting and Metamorphism* (ed.: M.P. Atherton & C.D. Gribble). Shiva Pub. Ltd., Cheshire: 277-285.
- WHALEN, J. B., CURRIE, K. L. AND CHAPPELL, B. W. (1987). A-type granites: geochemical characteristics, discrimination and petrogenesis. *Contrib. Mineral. Petrol.* 95: 407-419.