

DEGREE IN BIOLOGY

FINAL PROJECT

Effect of commercial activity of fisheries on the genetic variability of the spiny spider crab (*Maja brachydactyla*) in the north-eastern Atlantic



Lucía Gato Calvo
2013



UNIVERSIDADE DA CORUÑA

FACULDADE DE CIENCIAS

UNIVERSITY OF A CORUÑA

DEPARTMENT OF CELL AND MOLECULAR BIOLOGY

Area of Genetics

DEPARTMENT OF ANIMAL BIOLOGY, PLANT BIOLOGY AND ECOLOGY

Area of Zoology

EFFECT OF COMMERCIAL ACTIVITY OF FISHERIES ON THE
GENETIC VARIABILITY OF THE SPINY SPIDER CRAB (*Maja
brachydactyla*) IN THE NORTH-EASTERN ATLANTIC

Lucía Gato Calvo

2013

TRABALLO FIN DE GRAO

Dña Ana M González Tizón y D Luis Fernández Rodríguez autorizan a presentación do traballo de fin de grao “**Effect of comercial activity of fisheries on the genetic variability of the spiny spider crab (*Maja brachydactyla*) in the north-eastern Atlantic**” presentado por **Lucía Gato Calvo** para a súa defensa ante o tribunal cualificador.

En A Coruña, a 14 Junio de 2013

Asdo: Ana M González Tizón

Luis Rodríguez Fernández

This project is included in a collaborative project with the Department of Cell and Molecular Biology (area of Genetics) and the Department of Animal Biology, Plant Biology and Ecology (area of Zoology), which has been developed as a result of the concession of a Collaboration Grant at University Departments during the Academic Year 2012-2013 by the Ministry of Education, Culture and Sport.

INDEX

1. INTRODUCTION.....	p. 2
2. OBJECTIVES.....	p. 5
3. MATERIALS AND METHODS.....	p. 6
3.1. Sampling.....	p. 6
3.2. DNA extraction.....	p. 7
3.3. Quantification and quality assessment of the obtained DNA.....	p. 8
3.4. PCR amplification.....	p. 8
3.5. Sequencing and analysis of the obtained data.....	p. 11
4. RESULTS.....	p. 14
4.1. PCR amplification.....	p. 14
4.2. Analysis of the obtained sequences.....	p. 14
5. DISCUSSION.....	p. 17
6. REFERENCES.....	p. 19

1.- INTRODUCTION


The spiny spider crab *Maja brachydactyla* Balss, 1922, is a decapod crustacean which belongs to the family Majidae (Ng *et al.*, 2008). It can be found in the north-eastern Atlantic, from the British Isles and Ireland to the coasts of Senegal (*Picture 1*), reaching the archipelagos of Azores, Madeira, Canary Islands and Cape Verde (Sotelo *et al.*, 2008a). It was initially classified as *Maja squinado* Herbst, 1788, but it has recently been clarified that they are in fact two different species; morphological characteristics were used at first (Neumann, 1998) and then the difference was confirmed on the basis of molecular characteristics (Sotelo *et al.*, 2008a).



Picture 1. Spatial distribution of the spiny spider crab *Maja brachydactyla*

The life cycle of *Maja brachydactyla* (*Picture 2*) can be divided into three main stages: larval stage, juvenile stage and adult or reproductive stage (Corgos, 2004). Larvae are released in coastal areas, where they become juveniles in two to three weeks. During this period of time, they go through three larval stages separated by moults (two zoea stages and one megalopa stage). Once settlement has occurred, at the end of planktonic stage, juveniles remain in shallow waters (<15m) for two or three years (depending on the moment of the year when settlement takes place), until they undergo the terminal moult and reach sexual maturity to become adults. This usually happens at the end of summer or the beginning of autumn. Recent studies suggest that juvenile individuals frequently migrate between rocky and sandy bottoms, basically at night, in order to get food and refuges, respectively.

Once they have reached adulthood, spider crabs migrate to deeper waters (30-100m) to reproduce. Reproductive migrations begin in the autumn and ovigerous females return to shallow waters during the following spring to incubate their first spawning of the annual cycle. Incubation period lasts between 30 and 74 days, varying with water temperature, and sperm storage in spermathecae allows females to fertilize as many as three broods without mating in between (Corgos, 2004; González-Gurriarán *et al.*, 1998). Fertility is really high and each spawning can consist in more than 500.000 eggs in the largest females. As well, adults can live from two to three years, and that is the reason why spiny spider crab populations consist of several overlapping cohorts (Sotelo *et al.*, 2008b).



Picture 2. Life cycle of the spiny spider crab *Maja brachydactyla*

Spiny spider crab is an economically important fishing resource and it suffers an intensive exploitation pressure, especially in France and the United Kingdom (Freire *et al.*, 2002). In Galicia (north-west of Spain), a 90% of the new production of the population is annually extracted so the spiny spider crab has been classified as an overexploited resource. Nevertheless, genetic studies which have been carried out up to now do not show evidences of overexploitation. On the other hand, they do indicate the need to increase sampling sizes and include a higher number of sampling localities (Sotelo *et al.*, 2008b).

The intensive fishing activity for this resource could affect the genetic structure of its natural populations in two distinct ways:

1. Extraction and reduction of local genetic variants, which have associated genetic drift effects.
2. Export of spiny spider crabs to markets placed in other geographical localities, something which represents an artificial mechanism of genetic flow. This phenomenon could be occurring in Galicia as a consequence of the import and temporary storage of

spawning females in fisheries until sale. Eggs in an advanced developmental stage could hatch in the fishery and larvae could join local populations.

This study aims to contribute novel information on the possible homogenizing effect of the activity of fisheries on the genetic variability of population groups from different geographical localities.

2.- OBJECTIVES

This project has the objective of studying the effect of commercial activity on the genetic variability of the spiny spider crab *Maja brachydactyla* Balss, 1922, by means of molecular analysis of the mitochondrial genes for cytochrome c oxidase subunit I (COI) and 16S ribosomal subunit (16S).


3.- MATERIALS AND METHODS

3.1.- Sampling

Forty individuals from 3 localities in the north-eastern Atlantic region were analysed (*Picture 3*). Twenty five of them were from Lorient (France, $47^{\circ}44'45''\text{N}$ $3^{\circ}21'59''\text{W}$), 13 were from Saint-Malo (France, $48^{\circ}38'50''\text{N}$ $2^{\circ}00'32''\text{W}$) and 2 were from Cedeira (Spain, $43^{\circ}39'00''\text{N}$ $8^{\circ}03'00''\text{W}$). A pereopod or a pleopod (*Picture 4*) was obtained from each one and preserved in absolute ethanol until processing.



Picture 3. Sampling localities



Picture 4. Anatomy of decapod crustaceans. Pereiopods and pleopods are highlighted. Images were taken and modified from www.arthursclipart.org and www.pictospin.com

3.2.- DNA extraction

Genomic DNA of the individuals was extracted from pereiopods muscle tissue (with the exception of the individuals from Cedeira, in which the extraction was performed on pleopods) using RealPure extraction kit (REAL). Approximately 20mg of muscle tissue were dissected from each sample. They were added 600 μ l of lysis solution and were mechanically homogenized before the addition of 3 μ l of K proteinase in order to increase the surface exposed to enzymatic activity. After being inversion-mixed, samples were incubated overnight at 55°C with mechanical shaking.

Once cooled at room temperature, lysates were added 3 μ l of RNase, mixed by inversion and incubated at 37°C during 30min. Afterwards, 360 μ l of protein precipitation solution were added to the samples, which were also vigorously mixed by vortex and centrifuged at 14000g during 8min. This treatment resulted in the precipitation of a thick protein pellet at the bottom of the Eppendorf tubes.

While samples were being centrifuged, new Eppendorf tubes were disposed with 600 μ l of isopropanol, in which the supernatants containing the DNA were then poured. Later, samples were inversion-mixed and centrifuged again at 14000g during 3min. As a result, DNA formed a

thin pellet at the bottom of the tubes. Supernatants were discarded and DNA pellets were washed with 600µl of 70% ethanol. Then, samples were centrifuged at 14000g during 2min and ethanol was carefully removed. Tubes were inverted on clean absorbent paper and allowed to air-dry. Finally, DNA was re-suspended in 100µl of hydration solution and it was incubated overnight at room temperature to favour re-hydration.

3.3.- Quantification and quality assessment of the obtained DNA

The success of the DNA extraction procedure was assessed by means of a 1% agarose gel electrophoresis stained with RedSafe (Chembio). Once checking was done, the DNA quantity obtained in each case and its quality were determined with NanoDrop 1000 (Thermo scientific). Dilutions adjusted at 10ng/µl of concentration were then prepared (in the cases in which this was necessary) and stored at 4°C until they were used in the following molecular analysis. Original DNA extractions were preserved as a stock for future analysis at -80°C.

3.4.- PCR amplification

Amplification of 16S and COI mitochondrial genes was started as soon as genomic DNA samples were available. Primers 16Sar and 16Sbr employed in the studies of Sotelo *et al.* (2008a, b) and Palumbi *et al.* (1991) were used at first to amplify 16S gene, whereas LCO1490 and HCO2198 primers used in the studies of Sotelo *et al.* (2008a, b; 2009a, b) and Folmer *et al.* (1994) were used to amplify COI gene (*Table 1*).

Table 1. Sequences of the initial primers for the amplification of 16S and COI genes

Forward primers		Reverse primers	
Name	Sequence (5'→3')	Name	Sequence (5'→3')
16Sar	CGC CTG TTT ATC AAA AAC AT	16Sbr	CCG GTC TGA ACT CAG ATC ACG T
LCO1490	GGT CAA CAA ATC ATA AAG ATA TTG G	HCO2198	TAA ACT TCA GGG TGA CCA AAA AAT CA

A temperature gradient from 43°C to 54°C and two magnesium concentrations (1.5mM and 3mM) were tested to determine the best amplification conditions for each gene. Reactions were carried out in a final volume of 25µl, containing 25ng of DNA, PCR buffer (10mM HCl, 1.5mM MgCl₂, 50mM KCl, pH 8.3), dNTP mix (200µM), 0.5µM of each primer and 0.25U of BIOTAQ polymerase (Roche Applied Science). To test the 3mM magnesium concentration, 3.57µg of MgCl₂ were additionally added from a 25mM stock solution. Thermocycler (MyCycler Thermal Cycler, Bio-Rad Laboratories) was programmed as follows: 5min at 95°C; 35 cycles of 30s at 95°C, 30s at 43°C-54°C and 45s at 72°C; 5min at 72°C and 30min at 4°C (*Picture 5*). PCR products were visualized by being run in 1% agarose gels stained with RedSafe.

95°C 5min
35 cycles of:
95°C 30s
43°C-54°C 30s
72°C 45s
72°C 5min
4°C

Picture 5. Initial PCR profile for the amplification of 16S and COI genes

Another pair of primers employed by Sotelo *et al.* (2009a, b) was decided to be tested in the cases in which 16S gene amplification was not successful: 16L29 and 16HLeu (Schubart, 2009). Amplification conditions were the same as in the initial experiment; that is, a temperature gradient from 43°C to 54°C and magnesium concentrations of 1.5mM and 3mM were tested with the described PCR profile. PCR products were visualized in a 1% agarose gel electrophoresis and then they were sequenced. In the cases in which there was no amplification success, it was decided to request for 16L29 and 16HLeu primers again, but modified: M13 tails were added with the purpose of obtaining a proper sequence reading (Table 2).

Table 2. Sequences of the primers for the amplification of 16S gene

Forward primer		Reverse primer	
Name	Sequence (5'→3')	Name	Sequence (5'→3')
16L29M13	GTA AAA CGA CGG CCA GTY GCC TGT TTA TCA AAA AC	16HLeuM13	CAG GAA ACA GCT ATG ACC ATA TTA TCT GCC AAA AT

Furthermore, a touchdown PCR was programmed as to improve the specificity of the amplification reaction: 5min at 94°C; 5 cycles of 30s at 94°C, 90s at 45°C and 60s at 72°C; 5 cycles of 30s at 94°C, 90s at 65°C-60°C (with a decrease in temperature of 1°C per cycle) and 60s at 72°C; a final extension of 5min at 72°C and a cooling step at 4°C (Picture 6). Just as in the previous cases, PCR products were run in 1% agarose gels stained with RedSafe and they were sequenced when successfully amplified. Given the good performance which was obtained with this protocol, it was used in all the following amplifications of 16S gene.

94°C 5min	
5 cycles of:	94°C 30s 45°C 90s 72°C 60s
5 cycles of:	94°C 30s 65°C-60°C 90s (↓1°C/cycle) 72°C 60s
30 cycles of:	94°C 30s 60°C 90s 72°C 60s
72°C 5min	
4°C	

Picture 6. Touchdown PCR designed for the amplification of 16S gene with 16L29M13 and 16HLeuM13 primers

On the other hand, COI gene PCR products for which it was no possible to obtain the sequence were tried to be cloned by using the kit StrataClone PCR Cloning Kit (Agilent Technologies): 1.5µl of cloning buffer StrataClone were mixed with 1µl of PCR product and 0.5µl of vector mix StrataClone and the ligation reaction mixture was incubated at room temperature during 5min. Meanwhile, a tube of competent cells StrataClone SoloPack was thawed on ice. The ligation reaction was placed on ice when completed and 1µl of it was then mixed with half of the competent cells (25µl) without repeated pipetting. The resultant transformation mixture

was incubated on ice during 20min. Over this period of time, liquid LB medium was pre-warmed at 42°C (it had previously been prepared with 10g of NaCl, 10g of tryptone, 5g of yeast extract and distilled water in a final volume of 1 litre; pH had been adjusted to 7.0 with 5N NaOH and the mixture had been autoclave-sterilized). Once the incubation was finished, cells were heat-shocked at 42°C during 45s and they were then incubated on ice during 2min. Eventually, 250µl of pre-warmed LB medium were added to the transformation reaction mixture and 90µl of the resultant mixture were then plated on LB-ampicillin plates (they had previously been prepared following the same steps as with liquid LB medium, but adding 20g of agar and 100g ampicillin, too), over which it had previously been spread 40µl of 2% Xgal. Afterwards, these plates were incubated overnight at 37°C.

Later, white-coloured colonies were selected (they should contain a PCR product insert) and cell lines were plated on a new LB-ampicillin plate, which was subsequently incubated overnight at 37°C. Finally, several colonies of each cell line were randomly selected and they were individually placed into PCR tubes containing 10µl of sterile MilliQ water to induce cell lysis and then assess cloning success through PCR. Each tube was added PCR buffer (10mM Tris-HCl, 1.5mM MgCl₂, 50mM KCl, pH 8.3), dNTP mix (200mM), 0.8µM of each primer (M13F and M13R) and 0.25U of BIOTAQ polymerase (Roche Applied Science) in a final volume of 25µl. PCR was programmed as follows: 5min at 94°C; 35 cycles of 1min at 94°C, 1min at 55°C and 1min at 72°C; a final extension of 5min at 72°C and a cooling step at 4°C (*Picture 7*). PCR products were visualized in a RedSafe-stained 1% agarose gel electrophoresis.

94°C 5min
35 cycles of: 94°C 60s
55°C 60s
72°C 60s
72°C 5min
4°C

Picture 7. PCR profile for the amplification of the inserts

A literature search on PCR primers used up to now to amplify fragments of COI mitochondrial gene in decapod crustaceans was performed at last. The suitability of four primer pairs (*Table 3*) was decided to be tested.

Table 3. Sequences of the tested primer pairs for the amplification of COI gene

Forward primers		Reverse primers		References
Name	Sequence (5'→3')	Name	Sequence (5'→3')	
LCO1490M13	GTA AAA CGA CGG CCA GTG GTC AAC AAA TCA TAA AGA TAT TGG	HCO2198M13	GTA AAA CGA CGG CCA GTG GTC AAC AAA TCA TAA AGA TAT TGG	[1]
CrustDF1	GGT CWA CAA AYC ATA AAG AYA TTG G	CrustDR1	GGT CWA CAA AYC ATA AAG AYA TTG G	[2]
COL6	ACA AAT CAT AAA GAT ATY GG	COH6	ACA AAT CAT AAA GAT ATY GG	[3]
CrustF1M13	GTA AAA CGA CGG CCA GTT TTT CTA CAA ATC ATA AAG ACA TTG G	HCO2198M13	GTA AAA CGA CGG CCA GTG GTC AAC AAA TCA TAA AGA TAT TGG	[1], [4]

[1]: Matzen da Silva et al., 2011; [2]: Radulovici et al., 2009; [3]: Schubart, 2009; [4]: Costa et al., 2007

Each primer pair was tested at a temperature gradient of 50°C-56°C and two different magnesium concentrations (1.5mM and 3mM). The PCR profile was the same as initially

described and PCR products were run in a 1% agarose gel. COL6/COH6 was selected and a touchdown PCR was designed in order to improve specificity and guarantee a proper amplification of all samples, just as in the case of 16S gene. PCR profile was as follows: 5min at 94°C; 5 cycles of 30s at 94°C, 90s at 45°C and 60s at 72°C; 6 cycles of 30s at 94°C, 90s at 61°C-56°C (with a decrease in temperature of 1°C per cycle) and 60s at 72°C; 29 cycles of 30s at 94°C, 90s at 56°C and 60s at 72°C; a final extension of 5min at 72°C and a cooling step at 4°C (Picture 8). PCR products were run in 1% agarose gels and then they were sequenced. This procedure was used to amplify COI gene in all the following samples.

94°C 5min	
5 cycles of:	94°C 30s 45°C 90s 72°C 60s
6 cycles of:	94°C 30s 61°C-56°C 90s (↓1°C/cycle) 72°C 60s
29 cycles of:	94°C 30s 56°C 90s 72°C 60s
72°C 5min	
4°C	

Picture 8. Touchdown PCR designed for the amplification of COI gene with COL6 and COH6 primers

3.5.- Sequencing and analysis of the obtained data

Automatic sequencing of PCR products was carried out by the Molecular Biology Unit of the Research Support Services of the University of A Coruña. The genetic analysis system CEQTM 8000 (Beckman Coulter) and the genetic analysis system 3130xl (Applied Biosystems) were used for that purpose. The technique of automatic sequencing is based on the traditional Sanger sequencing method, but it employs fluorescent marking instead of radioactive marking; this allows the simultaneous reading of the products of four different reaction mixtures (a different reaction mixture is usually prepared for each triphosphate nucleotide, and each of them is marked with a different fluorochrome).

The identity of the sequences was confirmed using BLAST computer service, which is implemented in the National Centre for Biotechnology Information (NCBI). This means that each obtained sequence was aligned with the available sequences from GenBank database (also implemented in NCBI) and a signification value was calculated for the identified local similarities. So, sequences were checked out with the aim of assessing whether they really corresponded to 16S and COI genes of the spiny spider crab *Maja brachydactyla*.

COI sequences were aligned with the computer program ClustalX (Larkin *et al.*, 2007) and electropherograms were revised in order to correct the sequences by hand. The resultant alignment was saved as a .pir format document with the aim of transferring it to BioEdit v7.1.11 (Hall, 1999). Modification of the mitochondrial genetic code of *Escherichia coli* (which

is implemented by default) to convert it into the mitochondrial genetic code of decapod crustaceans was required to perform data analysis with this computer program. This implied the determination of the mitochondrial genetic code of decapod crustaceans through GenDecoder (Abascal *et al.*, 2006). The performed modifications were those represented in Table 4.

Table 4. Modifications performed in mitochondrial genetic code

Codon	Amino acid	
	<i>Escherichia coli</i>	Decapod crustaceans
TGA	Stop	Tryptophane
AGG	Arginine	Serine
AGA	Arginine	Serine
ATA	Isoleucine	Methionine

Once the modifications were done, the start codon was determined, taking into account the putative proteins which would be obtained by the translation in each of the six reading frames. Afterwards, nucleotide sequences were transformed into amino acid sequences and the latter were aligned. This amended alignment was transformed again into nucleotide sequences, which were saved as a *.pir* format document to be employed in the following analyses.

The amended alignment was analysed with DnaSP v5.10.01 (Librado & Rozas, 2009), in which the character of protein-coding sequence was specified and the mitochondrial genetic code of *Drosophila* was selected (previously, it had been determined to be equivalent to the mitochondrial genetic code of decapod crustaceans). Considering that the amplified COI region lacks introns, the whole length of the sequences was established as protein-coding and 3 sequence sets were defined: Lorient, Saint-Malo and Cedeira. Several variability parameters were then calculated for all the populations as a whole and for each single population: number of polymorphic sites (S), number of segregating sites (SS), number of synonymous changes (SC), nucleotide diversity (π , as well as its standard deviation), number of haplotypes (k), haplotype diversity (h, as well as its standard deviation), number of synonymous sites, number of non-synonymous sites, synonymous substitution rate (Ks), non-synonymous substitution rate (Ka) and Tajima's D (D, as well as its statistical significance). At last, the data file was saved as a *.meg* format document.

Finally, sequences were analysed with MEGA5 (Tamura *et al.*, 2011). The 3 mentioned data sets were defined again and the best-fit model of nucleotide substitution was estimated with the software designed by Posada & Crandall (1998), which is implemented in MEGA5. Inter-group and intra-group genetic distances were determined and a maximum likelihood phylogeny tree was built. The nucleotide composition of the sequences was also obtained.

On the other hand, the sequences of 16S gene were aligned with ClustalX (Larkin *et al.*, 2007) and corrected by hand through electropherograms revision, too. As it is not a protein-coding gene, these sequences were directly analysed with DnaSP v5.10.01 (Librado & Rozas, 2009). Likewise, 3 sets of sequences were defined (Lorient, Saint-Malo and Cedeira) and the following parameters of variability were computed: number of polymorphic sites (S), number of segregating sites (SS), nucleotide diversity (π , as well as its standard deviation), number of

haplotypes (k), haplotype diversity (h , as well as its standard deviation) and Tajima's D (D , as well as its statistical significance). The data file was converted into a *.meg* format document and MEGA5 was used to carry out a similar analysis to that described for 16S gene (definition of groups, estimation of the best-fit nucleotide substitution model, computing of inter-group and intra-group genetic distances, building of a maximum likelihood phylogeny tree and obtaining nucleotide composition).

Eventually, sequences of 16S and COI genes were concatenated in a new data file through DnaSP v5.10.01. It was saved as a *.meg* format document with the aim of building a phylogeny tree with a higher discrimination power through MEGA5.

4.- RESULTS

4.1.- PCR amplification

The primers which were initially tested for the amplification of 16S mitochondrial gene, namely 16Sar and 16Sbr (Palumbi *et al.*, 1991), did not generate amplification products in any case. Afterwards, 16L29 and 16HLeu primers (Schubart, 2009) did generate the expected amplification products and they were visualized in agarose gel electrophoresis. However, complete sequencing was not possible in any case and sequences with variable and shorter-than-expected lengths (approximately 700bp) were always obtained. Eventually, 16L29M13 and 16HLeuM13 showed an excellent yield and 16S gene was amplified and sequenced in every case.

On the other hand, the primers which were initially tested for the amplification of COI mitochondrial gene, namely LCO1490 and HCO2198 (Folmer *et al.*, 1994), did generate amplification products under the described conditions in every case. This was checked through electrophoresis gels but sequencing was not possible in any case. For this reason, some PCR products were tried to be cloned, but the ligation to the vector was not achieved in any case. As long as sequencing and cloning turned out to be impossible, the literature search which resulted in the selection of COL6 and COH6 primers (Schubart, 2009) was carried out. Every tested primer pair (*Table 3*) generated amplification products but the most suitable one was COL6/COH6 because of the high intensity of the bands in the electrophoresis gel and the absence of unspecific amplification products. The efficiency of this primer pair was excellent and COI gene was amplified and sequenced in every case.

4.2.- Analysis of the obtained sequences

The length of the amplified 16S mitochondrial gene fragment varied between 671bp and 694bp. The nucleotide composition averaged 35.5% thymine, 11.5% cytosine, 34.1% adenine and 18.9% guanine. The analysis with DnaSP v5.10.01 computer program resulted in the identification of 30 haplotypes (k) defined by 32 polymorphic sites, between which there were 17 segregating sites (SS) and 16 parsimony informative sites (S*) (*Table 5*); that is, one of the 17 segregating sites does not contribute information to the differentiation of localities. Likewise, nucleotide diversity (π) and haplotype diversity (h) were quite high; SD(π) and SD(h) respectively represent their standard deviations, which are remarkably high in the case of Cedeira population due to the reduced sampling size. Tajima's test (D represents the statistical parameter and P(D) its significance) had no significance in every case in which it could be applied. This points out the selectively neutral nature of the mutations which were found in the different localities. The statistical test could not be applied to Cedeira population, from which only 2 individuals were available (the test requires 4 at least).

Table 5. Genetic variability parameters for 16S gene in the different localities

Locality	n	S*	SS	π	SD(π)	k	h	SD(h)	D	P(D)
Lorient	25	27 (14+13)	12	0.01075	0.00169	19	0.960	0.027	-1.528	>0.10**
Saint Malo	13	29 (16+13)	15	0.01355	0.00167	13	1.000	0.030	-1.336	>0.10**
Cedeira	2	17 (17+0)	17	0.02707	0.01354	2	1.000	0.500	n/a	n/a
Total	40	32 (16+16)	17	0.00925	0.00114	30	0.962	0.020	-1.587	0.10>P>0.05**

*The number of polymorphic sites which appear only once and the number of polymorphic sites which are parsimony informative are given in brackets

**Not significant test for $\alpha=0.05$

The best-fit nucleotide substitution model was T92+G+I (Tamura, 1992), which was employed to estimate mean genetic distances inside localities (d, intra-group distances) and between different localities (inter-group distances) (Table 6). The former were strongly similar and low through the different localities and their standard deviations (SD(d)) were really small, something which indicates that the individuals belonging to each sampling locality are quite similar between them. Furthermore, inter-group genetic distances (values under the principal diagonal of the matrix on the right) were also very low and similar, with equally reduced standard deviations (values over the principal diagonal of the matrix). Therefore, there are not large differences between the localities of study.

Table 6. Intra-group and inter-group genetic distances for 16S gene

Locality	d	SD(d)	Lorient	Saint-Malo	Cedeira
Lorient	0.009	0.002		0.002	0.002
Saint-Malo	0.009	0.002	0.009		0.002
Cedeira	0.010	0.004	0.010	0.009	

T92+G+I model was also used to build a maximum likelihood phylogeny tree. In accordance with the obtained inter-group genetic distances, this tree did not show a clear differentiation between localities (data not shown).

On the other hand, the length of the amplified COI gene fragment varied between 555bp and 579bp. Mean nucleotide composition was 35.1% thymine, 20.3% cytosine, 26.3% adenine and 18.2% guanine. The analysis with DnaSP v5.10.01 resulted in the identification of 39 haplotypes (k) defined 69 polymorphic sites, between which 47 were segregating sites (SS) and 46 were parsimony informative sites (S*) (Table 7); this means that one out of 47 segregating sites does not contribute information to the differentiation of localities. Moreover, 10 out of 47 segregating sites represented synonymous changes (SC); that is, they were generally placed in the third position of a codon and they did not involve any change in the amino acid composition of cytochrome c oxidase subunit I. Nucleotide diversity (π) and haplotype diversity (h) were quite high and their standard deviations (SD(π) and SD(h)) were small, just as in the case of 16S gene; the only exception was Cedeira locality again, in which standard deviation was unusually high because of the small sampling size. Tajima's test was significant for Lorient locality and for all the samples taken as a whole, indicating that at least one of the identified mutations is not selectively neutral. The negative value of D suggests either a possible purifying selection or a population expansion, given the excess of low frequency polymorphisms which were found in relation to the expected results derived from the

hypothesis of the test. Tajima's test could not be applied to Cedeira locality because it did not have the minimum required number of sequences.

Table 7. Parameters of genetic variability for COI gene in the different localities

Localidad	n	S*	SS	SC	π	SD(μ)	k	h	SD(h)	D	P(D)
Lorient	25	65 (47+18)	45	10	0.01858	0.00454	25	1.000	0.011	-1.839	<0.05
Saint-Malo	13	52 (31+21)	20	1	0.02219	0.00260	13	1.000	0.030	-1.223	>0.10**
Cedeira	2	15 (15+0)	11	4	0.02273	0.01136	2	1.000	0.500	n/a	n/a
Total	40	69 (46+23)	47	10	0.01631	0.00291	39	0.999	0.006	-1.851	<0.05

*The number of polymorphic sites which appear only once and the number of polymorphic sites which are parsimony informative are given in brackets

**Not significant test for $\alpha=0.05$

Likewise, pair-wise comparisons resulted in the identification of 0 to 6 synonymous differences and 0 to 7 non-synonymous differences. Synonymous evolution rate (Ks) and non-synonymous evolution rate (Ka) varied from 0 to 0.047 and between 0 and 0.0182, respectively. Therefore, synonymous evolution rate is higher than non-synonymous evolution rate, just as it could be expected of a protein that is important for the organism.

The best-fit nucleotide substitution model was T92+G+I (Tamura, 1992), like in the case of 16S gene. This model was employed to estimate mean genetic distances inside localities (d, intra-group distances) and between different localities (inter-group distances) (Table 8). The results were almost the same as the ones obtained for 16S gene: intra-group distances were low and similar through the different localities again, with reduced standard deviations. This indicates that individuals from each locality are very similar between them. Also, inter-group genetic distances (values under the principal diagonal of the matrix in the right) were equally small and similar and had little standard deviations (values over the principal diagonal of the matrix). This suggests that there are not important differences between different localities.

Table 8. Intra-group and inter-group genetic distances for COI gene

Locality	d	SD(d)	Lorient	Saint-Malo	Cedeira
Lorient	0.017	0.003		0.003	0.003
Saint-Malo	0.016	0.003	0.017		0.003
Cedeira	0.018	0.006	0.016	0.017	

T92+G+I model was used to build a maximum likelihood phylogeny tree for COI gene. Similarly to the results obtained for 16S gene and in accordance with the discussed inter-group genetic distances, this tree did not allow to distinguish between different localities (data not shown).

Finally, the maximum likelihood phylogeny tree which was built with the concatenated sequences of 16S and COI genes did not discriminate different populations either (data not shown).

5.- DISCUSSION

The genetic variability of the spiny spider crab *Maja brachydactyla* was studied in three coastal localities of north-eastern Atlantic. The main conclusions which can be drawn from this study are, first, that the genetic variability of the spiny spider crab is quite high in spite of the high fishing pressure that it suffers, and, second, those localities seem to be genetically homogeneous.

The range of nucleotide diversity observed values in mitochondrial DNA ($\pi=0.00925-0.02273$) agrees with the mean values which have been reported for other crustaceans ($\pi>0.01$). The same is of application to haplotype diversity, which in this study varied between 0.960 and 1.000 and which has a mean value higher than 0.8 in other crustaceans (Khamnamtong *et al.*, 2009; Inoue *et al.*, 2007; Trontelj *et al.*, 2005; Stamatis *et al.*, 2004; Lavery *et al.*, 1996). This fact contradicts the results which were obtained in the studies of Sotelo *et al.* (2008b), in which nucleotide diversity is an order of magnitude lower ($\pi=0.003-0.005$), and implies that the spiny spider crab Atlantic population is quite more diverse than what was found initially. However, a haplotype diversity of 1.000 is too high and that could be due to the reduced sampling size (especially in the case of Cedeira) and/or to a random sampling mistake.

Anyway, the two French sampled localities have an important economic activity based in the spiny spider crab and they export huge quantities of this organism to Spain. Therefore, the strong exploitation pressure does not appear to have affected the genetic diversity of *Maja brachydactyla* in these areas. It would be interesting to have further individuals and sampling localities from Spanish Atlantic and Cantabric coasts, as the importation of French spiny spider crabs may be affecting the genetic structure of local communities. Unfortunately, the available data are not enough to obtain conclusive results for the moment.

On the other hand, further analysis on the geographic distribution of the identified haplotypes is needed. It could contradict the apparent genetic homogeneity which has been found to reveal some kind of population structure, just as it was reported by Sotelo *et al.* (2008b).

The remarkably usefulness of pleopods for the extraction of *Maja brachydactyla* genomic DNA has to be highlighted because they are not as relevant as pereopods to commercial activity. The performance of the DNA extraction protocol was very similar to that of pereopods and PCR amplification was equally successful. So, it is indeed possible to sample market spiny spider crabs without causing neither an extensive damage nor a reduction in price (which can derive from the lack of some of their valuable walking legs).

Finally, the failure of the amplification of 16S mitochondrial gene with the primers 16Sar and 16Sbr (Palumbi *et al.*, 1991) and the impossibility of neither sequencing nor cloning the PCR products of COI gene obtained with LCO1490 and HCO2198 primers (Folmer *et al.*, 1994) could be explained by the universal character of these primer pairs, which were originally developed for the amplification of 16S and COI genes in any kind of organism. The spiny spider crab might have some mutation in the annealing region which could impede the association between the primers and the DNA template, and this will explain the lack of amplification of 16S gene. In the case of COI gene, it is possible that the annealing specificity is not high enough and that,

consequently, primers are able to become associated with several regions along the DNA template (even with a pseudogene) to generate amplification products of different lengths and sequences. These unspecific products can severely hinder sequencing and so make it impossible to obtain the expected results. This problem was already discussed by Schubart (2009), who pointed out the fact that this primer pair is not optimized for the amplification of COI mitochondrial gene in decapod crustaceans.

In conclusion, the genetic variability of the spiny spider crab *Maja brachydactyla* seems to be higher than initially reported and its local populations along north-eastern Atlantic coast appear to be genetically homogeneous. Nevertheless, results are not conclusive and more detailed studies should be carried out in order to determine the effect of fishing activity on this resource with accuracy.

6.- REFERENCES

- Abascal, F., Zardoya, R., Posada, D. 2006. GenDecoder: genetic code prediction for metazoan mitochondria. *Nucl Acids Res* 34: W389-W393.
- Andrés, M., Estévez, A., Anger, K., Rotlland, G. 2008. Developmental patterns of larval growth in the edible spider crab, *Maja brachydactyla* (Decapoda: Majidae). *J Exp Mar Biol Ecol* 357: 35-40.
- Balss, H. 1922. Crustacea VII: Decapoda Brachyura (Oxyrhyncha und Brachyrhyncha) und geographische Übersicht über Crustacea Decapoda. In: W. Michaelsen (ed.), Beiträge zur Kenntnis der Meeresfauna Westafrikas, pp. 69-110. Friederichsen and Co., Hamburg.
- Corgos, A. 2004. Estrategia vital, estructura espacial y dinámica poblacional de la centolla, *Maja squinado* (Decapoda: Majidae). PhD Thesis. Departamento de Biología Animal, Biología Vegetal e Ecología. Universidade de A Coruña.
- Costa, F.O., deWaard, J.R., Boutillier, J., Ratnasingham, S., Dooh, R.T., Hajibabaei, M., Hebert, P.D.N. 2007. Biological identifications through DNA barcodes: the case of the Crustacea. *Can J Fish Aquat Sci* 64: 272-295.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotech* 3: 294-299.
- Freire, J., Bernárdez, C., Corgos, A., Fernández, L., González-Gurriarán, E., Sampedro, M.P., Verísimo, P. 2002. Management strategies for sustainable invertebrate fisheries in coastal ecosystems of Galicia (NW Spain). *Aquat Ecol* 36: 41-50.
- Freire, J., Carabel, S., Verísimo, P., Bernárdez, C., Fernández, L. 2009. Patterns of juvenile habitat use by the spider crab *Maja brachydactyla* as revealed by stable isotope analyses. *Sci Mar* 73: 39-49.
- González-Gurriarán, E., Fernández, L., Freire, J., Muiño, R. 1998. Mating and role of seminal receptacles in the reproductive biology of the spider crab *Maja squinado* (Decapoda: Majidae). *J Exp Mar Biol Ecol* 220: 269-285.
- Hall, T.A. 1999. Bioedit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser* 41: 95-98.
- Herbst, J.F. 1788. Versuch einer Naturgeschichte der Krabben und Krebse nebst einer systematischen Beschreibung ihrer verschiedenen Arten. Lange H.A. (Ed.). pp. 207-238. Berlin und Stralsund.
- Inoue, N., Watanabe, H., Kojima, S., Sekiguchi, H. 2007. Population structure of Japanese lobster *Panulirus japonicus* inferred by nucleotide sequence analysis of mitochondrial COI gene. *Fish Sci* 73: 550-556.

- Khamnamtong, B., Klinbunga, S., Menasveta, P. 2009. Genetic diversity and geographic differentiation of the giant tiger shrimp (*Penaeus monodon*) in Thailand analyzed by mitochondrial COI sequences. *Biochem Genet* 47: 42-55.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., López, R., Thompson, J.D., Gibson, T.J., Higgins, D.G. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23: 2947-2948.
- Lavery, S., Moritz, C., Fielder, D.R. 1996. Indo-Pacific population structure and evolutionary history of the coconut crab *Birgus latro*. *Mol Ecol* 5: 557-570.
- Librado, P., Rozas, J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25: 1451-1452.
- Matzen da Silva, J., Creer, S., dos Santos, A., Costa, A.C., Cunha, M.R., Costa, F.O., Carvalho, G.R. 2011. Systematic and evolutionary insights derived from mtDNA COI Barcode Diversity in the Decapoda (Crustacea: Malacostraca). *PLoS ONE* 6: e19449.
- Neumann, V. 1998. A review of the *Maja squinado* (Crustacea: Decapoda: Brachyura) species-complex with a key to the eastern Atlantic and Mediterranean species of the genus. *J Nat Hist* 32: 1667-1684.
- Ng, P.K.L., Guinot, D., Davie, P.J.F. 2008. Systema Brachyurorum: part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles B Zool* 17: 1-286.
- Palumbi, S.R., Martin, A., Romano, S., McMillan, W.O., Stice, L., Grabowski, G. 1991. The simple fool's guide to PCR, Version 2.0. Department of Zoology, University of Hawaii, Honolulu.
- Posada, D., Crandall, K.A. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Radulovici, A.E., Sainte-Marie, B., Dufresne, F. 2009. DNA barcoding of marine crustaceans from the Estuary and Gulf of St Lawrence: a regional-scale approach. *Mol Ecol Resour* 9: 181-187.
- Rotlland, G., González-Gurriarán, E., Fernández, L., Benhalima, K., Ribes, E. 2007. Ovarian maturation of the multi-spawning spider crab *Maja brachydactyla* (Decapoda: Majidae) with special reference to yolk formation. *Mar Biol* 152: 383-394.
- Schubart, C.D. 2009. Mitochondrial DNA and decapod phylogenies: the importance of pseudogenes and primer optimization. In: Martin JW, Crandall KA & Felder DL (Eds.), *Crustacean Issues 18: Decapod Crustacean Phylogenetics*, pp. 47-65. Boca Raton, Florida: Taylor & Francis/CRC Press.
- Sotelo, G., Morán, P., Posada, D. 2008a. Genetic identification of the northeastern Atlantic spiny spider crab as *Maja brachydactyla* Balss, 1992. *J Crustacean Biol* 28: 76-81.

- Sotelo, G., Morán, P., Fernández, L., Posada, D. 2008b. Genetic variation of the spiny spider crab *Maja brachydactyla* in the northeastern Atlantic. *Mar Ecol Prog Ser* 362: 211-223.
- Sotelo, G., Morán, P., Posada, D. 2009a. Molecular phylogeny and biogeographic history of the European *Maja* spider crabs (Decapoda, Majidae). *Mol Phylogenet Evol* 53: 314-319.
- Sotelo, G. 2009b. Phylogeography of the Galician crabs *Necora puber* and *Maja brachydactyla* in the NE Atlantic. PhD Thesis. Departamento de Bioquímica, Genética e Inmunología. Universidad de Vigo.
- Stamatis, C., Triantafyllidis, A., Moutou, K.A., Mamuris, Z. 2004. Mitochondrial DNA variation in northeast Atlantic and Mediterranean populations of Norway lobster, *Nephrops norvegicus*. *Mol Ecol* 13: 1377-1390.
- Tamura, K. 1992. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C content biases. *Mol Biol Evol* 9: 678-687.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S. 2011. MEGA5: Molecular Evolutionary Genetic Analysis using maximum likelihood, evolutionary distance and maximum parsimony methods. *Mol Biol Evol* 28: 2731-2739.
- Trontelj, P., Machino, Y., Sket, B. 2005. Phylogenetic and phylogeographic relationships in the crayfish genus *Austropotamobius* inferred from mitochondrial COI gene sequences. *Mol Phylogenet Evol* 34: 212-226.