



## OPEN ACCESS

## EDITED BY

Dan Lindholm,  
University of Helsinki, Finland

## REVIEWED BY

Mikko Hiltunen,  
University of Eastern Finland, Finland  
Vesa Olkkonen,  
Minerva Foundation, Finland

## \*CORRESPONDENCE

Jair C. Soares

✉ [Jair.C.Soares@uth.tmc.edu](mailto:Jair.C.Soares@uth.tmc.edu)

Chu-Huang Chen

✉ [cchen@texasheart.org](mailto:cchen@texasheart.org)

<sup>†</sup>These authors have contributed equally to this work

RECEIVED 10 August 2023

ACCEPTED 30 October 2023

PUBLISHED 16 November 2023

## CITATION

Akyol O, Akyol S, Chou M-C, Chen S, Liu C-K, Selele S, Soares JC and Chen C-H (2023) Lipids and lipoproteins may play a role in the neuropathology of Alzheimer's disease.

*Front. Neurosci.* 17:1275932.

doi: 10.3389/fnins.2023.1275932

## COPYRIGHT

© 2023 Akyol, Akyol, Chou, Chen, Liu, Selele, Soares and Chen. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Lipids and lipoproteins may play a role in the neuropathology of Alzheimer's disease

Omer Akyol<sup>1†</sup>, Sumeyya Akyol<sup>2†</sup>, Mei-Chuan Chou<sup>3</sup>, Shioulan Chen<sup>4</sup>, Ching-Kuan Liu<sup>5</sup>, Salih Selele<sup>6</sup>, Jair C. Soares<sup>6\*</sup> and Chu-Huang Chen<sup>1\*</sup>

<sup>1</sup>Molecular Cardiology, Vascular and Medicinal Research, The Texas Heart Institute, Houston, TX, United States, <sup>2</sup>NX Prenatal, Houston, TX, United States, <sup>3</sup>Department of Neurology, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan, <sup>4</sup>Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan, <sup>5</sup>Institute of Precision Medicine, College of Medicine, National Sun Yat-sen University, Kaohsiung, Taiwan, <sup>6</sup>Department of Psychiatry and Behavioral Sciences, UTHealth Houston McGovern Medical School, Houston, TX, United States

Alzheimer's disease (AD) and other classes of dementia are important public health problems with overwhelming social, physical, and financial effects for patients, society, and their families and caregivers. The pathophysiology of AD is poorly understood despite the extensive number of clinical and experimental studies. The brain's lipid-rich composition is linked to disturbances in lipid homeostasis, often associated with glucose and lipid abnormalities in various neurodegenerative diseases, including AD. Moreover, elevated low-density lipoprotein (LDL) cholesterol levels may be related to a higher probability of AD. Here, we hypothesize that lipids, and electronegative LDL (L5) in particular, may be involved in the pathophysiology of AD. Although changes in cholesterol, triglyceride, LDL, and glucose levels are seen in AD, the cause remains unknown. We believe that L5—the most electronegative subfraction of LDL—may be a crucial factor in understanding the involvement of lipids in AD pathology. LDL and L5 are internalized by cells through different receptors and mechanisms that trigger separate intracellular pathways. One of the receptors involved in L5 internalization, LOX-1, triggers apoptotic pathways. Aging is associated with dysregulation of lipid homeostasis, and it is believed that alterations in lipid metabolism contribute to the pathogenesis of AD. Proposed mechanisms of lipid dysregulation in AD include mitochondrial dysfunction, blood–brain barrier disease, neuronal signaling, inflammation, and oxidative stress, all of which lead ultimately to memory loss through deficiency of synaptic integration. Several lipid species and their receptors have essential functions in AD pathogenesis and may be potential biomarkers.

## KEYWORDS

Alzheimer's disease, lipids, cholesterol, electronegative LDL, LDLR, LOX-1

## 1. Introduction

Neurodegeneration refers to ongoing neuron and synapse loss in specific brain regions, leading to disorders like Alzheimer's disease (AD), Parkinson's disease, and Huntington's disease. AD is characterized by the presence of neurofibrillary tangles and senile plaques in the brain. Key factors in the pathogenesis of AD include  $\beta$ -amyloid protein ( $\beta$ ) buildup and

neuronal apoptosis. Current therapies cannot fully prevent progression due to permanent neuron loss (Citron, 2010) but can slow it down. A clear-cut relationship between lipids/lipoproteins/apolipoproteins and AD has not yet been established, even though there are signs of abnormalities in lipid metabolism. Despite extensive preclinical and clinical research over the past three decades, no disease-preventive or disease-modifying treatment alternatives have been identified for AD and late-onset clinical dementia.

In this article, we review and characterize the role of lipids and lipoproteins in the neuropathology of AD and have focused on their impact on amyloid precursor protein (APP) processing, tau phosphorylation, and A $\beta$  production. Lipids and lipoproteins play a key role in the processing of APP and the production of A $\beta$ . Abnormal phosphorylation of tau protein, also a hallmark feature of AD, can be affected by lipids and lipoproteins. Therefore, a better understanding of the mechanistic role of these molecules in AD pathology may provide avenues for therapeutic interventions.

## 2. Hypothesis

The pathophysiology of AD involves multiple genetic, environmental, developmental, social, biochemical, and anatomical factors. The speculative presumption has been that reactive oxygen species (ROS) play a key role in AD by oxidizing lipids, which leads to the modification of low-density lipoprotein (LDL) particles through various mechanisms, primarily oxidation. This modification results in the formation of a subclass of products known as negatively charged LDL. The major modifiable risk factors of AD, such as lifestyle (diet, alcohol, smoking, etc.), traumatic brain injury, type 2 diabetes, and vascular disease (Edwards Iii et al., 2019), are all linked to LDL metabolism and lipoprotein disruption. Our hypothesis is that LDL, especially its most electronegative subfraction called L5 (Figure 1), may play a key role in AD pathophysiology.

Vascular risk factors, such as type 2 diabetes, hyperlipidemia, smoking, obesity, high blood pressure, and physical inactivity, can reduce cerebral perfusion, activate neuroinflammatory responses, and increase ROS production. These risk factors may induce A $\beta$  production and cause intellectual disability in the form of cognitive decline. Elevated plasma LDL cholesterol levels are more likely to be related to early onset AD, and gene-based rare variant testing has shown that rare apoB-coding variants were significantly more abundant in early-onset AD (Wingo et al., 2019).

Human and animal investigations have demonstrated that the pathophysiology of AD is directly related to impaired lipid metabolism. Recent studies have shown that L5 has an important role in the initiation and evolution of atherosclerosis and in the increased interrelationship of vasoconstriction with A $\beta$ . The combined effects of A $\beta$  and ROS may lead to cognitive impairment through cerebrovascular dysfunction. Therefore, both ROS-mediated lipoprotein changes and L5, the major product of these oxidative changes, may be considered as potential biomarkers of metabolic stress and may be important in the pathogenesis of AD and other types of dementia (Akyol et al., 2020).

## 3. Supporting evidence

### 3.1. Contemporary pathophysiology of AD

There is increasing evidence that cerebrovascular disease (CVD) affects the pathogenesis of AD. Co-occurrence of CVD and AD pathology is common in older patients with dementia (Attems and Jellinger, 2014), and vascular pathology images in the brain are regularly monitored in MRI scans of AD patients (Staffaroni et al., 2017). Observational epidemiological investigations have found that lifestyle and cardiovascular risk factors are correlated with dementia risk, and targeting these risk factors may be a feasible prevention strategy for dementia (Norton et al., 2014). In two independent commissioned reports by the *Lancet* and *National Academy of Medicine*, encouraging results suggest that modifying risk factors including hypertension, diabetes, physical inactivity, and obesity could prevent up to 35% of cases of dementia (Livingston et al., 2017; National Academies of Sciences and Medicine, 2017).

In a recent meta-analysis by Bellenguez et al. a two-stage, genome-wide association study led to the validation of 33 previously identified loci. This groundbreaking research not only effectively doubled the overall number of genetic loci linked to AD risk, but also significantly contributed to our knowledge in the field by expanding our understanding of AD's underlying pathophysiology (Bellenguez et al., 2022). The most prominent gene sets uncovered in this study were closely tied to amyloid/tau, whereas several other gene sets with significant associations pertained to lipid metabolism, endocytosis, and immune processes, including the activation of macrophages and microglial cells. The studies have shown that cardiovascular-related loci elevate the risk of late-onset AD. The  $\epsilon$ 4 allele of apolipoprotein-E (APOE), a major risk factor for AD, encodes a lipid transporter required in the metabolism of cholesterol (Mahley, 2016). The FINGER study, using 11C-Pittsburgh compound B-PET imaging, found a link between early brain A $\beta$  accumulation and declining executive functions and a strong APOE4 association with A $\beta$  positivity (Kempainen et al., 2018). Genome-wide association studies have identified a single nucleotide polymorphism (SNP) in late-onset AD that is involved in lipid processing (Lambert et al., 2013; Reitz, 2013) and cholesterol metabolism (Carmona et al., 2018). Apart from APOE, 90 SNPs on 19 chromosomes elevated the risk of both AD and cardiovascular issues. Four novel genetic variants that increase AD risk were identified in a meta-analysis across three independent cohorts (Broce et al., 2019). In AD-affected brains, three genes showed differential expression changes, suggesting shared risk with cardiovascular disease (Broce et al., 2019).

Clinical, epidemiological, and animal studies suggest that elevated plasma lipids (i.e., hypercholesterolemia) are risk factors for AD (Xue-Shan et al., 2016). Neuronal cells are the target for cholesterol binding via LDL receptors (LDLR). Hypercholesterolemia can impair the blood brain barrier (BBB), leading to metabolic disintegration of cholesterol in the brain (Xue-Shan et al., 2016). Collectively, these findings indicate a common genetic relationship between AD and plasma lipids, with particular genes identified as potential drivers of this genetic link.

Individuals with AD may have both vascular and AD pathology, potentially worsening cognitive decline (Broce et al., 2019). Newly discovered genetic loci in MBLAC1, MINK1, and MTCH2/SPI1 may increase susceptibility to brain inflammation and vascular lesions,




FIGURE 1

Composition and structural characteristics of regular (L1) and electronegative (L5) LDL in humans. The apolipoprotein content specifies the structure, receptor interactions, assembly, and metabolism of these lipoproteins. The L5 subtype differs from regular LDL because of its additional apolipoproteins. Apo(a), apolipoprotein small (a); ApoA1, apolipoprotein-A1; ApoB100, Apolipoprotein-B100; ApoCIII, apolipoprotein-CIII; APOE, apolipoprotein-E.

leading to worse AD outcomes (Broce et al., 2019). Creating a multi-gene “hazard” model for cardiovascular pathways could help predict when older adults might develop AD (Desikan et al., 2017) and allow for targeted therapeutic intervention to alter its course.

### 3.1.1. Oxidative stress in linking L5 to AD

Several studies have revealed a significant correlation between inflammatory and vascular risk factors and AD, along with a strong association between dementia and factors such as nitrate and oxidative stress (Perry et al., 2002; Fitzpatrick et al., 2004). AD patients have elevated synaptic cholesterol release and neuronal degeneration, potentially contributing to the susceptibility of lipoproteins to ROS, leading to the generation of oxidized LDL (ox-LDL). This ox-LDL could serve as a stable biomarker for monitoring the inflammatory cascade in AD (Kankaanpaa et al., 2009).

A positive correlation between plasma NO concentration and oxidative stress marker 3-nitrotyrosine (3-NT) levels (Sinem et al., 2010) suggests insights into AD’s mechanisms and pathophysiology, potentially linked to ROS-induced stress. When the body is exposed to disease, infection, or certain environmental stressors, oxidation surpasses the level of antioxidants, and huge amounts of highly reactive ROS begin to accumulate (Zhao et al., 2014). AD patients show markers of lipid peroxidative oxidative stress in the hippocampus and cortex. Elevated NO levels in animal studies are associated with A $\beta$  neurotoxicity, potentially contributing to AD development (Luth et al., 2001). Additionally, plasma ox-LDL levels are higher in AD patients than in controls (Zhao et al., 2014), but they negatively correlate with Mini-Mental State Exam (MMSE)

scores. These findings may relate to increased cerebral cholesterol concentration and oxidative/nitrative stress. In another study, plasma ox-LDL levels were significantly higher in AD patients than in those with normal cognitive function (Dildar et al., 2010). However, an earlier study found little change in plasma ox-LDL levels in AD patients, particularly in severe cases (Zafrilla et al., 2006).

### 3.1.2. BBB disruption in AD

Impairment of the BBB is a significant factor in AD pathogenesis (Montagne et al., 2015; van de Haar et al., 2016). The damaged BBB allows proinflammatory and neurotoxic factors, such as thrombin, immunoglobulins, hemoglobin, bacterial breakdown products, iron, and fibrinogen, to disrupt CNS function and promote neuroinflammation and neurodegeneration (Zlokovic, 2011). Recent research suggests that BBB dysfunction is an early event leading to mild cognitive damage in AD (van de Haar et al., 2016), offering a potential target for intervention. However, despite increased understanding, clinical strategies to prevent BBB disruption in AD are still lacking. Vascular risk factors significantly impact the incidence of vascular dementia and AD (Gudala et al., 2013). A meta-analysis revealed a 73% increased risk of overall dementia, a 127% increased risk of vascular dementia, and a 56% increased risk of AD associated with diabetes (Gudala et al., 2013). Mid-life vascular risk factors, including diabetes, obesity, high cholesterol, and hypertension, have the most substantial impact on late-onset dementia due to AD (Reitz, 2013).



## 3.2. Dyslipidemia in AD

There is growing evidence of a strong connection between dementia and lipid metabolism, especially in AD, where alterations in lipid levels and distribution play a role in disease progression (Hottman et al., 2014). Multiple studies have linked dyslipidemia to AD, primarily because of elevated levels of total cholesterol and LDL-cholesterol (LDL-C) and reduced levels of HDL-cholesterol (HDL-C) (Launer et al., 2001; Wanamaker et al., 2015). Elevated circulatory HDL-C amounts are associated with a reduced chance of CVD (Miller and Miller, 1975). Furthermore, oxidation-derived stress in the CNS can lead to the oxidation of LDL-C (ox-LDL) and VLDL (ox-VLDL), which determine cytotoxicity of oxidized forms (Escargueil-Blanc et al., 1994). The incidence (Grossi et al., 2018) of the  $\epsilon 4$ -allele of the APOE gene is higher in AD patients than in controls, which indicates the importance of genetic factors in AD development.

The relationship between cholesterol and AD is well-established, yet the precise mechanisms of cholesterol's impact on AD pathophysiology remain unclear (Akyol et al., 2021). Brain cholesterol directly affects A $\beta$ -related processing. Reduced cholesterol in hippocampal neurons has been linked to a decrease in A $\beta$  oligomerization (Schneider et al., 2006), whereas elevated cholesterol levels in AD brains promote the activity of  $\gamma$ - and  $\beta$ -secretases (Xiong et al., 2008). In essence, cholesterol may enhance the degradation of  $\beta$ - and  $\gamma$ -secretases in AD brains, leading to elevated A $\beta$  levels. In mild-AD and AD patients, cholesterol ester (CE) levels were higher compared to controls (Akyol et al., 2021). Brain cholesterol levels are primarily regulated by *de novo* synthesis in astrocytes, oligodendrocytes, and neurons (Grimm et al., 2017).

The mechanism connecting AD to the  $\epsilon 4$ -allele is not yet completely understood. Researchers have proposed a connection between protease activities that degrade A $\beta$  in cerebellar vascular infarction and carriers of the  $\epsilon 4$  allele (Zhu et al., 2013). In a study conducted by Koffie et al., AD patients who had the APOE  $\epsilon 4$ -allele tended to show a significantly greater buildup of A $\beta$ -oligomers than those with the APOE  $\epsilon 3$  allele (Koffie et al., 2012). This heightened accumulation of A $\beta$ -oligomers ultimately leads to the loss of synapses.

### 3.2.1. Lipidomics studies and 27-hydroxycholesterol in AD

A recent study identified specific lipid subclasses associated with more severe AD pathology, including glycerolipids (diacylglycerols [DAG] and triacylglycerols), neutral lipids (CE), glycerophospholipids (phosphatidyl ethanolamine, phosphatidyl glycerol, phosphatidyl inositol, phosphatidylserine, and lysyl-phosphatidylglycerol), sphingolipids, ceramides (CER), dihydroceramides, hexosylceramides, and lactosylceramides (Akyol et al., 2021). Lipidomic studies have shown that AD patients have reduced levels of ether-sterols, PC, sphingomyelin, and phospholipids compared to controls (Oresic et al., 2011). Kim et al. observed a significant increase (more than 2-fold) in 14 lipids in the plasma of individuals with AD, including DAG, phosphatidyl ethanolamine, TG, and CER. Among these lipids PE, DAG, and TG correlated strongly with brain atrophy, making them potential markers for early detection of mild cognitive impairment when combined with MMSE scores (Kim et al., 2013). Trushina et al. used non-targeted metabolomics through liquid chromatography/mass spectrometry and found altered cholesterol and sphingolipid

transport in the cerebrospinal fluid and plasma of AD patients compared to cognitively normal individuals (Trushina et al., 2013).

In the periphery, cholesterol is oxidized through the steroid side-chain, resulting in the formation of a compound called 27-hydroxycholesterol (27-OHC). The enzyme CYP27A1, found in almost all cells of the body, is responsible for producing 27-OHC. This mechanism allows cells to mobilize cholesterol as an alternative to the traditional method of HDL-mediated reversed cholesterol transport (Babiker et al., 1997). Recent studies have shown that 27-OHC can cross the BBB and directly enter the cerebral circulation. In a FINGER study, a decrease in 27-OHC levels over a two-year multidomain lifestyle and vascular intervention, which included cardiovascular risk management, dietary modifications, physical activity, and cognitive training, was associated with improved cognitive performance, particularly in the realm of memory (Sandebring-Matton et al., 2021). Furthermore, levels of 27-OHC in the cerebrospinal fluid depend on the integrity and functioning of the BBB. Damage to this barrier increases the transport of 27-OHC from the circulation into the brain (Leoni et al., 2003). The enzyme CYP7B1, found in neurons, is believed to play a critical role in the metabolism of 27-OHC (Meaney et al., 2007). Conversely, the primary mechanism for eliminating cholesterol from the brain involves converting it into a compound called 24S-hydroxycholesterol (24S-OHC), or cerebrosterol, which has the ability to cross the BBB (Bjorkhem et al., 1997). The enzyme CYP46A1, predominantly found in neuronal cells, is responsible for producing 24S-OHC (Lund et al., 1999). There is a reciprocal relationship between these two oxysterols at the BBB, with fluxes occurring in opposite directions. Both fluxes are concentration-driven, with a higher concentration of 24S-OHC in the grey matter, its major site of production, and a lower concentration of 27-OHC in the same area, the main site of metabolism (Heverin et al., 2005). The balance between 24S-OHC and 27-OHC, as well as their metabolites, is hypothesized to be important in amyloidogenesis in AD (Bjorkhem, 2006). Studies have shown a significantly increased ratio of 27-OHC to 24S-OHC in the brains of AD patients, supporting this hypothesis. The flux of 27-OHC into the brain could potentially be the missing connection between hypercholesterolemia and AD.

### 3.2.2. VLDL and LDL involvement in AD pathology

VLDL receptor polymorphisms have been associated with the risk of dementia, particularly in vascular and mixed dementia (Helbecque et al., 2001). A prospective study revealed an inverse relationship between total VLDL-C and the risk of dementia (Tynkkynen et al., 2018). This may be attributed, at least in part, to the association between cardiovascular events and VLDL-C (Wurtz et al., 2015). Furthermore, one subclass of VLDL lipoproteins (triglyceride-to-total lipid ratio in very-large-VLDL) reduced the risk of dementia, whereas another (total cholesterol-to-total lipid ratio in very-large-VLDL) increased the risk of dementia (Tynkkynen et al., 2018). No significant associations were observed between apolipoproteins, cholesterol, fatty acids, phospholipids, triglycerides, ketone bodies, glycolysis-related metabolites, markers of inflammation, or fluid balance metabolites and dementia/AD (Tynkkynen et al., 2018). In contrast, previous research has indicated that specific phospholipids are linked to the progression of mild cognitive impairment to AD (Mapstone et al., 2014; Li et al., 2016), although the metabolomics platform used in Tynkkynen's study differs from those in the previous studies.

Linkage analysis, conducted on a genome-wide scale, has identified chromosome-12p as a region of interest for AD, encompassing LDL-receptor-related protein-1,  $\alpha$ 2-macroglobulin, and lectin-like ox-LDL receptor 1 (OLR-1) as a susceptible region for AD (Wu et al., 1998; Lee et al., 2008). OLR-1, initially discovered in endothelial cells as an ox-LDL receptor (Sawamura et al., 1997), is implicated in the pathogenesis of conditions like diabetes, atherosclerosis, and hypertension (Palmieri et al., 2013). This receptor is fully expressed in the brain and spinal cord (Yamanaka et al., 1998). Inflammatory cytokines, oxidative stress, and ox-LDL can induce its expression (Mehta et al., 2006), and these stimuli may be regarded as a hallmark of AD (Agostinho et al., 2010). OLR-1 functions through ligand interactions, promoting cytokine secretion in brain cells and mitigating the neurotoxic effects of amyloid peptides. Adequate OLR-1 expression may be a key factor in preventing AD. Moreover, it is highly expressed in vascular endothelial cells and may be involved in amyloid angiopathy in the brain, associated with AD progression (Shi et al., 2006). This hypothesis is supported by the research of Lambert et al. (2003), which noted significantly lower OLR-1 mRNA levels in AD patients than in healthy controls.

In healthy individuals, the accumulation of A $\beta$  fibers is effectively prevented by several organized removal mechanisms, including BBB transport, extracellular proteolysis, receptor-mediated endocytosis, and the efflux of soluble A $\beta$  into the peripheral circulation (Ramanathan et al., 2015). The LDL receptor related protein 1 (Lrp-1) plays a pivotal role in A $\beta$  catabolism, internalization, and production (Pietrzik et al., 2002; Deane et al., 2004). Lrp-1 is synthesized on the endothelial cell surface of the BBB, facilitating the rapid removal of A $\beta$  from brain tissue through endocytosis and degradation (Pflanzner et al., 2011). Reduced levels of Lrp-1 in the brains of patients of AD patients have been consistently observed (Shinohara et al., 2017). A $\beta$  forms complexes with lactoferrin, APOE, prion proteins, and activated  $\alpha$ 2-macroglobulin, which undergo Lrp-1 mediated endocytosis (Kang et al., 2000; Rushworth et al., 2013). The monomeric A $\beta$  also binds to the extracellular ligand-binding domain of Lrp-1, facilitating endocytosis (Deane et al., 2004). Microglial cells predominantly migrate to plaques through micropinocytosis, effectively clearing A $\beta$  (Lee and Landreth, 2010). Lrp-1-mediated A $\beta$  clearance from pericytes helps prevent A $\beta$  accumulation in cerebral vessels (Sagare et al., 2013). Soluble Lrp-1 is cleaved and released from  $\beta$ -secretase in the periphery, further enhancing its binding to A $\beta$  and facilitating its cerebral removal (Sagare et al., 2007). Additionally, A $\beta$  removal from the system is supported by the liver, spleen, and kidney via Lrp-1 (Ramanathan et al., 2015).

### 3.2.3. Cholesterol metabolism dysfunction in microglia

Emerging evidence suggests that disruptions in cholesterol metabolism, especially in high cholesterol environments, can harm microglia function. When microglia cannot maintain proper cholesterol metabolism and accumulate lipid droplets, it triggers a pro-inflammatory lipid profile (Loving et al., 2021). Excessive cholesterol exposure can negatively affect microglia through membrane protein mechanisms, contributing to various aspects of disease-related microglia dysfunction, including increased inflammation and a reduced ability to engulf particles, both of which are associated with AD. In summary, in high cholesterol settings, elevated cholesterol levels in neuronal cell membranes lead to greater production of A $\beta$  due to the proximity of key proteins involved in A $\beta$

production. This, in turn, burdens microglia with clearing more A $\beta$ . Simultaneously, high cholesterol impairs microglia's ability to clear A $\beta$  and enhances their inflammatory signaling and production of ROS. These factors drive A $\beta$  accumulation, leading to plaque formation and creating a pro-inflammatory environment that contributes to neurodegeneration. Statin drugs, commonly prescribed and generally safe, may offer a potential approach to regulate cholesterol and reduce neuroinflammation (Munoz Herrera and Zivkovic, 2022). Moreover, a high cholesterol (3%) diet induces a pro-inflammatory response in rodent microglia models by activating the inflammasome. While anti-inflammatory cytokines are also released as part of the response, the ultimate outcome is a compromised BBB (Chen et al., 2018). Moreover, as supported by the literature (Duong et al., 2021), cholesterol overload can lead to chronic inflammation in microglia. This inflammation is further induced by L5, which activates microglia through mechanisms involving an increase in nitrogen/oxygen radicals (peroxides, nitric oxide [NO], and ROS), elevated levels of TNF- $\alpha$ , reduced basal levels of interleukin (IL)-10, and increased production of proinflammatory proteins such as inducible NOS and cyclooxygenase 2 (Yu et al., 2017). L5 predominantly induces apoptosis in BV-2 microglial cells, and this effect is significantly interrupted by signaling pathway inhibitors. This suggests that L5 may interact with toll like receptor-4 to regulate various biopathways, including NF- $\kappa$ B, PI3K/Akt, and MAPKs. These interactions with L5, contributing to atherosclerosis formation, are also implicated in the development of neurodegeneration linked to neuroinflammation (Yu et al., 2017).

### 3.2.4. Comparing the impact and receptor preferences of L5 and LDL

Circulating LDLs are a group of lipoproteins that have various compositions, electrical charges, and densities. Electropositive LDL and L5 differ in lipid and protein composition, size, and density (Sanchez-Quesada et al., 2012). L5 has much higher aggregation and proteoglycan binding ability, which may be due to its phospholipolytic activity, than does the more positively charged LDL (LDL (+)) (Sanchez-Quesada et al., 2012). L5 promotes inflammatory responses in cells by increasing the expression and secretion of cytokines. L5-releasing inflammatory cytokines were first described in the endothelium (De Castellarnau et al., 2000). By promoting gene transcription, L5 elicits the release of inflammatory mediators like IL-6, IL-10, and monocyte chemoattractant protein-1 (MCP1) (Bancells et al., 2010).

In the PC12 rat cell line, L5 caused cell damage in a time- and concentration-dependent manner (Wang et al., 2017). In a neutral comet assay designed to examine whether L5-induced cell death involved DNA damage, the results showed that L5 and ox-LDL, but not L1 (a less negatively charged form of LDL), led to the formation of comet tails, indicating double-strand breaks in chromatin (Wang et al., 2017). At lower concentrations, L5 hindered nerve growth factor-related neuron cell differentiation by inhibiting the PI3K/Akt cascade through LOX-1-related mechanisms. In the same study, the researchers investigated whether L5 caused apoptotic cell death and activated apoptosis-related pathways in PC12 cells. They found that L5, but not L1, increased caspase-3 expression 6–12h after exposure and that this apoptotic pathway was blocked by a pancaspase inhibitor called ZVAD-FMK (Wang et al., 2017). Additionally, L5 treatment led to increased levels of phosphorylated-p53 just 10 min after exposure. The study showed that p53-dependent apoptosis is a primary

mechanism of cell death induced by L5 in PC12 cells. Furthermore, L5 inhibited differentiation of cultured PC12 cells induced by nerve growth factor, indicating its harmful effects on these cells. The results suggest that LDL(−), particularly the L5 subfraction, is toxic to PC12 cells, reducing their viability in a time- and concentration-dependent manner through LOX-1/p53/caspase-3 dependent apoptosis and ATM/H2AX-related DNA damage. Lowering plasma LDL(−) levels and preventing the electronegative modification of LDL might help prevent cognitive impairment associated with conditions like AD, where high LDL levels are linked to the disease's progression.

L5 induces cytokine release in monocytes through CD14 toll-like receptor 4 (TLR-4) pathway (Estruch et al., 2013), which is triggered by increased CER content in L5 (Estruch et al., 2014). This signaling pathway has also been associated with the release of proinflammatory cytokine IL-1 $\beta$  (Estruch et al., 2015). Studies have extensively investigated signaling pathways induced by minimally modified LDL and ox-LDL, including activation of PI3k/Akt-mediated p38 MAPK (Rios et al., 2012), ERK (Namgaladze et al., 2008), AP-1 (Choi et al., 2012), and NF- $\kappa$ B transcription factors (Janabi et al., 2000). However, there is limited research on L5-induced signaling pathways, mostly focusing on endothelial cells. In endothelial cells, L5 activates p38 MAPK (Chu et al., 2013; Chen et al., 2015) and transcription factors NF- $\kappa$ B and AP-1 (Chu et al., 2013) while inhibiting the PI3k/Akt pathway (Lu et al., 2009). In monocytes, L5 activated AP-1 and NF- $\kappa$ B (Estruch et al., 2013). Additionally, in one study, L5 facilitated p38 MAPK phosphorylation through the PI3k/Akt1 and TLR4 pathway (Estruch et al., 2016), leading to the activation of NF- $\kappa$ B, CREB, and AP-1, ultimately resulting in cytokine release.

### 3.2.5. Impact of L5 on endothelial cells and platelets

Endothelial cells control the recruitment of blood lymphocytes and monocytes to arterial walls, a crucial step in atherosclerosis. L5 attracts monocytes and lymphocytes to the endothelium (Yang et al., 2003), indicating its involvement in the early stages of atherosclerosis. It also triggers the release of chemokines and adhesion molecules (Figure 2), including vascular-cell adhesion molecules (Ziouzenkova et al., 2003). In human umbilical vein endothelial cells (HUVECs), L5 induced the release of IL-8 and MCP1 (De Castellarnau et al., 2000), which are key factors in recruiting T lymphocytes and monocytes to the endothelium. This effect has been observed in individuals with normal lipid levels (De Castellarnau et al., 2000), familial hypercholesterolemia (FH) (Sanchez-Quesada et al., 2003), and diabetes mellitus (Benitez et al., 2007). In individuals with diabetes and FH, who have higher levels of L5, the inflammatory impact of L5 is likely more pronounced compared to normolipidemic subjects. HUVEC studies have also revealed that L5 triggers the release of various cytokines associated with inflammation, including epithelial cell-derived neutrophil-activating peptide-78 (Abe et al., 2007), granulocyte/monocyte colony stimulating factor, growth-related oncogene, and IL-6 (Benitez et al., 2006). This effect has also been observed in arterial-derived cultured human endothelial cells (de Castellarnau et al., 2007). Lastly, L5 induced the release of matrix metalloproteinases and the expression of vascular-endothelial growth factor in bovine arterial endothelial cells (Tai et al., 2006).

Furthermore, research suggests that hypercholesterolemia, particularly with elevated L5 levels, can increase the risk of thrombosis and atherosclerosis (Wang and Tall, 2016; Figure 3). Platelet function and biosynthesis can be affected by hypercholesterolemia, implying that FH patients may not have

abnormal platelets but rather may have an issue related to the cholesterol environment where platelets operate. Lowering LDL-C levels can temporarily reduce platelet aggregation. However, it is important to note that this evidence may apply primarily to the specific study participants (Wang and Tall, 2016).

## 3.3. Association between atherosclerosis and AD

Vascular risk factors such as cardiovascular disease (Stampfer, 2006), hypertension (Qiu, 2012), diabetes (Wijesinghe et al., 2016), and CVD (Yarchoan et al., 2012) have emerged as important contributors to AD and have been extensively studied (Rius-Perez et al., 2018). Surprisingly, risk factors related to the vascular system can exacerbate the progression of AD (Li et al., 2010). Vascular issues can lead to reduced blood flow in the brain, hindering the clearance of A $\beta$  and potentially playing a significant role in the development of AD (Gupta and Iadecola, 2015). Atherosclerosis, a condition causing vascular problems, represents a crucial target for reducing the risk of AD risk through intervention and early screening (Lopez-Oloriz et al., 2013).

A growing body of evidence suggests a link between AD and atherosclerosis (Roher et al., 2011). Epidemiologic studies have found common vascular risk factors shared by both conditions, including obesity, smoking, older age, diabetes, hypercholesterolemia, hypertension, APOE4 subtype, and hyperhomocysteinemia (Gorelick et al., 2011). A postmortem study of atherosclerotic occlusion of the circle of Willis arteries revealed that AD patients had more stenosis of major brain vessels than did control subjects (Roher et al., 2011). Atherosclerosis may impede the hyperphosphorylation of tau protein and the accumulation of A $\beta$  by reducing cerebral blood flow (Hansson et al., 2018). Further support for this association comes from a meta-analysis showing a higher prevalence of atherosclerosis in AD patients compared to healthy controls (Xie et al., 2020).

Atherosclerosis can lead to cognitive decline and neuronal damage in several ways. The cerebral hypoperfusion and hypoxia associated with atherosclerosis accelerate the overproduction of A $\beta$  in AD models (Iadecola, 2004). In hypoxic conditions, splitting of A $\beta$  proteins from A $\beta$  precursor protein is markedly increased by over-expression of  $\gamma$ - and  $\beta$ -secretases and downregulation of  $\alpha$ -secretase, resulting in increased A $\beta$  production (Salminen et al., 2017). This excess A $\beta$  can harm cerebrovascular function (Iadecola, 2004) and reduce vascular compliance, affecting cerebral blood flow in a pernicious cycle. Atherosclerosis-induced neurovascular disturbance may lead to nitrosative damage and oxidative stress (Chaitanya et al., 2012), which may result in neuroinflammatory responses and metabolic toxicity. Reduced clearance of A $\beta$  coupled with overproduction of A $\beta$  and ROS-induced stress expedites the deposition of A $\beta$ , which eventually leads to cognitive decline (Dore et al., 2013).

## 3.4. Lipid biomarkers in AD

Biomarkers for AD identified through CSF and neuroimaging are primarily confined to scientific research due to high costs, safety concerns, and the need for specialized expertise for clinical application (Sperling et al., 2011). In contrast, blood-based biomarkers, like ApoJ and APOE found in the plasma lipoproteome,






FIGURE 2


Illustration showing L5 and LDL signaling pathways through vascular endothelial cell membrane receptors. Endothelial cells have receptors for oxidized lipoproteins and machinery for lipoprotein receptor-mediated endocytosis. L5 signals are taken up through LOX-1 and PAFR receptors, whereas regular LDL and its subtypes (L1–L4) are internalized by LDLR. LDLR, a transmembrane glycoprotein found on most cell surfaces, governs LDL uptake and its subsequent lysosomal degradation. LDL is transported through endothelial cells using a process called receptor-mediated transcytosis. This transportation involves receptors like SRB1, ALK1, or LDLR as well as a direct transcytosis mechanism. In contrast, signaling through PAFR and LOX-1 triggers PI3K-PKC activation leading to AKT phosphorylation and reduced cAMP levels, while facilitating granule release. LOX-1 receptors are notably abundant in the human aorta, especially in atherosclerosis-prone areas. Exposure of endothelial cells to L5 causes a three-fold increase in membrane-bound LOX-1 levels. Arrows in the figure indicate the direction or influence of signaling pathways. Green arrows represent stimulation or activation, and red arrows with an end bar signify inhibition or suppression. ALK1, activin receptor-like kinase 1; Akt, protein kinase-B; Bcl2, B-cell lymphoma 2 (antiapoptotic protein); Bcl-xL, B-cell lymphoma extra-large (antiapoptotic protein); CRP, C-reactive protein; EC, endothelial cell; eNOS, endothelial NOS; FGF2, fibroblast-growth factor-2 (basic); FGF2R, fibroblast-growth factor receptor-2; FOXO, forkhead box transcription factor subgroup O; L5, electronegative LDL; LDLR, LDL-receptor; LOX-1, lectin like oxidized LDL receptor 1; PAFR, platelet activating factor-receptor; PI3K, phosphoinositide-3-kinase; RAP, LDL receptor-associated protein; SR-B1, scavenger receptor B1; TNF-α, tumor necrosis factor-alpha; TNFR, TNFα receptor; TS20-TS92, LOX-1 neutralizing-antibodies.

offer a cost effective and minimally invasive alternative for early AD detection, making them suitable for widespread clinical use (Sperling et al., 2011). These biomarkers play a crucial role in the progression of AD, making them potentially useful for identifying AD through blood samples. High plasma ApoJ levels and low plasma APOE levels are related to hippocampal atrophy, cerebral amyloidosis, cognitive decline, and dementia, including AD (Song et al., 2012; Gupta et al., 2016). However, it is worth noting that while elevated plasma ApoJ is linked to AD, its diagnostic impact is limited. It improves accuracy only by 8% when combined with factors like APOE genotype, sex, and age (Gupta et al., 2016). Therefore, a methodological approach to enhance the diagnostic value of the plasma lipoproteome should be developed. Fractionating the plasma lipoproteome into distinct classes may reveal unique associations with AD not evident when examining the plasma lipoproteome as a whole. To generate a plasma lipoproteomic workflow, studies involving targeted proteomic methods have been performed based on selected reaction monitoring and liquid chromatography and tandem mass spectrometry as well as relative quantitative peptide analyses (Li et al., 2018).

## 4. Conclusion

AD is a well-recognized neurodegenerative disorder, and its incidence is increasing, positioning it as one of the most impactful diseases alongside cancer and CVD (Brunden et al., 2009). This study provides preliminary evidence to support the innovative idea that quantifying the lipoproteome in the plasma can improve the accuracy of AD diagnosis. Larger scale validation studies are required to test circulating individual lipoproteins in the plasma as potential biomarkers in AD. The current gold standard biomarkers for AD in the CSF, such as tau, phosphorylated tau (p-tau), and Aβ 1–42, require invasive lumbar punctures, making them less suitable for high throughput screening or patient follow-up (Blennow and Hampel, 2003). Thus, exploring blood-based biomarkers, as suggested in our study, holds great promise for improving the accessibility and convenience of AD diagnosis and management in the future.

In this comprehensive review, we have undertaken a detailed exploration of L5. We have shed light on our novel and intriguing hypothesis that L5 may play a pivotal role in initiating



neuroinflammation and oxidative stress, which could ultimately contribute to the development of AD. As our knowledge increases and technologies advance, elucidating the pathophysiological mechanisms of AD becomes more feasible in regard to lipoprotein metabolism, particularly the distinctive role of L5. Future research should prioritize the thorough profiling of L5 and the initiation of AD clinical trials designed to unravel the true risk that L5 poses in neurodegeneration. Moreover, our review helps in providing new tools and insights for the development of innovative diagnostics and treatment strategies for AD and other dementia-related diseases. If a strong correlation between L5 and AD pathophysiology is established, the prospect of anti-L5 therapy and antioxidant therapy emerges as a promising avenue for preventing AD symptoms. Anti-L5 therapy, by reducing plasma L5 and inhibiting the electronegative modification of LDL, could emerge as a vital preventive measure against cognitive impairment in AD, even in the absence of a definitive way to halt AD progression. This represents a beacon of hope in the quest to mitigate the devastating impact of AD on individuals and society as a whole.

## Author contributions

OA: Conceptualization, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. SA: Conceptualization, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. M-CC: Conceptualization, Methodology, Writing – review & editing. SC: Conceptualization,

Methodology, Writing – review & editing. C-KL: Conceptualization, Methodology, Writing – review & editing. SS: Conceptualization, Methodology, Supervision, Writing – review & editing. JCS: Conceptualization, Methodology, Supervision, Writing – review & editing. C-HC: Funding acquisition, Methodology, Resources, Supervision, Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. This study was supported by the Vascular and Medicinal Research Fund. The Texas Heart Institute, #765–64050.

## Acknowledgments

The authors thank Rebecca Bartow, PhD, of The Texas Heart Institute in Houston, Texas, for providing editorial assistance.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Abe, Y., Fornage, M., Yang, C. Y., Bui-Thanh, N. A., Wise, V., Chen, H. H., et al. (2007). L5, the most electronegative subfraction of plasma LDL, induces endothelial vascular cell adhesion molecule 1 and CXC chemokines, which mediate mononuclear leukocyte adhesion. *Atherosclerosis* 192, 56–66. doi: 10.1016/j.atherosclerosis.2006.06.012
- Agostinho, P., Cunha, R. A., and Oliveira, C. (2010). Neuroinflammation, oxidative stress and the pathogenesis of Alzheimer's disease. *Curr. Pharm. Des.* 16, 2766–2778. doi: 10.2174/138161210793176572
- Akyol, O., Chowdhury, I., Akyol, H. R., Tessier, K., Vural, H., and Akyol, S. (2020). Why are cardiovascular diseases more common among patients with severe mental illness? The potential involvement of electronegative low-density lipoprotein (LDL) L5. *Med. Hypotheses* 142:109821. doi: 10.1016/j.mehy.2020.109821
- Akyol, S., Ugur, Z., Yilmaz, A., Ustun, I., Gorti, S. K. K., Oh, K., et al. (2021). Lipid profiling of Alzheimer's disease brain highlights enrichment in glycerol(phospho)lipid, and sphingolipid metabolism. *Cells* 10:2591. doi: 10.3390/cells10102591
- Attems, J., and Jellinger, K. A. (2014). The overlap between vascular disease and Alzheimer's disease—lessons from pathology. *BMC Med.* 12:206. doi: 10.1186/s12916-014-0206-2
- Babiker, A., Andersson, O., Lund, E., Xiu, R. J., Deeb, S., Reshef, A., et al. (1997). Elimination of cholesterol in macrophages and endothelial cells by the sterol 27-hydroxylase mechanism. Comparison with high density lipoprotein-mediated reverse cholesterol transport. *J. Biol. Chem.* 272, 26253–26261. doi: 10.1074/jbc.272.42.26253
- Bancells, C., Sanchez-Quesada, J. L., Birkelund, R., Ordóñez-Llanos, J., and Benítez, S. (2010). Electronegative LDL induces Fas and modifies gene expression in mononuclear cells. *Front. Biosci. (Elite Ed.)* 2, 78–86. doi: 10.2741/e68
- Bellenguez, C., Kucukali, F., Jansen, I. E., Klei, E. D., Moreno-Grau, S., Amin, N., et al. (2022). New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat. Genet.* 54, 412–436. doi: 10.1038/s41588-022-01024-z
- Benítez, S., Camacho, M., Bancells, C., Vila, L., Sanchez-Quesada, J. L., and Ordóñez-Llanos, J. (2006). Wide proinflammatory effect of electronegative low-density lipoprotein on human endothelial cells assayed by a protein array. *Biochim. Biophys. Acta* 1761, 1014–1021. doi: 10.1016/j.bbali.2006.03.020
- Benítez, S., Perez, A., Sanchez-Quesada, J. L., Wagner, A. M., Rigla, M., Arcelus, R., et al. (2007). Electronegative low-density lipoprotein subfraction from type 2 diabetic subjects is proatherogenic and unrelated to glycemic control. *Diabetes Metab. Res. Rev.* 23, 26–34. doi: 10.1002/dmrr.643
- Bjorkhem, I. (2006). Crossing the barrier: oxysterols as cholesterol transporters and metabolic modulators in the brain. *J. Intern. Med.* 260, 493–508. doi: 10.1111/j.1365-2796.2006.01725.x
- Bjorkhem, I., Lutjohann, D., Breuer, O., Sakinis, A., and Wennmalm, A. (1997). Importance of a novel oxidative mechanism for elimination of brain cholesterol. Turnover of cholesterol and 24(S)-hydroxycholesterol in rat brain as measured with 18O<sub>2</sub> techniques in vivo and in vitro. *J. Biol. Chem.* 272, 30178–30184. doi: 10.1074/jbc.272.48.30178
- Blennow, K., and Hampel, H. (2003). CSF markers for incipient Alzheimer's disease. *Lancet Neurol.* 2, 605–613. doi: 10.1016/s1474-4422(03)00530-1
- Broce, I. J., Tan, C. H., Fan, C. C., Jansen, I., Savage, J. E., Witoelar, A., et al. (2019). Dissecting the genetic relationship between cardiovascular risk factors and Alzheimer's disease. *Acta Neuropathol.* 137, 209–226. doi: 10.1007/s00401-018-1928-6
- Brunden, K. R., Trojanowski, J. Q., and Lee, V. M. (2009). Advances in tau-focused drug discovery for Alzheimer's disease and related tauopathies. *Nat. Rev. Drug Discov.* 8, 783–793. doi: 10.1038/nrd2959
- Carmona, S., Hardy, J., and Guerreiro, R. (2018). The genetic landscape of Alzheimer disease. *Handb. Clin. Neurol.* 148, 395–408. doi: 10.1016/B978-0-444-64076-5.00026-0
- Chaitanya, G. V., Cromer, W., Wells, S., Jennings, M., Mathis, J. M., Minagar, A., et al. (2012). Metabolic modulation of cytokine-induced brain endothelial adhesion molecule expression. *Microcirculation* 19, 155–165. doi: 10.1111/j.1549-8719.2011.00141.x
- Chen, W. Y., Chen, F. Y., Lee, A. S., Ting, K. H., Chang, C. M., Hsu, J. F., et al. (2015). Sesamol reduces the atherogenicity of electronegative L5 LDL in vivo and in vitro. *J. Nat. Prod.* 78, 225–233. doi: 10.1021/np500700z
- Chen, Y., Yin, M., Cao, X., Hu, G., and Xiao, M. (2018). Pro- and anti-inflammatory effects of high cholesterol diet on aged brain. *Aging Dis.* 9, 374–390. doi: 10.14338/AD.2017.0706
- Choi, S. H., Wiesner, P., Almazan, F., Kim, J., and Miller, Y. I. (2012). Spleen tyrosine kinase regulates AP-1 dependent transcriptional response to minimally oxidized LDL. *PLoS One* 7:e32378. doi: 10.1371/journal.pone.0032378
- Chu, C. S., Wang, Y. C., Lu, L. S., Walton, B., Yilmaz, H. R., Huang, R. Y., et al. (2013). Electronegative low-density lipoprotein increases C-reactive protein expression in vascular endothelial cells through the LOX-1 receptor. *PLoS One* 8:e70533. doi: 10.1371/journal.pone.0070533
- Citron, M. (2010). Alzheimer's disease: strategies for disease modification. *Nat. Rev. Drug Discov.* 9, 387–398. doi: 10.1038/nrd2896
- de Castellarnau, C., Bancells, C., Benítez, S., Reina, M., Ordóñez-Llanos, J., and Sanchez-Quesada, J. L. (2007). Atherogenic and inflammatory profile of human arterial endothelial cells (HUAEC) in response to LDL subfractions. *Clin. Chim. Acta* 376, 233–236. doi: 10.1016/j.cca.2006.07.024
- De Castellarnau, C., Sanchez-Quesada, J. L., Benítez, S., Rosa, R., Caveda, L., Vila, L., et al. (2000). Electronegative LDL from normolipemic subjects induces IL-8 and monocyte chemotactic protein secretion by human endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* 20, 2281–2287. doi: 10.1161/01.atv.20.10.2281
- Deane, R., Wu, Z., Sagare, A., Davis, J., Du Yan, S., Hamm, K., et al. (2004). LRP/amyloid beta-peptide interaction mediates differential brain efflux of Abeta isoforms. *Neuron* 43, 333–344. doi: 10.1016/j.neuron.2004.07.017
- Desikan, R. S., Fan, C. C., Wang, Y., Schork, A. J., Cabral, H. J., Cupples, L. A., et al. (2017). Genetic assessment of age-associated Alzheimer disease risk: development and validation of a polygenic hazard score. *PLoS Med.* 14:e1002258. doi: 10.1371/journal.pmed.1002258
- Dildar, K., Sinem, F., Gokhan, E., Orhan, Y., and Filiz, M. (2010). Serum nitrosative stress levels are increased in Alzheimer disease but not in vascular dementia. *Alzheimer Dis. Assoc. Disord.* 24, 194–197. doi: 10.1097/WAD.0b013e3181c53d0a
- Dore, V., Villemagne, V. L., Bourgeat, P., Fripp, J., Acosta, O., Chételat, G., et al. (2013). Cross-sectional and longitudinal analysis of the relationship between A $\beta$  deposition, cortical thickness, and memory in cognitively unimpaired individuals and in Alzheimer disease. *JAMA Neurol.* 70, 903–911. doi: 10.1001/jamaneurol.2013.1062
- Duong, M. T., Nasrallah, I. M., Wolk, D. A., Chang, C. C. Y., and Chang, T. Y. (2021). Cholesterol, atherosclerosis, and APOE in vascular contributions to cognitive impairment and dementia (VCID): potential mechanisms and therapy. *Front. Aging Neurosci.* 13:647990. doi: 10.3389/fnagi.2021.647990
- Edwards Iii, G. A., Gamez, N., Escobedo, G., Calderon, O., and Moreno-Gonzalez, I. (2019). Modifiable risk factors for Alzheimer's disease. *Front. Aging Neurosci.* 11:146. doi: 10.3389/fnagi.2019.00146
- Escargueil-Blanc, I., Salvayre, R., and Negre-Salvayre, A. (1994). Necrosis and apoptosis induced by oxidized low density lipoproteins occur through two calcium-dependent pathways in lymphoblastoid cells. *FASEB J.* 8, 1075–1080. doi: 10.1096/fasebj.8.13.7926374
- Estruch, M., Bancells, C., Beloki, L., Sanchez-Quesada, J. L., Ordóñez-Llanos, J., and Benítez, S. (2013). CD14 and TLR4 mediate cytokine release promoted by electronegative LDL in monocytes. *Atherosclerosis* 229, 356–362. doi: 10.1016/j.atherosclerosis.2013.05.011
- Estruch, M., Rajamaki, K., Sanchez-Quesada, J. L., Kovanen, P. T., Oorni, K., Benítez, S., et al. (2015). Electronegative LDL induces priming and inflammasome activation leading to IL-1 $\beta$  release in human monocytes and macrophages. *Biochim. Biophys. Acta* 1851, 1442–1449. doi: 10.1016/j.bbali.2015.08.009
- Estruch, M., Sanchez-Quesada, J. L., Ordóñez-Llanos, J., and Benítez, S. (2014). Ceramide-enriched LDL induces cytokine release through TLR4 and CD14 in monocytes. Similarities with electronegative LDL. *Clin. Investig. Arterioscler.* 26, 131–137. doi: 10.1016/j.arteri.2013.12.003
- Estruch, M., Sanchez-Quesada, J. L., Ordóñez-Llanos, J., and Benítez, S. (2016). Inflammatory intracellular pathways activated by electronegative LDL in monocytes. *Biochim. Biophys. Acta* 1861, 963–969. doi: 10.1016/j.bbali.2016.05.010
- Fitzpatrick, A. L., Kuller, L. H., Ives, D. G., Lopez, O. L., Jagust, W., Breitner, J. C., et al. (2004). Incidence and prevalence of dementia in the cardiovascular health study. *J. Am. Geriatr. Soc.* 52, 195–204. doi: 10.1111/j.1532-5415.2004.52058.x
- Gorelick, P. B., Scuteri, A., Black, S. E., Decarli, C., Greenberg, S. M., Iadecola, C., et al. (2011). Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the American heart association/American stroke association. *Stroke* 42, 2672–2713. doi: 10.1161/STR.0b013e3182299496
- Grimm, M. O., Mett, J., Grimm, H. S., and Hartmann, T. (2017). APP function and lipids: a bidirectional link. *Front. Mol. Neurosci.* 10:63. doi: 10.3389/fnmol.2017.00063
- Grossi, M. F., Carvalho, M. D. G., Silveira, J. N., Gonçalves, G. S., Gomes, K. B., Bicalho, M. A., et al. (2018). OxLDL plasma levels in patients with Alzheimer's disease. *Arq. Neuropsiquiatr.* 76, 241–246. doi: 10.1590/0004-282x20180012

- Gudala, K., Bansal, D., Schifano, F., and Bhansali, A. (2013). Diabetes mellitus and risk of dementia: a meta-analysis of prospective observational studies. *J. Diabet. Investig.* 4, 640–650. doi: 10.1111/jdi.12087
- Gupta, V. B., Doecke, J. D., Hone, E., Pedrini, S., Laws, S. M., Thambisetty, M., et al. (2016). Plasma apolipoprotein J as a potential biomarker for Alzheimer's disease: Australian imaging, biomarkers and lifestyle study of aging. *Alzheimers Dement.* 3, 18–26. doi: 10.1016/j.dadm.2015.12.001
- Gupta, A., and Iadecola, C. (2015). Impaired A $\beta$  clearance: a potential link between atherosclerosis and Alzheimer's disease. *Front. Aging Neurosci.* 7:115. doi: 10.3389/fnagi.2015.00115
- Hansson, O., Palmqvist, S., Ljung, H., Cronberg, T., van Westen, D., and Smith, R. (2018). Cerebral hypoperfusion is not associated with an increase in amyloid  $\beta$  pathology in middle-aged or elderly people. *Alzheimers Dement.* 14, 54–61. doi: 10.1016/j.jalz.2017.06.2265
- Helbecque, N., Berr, C., Cottel, D., Fromentin-David, I., Sazdovitch, V., Ricolfi, F., et al. (2001). VLDL receptor polymorphism, cognitive impairment, and dementia. *Neurology* 56, 1183–1188. doi: 10.1212/wnl.56.9.1183
- Heverin, M., Meaney, S., Lutjohann, D., Diczfalusy, U., Wahren, J., and Bjorkhem, I. (2005). Crossing the barrier: net flux of 27-hydroxycholesterol into the human brain. *J. Lipid Res.* 46, 1047–1052. doi: 10.1194/jlr.M500024-JLR200
- Hottman, D. A., Chernick, D., Cheng, S., Wang, Z., and Li, L. (2014). HDL and cognition in neurodegenerative disorders. *Neurobiol. Dis.* 72, 22–36. doi: 10.1016/j.nbd.2014.07.015
- Iadecola, C. (2004). Neurovascular regulation in the normal brain and in Alzheimer's disease. *Nat. Rev. Neurosci.* 5, 347–360. doi: 10.1038/nrn1387
- Janabi, M., Yamashita, S., Hirano, K., Sakai, N., Hiraoka, H., Matsumoto, K., et al. (2000). Oxidized LDL-induced NF-kappa B activation and subsequent expression of proinflammatory genes are defective in monocyte-derived macrophages from CD36-deficient patients. *Arterioscler. Thromb. Vasc. Biol.* 20, 1953–1960. doi: 10.1161/01.atv.20.8.1953
- Kang, D. E., Pietrzik, C. U., Baum, L., Chevallier, N., Merriam, D. E., Kounnas, M. Z., et al. (2000). Modulation of amyloid beta-protein clearance and Alzheimer's disease susceptibility by the LDL receptor-related protein pathway. *J. Clin. Invest.* 106, 1159–1166. doi: 10.1172/JCI11013
- Kankaanpaa, J., Turunen, S. P., Moilanen, V., Horkko, S., and Remes, A. M. (2009). Cerebrospinal fluid antibodies to oxidized LDL are increased in Alzheimer's disease. *Neurobiol. Dis.* 33, 467–472. doi: 10.1016/j.nbd.2008.12.001
- Kemppainen, N., Johansson, J., Teuvo, J., Parkkola, R., Joutsa, J., Ngandu, T., et al. (2018). Brain amyloid load and its associations with cognition and vascular risk factors in FINGER study. *Neurology* 90, e206–e213. doi: 10.1212/WNL.0000000000004827
- Kim, W. S., Li, H., Ruberu, K., Chan, S., Elliott, D. A., Low, J. K., et al. (2013). Deletion of Abca7 increases cerebral amyloid- $\beta$  accumulation in the J20 mouse model of Alzheimer's disease. *J. Neurosci.* 33, 4387–4394. doi: 10.1523/JNEUROSCI.4165-12.2013
- Koffie, R. M., Hashimoto, T., Tai, H. C., Kay, K. R., Serrano-Pozo, A., Joyner, D., et al. (2012). Apolipoprotein E4 effects in Alzheimer's disease are mediated by synaptotoxic oligomeric amyloid- $\beta$ . *Brain* 135, 2155–2168. doi: 10.1093/brain/aws127
- Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C., et al. (2013). Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat. Genet.* 45, 1452–1458. doi: 10.1038/ng.2802
- Lambert, J. C., Luedeking-Zimmer, E., Merrot, S., Hayes, A., Thaker, U., Desai, P., et al. (2003). Association of 3'-UTR polymorphisms of the oxidized LDL receptor 1 (OLR1) gene with Alzheimer's disease. *J. Med. Genet.* 40, 424–430. doi: 10.1136/jmg.40.6.424
- Launer, L. J., White, L. R., Petrovitch, H., Ross, G. W., and Curb, J. D. (2001). Cholesterol and neuropathologic markers of AD: a population-based autopsy study. *Neurology* 57, 1447–1452. doi: 10.1212/wnl.57.8.1447
- Lee, J. H., Cheng, R., Rogaeva, E., Meng, Y., Stern, Y., Santana, V., et al. (2008). Further examination of the candidate genes in chromosome 12p13 locus for late-onset Alzheimer disease. *Neurogenetics* 9, 127–138. doi: 10.1007/s10048-008-0122-8
- Lee, C. Y., and Landreth, G. E. (2010). The role of microglia in amyloid clearance from the AD brain. *J. Neural Transm.* 117, 949–960. doi: 10.1007/s00702-010-0433-4
- Leoni, V., Masterman, T., Patel, P., Meaney, S., Diczfalusy, U., and Bjorkhem, I. (2003). Side chain oxidized oxysterols in cerebrospinal fluid and the integrity of blood-brain and blood-cerebrospinal fluid barriers. *J. Lipid Res.* 44, 793–799. doi: 10.1194/jlr.M200434-JLR200
- Li, D., Huang, F., Zhao, Y., Villata, P. W., Griffin, T. J., Zhang, L., et al. (2018). Plasma lipoproteome in Alzheimer's disease: a proof-of-concept study. *Clin. Proteomics* 15:31. doi: 10.1186/s12014-018-9207-z
- Li, D., Misialek, J. R., Boerwinkle, E., Gottesman, R. F., Sharrett, A. R., Mosley, T. H., et al. (2016). Plasma phospholipids and prevalence of mild cognitive impairment and/or dementia in the ARIC neurocognitive study (ARIC-NCS). *Alzheimers Dement.* 3, 73–82. doi: 10.1016/j.dadm.2016.02.008
- Li, J., Zhang, M., Xu, Z. Q., Gao, C. Y., Fang, C. Q., Deng, J., et al. (2010). Vascular risk aggravates the progression of Alzheimer's disease in a Chinese cohort. *J. Alzheimers Dis.* 20, 491–500. doi: 10.3233/JAD-2010-1383
- Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D., et al. (2017). Dementia prevention, intervention, and care. *Lancet* 390, 2673–2734. doi: 10.1016/S0140-6736(17)31363-6
- Lopez-Oloriz, J., Lopez-Cancio, E., Arenillas, J. F., Hernandez, M., Jimenez, M., Dorado, L., et al. (2013). Asymptomatic cervicocerebral atherosclerosis, intracranial vascular resistance and cognition: the AsIA-neuropsychology study. *Atherosclerosis* 230, 330–335. doi: 10.1016/j.atherosclerosis.2013.08.011
- Loving, B. A., Tang, M., Neal, M. C., Gorkhali, S., Murphy, R., Eckel, R. H., et al. (2021). Lipoprotein lipase regulates microglial lipid droplet accumulation. *Cells* 10:198. doi: 10.3390/cells10020198
- Lu, J., Yang, J. H., Burns, A. R., Chen, H. H., Tang, D., Walterscheid, J. P., et al. (2009). Mediation of electronegative low-density lipoprotein signaling by LOX-1: a possible mechanism of endothelial apoptosis. *Circ. Res.* 104, 619–627. doi: 10.1161/CIRCRESAHA.108.190116
- Lund, E. G., Guileyardo, J. M., and Russell, D. W. (1999). cDNA cloning of cholesterol 24-hydroxylase, a mediator of cholesterol homeostasis in the brain. *Proc. Natl. Acad. Sci. U. S. A.* 96, 7238–7243. doi: 10.1073/pnas.96.13.7238
- Luth, H. J., Holzer, M., Gartner, U., Staufenbiel, M., and Arendt, T. (2001). Expression of endothelial and inducible NOS-isoforms is increased in Alzheimer's disease, in APP23 transgenic mice and after experimental brain lesion in rat: evidence for an induction by amyloid pathology. *Brain Res.* 913, 57–67. doi: 10.1016/S0006-8993(01)02758-5
- Mahley, R. W. (2016). Central nervous system lipoproteins: ApoE and regulation of cholesterol metabolism. *Arterioscler. Thromb. Vasc. Biol.* 36, 1305–1315. doi: 10.1161/ATVBAHA.116.307023
- Mapstone, M., Cheema, A. K., Fiandaca, M. S., Zhong, X., Myhre, T. R., MacArthur, L. H., et al. (2014). Plasma phospholipids identify antecedent memory impairment in older adults. *Nat. Med.* 20, 415–418. doi: 10.1038/nm.3466
- Meaney, S., Heverin, M., Panzenboeck, U., Ekstrom, L., Axelsson, M., Andersson, U., et al. (2007). Novel route for elimination of brain oxysterols across the blood-brain barrier: conversion into 7 $\alpha$ -hydroxy-3-oxo-4-cholestenoic acid. *J. Lipid Res.* 48, 944–951. doi: 10.1194/jlr.M600529-JLR200
- Mehta, J. L., Chen, J., Hermonat, P. L., Romeo, F., and Novelli, G. (2006). Lectin-like, oxidized low-density lipoprotein receptor-1 (LOX-1): a critical player in the development of atherosclerosis and related disorders. *Cardiovasc. Res.* 69, 36–45. doi: 10.1016/j.cardiores.2005.09.006
- Miller, G. J., and Miller, N. E. (1975). Plasma-high-density-lipoprotein concentration and development of ischaemic heart-disease. *Lancet* 1, 16–19. doi: 10.1016/S0140-6736(75)92376-4
- Montagne, A., Barnes, S. R., Sweeney, M. D., Halliday, M. R., Sagare, A. P., Zhao, Z., et al. (2015). Blood-brain barrier breakdown in the aging human hippocampus. *Neuron* 85, 296–302. doi: 10.1016/j.neuron.2014.12.032
- Munoz Herrera, O. M., and Zivkovic, A. M. (2022). Microglia and cholesterol handling: implications for Alzheimer's disease. *Biomedicine* 10:3105. doi: 10.3390/biomedicines10123105
- Namgaladze, D., Kollas, A., and Brune, B. (2008). Oxidized LDL attenuates apoptosis in monocytic cells by activating ERK signaling. *J. Lipid Res.* 49, 58–65. doi: 10.1194/jlr.M700100-JLR200
- National Academies of Sciences and Medicine. (2017). *Preventing cognitive decline and dementia: A way forward*, Washington (DC): National Academies of Sciences and Medicine (2017).
- Norton, S., Matthews, F. E., Barnes, D. E., Yaffe, K., and Brayne, C. (2014). Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *Lancet Neurol.* 13, 788–794. doi: 10.1016/S1474-4422(14)70136-X
- Oresic, M., Hyotylainen, T., Herukka, S. K., Sysi-Aho, M., Mattila, I., Seppanan-Laakso, T., et al. (2011). Metabolome in progression to Alzheimer's disease. *Trans. Psychiatry* 1:e57. doi: 10.1038/tp.2011.55
- Palmieri, V. O., Coppola, B., Grattagliano, I., Casieri, V., Cardinale, G., Portincasa, P., et al. (2013). Oxidized LDL receptor 1 gene polymorphism in patients with metabolic syndrome. *Eur. J. Clin. Invest.* 43, 41–48. doi: 10.1111/eci.12013
- Perry, G., Taddeo, M. A., Nunomura, A., Zhu, X., Zenteno-Savin, T., Drew, K. L., et al. (2002). Comparative biology and pathology of oxidative stress in Alzheimer and other neurodegenerative diseases: beyond damage and response. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 133, 507–513. doi: 10.1016/S1532-0456(02)00119-9
- Pflanzner, T., Janko, M. C., Andre-Dohmen, B., Reuss, S., Weggen, S., Roebroek, A. J., et al. (2011). LRP1 mediates bidirectional transcytosis of amyloid- $\beta$  across the blood-brain barrier. *Neurobiol. Aging* 32, e2321–e2311. doi: 10.1016/j.neurobiolaging.2010.05.025
- Pietrzik, C. U., Busse, T., Merriam, D. E., Weggen, S., and Koo, E. H. (2002). The cytoplasmic domain of the LDL receptor-related protein regulates multiple steps in APP processing. *EMBO J.* 21, 5691–5700. doi: 10.1093/emboj/cdf568
- Qiu, C. (2012). Preventing Alzheimer's disease by targeting vascular risk factors: hope and gap. *J. Alzheimers Dis.* 32, 721–731. doi: 10.3233/JAD-2012-120922
- Ramanathan, A., Nelson, A. R., Sagare, A. P., and Zlokovic, B. V. (2015). Impaired vascular-mediated clearance of brain amyloid beta in Alzheimer's disease: the role, regulation and restoration of LRP1. *Front. Aging Neurosci.* 7:136. doi: 10.3389/fnagi.2015.00136
- Reitz, C. (2013). Dyslipidemia and the risk of Alzheimer's disease. *Curr. Atheroscler. Rep.* 15:307. doi: 10.1007/s11883-012-0307-3

- Rios, F. J., Koga, M. M., Ferracini, M., and Jancar, S. (2012). Co-stimulation of PAFR and CD36 is required for oxLDL-induced human macrophages activation. *PLoS One* 7:e36632. doi: 10.1371/journal.pone.0036632
- Rius-Perez, S., Tormos, A. M., Perez, S., and Talens-Visconti, R. (2018). Vascular pathology: cause or effect in Alzheimer disease? *Neurologia* 33, 112–120. doi: 10.1016/j.nrl.2015.07.010
- Roher, A. E., Tyas, S. L., Maarouf, C. L., Daus, I. D., Kokjohn, T. A., Emmerling, M. R., et al. (2011). Intracranial atherosclerosis as a contributing factor to Alzheimer's disease dementia. *Alzheimers Dement.* 7, 436–444. doi: 10.1016/j.jalz.2010.08.228
- Rushworth, J. V., Griffiths, H. H., Watt, N. T., and Hooper, N. M. (2013). Prion protein-mediated toxicity of amyloid- $\beta$  oligomers requires lipid rafts and the transmembrane LRP1. *J. Biol. Chem.* 288, 8935–8951. doi: 10.1074/jbc.M112.400358
- Sagare, A. P., Bell, R. D., Zhao, Z., Ma, Q., Winkler, E. A., Ramanathan, A., et al. (2013). Pericyte loss influences Alzheimer-like neurodegeneration in mice. *Nat. Commun.* 4:2932. doi: 10.1038/ncomms3932
- Sagare, A., Deane, R., Bell, R. D., Johnson, B., Hamm, K., Pendu, R., et al. (2007). Clearance of amyloid-beta by circulating lipoprotein receptors. *Nat. Med.* 13, 1029–1031. doi: 10.1038/nm1635
- Salminen, A., Kauppinen, A., and Kaarniranta, K. (2017). Hypoxia/ischemia activate processing of amyloid precursor protein: impact of vascular dysfunction in the pathogenesis of Alzheimer's disease. *J. Neurochem.* 140, 536–549. doi: 10.1111/jnc.13932
- Sanchez-Quesada, J. L., Camacho, M., Anton, R., Benitez, S., Vila, L., and Ordonez-Llanos, J. (2003). Electronegative LDL of FH subjects: chemical characterization and induction of chemokine release from human endothelial cells. *Atherosclerosis* 166, 261–270. doi: 10.1016/s0021-9150(02)00374-x
- Sanchez-Quesada, J. L., Villegas, S., and Ordonez-Llanos, J. (2012). Electronegative low-density lipoprotein. A link between apolipoprotein B misfolding, lipoprotein aggregation and proteoglycan binding. *Curr. Opin. Lipidol.* 23, 479–486. doi: 10.1097/MOL.0b013e328357c933
- Sandebing-Matton, A., Goikolea, J., Bjorkhem, I., Paternain, L., Kemppainen, N., Laatikainen, T., et al. (2021). 27-hydroxycholesterol, cognition, and brain imaging markers in the FINGER randomized controlled trial. *Alzheimers Res. Ther.* 13:56. doi: 10.1186/s13195-021-00790-y
- Sawamura, T., Kume, N., Aoyama, T., Moriwaki, H., Hoshikawa, H., Aiba, Y., et al. (1997). An endothelial receptor for oxidized low-density lipoprotein. *Nature* 386, 73–77. doi: 10.1038/386073a0
- Schneider, A., Schulz-Schaeffer, W., Hartmann, T., Schulz, J. B., and Simons, M. (2006). Cholesterol depletion reduces aggregation of amyloid-beta peptide in hippocampal neurons. *Neurobiol. Dis.* 23, 573–577. doi: 10.1016/j.nbd.2006.04.015
- Shi, J., Tian, J., Pritchard, A., Lendon, C., Lambert, J. C., Iwatsubo, T., et al. (2006). A 3'-UTR polymorphism in the oxidized LDL receptor 1 gene increases Abeta40 load as cerebral amyloid angiopathy in Alzheimer's disease. *Acta Neuropathol.* 111, 15–20. doi: 10.1007/s00401-005-1108-3
- Shinohara, M., Tachibana, M., Kanekiyo, T., and Bu, G. (2017). Role of LRP1 in the pathogenesis of Alzheimer's disease: evidence from clinical and preclinical studies. *J. Lipid Res.* 58, 1267–1281. doi: 10.1194/jlr.R075796
- Sinem, F., Dildar, K., Gokhan, E., Melda, B., Orhan, Y., and Filiz, M. (2010). The serum protein and lipid oxidation marker levels in Alzheimer's disease and effects of cholinesterase inhibitors and antipsychotic drugs therapy. *Curr. Alzheimer Res.* 7, 463–469. doi: 10.2174/156720510791383822
- Song, F., Poljak, A., Crawford, J., Kochan, N. A., Wen, W., Cameron, B., et al. (2012). Plasma apolipoprotein levels are associated with cognitive status and decline in a community cohort of older individuals. *PLoS One* 7:e34078. doi: 10.1371/journal.pone.0034078
- Sperling, R. A., Aisen, P. S., Beckett, L. A., Bennett, D. A., Craft, S., Fagan, A. M., et al. (2011). Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 7, 280–292. doi: 10.1016/j.jalz.2011.03.003
- Staffaroni, A. M., Elahi, F. M., McDermott, D., Marton, K., Karageorgiou, E., Sacco, S., et al. (2017). Neuroimaging in dementia. *Semin. Neurol.* 37, 510–537. doi: 10.1055/s-0037-1608808
- Stampfer, M. J. (2006). Cardiovascular disease and Alzheimer's disease: common links. *J. Intern. Med.* 260, 211–223. doi: 10.1111/j.1365-2796.2006.01687.x
- Tai, M. H., Kuo, S. M., Liang, H. T., Chiou, K. R., Lam, H. C., Hsu, C. M., et al. (2006). Modulation of angiogenic processes in cultured endothelial cells by low density lipoproteins subfractions from patients with familial hypercholesterolemia. *Atherosclerosis* 186, 448–457. doi: 10.1016/j.atherosclerosis.2005.08.022
- Trushina, E., Dutta, T., Persson, X. M., Mielke, M. M., and Petersen, R. C. (2013). Identification of altered metabolic pathways in plasma and CSF in mild cognitive impairment and Alzheimer's disease using metabolomics. *PLoS One* 8:e63644. doi: 10.1371/journal.pone.0063644
- Tynkkynen, J., Chouraki, V., van der Lee, S. J., Hernesniemi, J., Yang, Q., Li, S., et al. (2018). Association of branched-chain amino acids and other circulating metabolites with risk of incident dementia and Alzheimer's disease: a prospective study in eight cohorts. *Alzheimers Dement.* 14, 723–733. doi: 10.1016/j.jalz.2018.01.003
- van de Haar, H. J., Burgmans, S., Jansen, J. F., van Osch, M. J., van Buchem, M. A., Muller, M., et al. (2016). Blood-brain barrier leakage in patients with early Alzheimer disease. *Radiology* 281, 527–535. doi: 10.1148/radiol.2016152244
- Wanamaker, B. L., Swiger, K. J., Blumenthal, R. S., and Martin, S. S. (2015). Cholesterol, statins, and dementia: what the cardiologist should know. *Clin. Cardiol.* 38, 243–250. doi: 10.1002/clc.22361
- Wang, J. Y., Lai, C. L., Lee, C. T., and Lin, C. Y. (2017). Electronegative low-density lipoprotein L5 impairs viability and NGF-induced neuronal differentiation of PC12 cells via LOX-1. *Int. J. Mol. Sci.* 18:1744. doi: 10.3390/ijms18081744
- Wang, N., and Tall, A. R. (2016). Cholesterol in platelet biogenesis and activation. *Blood* 127, 1949–1953. doi: 10.1182/blood-2016-01-631259
- Wijesinghe, P., Shankar, S. K., Yasha, T. C., Gorrie, C., Amarantunga, D., Hulathduwa, S., et al. (2016). Vascular contributions in Alzheimer's disease-related neuropathological changes: first autopsy evidence from a south Asian aging population. *J. Alzheimers Dis.* 54, 1607–1618. doi: 10.3233/JAD-160425
- Wingo, T. S., Cutler, D. J., Wingo, A. P., Le, N. A., Rabinovici, G. D., Miller, B. L., et al. (2019). Association of early-onset Alzheimer disease with elevated low-density lipoprotein cholesterol levels and rare genetic coding variants of APOB. *JAMA Neurol.* 76, 809–817. doi: 10.1001/jamaneurol.2019.0648
- Wu, W. S., Holmans, P., Wavrant-DeVrieze, F., Shears, S., Kehoe, P., Crook, R., et al. (1998). Genetic studies on chromosome 12 in late-onset Alzheimer disease. *JAMA* 280, 619–622. doi: 10.1001/jama.280.7.619
- Wurtz, P., Havulinna, A. S., Soininen, P., Tynkkynen, T., Prieto-Merino, D., Tillin, T., et al. (2015). Metabolite profiling and cardiovascular event risk: a prospective study of 3 population-based cohorts. *Circulation* 131, 774–785. doi: 10.1161/CIRCULATIONAHA.114.013116
- Xie, B., Shi, X., Xing, Y., and Tang, Y. (2020). Association between atherosclerosis and Alzheimer's disease: a systematic review and meta-analysis. *Brain Behav.* 10:e01601. doi: 10.1002/brb3.1601
- Xiong, H., Callaghan, D., Jones, A., Walker, D. G., Lue, L. F., Beach, T. G., et al. (2008). Cholesterol retention in Alzheimer's brain is responsible for high beta- and gamma-secretase activities and Abeta production. *Neurobiol. Dis.* 29, 422–437. doi: 10.1016/j.nbd.2007.10.005
- Xue-Shan, Z., Juan, P., Qi, W., Zhong, R., Li-Hong, P., Zhi-Han, T., et al. (2016). Imbalanced cholesterol metabolism in Alzheimer's disease. *Clin. Chim. Acta* 456, 107–114. doi: 10.1016/j.cca.2016.02.024
- Yamanaka, S., Zhang, X. Y., Miura, K., Kim, S., and Iwao, H. (1998). The human gene encoding the lectin-type oxidized LDL receptor (OLR1) is a novel member of the natural killer gene complex with a unique expression profile. *Genomics* 54, 191–199. doi: 10.1006/geno.1998.5561
- Yang, C. Y., Raya, J. L., Chen, H. H., Chen, C. H., Abe, Y., Pownall, H. J., et al. (2003). Isolation, characterization, and functional assessment of oxidatively modified subfractions of circulating low-density lipoproteins. *Arterioscler. Thromb. Vasc. Biol.* 23, 1083–1090. doi: 10.1161/01.ATV.0000071350.78872.C4
- Yarchoan, M., Xie, S. X., Kling, M. A., Toledo, J. B., Wolk, D. A., Lee, E. B., et al. (2012). Cerebrovascular atherosclerosis correlates with Alzheimer pathology in neurodegenerative dementias. *Brain* 135, 3749–3756. doi: 10.1093/brain/aw271
- Yu, L. E., Lai, C. L., Lee, C. T., and Wang, J. Y. (2017). Highly electronegative low-density lipoprotein L5 evokes microglial activation and creates a neuroinflammatory stress via toll-like receptor 4 signaling. *J. Neurochem.* 142, 231–245. doi: 10.1111/jnc.14053
- Zafriilla, P., Mulero, J., Xandri, J. M., Santo, E., Caravaca, G., and Morillas, J. M. (2006). Oxidative stress in Alzheimer patients in different stages of the disease. *Curr. Med. Chem.* 13, 1075–1083. doi: 10.2174/092986706776360978
- Zhao, Z., Zhou, H., Peng, Y., Qiu, C. H., Sun, Q. Y., Wang, F., et al. (2014). Expression and significance of plasma 3-NT and ox-LDL in patients with Alzheimer's disease. *Genet. Mol. Res.* 13, 8428–8435. doi: 10.4238/2014.October.20.19
- Zhu, H., Bhadelia, R. A., Liu, Z., Vu, L., Li, H., Scott, T., et al. (2013). The association between small vessel infarcts and the activities of amyloid- $\beta$  peptide degrading proteases in apolipoprotein E4 allele carriers. *Angiology* 64, 614–620. doi: 10.1177/0003319712462125
- Ziuzenkova, O., Asatryan, L., Sahady, D., Orasanu, G., Perrey, S., Cutak, B., et al. (2003). Dual roles for lipolysis and oxidation in peroxisome proliferation-activator receptor responses to electronegative low density lipoprotein. *J. Biol. Chem.* 278, 39874–39881. doi: 10.1074/jbc.M306786200
- Zlokovic, B. V. (2011). Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat. Rev. Neurosci.* 12, 723–738. doi: 10.1038/nrn3114