



UNIVERSITY OF  
PLYMOUTH



School of Geography, Earth and Environmental Sciences  
Faculty of Science and Engineering

2024-03-08

## Worldwide Accumulation of Atmospheric Mercury in Glacier Cryoconite

Agnieszka Pasieka

Kamil Brudecki

Przemysław Niedzielski

Alexandra Proch

Nozomu Takeuchi

et al. *See next page for additional authors*

*Let us know how access to this document benefits you*



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

### General rights

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

### Take down policy

If you believe that this document breaches copyright please [contact the library](#) providing details, and we will remove access to the work immediately and investigate your claim.

Follow this and additional works at: <https://pearl.plymouth.ac.uk/gees-research>

---

### Recommended Citation

Pasieka, A., Brudecki, K., Niedzielski, P., Proch, A., Takeuchi, N., Ambrosini, R., Owens, P., Zawierucha, K., Baccolo, G., Clason, C., Beard, D., Yde, J., & Łokas, E. (2024) 'Worldwide Accumulation of Atmospheric Mercury in Glacier Cryoconite', Available at: <https://doi.org/10.5194/egusphere-egu24-8179>

This Conference Proceeding is brought to you for free and open access by the Faculty of Science and Engineering at PEARL. It has been accepted for inclusion in School of Geography, Earth and Environmental Sciences by an authorized administrator of PEARL. For more information, please contact [openresearch@plymouth.ac.uk](mailto:openresearch@plymouth.ac.uk).

---

## Authors

Agnieszka Pasieka, Kamil Brudecki, Przemysław Niedzielski, Alexandra Proch, Nozomu Takeuchi, Roberto Ambrosini, Philip N. Owens, K Zawierucha, Giovanni Baccolo, Caroline Clason, Dylan Beard, Jacob Clement Yde, and Edyta Łokas



PEARL

### **Worldwide Accumulation of Atmospheric Mercury in Glacier Cryoconite**

Pasieka, Agnieszka; Brudecki, Kamil; Niedzielski, Przemysław; Proch, Alexandra; Takeuchi, Nozomu; Ambrosini, Roberto; Owens, Philip N.; Zawierucha, K; Baccolo, Giovanni; Clason, Caroline; Beard, Dylan; Yde, Jacob Clement; Łokas, Edyta

**DOI:**

[10.5194/egusphere-egu24-8179](https://doi.org/10.5194/egusphere-egu24-8179)

**Publication date:**

2024

**Document version:**

Publisher's PDF, also known as Version of record

**Link:**

[Link to publication in PEARL](#)

**Citation for published version (APA):**

Pasieka, A., Brudecki, K., Niedzielski, P., Proch, A., Takeuchi, N., Ambrosini, R., Owens, P. N., Zawierucha, K., Baccolo, G., Clason, C., Beard, D., Yde, J. C., & Łokas, E. (2024).

*Worldwide Accumulation of Atmospheric Mercury in Glacier Cryoconite*. Abstract from EGU General Assembly 2024, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu24-8179>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

EGU24-8179, updated on 04 Jun 2024

<https://doi.org/10.5194/egusphere-egu24-8179>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



## Worldwide Accumulation of Atmospheric Mercury in Glacier Cryoconite

**Agnieszka Pasieka**<sup>1,2</sup>, Kamil Brudecki<sup>2</sup>, Przemysław Niedzielski<sup>3</sup>, Aleksandra Proch<sup>3</sup>, Nozomu Takeuchi<sup>4</sup>, Roberto Ambrosini<sup>5</sup>, Phil Owens<sup>6</sup>, Krzysztof Zawierucha<sup>7</sup>, Giovanni Baccolo<sup>8,9</sup>, Caroline Clason<sup>10</sup>, Dylan Beard<sup>11</sup>, Jacob Clement Yde<sup>12</sup>, and Edyta Łokas<sup>2</sup>

<sup>1</sup>Department of Radiation Protection, Institute of Nuclear Physics, Polish Academy of Sciences, Krakow, Poland (agnieszka.pasieka@ifj.edu.pl)

<sup>2</sup>Department of Mass Spectrometry, Institute of Nuclear Physics, Polish Academy of Sciences, Krakow, Poland

<sup>3</sup>Department of Analytical Chemistry, Faculty of Chemistry, Adam Mickiewicz University, Poznań, Poland

<sup>4</sup>Department of Earth Sciences, Chiba University, Chiba, Japan

<sup>5</sup>Department of Environmental Science and Policy, University of Milano, Milan, Italy

<sup>6</sup>Department of Geography, Earth and Environmental Sciences, University of Northern British Columbia, Prince George, Canada

<sup>7</sup>Department of Animal Taxonomy and Ecology, Faculty of Biology, Adam Mickiewicz University, Poznań, Poland

<sup>8</sup>Laboratory of Environmental Chemistry, Paul Scherrer Institut, Villigen, Switzerland

<sup>9</sup>Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

<sup>10</sup>Department of Geography, Durham University, Durham, UK

<sup>11</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, UK

<sup>12</sup>Department of Civil Engineering and Environmental Sciences, Western Norway University of Applied Sciences, Sogndal, Norway

Mercury (Hg) is a chemical element recognized as one of the most toxic among all naturally occurring elements, with health risks depending on its form, concentration, route and time of exposure. Mercury appears in the environment as a result of human activities, which include burning coal or lignite, improper waste disposal, oil refining, use of mercury-containing pesticides and fertilizers, and industrial development such as mining, chemical, pharmaceutical and paper industries. The element also appears in the environment as a consequence of natural phenomena, among which are volcanic emissions, rock erosion, biomass burning and geothermal processes, but also as a result of re-emissions. Mercury can persist in the atmosphere for up to several months, which promotes the transfer of the element to areas far from the emitting source.

Cryoconite, a sediment accumulating on the surface of glaciers, is known to accumulate atmospheric contaminants such as Hg likely due to biofilm producing extracellular polymeric substances. Mercury is a contaminant of primary concern in the global environment, including cryosphere environments such as glaciers, due to its high toxicity to biota. This study, for the first time, presents a comprehensive global analysis of the variation in Hg concentrations, observed in cryoconite holes and deposits from the surface of 27 glaciers in both hemispheres, comprising 105 samples in total. Concentrations of Hg were determined through ICP-MS/MS.

The results indicate a higher Hg content in cryoconite from glaciers located in the Northern Hemisphere, which can be linked to the proximity of highly industrialized areas, which contrasts to glaciers located in the Southern Hemisphere. The highest Hg content was measured in cryoconite located in Norway and Alaska (up to 0.7 ppm), and the Alps (close to 0.5 ppm), correlated with the levels of industrialization in these regions. Our results reveal a broad pattern of reduction in Hg concentrations in cryoconite with altitude, which may be related to the topographical relief affecting the transport of contaminants from higher altitudes to lower.

As a result of global warming, the majority of glaciers are retreating. The accumulated Hg in the cryoconite can be released during melting of glaciers and thus may also contribute to contamination of the downstream ecosystems and local communities through consumption of contaminated food and water in polar and alpine regions. Therefore, studies like this are needed to monitor the levels and fate of Hg in glaciers and ice caps.