

This is the peer reviewed version of the following article:

Outcome of 421 adult patients with Philadelphia-negative acute lymphoblastic leukemia treated under an intensive program inspired by the GIMEMA LAL1913 clinical trial: a Campus ALL study / Lazzarotto, Davide; Cerrano, Marco; Papayannidis, Cristina; Chiaretti, Sabina; Mosna, Federico; Fracchiolla, Nicola; Zappasodi, Patrizia; Imbergamo, Silvia; Del Principe, Maria Ilaria; Lunghi, Monia; Lussana, Federico; Piccini, Matteo; Fumagalli, Monica; Dargenio, Michelina; Salutari, Prassede; Forghieri, Fabio; Da Molin, Teresa Giulia; Bonifacio, Massimiliano; Olivi, Matteo; Giglio, Fabio; Trappolini, Silvia; Leoncin, Matteo; Mule, Antonino; Delia, Mario; Pasciolla, Crescenza; Grimaldi, Francesco; Cambo, Benedetta; Santoro, Lidia; Guolo, Fabio; Minetto, Paola; Defina, Marzia; Chiusolo, Patrizia; Fanin, Matteo; Mauro, Endri; Aprile, Lara; Mazzone, Carla; Trastulli, Fabio; Ciccone, Maria; De Gobbi, Marco; Cignetti, Alessandro; De Bellis, Eleonora; Mancini, Valentina; Piciocchi, Alfonso; Vignetti, Marco; Marsili, Giovanni; Starza, Irene Della; Fanin, Renato; Luppi, Mario; Ferrara, Felicetto; Pizzolo, Giovanni; Bassan, Renato; Foa, Robin; Candoni, Anna. - In: HAEMATOLOGICA. - ISSN 1592-8721. - (2024), pp. 1-35. [10.3324/haematol.2024.285638]

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

27/10/2024 23:34

(Article begins on next page)

27/10/2024 23:34

Outcome of 421 adult patients with Philadelphia-negative acute lymphoblastic leukemia treated under an intensive program inspired by the GIMEMA LAL1913 clinical trial: a Campus ALL study

by Davide Lazzarotto, Marco Cerrano, Cristina Papayannidis, Sabina Chiaretti, Federico Mosna, Nicola Fracchiolla, Patrizia Zappasodi, Silvia Imbergamo, Maria Iliara Del Principe, Monia Lunghi, Federico Lussana, Matteo Piccini, Monica Fumagalli, Michelina Dargenio, Prassede Salutati, Fabio Forghieri, Teresa Giulia Da Molin, Massimiliano Bonifacio, Matteo Olivi, Fabio Giglio, Silvia Trappolini, Matteo Leoncin, Antonino Mulè, Mario Delia, Crescenza Pasciolla, Francesco Grimaldi, Benedetta Cambò, Lidia Santoro, Fabio Guolo, Paola Minetto, Marzia Defina, Patrizia Chiusolo, Matteo Fanin, Endri Mauro, Lara Aprile, Carla Mazzone, Fabio Trastulli, Maria Ciccone, Marco De Gobbi, Alessandro Cignetti, Eleonora De Bellis, Valentina Mancini, Alfonso Picocchi, Marco Vignetti, Giovanni Marsili, Irene Della Starza, Renato Fanin, Mario Luppi, Felicetto Ferrara, Giovanni Pizzolo, Renato Bassan, Robin Foà, and Anna Candoni

Received: May 15, 2024.

Accepted: August 6, 2024.

Citation: Davide Lazzarotto, Marco Cerrano, Cristina Papayannidis, Sabina Chiaretti, Federico Mosna, Nicola Fracchiolla, Patrizia Zappasodi, Silvia Imbergamo, Maria Iliara Del Principe, Monia Lunghi, Federico Lussana, Matteo Piccini, Monica Fumagalli, Michelina Dargenio, Prassede Salutati, Fabio Forghieri, Teresa Giulia Da Molin, Massimiliano Bonifacio, Matteo Olivi, Fabio Giglio, Silvia Trappolini, Matteo Leoncin, Antonino Mulè, Mario Delia, Crescenza Pasciolla, Francesco Grimaldi, Benedetta Cambò, Lidia Santoro, Fabio Guolo, Paola Minetto, Marzia Defina, Patrizia Chiusolo, Matteo Fanin, Endri Mauro, Lara Aprile, Carla Mazzone, Fabio Trastulli, Maria Ciccone, Marco De Gobbi, Alessandro Cignetti, Eleonora De Bellis, Valentina Mancini, Alfonso Picocchi, Marco Vignetti, Giovanni Marsili, Irene Della Starza, Renato Fanin, Mario Luppi, Felicetto Ferrara, Giovanni Pizzolo, Renato Bassan, Robin Foà, and Anna Candoni. Outcome of 421 adult patients with Philadelphia-negative acute lymphoblastic leukemia treated under an intensive program inspired by the GIMEMA LAL1913 clinical trial: a Campus ALL study. *Haematologica*. 2024 Aug 15. doi: 10.3324/haematol.2024.285638 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication.

E-publishing of this PDF file has been approved by the authors.

After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal.

All legal disclaimers that apply to the journal also pertain to this production process.

Outcome of 421 adult patients with Philadelphia-negative acute lymphoblastic leukemia treated under an intensive program inspired by the GIMEMA LAL1913 clinical trial: a Campus ALL study

RUNNING HEADS: Real-life study on 421 adult Ph-neg ALL treated with LAL1913 program

Davide Lazzarotto¹, Marco Cerrano², Cristina Papayannidis³, Sabina Chiaretti⁴, Federico Mosna⁵, Nicola Fracchiolla⁶, Patrizia Zappasodi⁷, Silvia Imbergamo⁸, Maria Ilaria Del Principe⁹, Monia Lunghi¹⁰, Federico Lussana¹¹, Matteo Piccini¹², Monica Fumagalli¹³, Michelina Dargenio¹⁴, Prassede Salutari¹⁵, Fabio Forghieri¹⁶, Teresa Giulia Da Molin^{17,18}, Massimiliano Bonifacio^{17,18}, Matteo Olivi¹⁹, Fabio Giglio²⁰, Silvia Trappolini²¹, Matteo Leoncin²², Antonino Mulè²³, Mario Delia²⁴, Crescenza Pasciolla²⁵, Francesco Grimaldi²⁶, Benedetta Cambò²⁷, Lidia Santoro²⁸, Fabio Guolo^{29,30}, Paola Minetto^{29,30}, Marzia Defina³¹, Patrizia Chiusolo^{32,33}, Matteo Fanin^{1,34}, Endri Mauro³⁵, Lara Aprile³⁶, Carla Mazzone³⁷, Fabio Trastulli³⁸, Maria Ciccone³⁹, Marco De Gobbi⁴⁰, Alessandro Cignetti⁴¹, Eleonora De Bellis⁴², Valentina Mancini⁴³, Alfonso Piciocchi⁴⁴, Marco Vignetti⁴⁴, Giovanni Marsili⁴⁴, Irene Della Starza^{4,44}, Renato Fanin^{1,34}, Mario Luppi¹⁶, Felicetto Ferrara³⁸, Giovanni Pizzolo^{17,18}, Renato Bassan²², Robin Foà⁴, Anna Candoni^{1,16}

¹ Clinica Ematologica - Centro Trapianti e Terapie Cellulari, Azienda Sanitaria Universitaria Friuli Centrale, Udine, Italy

² S.C. Ematologia, AOU Città della Salute e della Scienza - Presidio Molinette, Torino, Italy

³ IRCCS Azienda Ospedaliero-Universitaria di Bologna, Istituto di Ematologia "Seràgnoli", Bologna, Italy

⁴ Ematologia, Dipartimento di Medicina Traslazionale e di Precisione, "Sapienza" Università di Roma, Roma, Italy

⁵ Ematologia e Centro Trapianti di Midollo Osseo, Azienda Sanitaria dell'Alto Adige, Bolzano, Italy

⁶ U.O. Ematologia, IRCCS Ca' Granda Ospedale Maggiore Policlinico di Milano, Italy

⁷ Division of Hematology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

⁸ Dipartimento Strutturale Aziendale Medicina, University of Padova, Padova, Italy

- ⁹ Department of Biomedicine and Prevention, University of Rome Tor Vergata, Roma, Italy
- ¹⁰ Division of Hematology, Department of Translational Medicine, AOU Maggiore della Carità, Università del Piemonte Orientale, Novara, Italy
- ¹¹ Hematology and Bone Marrow Transplantation Unit, Azienda Socio-Sanitaria Territoriale Papa Giovanni XXIII, Bergamo, Italy
- ¹² SODc Ematologia, Azienda Ospedaliera Universitaria Careggi, Firenze, Italy
- ¹³ Ematologia, Ospedale San Gerardo, ATS Brianza, Monza, Italy
- ¹⁴ S.C. Ematologia, Ospedale Vito Fazzi, Lecce, Italy
- ¹⁵ Hematology Unit, Ospedale Civile Santo Spirito, Pescara, Italy
- ¹⁶ Clinica Ematologica, Azienda Ospedaliero Universitaria di Modena, Dip. Scienze Mediche e Chirurgiche, UNIMORE, Modena, Italy
- ¹⁷ Department of Engineering for Innovation Medicine, Section of Hematology, University of Verona, Verona, Italy
- ¹⁸ Department of Medicine, Azienda Ospedaliera Universitaria Integrita di Verona, Verona, Italy
- ¹⁹ Department of Oncology and Hematology, A.O.U. Città della Salute e della Scienza di Torino, Torino, Italy
- ²⁰ Unità Di Ematologia E Trapianto Di Midollo Osseo, IRCCS Ospedale San Raffaele di Milano, Milano, Italy
- ²¹ S.O.D. Clinica Ematologica, Azienda Ospedaliero-Universitaria Ospedali Riuniti Umberto I, Ancona, Italy
- ²² UOC Ematologia, Azienda Ulss3 Serenissima, Ospedale dell'Angelo, Mestre, Italy
- ²³ Divisione di Ematologia ad indirizzo oncologico, A.O. Ospedali Riuniti Villa Sofia-Cervello, Palermo, Italy
- ²⁴ UO Ematologia con Trapianto - Azienda Ospedaliero-Universitaria-Consortiale Policlinico di Bari, Bari, Italy
- ²⁵ U.O. di Ematologia, IRCCS Istituto Tumori Giovanni Paolo II, Bari, Italy
- ²⁶ Dipartimento di Medicina Clinica e Chirurgia, AOU Federico II, Napoli, Italy
- ²⁷ Hematology and BMT Unit, Department of Clinical and Experimental Medicine, University of Parma, Parma, Italy
- ²⁸ U.O.C. Ematologia e Trapianto di Midollo Osseo, Avellino, Italy
- ²⁹ Clinica Ematologica, Dipartimento Di Medicina Interna (DiMI), Università Degli Studi di Genova, Genova, Italy
- ³⁰ IRCCS Ospedale Policlinico San Martino, Genova, Italy
- ³¹ UOC Ematologia, Azienda Ospedaliero Universitaria Senese, Siena, Italy
- ³² Dipartimento di Diagnostica per Immagini, Radioterapia Oncologica ed Ematologia, Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italy

³³ Sezione di Ematologia, Dipartimento di Scienze Radiologiche ed Ematologiche, Università Cattolica del Sacro Cuore, Roma, Italy

³⁴ Dipartimento di Medicina, Università degli Studi di Udine, Udine, Italy

³⁵ Hematology Unit, Santa Maria di Ca' Foncello Hospital, Treviso, Italy

³⁶ S.C. Ematologia, Ospedale S.G. Moscati, Taranto, Italy

³⁷ Haematology, Department of Medical Area, St. Eugenio Hospital, Roma, Italy

³⁸ Hematology, Hospital "Antonio Cardarelli", Napoli, Italy

³⁹ UO Ematologia, Dipartimento di Medicine Specialistiche, Azienda Ospedaliero-Universitaria Arcispedale S Anna, Ferrara, Italy

⁴⁰ Department of Clinical and Biological Sciences, San Luigi Gonzaga Hospital, University of Turin, Orbassano, Italy

⁴¹ SCU Ematologia e Terapie Cellulari, AO Ordine Mauriziano, Torino, Italy

⁴² UCO Ematologia, Azienda Sanitaria Universitaria Giuliano Isontina, Trieste, Italy

⁴³ ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy

⁴⁴ Centro Dati Fondazione GIMEMA Franco Mandelli, Roma, Italy

Authors' contributions: A.C. designed the study (and she is the Principal Investigator), wrote and revised the paper. D.L. performed the analysis, wrote the paper and collected data. R.F., C.P., M.C., S.C., collected data and contributed to the revision of paper. All other authors collected data and contributed to the final approval of the manuscript.

Corresponding authors:

Davide Lazzarotto, MD and Anna Candoni, MD

Clinica Ematologica - Centro Trapianti e Terapie Cellulari, Azienda Sanitaria Universitaria Friuli Centrale, Udine.

E-mail: davide.lazzarotto@asufc.sanita.fvg.it; acandoni@unimore.it

Data sharing statement: Data regarding this research cannot be shared openly to protect study participant privacy.

Acknowledgments. We would like to thank all the members of the Campus ALL-Italy network.

Funding. The work was partly supported by Associazione Italiana per la Ricerca sul Cancro (AIRC), 5x1000 Metastases Special Program, N° 21198, Milan, Italy (RF)

Disclosures: all the Authors declared no conflict of interest related to the submitted work

ABSTRACT

The introduction of pediatric-inspired regimens in adult Philadelphia-negative acute lymphoblastic leukemia (Ph- ALL) has significantly improved patients' prognosis. Within the Campus ALL network we analyzed the outcome of adult Ph- ALL patients treated according to the GIMEMA LAL1913 protocol outside the clinical trial, to compare the real-life data with the study results. We included 421 consecutive patients, with a median age of 42 years. The complete remission (CR) rate after the first course of chemotherapy was 94% and a measurable residual disease (MRD) negativity after the third course was achieved in 72% of patients. The 3-year overall survival (OS) and disease-free survival (DFS) were 67% and 57%, respectively. In a multivariate analysis, MRD positivity negatively influenced DFS. In a time-dependent analysis including only very high risk (VHR) and MRD positive cases, transplanted (HSCT) patients had a significantly better DFS than non-HSCT ones ($P=0.0017$). During induction, grade ≥ 2 pegaspargase-related hepato-toxicity was observed in 25% of patients (vs 12% in the GIMEMA LAL1913 trial, $P=0.0003$).

In this large real-life cohort of Ph- ALL, we confirmed the very high CR rate and a superimposable OS and DFS compared to the GIMEMA LAL1913 clinical trial: CR rate after C1 94% vs 85%, $P=0.0004$; 3-year OS 67% vs 67%, $P=0.94$; 3-year DFS 57% vs 63%, $P=0.17$. HSCT confirms its important role in VHR and MRD-positive patients. The rate of pegaspargase-related toxicity was significantly higher in the real-life setting, emphasizing the importance of dose adjustment in the presence of risk factors to avoid excessive toxicity.

INTRODUCTION

No definitive consensus exists on the optimal treatment regimen for adult Philadelphia chromosome-negative acute lymphoblastic leukemia (Ph- ALL) to optimally balance efficacy and toxicity, as shown by the different treatment backbones employed by cooperative study groups.¹⁻⁷ Nonetheless, in the last years numerous phase 2 and phase 3 clinical trials from different countries have been associated with rather favorable results compared to previous experiences.¹⁻⁷ These improvements have been achieved using intensive pediatric-inspired protocols, new formulations of asparaginase and revised stratification models which included measurable residual disease (MRD) monitoring, in addition to baseline risk factors.⁸⁻¹⁰ However, data on the real-life applicability of therapeutic regimens tested in clinical trials, which inherently enroll selected patient populations, are very limited.

Recently, the Gruppo Italiano Malattie EMatologiche dell'Adulto (GIMEMA) published the results of the LAL1913 clinical trial, which included 203 homogeneously treated adult Ph- ALL patients with a pediatric-inspired protocol.¹ After the completion of this study, most Italian hematology centers used the same therapeutic program in their clinical practice while the new protocol for Ph- ALL was under discussion. In this paper, we report the efficacy and safety data of a chemotherapy program performed according to the GIMEMA LAL1913 protocol in adult patients with Ph- ALL treated outside the clinical trial, in a real-life setting.

METHODS

Patients and objectives of the study

We included 421 consecutive adult patients with newly diagnosed Ph- ALL or lymphoblastic lymphoma (LL, with <20% bone marrow blasts) treated according to the GIMEMA LAL1913 protocol,¹ outside the clinical trial, between September 2016 and December 2022. The data were collected from 39 hematology centers that are part of the Campus ALL network in Italy.

The main objectives of the study were to compare the complete remission (CR) rate, the overall survival (OS) and the disease-free survival (DFS) between the real-life cohort (421 cases) and the GIMEMA LAL1913 clinical trial population (203 cases). Secondary endpoints included the evaluation of the treatment toxicity and the allogeneic stem cell transplantation (HSCT) rate according to the risk-group at diagnosis.

Diagnostic procedures such as immunophenotyping, cytogenetics and molecular studies were carried out according to the GIMEMA LAL1913 protocol indications.^{1,2} The Philadelphia-like signature was not routinely tested in this real-life population.

In line with the GIMEMA LAL1913 trial, three risk classes were defined at diagnosis (as reported in the supplementary section).

This observational study was approved by the Ethics Committee of Friuli Venezia Giulia, Italy (ethical approval number CEUR-2022-Os-03) and conducted in accordance with the 2008 revision of the Helsinki Declaration.

Treatment protocol

All patients were treated according to the GIMEMA LAL1913 protocol as described by Bassan R et al. and detailed in **Supplemental Table 1**.¹

Antibiotic, antimycotic and antiviral prophylaxis, and pegaspargase toxicity management were administered according to the policy of each center.

Treatment-related toxicity was evaluated according to the Common Terminology Criteria for Adverse Events (CTCAE), version 4.0.

MRD analysis

MRD analysis was carried out on bone marrow samples through real-time quantitative (RTq) -PCR for immunoglobulin (IG) or T-cell receptor (TR) gene rearrangements following the EuroMRD guidelines¹¹ in 3 reference laboratories (as in the GIMEMA LAL1913 trial), or locally through multiparameter flow cytometry (MFC) targeting leukemia associated immunophenotype in patients lacking suitable molecular probes. Similarly to the GIMEMA LAL1913 trial, data on MRD were collected at four specific time points: end of induction week 4 (TP1), weeks 10 (end of course 3, TP2), 16 (end of course 5, TP3) and 22 (end of course 7, TP4). Patients with low positive ($\leq 10^{-4}$) or negative TP2-3 and negative TP4 (or negative TP2-3 when TP4 was missing) were defined as MRD-negative (MRD-neg), while those with TP2-3 $\geq 10^{-4}$ and/or positive TP4 were defined as MRD-positive (MRD-pos), according to the LAL1913 clinical trial.

Statistical analysis

The comparison between baseline characteristics among subgroups was obtained using the Fisher's exact or Chi-squared test for categorical variables, the Student's t-test for normally distributed variables and the Mann-Whitney test for non-normally distributed variables. Logistic regression was used to study variables influencing the obtainment of MRD-negativity at TP2. Median follow-up time was calculated among survivors and was last updated in June 2023.

The response evaluation criteria are reported in the supplementary section.

OS was calculated from the date of diagnosis to the date of the last follow-up or to the date of death by any cause. DFS was calculated from the date of the first CR achievement to the date of the last follow-up, relapse or death by any cause. DFS stratification for MRD followed the definition for MRD-neg and MRD-pos described in the "MRD analysis" section using the available time-points for each patient. OS and DFS were estimated according to the Kaplan-Meier method and the

differences between groups were compared with the log-rank test. Univariate and multivariate analyses were carried out by Cox regression for OS and DFS.

Simon-Makuch plot was used to assess the time-dependent effects of HSCT and Mantel-Byar test was used for comparison of survival curves.

The same descriptive statistics were used to compare the characteristics of the real-life and the LAL1913 clinical trial populations. To compare OS and DFS, a subclass matching propensity score was performed (5 quantile classes) considering the following variables: age, sex, risk, lineage, and transplant. All 602 observations were matched, and the real-life data were weighted according to subclassification. Propensity score estimates were calculated using a logistic regression model. A summary of the characteristics of the patients in the propensity score matching is reported in **Supplemental Table 2**.

Statistical significance in all cases was considered for a P value less than 0.05.

RESULTS

Patients' characteristics

The main characteristics of the 421 patients are summarized in **Table 1**. Median age was 42 years (range 18-80) and was significantly lower in T-ALL/LL patients (38.5 vs 45, $P=0.0009$); 23% ($n=97$) were older than 55 years; 52.5% ($n=221$) had B-ALL and 12% ($n=50$) had LL (of which $n=45$ T-lineage, $P<0.0001$).

The median WBC count was significantly higher in T-ALL ($P<0.0001$), as also the involvement of lymph nodes and mediastinum (42% and 47% of patients, respectively). Central nervous system (CNS) involvement was documented in 9% ($n=37$) of patients at disease onset (more frequently in T-ALL/LL, 12.5% vs 5%, $P=0.0149$).

As for cytogenetics/genetics (evaluable in 81% of patients, n=342), 15 patients had a *KMT2A*;11q23 rearrangement, 45 had other adverse karyotypes, while a t(1;19)/*TCF3*::*PBX1* translocation was detected in 5 patients and a hyperdiploidy in 15. As specified in the “Methods” section, the Philadelphia-like signature was not routinely tested.

Overall, 49% of patients were standard risk (SR), 10% high risk (HR) and 41% very high risk (VHR). T-ALL/LL patients displayed VHR features more frequently (52% vs 30% of B-ALL/LL, P<0.0001).

The median follow-up of the entire population was 24.6 months. At the last follow-up, 306 patients (73%) were alive (251/306 - 82% in CR1) and 115 (27%) had died (64/115 - 56% due to underlying disease, 24/115 - 21% due to transplant-related mortality, 9/115 - 8% deaths during induction, 5/115 - 4% deaths in CR during subsequent courses of chemotherapy, 13/115 - 11% due to other causes).

Treatment and response

All 421 patients received the first course of therapy (C1) and 358 (85%) of them were able to continue the treatment up to the third course (C3). Prior to C3 we recorded 15 deaths, 9 during induction (2%), 3 during consolidation (7 of which due to infection) and 3 unrelated to disease or therapy, while 40 patients switched to an alternative treatment, 26 (65%) due to refractoriness or early progression (14 after C1 and 12 after C2) and 14 (35%) due to adverse events (10 after C1 and 4 after C2). Eight patients had a short follow up (too early) and did not undergo C3 at data cut off. Overall, only 6% of the entire patient population (26/421) was refractory after C2.

The morphologic CR rate after C1 was 94% (356/379) and after C2 95% (329/347) of evaluable patients (not evaluable patients were those with LL without marrow involvement and those in which the bone marrow study was not performed). The early death rate was 3% (n=12) of the whole population.

After C3, 146 patients (35%) underwent a HSCT in first line; in 16% of patients (n=24), the procedure was preceded by immunotherapy for MRD persistence (Blinatumomab in 22 cases, Inotuzumab in 2 cases). The two main indications for HSCT were: VHR disease (70.5%, n=103) and MRD positivity (21%, n=31). HSCT was more frequently carried out in T-ALL patients (47% vs 31%, P=0.002). Overall, 129 SR-MRD-neg patients were able to proceed to maintenance. Globally, 39 patients were treated with immunotherapy (35 Blinatumomab, 3 Inotuzumab, 1 Daratumumab) for MRD persistence after first-line chemotherapy.

MRD study was available in 381 patients (90.5%); 71% (n=269) were monitored by RTq-PCR for Ig/TR gene rearrangements, and the remaining 29% (n=112) by MFC. The rates of MRD negativity at TP1 and TP2 were, respectively, 46% and 67% of the evaluable patients (72% when excluding LL patients without MRD study on bone marrow).

A summary of the MRD response at the different time points is provided in **Table 2**, and did not differ between B-ALL and T-ALL patients. A multivariate logistic regression analysis including age, risk category, lineage, ECOG score, and CNS involvement was carried out to study variables influencing the obtainment of MRD negativity at TP2, and we found that the presence of a HR or a VHR risk class was the only factor associated with failure to achieve MRD negativity (OR 0.38, CI 0.22 – 0.64, P=0.0003).

Side effects and toxicities

Chemotherapy dose reductions beyond those established by the LAL1913 protocol in patients >55 years were required during C1 in 118 patients (28%), due in 50% of patients (59) to either toxicity or infection.

Table 3 summarizes the pegaspargase-related toxicity. During C1, 382 patients (91%) received pegaspargase, and 49% of them (189) developed a grade ≥ 2 related toxicity (mainly hepatic toxicity)

while thrombosis, pancreatic toxicity and hypersensitivity reaction were rare (as reported in **Table 3**). The global rate of grade ≥ 2 pegaspargase-related toxicity at C2 was 32% (101/314). Pegaspargase was not administered at C2 in 12% of patients (47/382) due to previous related toxicity at C1. In addition, a drug dose reduction was required during C2 in 27% of patients receiving pegaspargase (86/314). During C5 and C6, the global rate of grade ≥ 2 pegaspargase-related toxicity was 38% and 30%, respectively. A pegaspargase dose reduction at C5 and C6 was required in 28% (50/177) and 35% (45/129) of cases, respectively. The drug was omitted at C5, due to the previous related toxicity, in 9% of patients and in 19% of patients at C6 (**Table 3**).

Pegaspargase-free courses (C3, C4, C7 and C8) were administered at the programmed full doses of chemotherapy in 93%, 93%, 89% and 87% of patients, respectively.

Infectious complications were more frequently recorded during C1. Bacteremia/sepsis, being the most common infection, was observed in 14% of patients (59), followed by pneumonia in 11% (45); 20 cases of pneumonia (5% of the whole population) were mycotic. Moreover, during C1, 22% (91) of patients developed febrile neutropenia. In the following courses the number of patients developing bacteremia/sepsis was lower (between 1% and 9%); the courses with the highest number of events observed were C3, C6 and C7 (respectively, 7%, 7% and 9%). Also, the number of patients developing pneumonia was lower (between 0% and 4%) with similar percentages in the different courses. The number of patients developing febrile neutropenia beyond C1 ranged between 3% and 21% of patients, and again a higher number of events was observed at C3, C6 and C7 (respectively, 17%, 18% and 21%).

Survival analysis and prognostic factors

The 3-year OS probability was 67% (median not reached), without significant differences between patients aged ≤ 40 and those aged 41-55 (76% vs 63%, $P=0.28$). However, both these groups had a

significantly higher 3-year OS than patients aged >55 (55%, Logrank test $P=0.0007$ vs patients aged ≤ 40 and $P=0.041$ vs patients aged 41-55), as shown in **Figures 1A** and **1C**.

The 3-year DFS probability was 57% (median not reached), without significant differences between patients aged ≤ 40 and patients aged 41-55 (61% vs 60%, $P=0.77$). Again, both these groups had a significantly higher 3-year DFS than patients aged >55 (46%, Logrank test $P=0.011$ vs patients aged ≤ 40 and $P=0.050$ vs patients aged 41-55), as shown in **Figures 1B** and **1D**.

Figure 2A shows the DFS curves for MRD-neg and MRD-pos patients. The 3-year DFS was 67% in MRD-neg patients vs 32% in MRD-pos ones (Logrank test $P < 0.0001$), respectively.

In univariate analysis, a younger age predicted a better OS, while CNS involvement and MRD-pos predicted a worse OS. A younger age (≤ 55 years) also predicted a better DFS in univariate analysis, while MRD-pos, CNS involvement, high leucocyte count ($>30 \times 10^9/l$), adverse cytogenetics, the presence of a *KMT2A* rearrangement and the VHR risk class per se predicted a worse DFS. In multivariate analysis for OS and DFS, significance was retained only for MRD-pos (**Figure 3**).

To better analyze the effect of HSCT, a time-dependent analysis was performed for DFS. HSCT did not show a benefit when considering the whole population, but when we considered just VHR or MRD-pos patients, i.e. those who were transplant candidate according to the GIMEMA LAL1913 protocol, the impact of HSCT was significant (Mantel-Byar $P=0.0017$). In **Figure 2B** we report the Simon-Makuch plot for DFS of VHR and MRD-pos patients according to HSCT.

Comparison with the results of the GIMEMA LAL 1913 trial

As shown in **Table 4**, we compared the most important findings of this real-life observational study (including 421 cases) and the results of the GIMEMA LAL1913 clinical trial (including 203 cases).¹ The real-life population was slightly older, although the difference was not significant (median age 42 vs 40 years, $P=0.5$, with patients older than 55: 23% vs 19%, $P=0.33$) and included a higher

number of T-ALL/LL (47.5% vs 31.5%, $P=0.0002$). Moreover, in the real-life population cohort we observed a non-significantly higher proportion of HR+VHR patients (51% vs 43%, $p=0.09$).

The CR rate at TP1 was higher in the real-life population (94% vs 85%, $P=0.0004$), but the rate of MRD negativity at both TP1 and TP2 was lower (respectively 46% vs 56%, $P=0.04$, and 72% vs 80%, $P=0.04$).

Importantly, OS and DFS were similar in the two studies, with a 3-year OS of 67% vs 67%, $P=0.94$, and a 3-year DFS of 57% vs 63%, $P=0.17$, respectively (**Figure 4**). When weighed according to the propensity score performed, the 3-year OS and DFS were 67% and 55%, respectively (with $P=0.94$ and $P=0.17$, compared to the GIMEMA LAL1913 trial data). The rate of HSCT in first line was higher in the real-life setting (35% vs 28%), though without reaching a significant difference ($P=0.09$).

Finally, we compared pegaspargase-related adverse events during C1, and we observed a higher rate of grade ≥ 2 hepatic toxicity in patients treated in the real-life setting compared to those included in the LAL1913 trial (25% vs 12%, $P=0.0003$), while the rates of grade 3 pancreatic toxicity and the thrombotic events were similar in the two cohorts (3% vs 1%, $P=0.26$, and 2% vs 2%, $P=1.00$).

DISCUSSION

Pediatric-inspired protocols have improved the outcome of Ph- ALL in adults,⁸ as demonstrated by several trials yielding comparable results, with CR rates around 90% and OS and DFS rates above 60% at 3-5 years, despite the differences in trial design and asparaginase formulations and dosage.¹⁻

⁷ In trials including older adults (>55 years), this patient population witnesses worse results, with inferior CR rates and survival.¹⁻⁴ This is likely due to the difficulties in delivering optimal

chemotherapy doses, increased complication rates, and a different disease biology compared to younger patients.^{3,4,12}

Despite the significant number of clinical trials using pediatric-inspired protocols in adult Ph- ALL, very limited data are available on the feasibility, toxicities and outcome of these protocols in the context of the daily clinical practice outside of clinical trials.¹³⁻¹⁶ Since the completion of the GIMEMA LAL1913 trial, the results of which have been recently published,¹ most Italian hematology centers have adopted this pediatric-inspired therapeutic program as the standard of care for the clinical management of newly diagnosed adult Ph- ALL. The current study conducted within the Campus ALL network and involving 39 hematology centers in Italy was thus aimed at analyzing the feasibility and performance of the LAL1913 program in the real-life in terms of tolerability and outcome, and to compare these results with those obtained in the original clinical trial.¹ To our knowledge, this multicenter real-life study that included 421 adult Ph-ALL patients homogeneously treated according to a pediatric-inspired protocol (GIMEMA LAL1913)¹ is the largest so far available. Some differences emerged between the characteristics of the real-life population compared to the clinical trial one. The real-life cohort included more T-ALL (47.5% vs 31.5%) and a higher proportion of HR+VHR patients, albeit the difference was not significant (51% vs 43%). The median age and the proportion of patients aged >55 were comparable in the two cohorts (23% in real-life vs 19% in clinical trial, P=0.33).

In consideration of the abovementioned differences, a high CR rate after C1 (94%) was observed in the real-life setting, resulting even higher than the rate reported in the GIMEMA LAL1913 clinical trial (85%, p=0.0004). However, the rate of MRD-neg patients at both TP1 and TP2 was lower in the real-life cohort compared to the clinical trial results (respectively 46% vs 56%, P=0.04, and 72% vs 80%, P=0.04), and this can be explained by the higher number of HR and VHR patients included in our study, considering that this was the only variable that significantly influenced the achievement

of a MRD negativity a TP2. It is noteworthy that the OS and DFS observed in our real-life population were comparable to the results reported in the GIMEMA LAL1913 clinical trial (3-year OS 67% vs 67%, P=0.94; 3-year DFS 57% vs 63%, P=0.17). The CR and MRD-neg rates, OS and DFS were also in line with other published prospective clinical trials.^{3,4,6}

Similarly to other studies, age had an impact on OS in this real-life analysis,^{1,4,14-16} but unlike the GIMEMA LAL1913 clinical trial, patients aged 41-55 years fared the same as patients aged ≤ 40 years, and only patients aged >55 years showed a significantly reduced survival.¹

Besides, this study confirmed that biological features of the disease at diagnosis (cytogenetics, leukocytosis, *KMT2A* rearrangements) played an important role in DFS in univariate analysis, contributing to the definition of HR and VHR classes. This effect was not evident in the GIMEMA LAL1913 clinical trial, in which only patients with *KMT2A* rearranged ALL showed a significantly worse outcome.¹

We also confirmed the crucial role of MRD monitoring in the clinical practice and its important prognostic impact on OS and DFS as observed in many clinical trials.^{1,2,14,17-19}

Our real-life data also highlighted the prognostic impact of first-line HSCT in patients with unfavorable risk factors (VHR and/or MRD-pos), with results comparable to those reported in the GIMEMA LAL1913 clinical study. In our population, a proportion of patients was bridged to HSCT procedure with immunotherapy for MRD positivity, an option that was not available in the GIMEMA LAL1913 trial, but given that this approach is becoming standard in the clinical practice and that several trials (such as the ongoing GIMEMA LAL2317) are exploring a sequential chemo-immunotherapy approach, the role of HSCT may change in the near future.^{17,20,21}

In terms of tolerability, the highest rate of pegaspargase-related toxicity in our study was observed at C1, with 49% of patients experiencing at least one grade ≥ 2 adverse event. In particular, during C1 25% of patients in this study developed grade ≥ 2 hepatic toxicity compared to 12% of patients in the

GIMEMA LAL1913 clinical trial (P=0.0003). Overall, the pegaspargase-related toxicity observed compares favorably to other international reports in the literature.^{22,23} This finding may reflect less attention to risk factors for pegaspargase-related toxicity (such as obesity, hepatopathies) and/or a less stringent patient selection in the real-life setting than in the GIMEMA LAL1913 clinical trial.

Infectious complications are a significant concern during the management of Ph- ALL patients.²⁴ In our analysis, infectious complications mainly occurred at C1, with 14% of patients developing bacteremia/sepsis and 11% of patients developing pneumonia (mycotic in nearly half of the cases). While pneumonia represents an uncommon event in subsequent courses, the number of patients who developed bacteremia/sepsis reached 9% of cases, in line with previous studies involving Ph- ALL patients treated with intensive protocols.^{6,7,13} Despite the low rate of early mortality observed in our study (3%, 12/421), infections were the main cause of death, affecting more than half of cases. This suggests the need to improve infection surveillance, prophylactic measures, and antimicrobial therapy.

In summary, our study demonstrates the feasibility and favorable outcome of a pediatric-inspired therapeutic regimen in a large real-world setting with CR rates and OS and DFS similar to those reported in the originator clinical trial (GIMEMA LAL1913). Moreover, our analysis confirms the important role of HSCT in patients with high risk factors or MRD positivity. Therefore, outside of clinical trials efforts should be made to obtain a detailed disease characterization, keeping up with recently identified molecular subgroups, and to strictly monitor MRD at the appropriate time points to better identify patients with risk factors for early HSCT referral.⁹⁻¹⁰

A limitation of our study is the lack of availability of the Ph- like signature for the majority of the patients, that should be more widely tested in the common clinical practice.²⁵⁻²⁸ Besides, the widespread use of immunotherapy in patients with pre-transplant MRD persistence could improve the outcome of transplant, and we are waiting for the results of the studies testing these

approaches in patients with baseline high risk features²⁹ or in all cases, including MRD-negative patients, where blinatumomab also appears to be effective.³⁰

The tolerability of pegaspargase in the real-life setting remains an important concern, given the cornerstone role of this drug in the therapy regimen, and indeed our study is limited by the lack of a precise correlation analysis between pegaspargase dosage and response. Future studies are needed to individualize the dosage of the drug to be given to each patient with a tolerated dose while remaining within the efficacy range.^{8,20,22,31,32} Probably, the widespread availability of asparaginase activity monitoring could be useful to optimize dose calculation.³³

A detailed analysis of infectious complications was beyond the purpose of this study (and will be detailed in a subsequent report), but this remains an important issue to address, as infections are an important cause of morbidity and mortality, and efforts should be made to uniform anti-infectious prophylaxis, especially on the anti-mycotic front.^{24,34}

Finally, elderly patients still show inferior outcomes compared to younger patients, even observing a median OS of 45 months, and future studies should aim at identifying the optimal age cut-off to define the “elderly” and to design better and tailored induction therapies incorporating frontline immunotherapy to reduce toxicity and improve the outcome.³⁵⁻³⁷

REFERENCES

1. Bassan R, Chiaretti S, Della Starza I, et al. Pegaspargase-modified risk-oriented program for adult acute lymphoblastic leukemia: results of the GIMEMA LAL1913 trial. *Blood Adv.* 2023;7(16):4448-4461
2. Bassan R, Pavoni C, Intermesoli T, et al. Updated risk-oriented strategy for acute lymphoblastic leukemia in adult patients 18-65 years: NILG ALL 10/07. *Blood Cancer J.* 2020;10(11):119.
3. Huguet F, Leguay T, Raffoux E, et al. Pediatric-inspired therapy in adults with Philadelphia chromosome negative acute lymphoblastic leukemia: the GRAALL-2003 study. *J Clin Oncol.* 2009;27(6):911-918.
4. Huguet F, Chevret S, Leguay T, et al. Intensified therapy of acute lymphoblastic leukemia in adults: report of the randomized GRAALL-2005 clinical trial. *J Clin Oncol.* 2018;36(24):2514-2523.
5. Toft N, Birgens H, Abrahamsson J, et al. Results of NOPHO ALL2008 treatment for patients aged 1-45 years with acute lymphoblastic leukemia. *Leukemia.* 2018;32(3):606-615.
6. Stock W, Luger SM, Advani AS, et al. A pediatric regimen for older adolescents and young adults with acute lymphoblastic leukemia: results of CALGB 10403. *Blood.* 2019;133(14):1548-1559.
7. Testi AM, Canichella M, Vitale A, et al. Adolescent and young adult acute lymphoblastic leukemia. Final results of the phase II pediatric-like GIMEMA LAL-1308 trial. *Am J Hematol.* 2021;96(3):292-301.
8. Siegel SE, Stock W, Johnson RH, et al. Pediatric-inspired treatment regimens for adolescents and young adults With Philadelphia chromosome-negative acute lymphoblastic leukemia: a review. *JAMA Oncol.* 2018;4(5):725-734.
9. Gökbuget N, Boissel N, Chiaretti S, et al. Diagnosis, Prognostic Factors and Assessment of ALL in Adults: 2023 ELN Recommendations from a European Expert Panel. *Blood.* 2024;143(19):1891-1902.
10. Gökbuget N, Boissel N, Chiaretti S, et al. Management of ALL in Adults: 2023 ELN Recommendations from a European Expert Panel. *Blood.* 2024;143(19):1903-1930.
11. van der Velden VH, Cazzaniga G, Schrauder A, et al. Analysis of minimal residual disease by Ig/TCR gene rearrangements: guidelines for interpretation of real-time quantitative PCR data. *Leukemia.* 2007;21(4):604-611.
12. Gökbuget N. How I treat older patients with ALL. *Blood.* 2013;122(8):1366-1375.

13. Hanbali A, Kotb A, Fakhri RE, et al. Improved survival in adolescents and young adults (AYA) patients aged 14-55 years with acute lymphoblastic leukemia using pediatric-inspired protocol - a retrospective analysis of a real-world experience in 79 of patients treated at a national tertiary care referral center. *Leuk Res Rep.* 2021;16:100270.
14. Ferrari LC, Rivas MM, Navickas AB, et al. PH negative acute lymphoblastic leukemia in adolescents and young adults treated according a MRD adapted BFM ALL IC 2009 protocol: Argentine real-world data on 171 patients. *Ann Hematol.* 2023;102(5):1087-1097.
15. Reed DR, Wooster M, Isom S, et al. Real-world outcomes of adult patients with acute lymphoblastic leukemia treated with a modified CALGB 10102 regimen. *Ann Hematol.* 2023;102(4):897-906.
16. Oravcova I, Lukas J, Cingelova S, et al. Treatment of Adults and Young Adults with Acute Lymphoblastic Leukemia: Real Life Data from Two Centers in Slovakia. *Clin Lymphoma Myeloma Leuk.* 2021;21(10):e782-e791.
17. Gökbuğet N, Kneba M, Raff T, et al. Adult patients with acute lymphoblastic leukemia and molecular failure display a poor prognosis and are candidates for stem cell transplantation and targeted therapies. *Blood.* 2012;120(9):1868-1876.
18. Beldjord K, Chevret S, Asnafi V, et al. Oncogenetics and minimal residual disease are independent outcome predictors in adult patients with acute lymphoblastic leukemia. *Blood.* 2014;123(24):3739-3749.
19. Dhédin N, Huynh A, Maury S, et al. Role of allogeneic stem cell transplantation in adult patients with Ph-negative acute lymphoblastic leukemia. *Blood.* 2015;125(16):2486-2496.
20. Bassan R, Chiaretti S, Della Starza I, et al. Preliminary results of the GIMEMA LAL2317 sequential chemotherapy-blinatumomab frontline trial for newly diagnosed adult Ph-negative B-lineage ALL patients. *Hemasphere.* 2021;5(8):S114.
21. Ribera JM, Morgades M, Ciudad J, et al. Chemotherapy or allogeneic transplantation in high-risk Philadelphia chromosome-negative adult lymphoblastic leukemia. *Blood.* 2021;137(14):1879-1894.

22. Stock W, Douer D, DeAngelo DJ, et al. Prevention and management of asparaginase/pegasparaginase-associated toxicities in adults and older adolescents: recommendations of an expert panel. *Leuk Lymphoma*. 2011;52(12):2237-2253.
23. Aldoss I, Douer D. How I treat the toxicities of pegasparaginase in adults with acute lymphoblastic leukemia. *Blood*. 2020;135(13):987-995.
24. Mariette C, Tavernier E, Hocquet D, et al. Epidemiology of invasive fungal infections during induction therapy in adults with acute lymphoblastic leukemia: a GRAALL-2005 study. *Leuk Lymphoma*. 2017;58(3):586-593.
25. Moorman AV. New and emerging prognostic and predictive genetic biomarkers in B-cell precursor acute lymphoblastic leukemia. *Haematologica*. 2016;101(4):407-416.
26. Paietta E, Roberts KG, Wang V, et al. Molecular classification improves risk assessment in adult BCR-ABL1-negative B-ALL. *Blood*. 2021;138(11):948-958.
27. Moorman AV, Barretta E, Butler ER, et al. Prognostic impact of chromosomal abnormalities and copy number alterations in adult B-cell precursor acute lymphoblastic leukaemia: a UKALL14 study. *Leukemia*. 2022;36(3):625-636.
28. Chiaretti S, Messina M, Foà R. BCR/ABL1-like acute lymphoblastic leukemia: How to diagnose and treat? *Cancer*. 2019;125(2):194-204.
29. Boissel N, Huguet F, Leguay T, et al. Blinatumomab during Consolidation in High-Risk Philadelphia Chromosome (Ph)-Negative B-Cell Precursor (BCP) Acute Lymphoblastic Leukemia (ALL) Adult Patients: A Two-Cohort Comparison within the Graall-2014/B Study. *Blood*. 2022;140(Supplement 1):507-509
30. Litzow MR, Sun Z, Paietta E, et al. Consolidation Therapy with Blinatumomab Improves Overall Survival in Newly Diagnosed Adult Patients with B-Lineage Acute Lymphoblastic Leukemia in Measurable Residual Disease Negative Remission: Results from the ECOG-ACRIN E1910 Randomized Phase III National Cooperative Clinical Trials Network Trial. *Blood*. 2022;140(Supplement 2):LBA-1.

31. Lussana F, Minetto P, Ferrara F, Chiaretti S, Specchia G, Bassan R. National Italian Delphi panel consensus: which measures are indicated to minimize pegylated-asparaginase associated toxicity during treatment of adult acute lymphoblastic leukemia? *BMC Cancer*. 2020;20(1):956.
32. Derman BA, Streck M, Wynne J, et al. Efficacy and toxicity of reduced vs. standard dose pegylated asparaginase in adults with Philadelphia chromosome-negative acute lymphoblastic leukemia. *Leuk Lymphoma*. 2020;61(3):614-622.
33. Schore RJ, Devidas M, Bleyer A, et al. Plasma asparaginase activity and asparagine depletion in acute lymphoblastic leukemia patients treated with pegaspargase on Children's Oncology Group AALL07P4. *Leuk Lymphoma*. 2019;60(7):1740-1748.
34. Keng MB, Keng HC, Tan BH, Wong GC. High risk of invasive fungal infections in adult acute lymphoblastic leukemia patients receiving induction and salvage chemotherapy. *Leuk Lymphoma*. 2017;58(8):2017-2018.
35. Luskin MR. Acute lymphoblastic leukemia in older adults: curtain call for conventional chemotherapy? *Hematology Am Soc Hematol Educ Program*. 2021;2021(1):7-14.
36. Stelljes M, Raffel S, Alakel N, et al. Inotuzumab Ozogamicin as Induction Therapy for Patients Older Than 55 Years With Philadelphia Chromosome-Negative B-Precursor ALL. *J Clin Oncol*. 2024;42(3):273-282.
37. Goekbuget N, Schwartz S, Faul C, et al. Dose Reduced Chemotherapy in Sequence with Blinatumomab for Newly Diagnosed Older Patients with Ph/BCR::ABL Negative B-Precursor Adult Lymphoblastic Leukemia (ALL): Preliminary Results of the GMALL Bold Trial. *Blood*. 2023;142(Supplement 1):964.

TABLE 1. Characteristics of the 421 study patients.

	All patients (n=421)	B-ALL/LL (n=221)	T-ALL/LL (n=200)
Age (years), median (range)	42.0 (18-80)	45.0 (18-80)	38.5 (18-72)
≤ 40, n (%)	199 (47)	87 (39)	112 (56)
>40-55, n (%)	125 (30)	66 (30)	59 (29.5)
>55, n (%)	97 (23)	68 (31)	29 (14.5)
Gender (male), n (%)	248 (59)	111 (50)	137 (68.5)
Diagnosis , n (%)			
ALL	371 (88)	216 (98)	155 (77.5)
LL	50 (12)	5 (2)	45 (22.5)
ECOG PS , n			
0;1;2;3;4;NA	212;147;42;15;3;2	122;77;19;3;0;0	90;70;23;12;3;2
Hemoglobin (g/dl), median (range)	10.7 (4.1-17.2)	9.3 (4.1-15.1)	12.3 (4.2-17.2)
WBC (10 ⁹ /l), median (range)	9.9 (0.2-626.9)	6.9 (0.2-626.9)	12.2 (0.5-538.0)
≤ 30 (%)	304 (72)	170 (77)	134 (67)
>30-100 (%)	66 (16)	31 (14)	35 (17.5)
>100 (%)	51 (12)	20 (9)	31 (15.5)
PB blasts (%), median (range)	38.0 (0.0-100.0)	40.5 (0.0-100.0)	30.0 (0.0-100.0)
BM blasts (%), median (range)	80.0 (0.0-100.0)	85.0 (0.0-100.0)	73.0 (0.0-100.0)
Platelets (10 ⁹ /l), median (range)	78 (4-753)	63.5 (8-400)	115 (4-753)
Hepatomegaly , n (%)	27 (6)	12 (5)	15 (7.5)
Splenomegaly , n (%)	44 (10.5)	18 (8)	26 (13)
Lymphadenopathy , n (%)	110 (26)	26 (12)	84 (42)
Mediastinal mass , n (%)	96 (23)	2 (1)	94 (47)
CNS involvement , n (%)	37 (9)	12 (5)	25 (12.5)
Other involved sites , n (%)	5		
testis/ovary:skin	1:4	-	-
Immunophenotype , n (%)			
B : pro, common, pre, MPAL, UND	221 (52.5)	39,130,23,5,24	
T : ETP, pro, pre, cortical, mature, MPAL, UND	200 (47.5)		40,15,23,57,13,4,48
Citogenetics/genetics , n (%)			
Evaluable, n (%)	342 (81.2)	191 (86.4)	151 (75.5)
Normal	174 (51)	78	96
Adverse	60 (17)	45	15
- t(4;11)/KMT2A::AFF4, t(11;19)	15	15	-
- Other ^a	45	30	15
Non-adverse	108 (32)	67	40
- t(1;19)/E2A::PBX1	5	5	-
- Hyperdiploid	15	15	0
- Other non-adverse	88	48	40
Risk stratification , n (%)			
Evaluable 420/421 (99.8%)			
Standard risk (SR)	207 (49)	120 (54)	87 (44)
High risk (HR)	42 (10)	34 (15)	8 (4)
Very high risk (VHR)	171 (41)	67 (30)	104 (52)

Abbreviations: ALL, acute lymphoblastic leukemia; LL, lymphoblastic lymphoma; ECOG PS, Eastern Cooperative Oncology Group performance status; NA, not available; WBC, white blood cells; PB, peripheral blood; BM, bone marrow; CNS, central nervous system; UND, undefined; ETP: early T precursor

^a other than t(4;11)/KMT2A rearrangement: 11q23, +8, -7, del6q, t(8;14) abnormalities, low hypodiploidy (30-39 chromosomes), near triploidy (60-78 chromosomes) or complex karyotype with ≥5 unrelated anomalies

TABLE 2. Summary of response according to different time points.

	All patients (n=421)	B-ALL (n=216)	T-ALL (n=155)	LL (n=50)
Response at TP1 (in available cases)	379	209	147	23
MRD-pos or unk CR (%)	181 (48)	108 (52)	66 (45)	7 (30)
MRD-neg CR (%)	175 (46)	91 (43.5)	71 (48)	13 (57)
No CR (%)	23 (6)	10 (5)	10 (7)	3 (13)
Response at TP2 (in available cases), n (%)	344*	169	128	47*
MRD-pos CR (%)	70 (20)	44 (26)	25 (19.5)	1 (2)
MRD-neg CR (%)	231 (67)	115 (68)	96 (75)	20 (43)
MRD-unk CR (%)	23 (7)*	4 (2)	2 (2)	17 (36)*
No CR (%)	20 (6)*	6 (4)	5 (4)	9 (19)*
Response at TP3 (in available cases), n (%)	168	89	64	15
MRD-pos CR (%)	15 (9)	7 (8)	8 (12.5)	0
MRD-neg CR (%)	144 (86)	77 (86.5)	52 (81)	15 (100)
MRD-unk CR (%)	4 (2)	4 (4.5)	0	0
No CR (%)	5 (3)	1 (1)	4 (6)	0
Response at TP4 (in available cases), n (%)	141**	69	43	29**
MRD-pos CR (%)	7 (5)	2 (3)	4 (9)	1 (3)
MRD-neg CR (%)	112 (79)	61 (88)	39 (91)	12 (41)
MRD-unk CR (%)	19 (13.5)**	5 (7)	0	14 (48)**
No CR (%)	3 (2)**	1 (1.5)	0	2 (7)**

Abbreviations: ALL, acute lymphoblastic leukemia; LL, lymphoblastic lymphoma; MRD-pos, measurable residual disease positive; MRD-neg, measurable residual disease negative; MRD-unk, measurable residual disease unknown; CR, complete remission

*including 23 patients evaluated only with PET scan (17 CR, 6 not CR) with bone marrow MRD not evaluable

** including 16 patients evaluated only with PET scan (14 CR, 2 not CR) with bone marrow MRD not evaluable

TABLE 3. Summary of pegaspargase-related toxicity

Study parameter	Course 1	Course 2	Course 5	Course 6
Patients	421	382	203	167
Had pegaspargase, n (%)	382/421 (91)	314/382 (82)	177/203 (87)	129/167 (77)
Not administered for previous pegaspargase toxicity, n (%)	NA	47/382 (12)	19/203 (9)	31/167 (19)
Pegaspargase dosing, n (%)				
• Reduced	61/382 (16)	86/314 (27)	50/177 (28)	45/129 (35)
• Full Dose	321/382 (84)	228/314 (73)	127/177 (72)	84/129 (65)
Had pegaspargase-related toxicity grade ≥ 2 , n (%)	189/382 (49)	101/314 (32)	67/177 (38)	39/129 (30)
• Hepatobiliary $G_{\geq 2}/\geq 3$, n (%)	96 (25) / 59 (15)	47 (15) / 15 (5)	36 (20) / 17 (10)	19 (15) / 10 (8)
• Pancreatic $G_{\geq 2}/\geq 3$, n (%)	21 (6) / 11 (3)	1 (0.5) / 1 (0.5)	4 (2) / 3 (2)	0 / 0
• Thrombosis $G_{\geq 2}/\geq 3$, n (%)	7 (2) / 6 (2)	5 (2) / 4 (1)	0 / 0	0 / 0
• Coagulopathy $G_{\geq 2}/\geq 3$, n (%)	81 (21) / 24 (6)	57 (18) / 13 (4)	37 (21) / 10 (6)	22 (17) / 6 (5)
• Metabolic $G_{\geq 2}/\geq 3$, n (%)	13 (3) / 9 (2)	7 (2) / 4 (1)	6 (3) / 4 (2)	2 (2) / 1 (1)

Abbreviations: NA, not applicable

TABLE 4. Comparison between the study results and the GIMEMA LAL1913 clinical trial.

	Campus ALL study (421 patients)	GIMEMA LAL1913 trial (203 patients)	P
Age (years), median (range)	42.0 (18-80)	39.8 (18-65)	0.5000
≤40, n (%)	199 (47)	103 (51)	0.4773
>40-55, n (%)	125 (30)	61 (30)	0.9867
>55, n (%)	97 (23)	39 (19)	0.3300
Diagnosis , n (%)			
ALL	371 (88)	183 (90)	0.5702
LL	50 (12)	20 (10)	0.5200
B-ALL/T-ALL	221/200	139/64	0.0002
WBC (10 ⁹ /l), median (range)	9.9 (0.2-626.9)	7.1 (1.5-347.3)	
≤30 (%)	304 (72)	159 (78)	0.1251
>30-100 (%)	66 (16)	31 (15)	0.9910
>100 (%)	51 (12)	13 (6)	0.0395
CNS involvement , n (%)	37 (9)	19 (9)	0.9559
Risk stratification , n (%)			
Standard risk (SR)	207 (49)	115 (58)	0.0994
High risk (HR)	42 (10)	20 (10)	0.9175
Very high risk (VHR)	171 (41)	68 (33)	0.0951
CR at TP1 (%)	94	85	0.0004
MRD negativity at TP2 (cases available) (%)	72	80	0.0401
Refractory patients before course 3 (%)	6	3	0.2030
First line HSCT rate (%)	35	28	0.0921
Median follow-up (months)	25	39	
3-year OS (%)	67	67	0.9400
3-year DFS (%)	57	63	0.1700
Pegaspargase toxicity (%)			
Grade ≥2 hepatic toxicity at C1	25	12	0.0003
Grade ≥3 pancreatic toxicity at C1	3	1	0.2592
Grade ≥3 thrombosis at C1	2	2	0.9988

Abbreviations: ALL, acute lymphoblastic leukemia; LL, lymphoblastic lymphoma; WBC, white blood cells; CNS, central nervous system; CR, complete remission; MRD, measurable residual disease; HSCT, allogeneic stem cell transplantation; OS, overall survival; DFS, disease free survival

Figure legends

FIGURE 1. Overall survival (OS) and disease-free survival (DFS). A and B) OS and DFS of the entire population. C and D) OS and DFS stratified for age (≤ 40 , 41-55, > 55).

Abbreviations: NR, not reached

FIGURE 2. Disease-free survival (DFS) according to MRD and transplant (HSCT) realization. A) DFS stratified for MRD status. B) Simon-Makuch plot of DFS of VHR and MRD-positive patients according to HSCT


Abbreviations: MRD, measurable residual disease; HSCT, allogeneic stem cell transplantation; VHR, very high risk

FIGURE 3. Univariate and multivariate analysis. A) Univariate for OS. B) Univariate analysis for DFS. C) Multivariate analysis for OS. D) Multivariate analysis for DFS

Abbreviations: OS, overall survival; DFS, disease-free survival; ECOG PS, Eastern Cooperative Oncology Group performance status; LL, lymphoblastic lymphoma; ALL, acute lymphoblastic leukemia; WBC, white blood cell; CNS, central nervous system; Cyto, cytogenetics; MRD, measurable residual disease; SR, standard risk; HR, high risk; VHR, very high risk


FIGURE 4. Comparison of overall survival (OS) and disease-free survival (DFS) between the study population (421 cases) and the GIMEMA LAL1913 clinical trial population (203 cases).

Abbreviations: RL-LAL1913, real life population; LAL1913, GIMEMA LAL1913 clinical trial





Number at risk

421	281	185	124	83	32	7	0
-----	-----	-----	-----	----	----	---	---





Number at risk

392	209	135	90	59	27	3	0
-----	-----	-----	----	----	----	---	---




Number at risk

Group: ≤40							
199	139	88	55	36	12	3	0
Group: 41-55							
125	89	64	43	32	13	3	0
Group: >55							
97	53	33	26	15	7	1	0




Number at risk

Group: ≤40							
190	103	62	40	23	10	0	0
Group: 41-55							
117	67	49	35	27	12	2	0
Group: >55							
85	39	24	15	9	5	1	0



Number at risk

Group: MRDneg	254	156	106	71	49	22	2	0
Group: MRDpos	116	42	23	14	7	4	1	0




Number at risk

No HSCT	212	14	5	1
HSCT	0	65	28	8

A**Univariate OS**


age (18-40 vs 40+)
 age (18-55 vs 55+)
 female vs male
 ECOG PS 0-1 vs >1
 LL vs ALL
 Lineage T vs B
 WBC >30 vs ≤30
 CNS yes vs no
 Cyto non adv vs adverse
 Cyto normal vs adverse
 KMT2A-r pos vs neg
 MRD pos vs neg
 risk HR vs SR
 risk VHR vs SR



HR	CI	P
0,6205	0,4251-0,9056	0,0138
0,5260	0,3563-0,7766	0,0013
0,6863	0,4649-1,0131	0,0594
0,9677	0,5926-1,5804	0,8963
0,7878	0,4345-1,4283	0,4343
1,1647	0,8091-1,6766	0,4145
1,3746	0,9336-2,0238	0,1087
1,9909	1,1758-3,3711	0,0108
0,6329	0,3917-1,0228	0,0631
0,6243	0,3733-1,0441	0,0740
1,4445	0,5878-3,5500	0,4251
3,0551	2,0210-4,6182	<0,0001
1,3019	0,7100-2,3871	0,3962
1,2843	0,8728-1,8898	0,2065

B**Univariate DFS**


age (18-40 vs 40+)
 age (18-55 vs 55+)
 female vs male
 ECOG PS 0-1 vs >1
 LL vs ALL
 Lineage T vs B
 WBC >30 vs ≤30
 CNS yes vs no
 Cyto non adv vs adverse
 Cyto normal vs adverse
 KMT2A-r pos vs neg
 MRD pos vs neg
 risk HR vs SR
 risk VHR vs SR



HR	CI	P
0,7724	0,5515-1,0819	0,1315
0,6133	0,4226-0,8899	0,0104
0,7662	0,5394-1,0884	0,1390
0,7593	0,4391-1,1690	0,2134
0,5939	0,3129-1,1270	0,1127
1,0009	0,7169-1,3975	0,9958
1,7432	1,2329-2,4646	0,0018
1,7000	1,0107-2,8596	0,0466
0,5610	0,3589-0,8769	0,0116
0,5709	0,3549-0,9183	0,0215
2,4052	1,1722-4,9353	0,0173
3,3633	2,3864-4,7000	<0,0001
0,9703	0,5252-1,8001	0,9292
1,4804	1,0442-2,0988	0,0284

C**Multivariate OS**


age (18-40 vs 40+)
 MRD pos vs neg
 risk VHR+HR vs SR





HR	CI	P
0,7190	0,4528-1,1417	0,1641
2,9925	1,9599-4,5693	<0,0001
0,9486	0,6222-1,4461	0,8071

D**Multivariate DFS**

age (18-40 vs 40+)
 MRD pos vs neg
 risk VHR+HR vs SR



HR	CI	P
0,7531	0,5108-1,1103	0,1543
3,2157	2,2671-4,5612	<0,0001
1,1246	0,7931-1,5948	0,5119



SUPPLEMENTAL METHODS

Risk classification according to the GIMEMA LAL1913 clinical trial:

- Very high risk (VHR): WBC count $>100 \times 10^9/L$ or adverse cytogenetics/molecular biology such as t(4;11)/MLL rearrangement at 11q23, +8, -7, del6q, t(8;14), low hypodiploidy with 30-39 chromosomes, near triploidy with 60-78 chromosomes, karyotype with >5 unrelated anomalies, or an early/late non-cortical immunophenotype EGIL T-I/II/IV (CD1a negative) for T-precursor ALL
- High risk (HR): complete remission after the second cycle or, for B precursor ALL, a WBC count $>30 \times 10^9/L$ or a pro-B immunophenotype.
- Standard risk (SR): for B-precursor ALL a WBC count $<30 \times 10^9/L$, for T-precursor ALL a WBC count $<100 \times 10^9/L$ and a cortical immunophenotype EGIL T-III (CD1a+).

Response evaluation criteria: the CR was defined as the disappearance of clinical and laboratory signs of ALL/LL, including extramedullary disease if previously detected; a transfusion-free status with neutrophils $>1.0 \times 10^9/l$ and platelets $>100 \times 10^9/l$; and a normocellular or regenerating bone marrow with blast cell content $<5\%$. A recurrence was defined as the reappearance of $>5\%$ marrow leukemic cells and/or an extramedullary involvement. In cases of LL without marrow involvement and therefore MRD monitoring, the evaluation of response was carried out with PET scans at TP2 and TP4. Early death was defined as death not due to disease occurring before the third course of CHT.

SUPPLEMENTAL TABLE 1. Treatment protocol.

Treatment phase	Drugs	Dosing	Days
Prephase	Prednisone Cyclophosphamide	20 mg/m ² q12h 300 (200 if age >55) mg/m ²	-5 to -1 -3 to -1
Course 1 (C1)	Idarubicin Vincristine Dexamethasone Pegaspargase IT prophylaxis	12 (9 if age >55) mg/m ² 1.4 mg/m ² (max. 2 mg) 5 mg/m ² q12h 2000 (1000 if age >55) UI/ m ²	1,2 1,8,15,22 1-5, 15-19 10 1,15
Course 2,4,6 (C2, C4, C6)	Vincristine Idarubicin Cyclophosphamide Dexamethasone Cytarabine Pegaspargase Mercaptopurine IT prophylaxis	1.4 mg/m ² (max. 2 mg) 12 (9 if age >55) mg/m ² 1000 mg/m ² 5 mg/m ² q12h 75 mg/m ² 2000 (1000 if age >55) UI/ m ² 60 mg/m ²	1,8 (no course 2) 1 1 1-5 2-5 8 (no course 4) 1-10 1 (and 15, course 2)
HD courses 3,7 (C3, C7)	Methotrexate Cytarabine	2500 (B), 5000 (T), 1500 (if age >55) mg/m ² over 24 hours 2000 mg/m ²	1 3,4
HD course 5 (C5)	Methotrexate Pegaspargase Mercaptopurine	2500 (B), 5000 (T), 1500 (if age >55) mg/m ² over 24 hours 2000 (1000 if age >55) UI/ m ² 25 mg/m ²	1 3 8-18
Course 8 (C8)	Vincristine Idarubicin Dexamethasone Cyclophosphamide Prednisone IT prophylaxis	1.4 mg/m ² (max. 2 mg) 10 (7.5 if age >55) mg/m ² 5 mg/m ² q12h 300 (200 if age >55) mg/m ² 20 mg/m ² q12h	1,8 1,8 1-5 1-3 8-12 1,15
Maintenance courses M1,3,5,7,9,11	Cyclophosphamide Mercaptopurine Methotrexate IT prophylaxis	100 mg/m ² 75 mg/m ² 15 mg/m ²	1-4 8-28 8,15,22 1 (courses 3,5)
Maintenance courses M2,4,6,8,10,12	Vincristine Prednisone Mercaptopurine Methotrexate IT prophylaxis	1 mg/m ² (max. 2 mg) 20 mg/m ² q12h 75 mg/m ² 15 mg/m ²	1 1-5 8-28 8,15,22 1 (courses 2,4)
Maintenance courses M13-24	Mercaptopurine Methotrexate	75 mg/m ² 15 mg/m ²	1-28 1,8,15,22

Abbreviations: IT, intrathecal

SUPPLEMENTAL TABLE 2. Characteristics of the patients in the propensity score matching

Characteristic	Overall, N = 602	RL-LAL1913, N = 419	LAL1913, N = 183	p-value
Age, median (range)	42 (18 - 80)	42 (18 - 80)	41 (18 - 65)	0.5
Sex, n (%)				0.6
<i>Male</i>	349 (58%)	246 (59%)	103 (56%)	
<i>Female</i>	253 (42%)	173 (41%)	80 (44%)	
WBC, median (range)	9 (0 - 627)	10 (0 - 627)	6 (0 - 347)	0.001
<i>Unknown</i>	1	1	0	
Lineage, n (%)				<0.001
<i>B</i>	359 (60%)	221 (53%)	138 (75%)	
<i>T</i>	243 (40%)	198 (47%)	45 (25%)	
Risk, n (%)				0.6
<i>SR</i>	302 (50%)	207 (49%)	95 (52%)	
<i>HR</i>	63 (10%)	42 (10%)	21 (11%)	
<i>VHR</i>	237 (39%)	170 (41%)	67 (37%)	
Cytogenetics, n (%)				0.026
<i>Normal</i>	219 (48%)	173 (51%)	46 (39%)	
<i>Non adverse</i>	148 (32%)	108 (32%)	40 (34%)	
<i>Adverse</i>	93 (20%)	60 (18%)	33 (28%)	
<i>Unknown</i>	142	78	64	
CR, n (%)				0.2
<i>CR</i>	556 (92%)	391 (93%)	165 (90%)	
<i>no CR</i>	46 (7.6%)	28 (6.7%)	18 (9.8%)	
Transplant, n (%)	200 (33%)	144 (34%)	56 (31%)	0.4
Time to transplant, median (range)	6.12 (2.93 - 28.88)	6.04 (2.93 - 28.88)	6.41 (3.72 - 12.20)	0.2
<i>Unknown</i>	402	275	127	