

LCC: QA1-939
UDC: 517.9

MODULATED CHAOS AS A SOURCE OF IMAGES FOR NUMBERS

J.C. Tiago de Oliveira

Department of Mathematics, University of Évora
Institute of Scientific Research Bento da Rocha Cabral (Portugal)
E-mail: zct@uevora.pt

ABSTRACT

Mario Markus, a Chilean scientist and artist from Dortmund Max Planck Institute, has exposed a large set of images of Lyapunoff exponents for the logistic equation modulated through rhythmic oscillation of parameters. The pictures display features like foreground/background contrast, visualizing superstability, structural instability and, above all, multistability, in a way visually analogous to three-dimensional representation.

See, for instance, <http://www.mariomarkus.com/hp4.html>.

The present papers aims to classify, through codification of numbers in the unit interval, the ensemble of images thus generated. The above is intended as a part of a still unfulfilled work in progress, the classification of style in visual fractal images-a common endeavour to Art and Science.

Key words: chaos, fractals, pictures of Lyapunoff

1. INTRODUCTIONS

The pictures of Lyapunoff exponents, published by Markus et al (1), corresponding to symbolic sequences which control a quadratic – or topologically conjugate – iteration, is, through the alternative nature of parameters A and B, equivalent to a succession of zeros and ones. For such a symbolic sequence, the natural topology is the one of a Cantos set.

Now, it is also true that the infinite ensemble generated by sequencing zero and one is the transcription of the unit interval in basis 2; to what conditions is such a sequence bound to obey, in order to designate a generic real number, instead of a point in a meagre, zero – measure set such as a Cantor set?

Are such conditions compatible with the hypothesis basilar to the works of Markus?

If so it was, this would be another instantiation of a new paradigm – constructing fractals as number images.

A procedure developed in divergent ways by a few authors:

- **Jenny Harrison** (2), through a construction inspired by Denjoy and leading into continued fractals
- **Michel Mendes-France** (3), when he considered number-theoretic properties of sequences of coefficients for trigonometric sums.
- The early departed **Mário Sarreira and José Sousa Ramos** (4) in their codification of a cubic Mandelbrot set.

2. DEVELOPMENT

(1) $0.111111\dots = 1.000\dots$ is the constitutive equation of the binary number system. For a radix $b > 2$, it will not happen. $0.111\dots = \frac{1}{b} + \frac{1}{b^2} + \dots < 1.000\dots$; therefore, the free semigroup generated by the two symbols 0 and 1 will define a Cantos set.

The inverse inequality ($0.1111\dots > 1.00\dots$) is accomplished for $b < 2$, were a multiplicity of developments, with a magnitude dependent on the location, is present for each point of the unit segment, thus structuring it as a multifractal.

Equation (1) fills the same role as the “just-touching collage” in Barnsley (5); similar conditions may be found in Williams, Dubuc, Hata (6, 7, 8), along with iterated function systems, whose sum is the unit matrix, giving rise to connected sets.

The apparent tridimensionality of the pictures in (1) is a consequence of the difference between the basins of attraction, the cardinality of which is conjectured to be equal to the periodicity in the control signal (the A B sequence).

Such a conjecture would validate the following corollaries:

- a)The existence of a shift symmetry in the control sequence, with the same period as it
- b)The irrelevance of the first N terms of the control sequence, for whatever N, in periodic A B sequences

$$\begin{array}{lcl}
 \text{(duality =} & 1 \rightarrow 2 \rightarrow 4 \rightarrow 8 & \text{(shift = product by 2)} \\
 (2^4 - 1) - X = & \downarrow \downarrow \downarrow \downarrow & \\
 = 15 - X & 14 \quad 13 \quad 11 \quad 7 &
 \end{array}$$

As well as

$$\begin{array}{lcl}
 3 \rightarrow 6 \rightarrow 12 & & \\
 \downarrow \downarrow \downarrow & & \\
 12 \quad 9 \quad 3 & &
 \end{array}$$

As a convention, we shall designate them as the 8 and 12 sequences, in order to enhance the leading digit 1

$$\begin{array}{lcl}
 8 & : & 1000 \\
 12 & : & 1100
 \end{array}$$

Period 5:

- Exclusion of 00000 and 11111
 - No repeating of sub multiple sub sequences as 5 is a prime number
 - Generation of their respective sequences
- Shift = 2x, Duality = 31-x

$$\begin{array}{lcl}
 1 \rightarrow 2 \rightarrow 4 \rightarrow 8 \rightarrow 16 & & \\
 \downarrow \downarrow \downarrow \downarrow \downarrow & & \\
 30 \quad 29 \quad 27 \quad 23 \quad 15 & &
 \end{array}$$

Also, there is

$$\begin{array}{lcl}
 3 \rightarrow 6 \rightarrow 12 \rightarrow 24 & & \\
 \downarrow \downarrow \downarrow \downarrow & & \\
 28 \quad 25 \quad 19 \quad 7 & &
 \end{array}$$

Which merges with

$$\begin{array}{lcl}
 14 \rightarrow 28 & & \\
 \downarrow \downarrow & & \\
 17 \quad 3 & &
 \end{array}$$

In fact, the numbers subsuming them (24,17) represent the same pattern, a just from a shift:

$$\begin{array}{lcl}
 24 & : & 11000 \\
 17 & : & 10001
 \end{array}$$

The same phenomenon occurs between

$$\begin{array}{lcl}
 5 \rightarrow 10 \rightarrow 20 & & 9 \rightarrow 18 \\
 \downarrow \downarrow \downarrow & \text{and} & \downarrow \downarrow \\
 26 \quad 21 \quad 11 & & 22 \quad 13
 \end{array}$$

$$20 \quad : \quad 10100 \quad \text{being equivalent by 2 Shift to} \quad 18 \quad : \quad 10010$$

Therefore, we shall select, as tokens for periods sequences:

$$\begin{array}{lcl}
 16 & : & 10000 \\
 17 & : & 10001 \\
 18 & : & 10010
 \end{array}$$

Period 6: 64 sequences; elimination of 0 and 1, as usual

6 being a multiple of 2 and 3, we are bound to eliminate repetitions of segments 10 and 100, live





$$\begin{array}{lcl}
 101010 & : & 42 \\
 100100 & : & 36
 \end{array}$$

But it proves better to start by their duals

$$\begin{array}{lcl}
 010101 & : & 21 \\
 011011 & : & 27
 \end{array}$$










54 numbers remain, to be distributed.



Therefore the sequences will be codified as:


- 32 : 100000
- 33 : 100001
- 34 : 100010 (or 56 = 111000)
- 35 : 100011
- 36 : 100101

Period 7: excluding 0 and 1, and there being no submultiples, we need to explore 126 numbers



15 and 17 are not suitable generators, as previously they were found. We might remark that these sequences always have 14 elements; therefore, we need 2 more ($7 + 2 = 9$, $9 \times 14 = 126$).

The last elements in the 1st line grow according to arithmetic progression (647,70).
 71 is present in one previous sequence, as is 72.
 So, 73 and 74 should be good generators.



Until now, we have the followings symbolic sequences

Periods

| 1, 2, 3 | 4 | 5 | 6 | 7 |
|----------|-----------|----------|----|----|
| 1 (= 1 | 8 (=1000 | 16-10000 | 32 | 64 |
| 2 (= 10 | 9 (= 1001 | 17-10001 | 33 | 65 |
| 4 (= 100 | | 18-10010 | 34 | 66 |
| | | | 35 | 67 |
| | | | 37 | 68 |
| | | | | 69 |
| | | | | 70 |
| | | | | 73 |
| | | | | 74 |

5. GENERALIZING

If P is a prime number, this having no sub-multiples, the only excluded sequences are 0 and $2^P - 1$ (0000... and 1111...).
 Therefore, if every single set of isomorphic numbers (in terms of duality + shift) has the same quantity of terms, as they did until now, their quality must depend on the factorization of $2^P - 2 = 2(2^P - 1)$.
 Let us try it for the next prime numbers: 11, 13, 17.

11: $2^{11} = 2048$; $2^{11} - 2 = 2046 = 2 \times 1023 = 6 \times 341 = 6 \times 31 \times 11 = 11 \times 186 = 22 \times 93$


Therefore, there should be 93 ensembles of 22 equivalent numbers.

13: $2^{13} - 2 = 8190 = 10 \times 819 = 13 \times 10 \times 63 = 26 \times 315$
 315 equivalent segments

17: $(2^{(17-1)}-1)$ being divisible by 17, this quotient should be the member of different sequences.

For composite numbers, the problem is my difficult, we should eliminate the instantiation of sub-multiples. For instance, in the period 8 ($2^8 = 256$) we eliminate:

- 0 and 255
- The translates of period 2 (10101010 and 01010101)
- The duals and translate of 8 and 9



9, along with fifteen others.
 Therefore, $256 - 2 - 2 - 16 - 16 = 220$ will remain, to be distributed through 16 – long ensembles.

6. CONCLUSIONS

The number of image-generations sequences grows with N in a was majorated by $(2^{(P-1)} - 1) / P$, for prime p; which means, exponentially in N.
 Therefore, it is not a denumerable set. Is it a continuation, or a Cantor set, or something in between (a “fat” Cantor set) ?

As a tentative answer, let us be inspired by the following:

- A number such as 0.897897897... in decimal radix expansion is $= 897.0.001001001... = 897 / 999$

Therefore, in fractal number system, the corresponding numbers are to be

$1 / 1, 2 / 3, 4 / 7, 8 / 19, 9 / 15 = 3 / 5$

$16 / 31, 17 / 31, 18 / 31, 32 / 63, 33 / 63 = 11 / 21$

$34 / 63, 35 / 63, 37 / 63, 64 / 127, 65 / 127, 66 / 127, 67 / 127, 68 / 127, 69 / 127, 70 / 127, 73 / 127, 74 / 127$

How for did such a sequence fill the $[\frac{1}{2}, 1]$ interval? Is it mifuncly deuse? Fat or meagre? What about its Hausdorff dimension?

Sequences, essentially because shift symmetric fails (it was a consequence of periodicity); therefore applet tridimensionality (multimodality) is lost.

However, duality, as well as irrelevance of transients, are maintained.

We might understand for the above that, restrictions of the a periodic, being less severe, on stating hypothesis – Markus pictures as images for numbers – which was not refuted in the realm of dyadic rrationals, maintains the pertinence of a conjecture.

REFERENCES

1. **Rosler Kiwi Hess Markus** (1989) Modulated Non-linear Processes and a Novel Mechanism to Induce Chaos Phiss. Rev. A, 39, 11
2. **Harrison** (1989) In R.L. Devanel ed, Chaos and Fractals, AMS
3. **Mendes France** (1991) La Constante de Plack d'une Comba Fractal Geometry and analysis
4. **Sarreira, Sousa Ramos** (1991) Symbolic Dynamics of Hercition of Cubic Complex Maps
5. **European Conference on Heration Theory**
6. **Barnsley** (1989) Fractals Everywhere
7. **Williams** (1992) The composition of Constructive Transposition Boletim Sociedade Matemática-Brasil
8. **Dahac** (1987) In Bull. Soc. Math. Quebec
9. **Hata** (1991) In Fractal Geometry and Analysis