

Available online at www.sciencedirect.com

Tectonophysics 461 (2008) 114–129

TECTONOPHYSICS

www.elsevier.com/locate/tecto

New insights from U–Pb zircon dating of Early Ordovician magmatism on the northern Gondwana margin: The Urra Formation (SW Iberian Massif, Portugal)

A.R. Solá ^{a,*}, M.F. Pereira ^b, I.S. Williams ^c, M.L. Ribeiro ^a, A.M.R. Neiva ^d, P. Montero ^e, F. Bea ^e, T. Zinger ^f^a Departamento de Geologia, INETI, Apt. 7586, 2720-866 Alfragide, Portugal^b Centro Geofísica Évora, Departamento Geociências, Universidade de Évora, Apt 94, 7002-554 Évora, Portugal^c Research School of Earth Sciences, Australian National University, Canberra, ACT, 0200, Australia^d Departamento de Ciências da Terra, Universidade de Coimbra, 3000-272 Coimbra, Portugal^e Departamento de Mineralogía y Petrología, Universidad de Granada, Campus Fuentenueva, 18002 Granada, Spain^f Institute Precambrian Geology and Geochronology, RAS, Makarova, Emb.2, 199034, St. Petersburg, Russia

Received 12 May 2007; received in revised form 20 November 2007; accepted 27 January 2008

Available online 7 February 2008

Abstract

The Central Iberian–Ossa-Morena transition zone (SW Iberian Massif) represents a segment of the northern Gondwana margin with a long geodynamic evolution, characterized by the superposition of Cadomian and Variscan events. The Early Ordovician is mainly represented by porphyritic felsic volcaniclastic rocks (the Urra Formation) that pass up into a siliciclastic sediments typical of the Central Iberian Zone (Lower Ordovician Armorican Quartzite Formation). The Urra Formation unconformably overlies the previously deformed and metamorphosed Ediacaran sediments of the Série Negra (with Ossa-Morena Zone paleogeographic affinity). New SHRIMP zircon data obtained from the Urra Formation volcaniclastic rocks indicate an Early Ordovician age ($^{206}\text{Pb}/^{238}\text{U}$ ages ranging from 494.6 ± 6.8 Ma to 488.3 ± 5.2 Ma) for this magmatic event. The inherited zircon cores indicate the presence of multicycle protoliths with different Precambrian ages: Neoproterozoic (698–577 Ma), Paleoproterozoic (2.33 Ga) and Paleoarchean (3.2–3.3 Ga). There is a noticeable lack of Meso- to Neoproterozoic ages. The data support the hypothesis that the volcaniclastic rocks were derived by partial melting of Cadomian basement (linked to a West African Craton provenance). The Urra Formation volcaniclastic rocks have rhyolitic to dacitic compositions, are peraluminous and similar to calc-alkaline high-K series suites elsewhere. Isotopic signatures present a wide range of values ($^{87}\text{Sr}/^{86}\text{Sr}_i = 0.7085\text{--}0.7190$, more restricted ϵNd_i (–2.65 to –0.35) and $\delta^{18}\text{O} = 9.63\text{--}10.34\text{‰}$, compatible with magmas derived from crustal rocks, including portions of the lower crust. Some samples show disturbance of the Rb–Sr system as shown by unrealistic values for ($^{87}\text{Sr}/^{86}\text{Sr}_i < 0.703$, probably due to Variscan deformation and metamorphism. The volcaniclastic rocks with a significant sedimentary contribution (upper unit) are distinguished from the others by the lowest values of ϵNd_i (–5.53 to –4.85). The geochemical data are compatible with an orogenic geodynamic environment. However, the “orogenic” signature can be considered to represent, in part, an inherited feature caused by melting of the Cadomian basement which also has calc-alkaline affinities. The Early Ordovician crustal growth and associated magmatism, represented by the Urra felsic volcaniclastic rocks and associated calc-alkaline granitoids, diorites and gabbros, can be interpreted in terms of the underplating and temporal storage of mantle-derived magmas as the potential source for the “orogenic melts” that were intruded during Early Paleozoic extension. This record of Early Ordovician magmatism has striking similarities with other correlatives from the Iberian, Bohemian and Armorican massifs that are discussed in this paper. This comparison reinforces the probable existence of a large-scale crustal melting process linked to a significant episode of extension on the northern Gondwana margin that probably resulted in the birth of the Rheic Ocean.

© 2008 Elsevier B.V. All rights reserved.

Keywords: Early Paleozoic rifting; Calc-alkaline magmatism; Central Iberian–Ossa-Morena transition zone; Northern Gondwana; Rheic Ocean

* Corresponding author. Tel.: +351 214705455; fax: +351 214718941.

E-mail address: rita.sola@ineti.pt (A.R. Solá).