

Find out how to access preview-only content

Look Inside Get Access

Close

European Journal of Wildlife Research

October 2008, Volume 54, Issue 4, pp 675-680

A half a century of measuring ungulate body condition using indices: is it time for a change?

Abstract

From a literature review of five wildlife ecology journals since 1937, we document how using indices to monitor ungulate body condition is common practice, with the kidney fat index (KFI = weight of fat around the kidneys/weight of kidneys without fat \times 100) as the favoured tool (82% of studies). In this context, we highlight the problems of using indices when underlying statistical assumptions are not met (isometry, parallel slopes between treatments). We show, with real and simulated data for two cervids with contrasting fat storage strategies, how results from analysis of variance of KFI values differ from analysis of covariance (ANCOVA) of raw data. We conclude that the KFI is affected by the restrictions typically associated with derived index values, and as a consequence, statistical analysis of the KFI could generate spurious results leading to erroneous interpretations concerning variation in body condition of ungulate populations. Thus, we recommend analysing fat weight as an untransformed variable in ANCOVA (kidney weight as covariate) to describe body condition variation in ungulates.

Communicated by H. Kierdorf



Within this Article:

1. Introduction
2. Materials and methods
3. Results
4. Discussion
5. References
6. References

Related Content



Supplementary Material (1)

References (24)

About this Article

Title

A half a century of measuring ungulate body condition using indices: is it time for a change?

Journal

European Journal of Wildlife Research
Volume 54, Issue 4 , pp 675-680

Cover Date

2008-10-01

DOI

10.1007/s10344-008-0194-7

Print ISSN

1612-4642

Online ISSN

1439-0574

Publisher

Springer-Verlag

Additional Links

- [Register for Journal Updates](#)
- [Editorial Board](#)
- [About This Journal](#)
- [Manuscript Submission](#)

Topics

- [Animal Ecology](#)
- [Zoology](#)

Keywords

- ANCOVA
- *Capreolus capreolus*
- *Dama dama*
- Kidney fat index
- Ratios
- Residuals
- Proportions

Authors

- Emmanuel Serrano ⁽¹⁾
- Russell Alpizar-Jara ⁽²⁾
- Nicolas Morellet ⁽¹⁾
- Aidan Jonathan Mark Hewison ⁽¹⁾

Author Affiliations

- 1. Equipe Ecologie des Populations, Laboratoire de Comportement et Ecologie de la Faune Sauvage, Institut National de la Recherche Agronomique (CEFS-INRA), Chemin de Borde-Rouge-Auzeville, B.P. 52627, Castanet-Tolosan Cedex, 31326, France
- 2. Departamento de Matemática e Centro de Investigação em Matemática e Aplicações (CIMA-U.E.), Universidade de Évora, Rua Romão Ramalho, 7000-671, Évora, Portugal

6.031.506 scientific documents at your fingertips
© Springer, Part of Springer Science+Business Media

You have been redirected to our new and improved site.

More info [I'm good, don't tell me again](#)
.springer.com