
On: 28 December 2014, At: 13:01 Publisher: Routledge Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



International Journal of Psychology

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/pijp20</u>

Invited Address, IUPsyS Invited Symposium, Invited Symposium, Symposium, Paper Session, Poster Session Published online: 02 Nov 2010.

To cite this article: (2008) Invited Address, IUPsyS Invited Symposium, Invited Symposium, Symposium, Paper Session, Poster Session, International Journal of Psychology, 43:3-4, 528-707, DOI: <u>10.1080/00207594.2008.10108486</u>

To link to this article: http://dx.doi.org/10.1080/00207594.2008.10108486

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Mexican elementary school 6-years-old children (N= 1500), from medium and low socioeconomic levels. It also identifies preferential playing games at home and at school, by boys and girls. Results of a questionnaire and the corresponding quantitative analysis are presented (SPSS, 12.0), showing favorite play places (at school, at home or outdoors), preferences and playing styles.

School failure in Mexican children: Effects of preacademic and linguistic behavioral levels

Guevara, Yolanda Dept. of Psychology, National University Mexico, Mexico City, Mexico Lopez, Alfredo Dept. of Psychology, National University Mexico, Mexico City, Mexico Huitrón, Blanca Dept. of Psychology, National University Mexico, Mexico City, Mexico Delgado, Ulises Dept. of Psychology, National University Mexico, Mexico City, Mexico Hermosillo, Angela Dept. of Psychology, National University Mexico, Mexico City, Mexico

Several studies have proved that, adequate levels of preacademic and linguistic behaviors (Previous knowledge or entry cognitive behaviors) are necessary to avoid school failure. Our research had two phases. 1): 262 children from low socioeconomic status were assessed through behavioral tests to determine linguistic and preacademic skills level, and reading, writing, and math skills, at the beginning, in the middle and at the end of the first school year. Results show a narrow relationship between entry repertoires and level of developed academic skills, with academic problems related to preacademic and linguistic skills. 2): Ten such children were trained in linguistic and preacademic skills, with encouraging results in school achievement.

Treatment of school failure with a program of discipline in the classroom

Vidal Lucena, Margarita Dept. de Investigación, Instituto Calasanz Educación, Madrid, Spain

School failure has been related to problems of conduct, among other causes. In this paper we present the first results of a program of discipline in the classroom, that has been designed on the basis of several pilot studies carried on in a Spanish school. This program is based on cognitive theory; it introduces explicit, simple, contextualized rules, according to a series of sequenced levels at school. The goal is to reduce the incidence of factors that are directly related to school failure, by introducing rules of behaviour that decrease the time of conflict and increase instructional time.

Treatment of school failure with behavioral techniques

Galindo, Edgar Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal

School failure, a main problem in Portugal (http:// www.min-edu.pt), is due to many factors, including psychosocial deficiencies. Behavioral procedures have been widely applied to help deficient persons, like Galindo (1999, 2001) with children in Latin-American slums. This experience is here applied in Portugal to 6 - 12 years old children, failing school because of family problems or social exclusion (poverty or ethnic minority). First, a quasi-experimental study (Multiple Baseline) was carried out, with 8 children. Results are evaluated in terms of % of attained objectives, time, and school satisfaction. Procedures were later applied to 20 children.

Treatment of school failure in Portuguese children: Case studies

Marcelino, Lilia Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal Galindo, Edgar Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal Selected case studies of 6-12 year old children living in Lisbon slums, participating in the project "Treatment of school failure with behavioral techniques" are presented. Behavioral diagnostic and treatment procedures were designed according to teachers' aims and consistent with the particular social and/or learning problems of each child. Causes of failure are multiple: family instability, child neglect, lack of Portuguese language (immigrants) or social skills. Results are evaluated by % of attained behavioral objectives, time of training and degree of school/family satisfaction. In most cases, treatment was successful. Nevertheless, remarkable differences were found between objective criteria and the degree of satisfaction.

School failure in Angola, Mozambique and Cape Verde: A project for Africa

Guerra Marques, Sónia Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal Sardinha, Erika Dada, Catia Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal Centeio, Denise Dept. of Psychology, Universidade Lusofona, Lisboa, Portugal African students participating in the project "Treatment of school failure with behavioral techniques' analyze the problem in their countries, where it is associated with mental, sensorial, motor or psychosocial deficiencies and the consequences of war. Percentages of primary school repeaters and survival rate to grade 5 are respectively 10% - 62 % in Mozambique, 15% -93% in Cape Verde and 29%-(unknown) in Angola, a country where additionally about 1,5 million children live in difficult conditions (3,000 on the streets) and 750,000 (310,784 in Mozambique) suffer deficiencies, mostly without attention. A project is presented, which includes a "Training Center for Children at Risk" managed by parents and depending from the local University, whose students apply behavioral techniques.

S-221: Dementia therapy: What can psychology contribute?

Gabriele Wilz, Katja Werheid (chair)

Due to the raising prevalence of dementia worldwide, there is a pressing need to develop supportive intervention strategies for patients affected by dementing disorders and their caregivers. However, research on clinical application and efficacy of psychological interventions in these populations is scarce. The present symposium gathers evidence on a variety of psychological treatments for patients with dementia as well as caregivers, ranging from psychotherapy to focused psychological and social interventions. Scientists from different European countries will present latest evidence on the evaluation of psychological treatments and discuss future developments in this area.

Description and outcomes of a psychological and occupational therapy intervention for dementia family caregivers

Losada, Andres Dept. of Psychology, Universidad Rey Juan Carlos, Madrid, Spain Márquez González, María Biological and Health Psycholo, Universidad Autónoma de Madrid, Madrid, Spain Peñacoba, Cecilia Dept. of Psychology, Universidad Rey Juan Carlos, Madrid, Spain Gallagher Thompson, Dolores School of Medicine, Stanford University, San Francisco, USA Knight, Bob G. Clinical Psychology, University of Southern Califor, Los Angeles, USA Cigarán, Margarita Dept. of Psychology, Universidad Rey Juan Carlos, Madrid, Spain Moreno, Ricardo Dept. of Psychology, Universidad Rey Juan Carlos, Madrid. Spain

Preliminary results of a multicomponent and interdisciplinary (cognitive-behaviour-therapy and occupational-therapy) intervention for dementia caregivers are presented. Caregivers were randomly assigned to an intervention condition (n = 40) or a control condition (n = 50), and blind interviews were conducted. The results show a significant reduction from pre-intervention to post-intervention of depression (p < .01), anxiety (p < .05) and dysfunctional thoughts (p < .01), and a significant increase in pleasurable activities (p < .01) and satisfaction with these activities (p < .01) in those caregivers that participated in the intervention condition. These changes were not significant in the control group.

Evaluation of health effects of assisted vacations for persons with dementia and their spouses

Wilz, Gabriele Inst. für Klin. Psychologie, Technische Universität Berlin, Berlin, Germany Fink-Heitz, Margit Institute of Public Health Res, Technische Universität München, Munich, Germany

Objective: An evaluation of assisted vacations for persons with dementia and their caregivers was conducted for the first time in caregiving research. Method: A quasi-experimental, 2-group design with two measuring times was used to examine whether assisted vacations lead to a reduction in physical complaints and depression in family caregivers (N=29). Results: The overall emotional and physical state of the participants in the intervention group showed significant improvements in comparison with the control group three months after the first interview. Conclusions: Assisted vacations can be seen as a way of diminishing the risk of stress disorders for family caregivers.

Dyadic exchange and well-being in couples with one spouse suffering from dementia

Braun, Melanie Sozial.- und Gesundheitspsy., Universität Zürich, Zürich, Switzerland

Objective: Dementia prevalence increases due to growing life expectancy. Equity and a balanced exchange predict relationship quality and wellbeing. In couples with dementia, equity should be impaired by a modified dyadic exchange. Method: Thirty couples (N=60) with a demented husband participated in the study. Social exchange, equity, and well-being are assessed longitudinally. Objective and subjective methods are combined to analyse dyadic processes. Results: Caregiving wives experience more depression and inequity, and lower satisfaction with life than their husbands. Despite their impairment, the demented subjects participated actively. Conclusions: Relations between exchange, well-being, and health decline comprise essential information for couple interventions.

Evaluation of a cognitive behavioural group intervention program for caregivers

Kalytta, Tanja Fachbereich I, Techn. Fachhochschule Berlin, Berlin, Germany Wilz, Gabriele Inst. für Klin. Psychologie, Technische Universität Berlin, Berlin, Germany

Objectives: The effect of a cognitive behavioural group program was examined in caregivers of dementia patients. Methods: The participants were 71 caregivers who were assigned to either a group receiving treatment or a group receiving usual care (control group). We measured caregivers' resources, body complaints, and depressive symptoms. MAN-OVAs allowed for examination of short-term effects of the intervention. Results: Caregivers who attended the intervention report a raised awareness and understanding of the illness. The participation at the intervention program was significantly associated with in changes body complaints. Conclusions: The findings confirm the importance of interventions for caregivers of dementia patients.

Effective factors in psychosocial interventions in dementia care

Vernooij-Dassen, Myrra Dept. of Clinical Psychology, University of Nijmegen, Nijmegen, Netherlands Many psychosocial interventions in dementia care have been developed over the last decades. Metaanalyses provide evidence on effectiveness of some interventions. No single intervention can address the complexity of needs of dementia patients and their carers. Factors that contribute to effectiveness