



## Green Processing and Synthesis

Editor-in-Chief: Hessel, Volker

[Get eTOC Alert >](#)

[Get New Article Alerts >](#)

[SEE ALL FORMATS AND PRICING](#)

Volume	Issue	Page
<input type="text"/>	<input type="text"/>	<input type="text"/>

### Volume 4, Issue 2 (Apr 2015)

[Previous Article](#) [Next Article](#)

30,00 € / \$42.00 / £23.00

#### ISSUES

#### VOLUME 4 (2015)

Issue 5 (Oct 2015), pp. 341-454  
Special issue: NaMiBIT...

Issue 4 (Aug 2015), pp. 257-339

Issue 3 (Jun 2015), pp. 165-255

Issue 2 (Apr 2015), pp. 67-163  
Special issue: 1st Int...

Issue 1 (Jan 2015), pp. 1-66

#### VOLUME 3 (2014)

Issue 6 (Dec 2014), pp. 397-501  
Special issue: 4th Int...

Issue 5 (Oct 2014), pp. 307-395  
Special issue: 4th Int...

Issue 4 (Aug 2014), pp. 257-306

Issue 3 (Jun 2014), pp. 187-

#### MOST DOWNLOADED ARTICLES

1. [Phytoextraction as a tool for green chemistry](#) by [Hunt, Andrew J./Anderson, Christopher W.N./Bruce, Neil/García, Andrea Muñoz/ Graedel, Thomas E./Hodson, Mark/Meech, John A./Nassar, Nedal T./Parker, Helen L./Ryott,](#)

## Can mitochondrial malondialdehyde content be a useful tool to evaluate sea lamprey juveniles' capacity to seawater acclimatization?

Marta Candeias<sup>1</sup> / Isabel Alves-Pereira<sup>1, 2</sup> / Maria João Lança<sup>1, 3</sup> / Ana Filipa Ferreira<sup>4</sup> / Bernardo R. Quintella<sup>4, 5</sup> / Pedro R. Almeida<sup>4, 6</sup> / Rui Ferreira<sup>1, 2</sup>

<sup>1</sup> ICAAM-Instituto de Ciências Agrárias e Ambientais Mediterrânicas, Universidade de Évora, Évora, Portugal

<sup>2</sup> Departamento de Química, ECT, Universidade de Évora, Évora, Portugal

<sup>3</sup> Departamento de Zootecnia, ECT, Universidade de Évora, Évora, Portugal

<sup>4</sup> MARE – Centro de Ciências do Mar e do Ambiente, Lisboa, Portugal

<sup>5</sup> Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Portugal

<sup>6</sup> Departamento de Biologia, ECT, Universidade de Évora, Évora, Portugal

Corresponding author: Rui Ferreira, ICAAM-Instituto de Ciências Agrárias e Ambientais Mediterrânicas, Universidade de Évora, Évora, Portugal; and Departamento de Química, ECT, Universidade de Évora, Évora, Portugal, e-mail: [\(email\)](#).

**Citation Information:** Green Processing and Synthesis. Volume 4, Issue 2, Pages 133–139, ISSN (Online) 2191-9550, ISSN (Print) 2191-9542, DOI: [10.1515/gps-2014-0087](https://doi.org/10.1515/gps-2014-0087), April 2015

[Request Permissions](#)

### Publication History

Received: 2014-10-31

Accepted: 2015-02-16

Published Online: 2015-04-02

### Abstract

The sea lamprey is an anadromous species that migrates twice during its life cycle between freshwater and seawater. Microphagous larvae generally spend 4–5 years burrowed in the substrate of rivers and streams before undergoing metamorphosis that ends with the beginning of the juvenile trophic migration. Once metamorphosis is complete, sea lamprey juvenile downstream migrants are fully tolerant to seawater salinity. Pollution resulting from industrial effluents may disturb the seawater acclimatization causing oxidative damage, and ultimately may lead to a decrease of sea lamprey population. The aim of this study was to compare salt acclimation of sea lamprey juveniles captured in river basins with different levels of aquatic pollution, using mitochondrial glutathione (GSH) and malondialdehyde (MDA) of gills and liver as markers of physiological stress and cell damage. The results showed that juveniles from the Lima basin exhibited the highest levels of mitochondrial MDA in gills, even though significant changes in the stress markers of mitochondrial gills of all animals subject to salt acclimation were not detected. In addition, an increase in the oxidative damage of hepatic mitochondria of macrophthalmia from the Vouga basin suggests the occurrence of metabolic failures with the potential to disturb the capacity to adaptation to the marine environment.

**Keywords:** cell damage; oxidative stress; *Petromyzon marinus*; seawater acclimation