


Contents lists available at [ScienceDirect](http://ScienceDirect)

## Archives of Gerontology and Geriatrics

journal homepage: [www.elsevier.com/locate/archger](http://www.elsevier.com/locate/archger)

# Role of physical activity, physical fitness, and chronic health conditions on the physical independence of community-dwelling older adults over a 5-year period



Catarina Pereira<sup>a,\*</sup>, Fátima Baptista<sup>b</sup>, Ana Cruz-Ferreira<sup>a</sup>

<sup>a</sup> Departamento de Desporto e Saúde, Escola de Ciências e Tecnologia, Universidade de Évora; Research Centre in Sports Sciences, Health Sciences and Human Development (CIDESD), Portugal

<sup>b</sup> Exercise and Health Laboratory, Interdisciplinary Center for the Study of Human Performance, Faculty of Human Kinetics, University of Lisbon, Portugal

## ARTICLE INFO

## Article history:

Received 27 July 2015

Received in revised form 2 February 2016

Accepted 4 February 2016

Available online 17 February 2016

## Keywords:

Physical functioning

Body composition

Metabolic expenditure

Fitness

Aging

## ABSTRACT

The variability in the individual characteristics and habits could help determine how older adults maintain independence. The impact of the variability in physical activity, physical fitness, body composition, and chronic health conditions (co-morbidities) on the independence of older adults, especially over time, is seldom examined. This study aims to analyze quantitatively the impact of baseline values and changes in physical activity, physical fitness, body composition, and co-morbidities on the physical independence of community-dwelling, older adults over a 5-year period. Data from 106 and 85 community-dwelling adults ( $\geq 60$  years) were collected at baseline and after five years, respectively. Linear regression selected the main predictors of changes in physical independence as follows: the baseline physical independence ( $\beta = 0.032$ ,  $R^2 = 9.9\%$ ) and co-morbidities ( $\beta = -0.191$ ,  $R^2 = 6.3\%$ ) and the changes in co-morbidities ( $\beta = -0.244$ ,  $R^2 = 10.8\%$ ), agility ( $\beta = -0.288$ ,  $R^2 = 6.7\%$ ), aerobic endurance ( $\beta = 0.007$ ,  $R^2 = 3.2\%$ ), and walking expenditure ( $\beta = 0.001$ ,  $R^2 = 5.1\%$ ) ( $p < 0.05$ ). In conclusion, baseline physical independence, baseline co-morbidities, and changes in co-morbidities, walking, agility, and aerobic endurance predicted physical independence over five years regardless of age and gender. Gains of up to 8.3% in physical independence were associated with improvements in these variables, which corresponds to regaining independence for performing one or two activities of daily living.

© 2016 Published by Elsevier Ireland Ltd.

## 1. Introduction

Life expectancies continue to rise worldwide (WHO, 2014). Consequently, more people are likely to live longer with infirmities and disabilities, and therefore, they are likely to lose their physical independence (Crimmins, Hayward, Hagedorn, Saito, & Brouard, 2009; Seeman, Merkin, Crimmins, & Karlamangla, 2010). Nonetheless, an early diagnosis of poor health conditions for which disability preventive measures are available (Crimmins et al., 2009) may also result in increased disability-free life expectancy (Crimmins et al., 2009; Hashimoto et al., 2010). As educated older adults generally adhere to preventive measures (Van Oyen et al., 2011) physical dependence may in fact decrease (Paterson & Warburton, 2010).

Physical dependence indicates a reduced physical capacity to independently perform common activities of daily living (ADLs). ADLs include the basic activities or self-care activities (BADLs); instrumental activities that allow an individual to function independently in a community (IADLs); and advanced or challenging activities that enrich life such as social, religious, and leisure activities (AADLs) (Fujiwara et al., 2008; Rikli & Jones, 1998, 2013). A functional decline in more demanding ADLs predicts future declines in less demanding ADLs, institutionalization, and death (Fujiwara et al., 2008; Vermeulen, Neyens, van Rossum, Spreuwerberg, & de Witte, 2011).

Most factors affecting the ability to perform ADLs involve chronic health conditions, physical activity and fitness as well as cognition status in frail and often institutionalized populations (Cahn-Weiner et al., 2007; Fujiwara et al., 2008; Paterson & Warburton, 2010; Vasquez, Batsis, Germain, & Shaw, 2014). These and many other studies that focus on disability and independence and that examine the impact of factors related to aging and physical independence have provided cross-sectional results, largely providing measures of central tendency and normative

\* Corresponding author at: Departamento de Desporto e Saúde, Escola de Ciências e Tecnologia, Universidade de Évora, Pavilhão Gimnodesportivo, Rua Reguengos de Monsaraz, 44, 7000-727 Évora, Portugal.

E-mail addresses: [clnp@uevora.pt](mailto:clnp@uevora.pt) (C. Pereira), [fbaptista@fmh.ulisboa.pt](mailto:fbaptista@fmh.ulisboa.pt) (F. Baptista), [anacf@uevora.pt](mailto:anacf@uevora.pt) (A. Cruz-Ferreira).