

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

PENALIZED LEAST SQUARES FITTING

Manfred von Golitschek and Larry L. Schumaker

Communicated by P. P. Petrushev

**Dedicated to the memory of our colleague Vasil Popov
January 14, 1942 – May 31, 1990**

ABSTRACT. Bounds on the error of certain penalized least squares data fitting methods are derived. In addition to general results in a fairly abstract setting, more detailed results are included for several particularly interesting special cases, including splines in both one and several variables.

1. Introduction. We begin with an abstract definition of what we mean by a penalized least squares fit. Suppose X , Y and S are linear spaces over \mathbb{R} , where $S \subseteq Y \subseteq X$. Let $\|\cdot\|_X : X \rightarrow \mathbb{R}$ and $\|\cdot\|_Y : Y \rightarrow \mathbb{R}$ be semi-norms induced by semi-definite inner products $\langle \cdot, \cdot \rangle$ on X and $[\cdot, \cdot]$ on Y , respectively. Given $f \in X$ and $\lambda > 0$, suppose there exists $s_\lambda(f, S)$ in S such that

$$\Phi(s_\lambda(f, S)) = \min_{u \in S} \Phi(u),$$

2000 *Mathematics Subject Classification*: Primary 41A15; Secondary 41A25.

Key words: Penalized least squares, Tikhonov regularization, Spline approximation.

*Supported by the Army Research Office under grant DAAD-19-02-10059.

where

$$\Phi(u) := \|f - u\|_X^2 + \lambda \|u\|_Y^2.$$

Then we call $s_\lambda(f, S)$ a penalized least squares fit of f corresponding to λ . If there exists a unique minimizer of Φ in S for each $f \in X$, then $Q_\lambda : X \rightarrow S$ defined by $Q_\lambda f := s_\lambda(f, S)$ defines a linear operator which is not in general a linear projection. Our aim in this paper is to investigate the behavior of $\|f - Q_\lambda f\|_X$ as a function of both λ and the approximation properties of S .

The paper is organized as follows. In Section 2 we give a detailed treatment of the penalized least squares problem for approximation by trigonometric polynomials. The next section is devoted to Tikhonov regularization (with weights) for univariate functions defined on an interval. The results include error bounds for certain classes of functions. In Section 4 we derive an explicit formula for the Tikhonov regularization of arbitrary functions in $C[-\pi, \pi]$. Our main L_2 error bounds for penalized least squares fits of univariate functions are contained in Section 5. General penalized least squares is treated in Section 6. We conclude the paper by outlining two typical applications: univariate splines and bivariate splines on triangulations.

2. Penalized Fourier series. Throughout this section, we take $X = L_2[-\pi, \pi]$ with the inner product $\langle f, g \rangle = \int_{-\pi}^{\pi} fg$, and set $Y = W_2^2[\mathbb{T}]$ with the semi-definite inner product $[f, g] = \int_{-\pi}^{\pi} f''g''$, where \mathbb{T} denotes the circle. We let $S = \mathcal{T}_n$ be the set of all real-valued trigonometric polynomials of degree at most n , with $n \in \mathbb{N}$ fixed. Clearly, for each $f \in L_2[-\pi, \pi]$ and $\lambda > 0$, there exists a unique minimal solution $s_{\lambda, n} := s_{\lambda, n}(f)$ of the penalized least squares problem

$$\min_{T \in \mathcal{T}_n} \left\{ \int_{-\pi}^{\pi} (f - T)^2 dx + \lambda \int_{-\pi}^{\pi} (T'')^2 dx \right\}.$$

Our first theorem compares $s_{\lambda, n}$ with the n -th partial sum $s_n(f) := s_{0, n}(f)$ of the Fourier series of f .

Theorem 2.1. *Suppose $f \in L_2[-\pi, \pi]$, and that*

$$f(x) = a_0 + \sum_{k=1}^{\infty} (a_k \cos(kx) + b_k \sin(kx))$$

is its Fourier series. Then for each $\lambda > 0$ and $n \in \mathbb{N}$, $s_{\lambda,n} = s_\lambda(f, \mathcal{T}_n)$ is given by

$$(2.1) \quad s_{\lambda,n}(x) = a_0 + \sum_{k=1}^n \frac{1}{1 + \lambda k^4} (a_k \cos(kx) + b_k \sin(kx)),$$

or equivalently,

$$(2.2) \quad s_n(f) = s_{\lambda,n} + \lambda s_{\lambda,n}^{(4)}.$$

Proof. The minimal solution $s_{\lambda,n}$ is characterized by the orthogonality relations

$$(2.3) \quad \int_{-\pi}^{\pi} (f - s_{\lambda,n})T = \lambda \int_{-\pi}^{\pi} s_{\lambda,n}'' T'', \quad \text{for all } T \in \mathcal{T}_n.$$

A simple computation shows that the function $s_{\lambda,n}$ given by (2.1) is the only function in \mathcal{T}_n which satisfies (2.3). This implies that $s_{\lambda,n}$ is the unique minimal solution. The proof of (2.2) is equally simple. \square

Taking the limit as $n \rightarrow \infty$ in Theorem 2.1, we get

Corollary 2.2. *Let f be as in Theorem 2.1. Then for each $\lambda > 0$, the function*

$$s_{\lambda,\infty}(x) = a_0 + \sum_{k=1}^{\infty} \frac{1}{1 + \lambda k^4} (a_k \cos(kx) + b_k \sin(kx)), \quad x \in \mathbb{R},$$

lies in $W_2^4(\mathbb{T})$, and

$$f = s_{\lambda,\infty} + \lambda s_{\lambda,\infty}^{(4)}.$$

Moreover, $s_{\lambda,\infty}$ is the unique solution of the minimum problem

$$\min \left\{ \int_{-\pi}^{\pi} (f - g)^2 dx + \lambda \int_{-\pi}^{\pi} (g'')^2 dx, \quad g \in W_2^2(\mathbb{T}) \right\}.$$

Let $\|\cdot\|_2$ and $\|\cdot\|_\infty$ denote the L_2 and uniform norms on $[-\pi, \pi]$, respectively. Since $\|f - s_n(f)\|_2 \leq \|f - T\|_2$ for all $T \in \mathcal{T}_n$, we have $\|f - s_{\lambda,n}\|_2 \geq \|f - s_n(f)\|_2$. In the next theorem we will see that the L_2 -norm $\|f - s_{\lambda,n}(f)\|_2$ changes very little for increasing $n > \lambda^{-1/4}$. Hence, for numerical purposes, it is not worthwhile to compute $s_{\lambda,n}(f)$ for n larger than $\lambda^{-1/4}$.

Theorem 2.3. *Let $f \in L_2[-\pi, \pi]$ and $0 < \lambda \leq 1$. Let $N_\lambda = \lfloor \lambda^{-1/4} \rfloor$ be the largest integer smaller than or equal to $\lambda^{-1/4}$. Then*

$$(2.4) \quad \frac{1}{2}\sqrt{\varepsilon_{\lambda,n}} \leq \|f - s_{\lambda,n}(f)\|_2 \leq \sqrt{\varepsilon_{\lambda,n}},$$

where $\varepsilon_{\lambda,n} := \varepsilon_{\lambda,n}(f)$ is defined by

$$\varepsilon_{\lambda,n} := \begin{cases} \|f - s_n(f)\|_2^2 + \lambda^2 \|s_n^{(4)}(f)\|_2^2, & \text{if } n \leq N_\lambda, \\ \varepsilon_{\lambda,N_\lambda}, & \text{if } n > N_\lambda. \end{cases}$$

Proof. We apply the orthogonality relation

$$\int_{-\pi}^{\pi} (f - s_n(f))T = 0, \quad T \in \mathcal{T}_n,$$

for $T = s_{\lambda,n}^{(4)} = s_{\lambda,n}^{(4)}(f)$. This gives $\int_{-\pi}^{\pi} (f - s_n(f))s_{\lambda,n}^{(4)} = 0$, and (2.2) implies

$$(2.5) \quad \|f - s_{\lambda,n}\|_2^2 = \|f - s_n(f) + \lambda s_{\lambda,n}^{(4)}\|_2^2 = \|f - s_n(f)\|_2^2 + \lambda^2 \|s_{\lambda,n}^{(4)}\|_2^2.$$

It follows from (2.1) that

$$\|s_{\lambda,n}^{(4)}\|_2^2 = \left\| \sum_{k=1}^n \frac{k^4}{1 + \lambda k^4} (a_k \cos(kx) + b_k \sin(kx)) \right\|_2^2 = \pi \sum_{k=1}^n \frac{k^8(a_k^2 + b_k^2)}{(1 + \lambda k^4)^2}.$$

If $n \leq N_\lambda$, this implies

$$\begin{aligned} \|s_{\lambda,n}^{(4)}\|_2^2 &\leq \pi \sum_{k=1}^n k^8(a_k^2 + b_k^2) = \|s_n^{(4)}\|_2^2 \\ \|s_{\lambda,n}^{(4)}\|_2^2 &\geq \frac{\pi}{4} \sum_{k=1}^n k^8(a_k^2 + b_k^2) = \frac{1}{4} \|s_n^{(4)}\|_2^2. \end{aligned}$$

If $n > N_\lambda$, then

$$\begin{aligned} \|s_{\lambda,n}^{(4)}\|_2^2 &= \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \pi \sum_{k=N_\lambda+1}^n \frac{k^8(a_k^2 + b_k^2)}{(1 + \lambda k^4)^2} \\ &\leq \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \pi \sum_{k=N_\lambda+1}^n \frac{a_k^2 + b_k^2}{\lambda^2} \\ &= \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \lambda^{-2} \left(\|f - s_{N_\lambda}\|_2^2 - \|f - s_n\|_2^2 \right). \end{aligned}$$

Moreover, if $n > N_\lambda$, then

$$\begin{aligned} \|s_{\lambda,n}^{(4)}\|_2^2 &= \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \pi \sum_{k=N_\lambda+1}^n \frac{k^8(a_k^2 + b_k^2)}{(1 + \lambda k^4)^2} \\ &\geq \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \pi \sum_{k=N_\lambda+1}^n \frac{a_k^2 + b_k^2}{4\lambda^2} \\ &= \|s_{\lambda,N_\lambda}^{(4)}\|_2^2 + \frac{1}{4\lambda^2} (\|f - s_{N_\lambda}\|_2^2 - \|f - s_n\|_2^2). \end{aligned}$$

Inserting these estimates into (2.5) yields (2.4). \square

3. Tikhonov regularization on $[a, b]$ with weights.

Suppose $w : [a, b] \rightarrow \mathbb{R}$ is a piecewise continuous nonnegative function on $[a, b]$ with $\int_a^b w > 0$.

In this section we take $X = L_2(a, b)$ with the inner product $\langle f, g \rangle = \int_a^b wfg$, and set $Y = S = W_2^2[a, b]$ with the semi-definite inner product $[f, g] = \int_a^b f''g''$.

Definition 3.1. Let $f \in L_2(a, b)$ and $\lambda > 0$. We call $\phi_{\lambda,w} = \phi_\lambda(f, w) \in W_2^2[a, b]$ the (nonperiodic) Tikhonov regularization of f corresponding to λ and w provided it minimizes

$$(3.1) \quad \int_a^b w(x)(f - u)^2 dx + \lambda \int_a^b (u'')^2 dx$$

with respect to all $u \in W_2^2[a, b]$.

For the weight $w = 1$, Ragozin [5] proved that the Tikhonov regularization ϕ_λ satisfies

$$\|f^{(q)} - \phi_\lambda^{(q)}\|_2 \leq C_q \lambda^{(2-q)/4} \|f''\|_2, \quad q = 0, 1, 2,$$

for some absolute constants $C_q > 0$. The next theorem states, among other things, that $C_0 \leq 1$ and $C_2 \leq 2$ for the weight $w = 1$.

Theorem 3.2. Let $f \in C[a, b]$ and $\lambda > 0$. Let $\phi_\lambda := \phi_\lambda(f, w)$ be the Tikhonov regularization of f corresponding to λ, w . Then $\phi_\lambda \in C^3[a, b]$, $\phi_\lambda^{(4)}$ is piecewise continuous on $[a, b]$, and

1) $w(x)[f(x) - \phi_\lambda(x)] = \lambda\phi_\lambda^{(4)}(x)$, for all $x \in [a, b]$ where w is continuous,

2) $(\phi_\lambda)''(a) = (\phi_\lambda)''(b) = \phi_\lambda^{(3)}(a) = \phi_\lambda^{(3)}(b) = 0$.

If $f \in W_2^2[a, b]$, then $\|(\phi_\lambda)''\|_2 \leq \|f''\|_2$ and

$$\left(\int_a^b w(f - \phi_\lambda)^2 \right)^{1/2} \leq \lambda^{1/2} \|f''\|_2.$$

Proof. Let $f \in C[a, b]$. The orthogonality relations for (3.1) are

$$(3.2) \quad \int_a^b w(f - \phi_\lambda)u - \lambda \int_a^b (\phi_\lambda)''u'' = 0, \quad \text{for all } u \in W_2^2[a, b].$$

We define the function F by

$$F(x) := \int_a^x \int_a^t w(s) (f(s) - \phi_\lambda(s)) ds dt, \quad a \leq x \leq b.$$

Then, $F(a) = F'(a) = 0$. Applying (3.2) for $u = 1$ and $u(x) = x$ yields

$$\int_a^b w(f - \phi_\lambda) dx = \int_a^b (f(x) - \phi_\lambda(x)) x dx = 0,$$

and thus $F'(b) = 0$ and $F(b) = 0$. Therefore, integrating by parts gives

$$\int_a^b w(f - \phi_\lambda)u = \int_a^b Fu''.$$

Hence (3.2) is equivalent to

$$\int_a^b (F - \lambda(\phi_\lambda)''u)u'' = 0, \quad \text{for all } u \in W_2^2[a, b],$$

and thus

$$\int_a^b (F - \lambda(\phi_\lambda)''g)g = 0, \quad \text{for all } g \in L_2[a, b].$$

This implies that

$$F(x) = \lambda(\phi_\lambda)''(x), \quad a \leq x \leq b.$$

Differentiating twice yields

$$w(x)(f(x) - \phi_\lambda(x)) = \lambda\phi_\lambda^{(4)}(x)$$

at all points $x \in [a, b]$ where w is continuous. Then $F = \lambda(\phi_\lambda)''$ and $F(a) = F(b) = 0$ imply $(\phi_\lambda)''(a) = (\phi_\lambda)''(b) = 0$, while $F' = \lambda\phi_\lambda^{(3)}$ and $F'(a) = F'(b) = 0$ imply $\phi_\lambda^{(3)}(a) = \phi_\lambda^{(3)}(b) = 0$.

Let $f \in W_2^2[a, b]$. We denote the minimal value in (3.1) by M . Then (3.1) and (3.2) imply that

$$M = \int_a^b w(f - \phi_\lambda)(f - u) + \lambda \int_a^b (\phi_\lambda)'' u'', \quad \text{for all } u \in W_2^2[a, b].$$

Taking $u = f$, it follows that

$$M = \lambda \int_a^b (\phi_\lambda)'' f'' \leq \lambda \|(\phi_\lambda)''\|_2 \|f''\|_2.$$

Therefore,

$$\int_a^b w(f - \phi_\lambda)^2 + \lambda \|(\phi_\lambda)''\|_2^2 \leq \lambda \|(\phi_\lambda)''\|_2 \|f''\|_2,$$

and thus

$$\|(\phi_\lambda)''\|_2 \leq \|f''\|_2, \quad \|\sqrt{w}(f - \phi_\lambda)\|_2 \leq \lambda^{1/2} \|f''\|_2.$$

This concludes the proof. \square

Theorem 3.3. *Let $w = 1$. Let $f \in W_2^4[a, b]$ and $f''(a) = f''(b) = f^{(3)}(a) = f^{(3)}(b) = 0$. Then the Tikhonov regularization ϕ_λ of f corresponding to $\lambda > 0$ satisfies*

$$\begin{aligned} \|f - \phi_\lambda\|_2 &\leq \lambda \|f^{(4)}\|_2, \\ \|f'' - (\phi_\lambda)''\|_2 &\leq \lambda^{1/2} \|f^{(4)}\|_2, \\ \|\phi_\lambda^{(4)}\|_2 &\leq \|f^{(4)}\|_2. \end{aligned}$$

Proof. We write $U := \{u \in W_2^4[a, b] : u''(a) = u''(b) = u^{(3)}(a) = u^{(3)}(b) = 0\}$. Clearly, $f \in U$ by assumption and $\phi_\lambda \in U$ by Theorem 3.2. For each $u \in U$ we obtain by integration by parts that

$$\int_a^b (f - \phi_\lambda)u^{(4)} = \int_a^b (f'' - (\phi_\lambda)'')u''.$$

Inserting the identity $f - \phi_\lambda = \lambda\phi_\lambda^{(4)}$, it follows that

$$(3.3) \quad \int_a^b (f'' - (\phi_\lambda)'')u'' - \lambda \int_a^b \phi_\lambda^{(4)}u^{(4)} = 0, \quad \text{for all } u \in U.$$

But (3.3) are the orthogonality relations for the minimum problem

$$(3.4) \quad M^* := \min_{u \in U} \|f'' - u''\|_2^2 + \lambda \|u^{(4)}\|_2^2,$$

and so ϕ_λ is the minimal solution of (3.4). Since $f \in U$, it follows that $M^* \leq \lambda \|f^{(4)}\|_2^2$, and thus

$$\|\phi_\lambda^{(4)}\|_2 \leq \|f^{(4)}\|_2, \quad \|f'' - (\phi_\lambda)''\|_2 \leq \lambda^{1/2} \|f^{(4)}\|_2.$$

Finally, we have

$$\|f - \phi_\lambda\|_2 = \lambda \|\phi_\lambda^{(4)}\|_2 \leq \lambda \|f^{(4)}\|_2,$$

which concludes the proof of the theorem. \square

4. An explicit formula for the Tikhonov regularization In this section we consider the inner product spaces $X = L_2(a, b)$ and $Y = S = W_2^2[a, b]$ of Section 3 with the weight function $w = 1$ on $[a, b]$, that is, $\langle f, g \rangle = \int_a^b fg$, $[f, g] = \int_a^b f''g''$. In Theorem 2.1 we have given an explicit formula for the periodic Tikhonov regularization. In this section we want to apply the methods of Section 2 to the nonperiodic case. We may assume for convenience that the interval is $[a, b] = [-\pi, \pi]$. Otherwise, under the linear transformation

$$x(t) = a + \frac{(b-a)(t+\pi)}{2\pi}, \quad -\pi \leq t \leq \pi,$$

of the interval $[-\pi, \pi]$ onto $[a, b]$, the Tikhonov regularizations $\phi_\lambda(x)$ on $[a, b]$ of f and $\psi_\mu(t)$ on $[-\pi, \pi]$ of $g(t) := f(x(t))$ are related by

$$(4.1) \quad \psi_\mu(t) = \phi_\lambda(x(t)), \quad \lambda = \left(\frac{b-a}{2\pi}\right)^4 \mu.$$

To prove (4.1), we simply compare the orthogonality relations

$$(4.2) \quad \int_a^b (f(x) - \phi_\lambda(x))u(x) dx = \lambda \int_a^b (\phi_\lambda)''(x)u''(x) dx, \quad u \in W_2^2[a, b],$$

and

$$(4.3) \quad \int_{-\pi}^{\pi} (g(t) - \psi_{\mu}(t))v(t) dt = \mu \int_{-\pi}^{\pi} \psi_{\mu}''(t)v''(t) dt, \quad v \in W_2^2[-\pi, \pi],$$

where the equivalence of (4.2) and (4.3) follows from the bijection $v(t) = u(x(t))$ of $W_2^2[a, b]$ onto $W_2^2[-\pi, \pi]$.

The main idea of this section is contained in the next lemma.

Lemma 4.1 *Each function $v \in C^1[-\pi, \pi]$ has a unique representation of the form*

$$(4.4) \quad v(t) = c_1 t + c_2 t^2 + w(t), \quad c_1 \in \mathbb{R}, \quad c_2 \in \mathbb{R}, \quad w \in C^1(\mathbb{T}).$$

In other words, (4.4) defines a bijection from $\mathbb{R}^2 \times C^1(\mathbb{T})$ onto $C^1[-\pi, \pi]$.

Proof. Given $c_1, c_2 \in \mathbb{R}$ and $w \in C^1(\mathbb{T})$, the function v in (4.4) lies in $C^1[-\pi, \pi]$. Conversely, for a function $v \in C^1[-\pi, \pi]$, the condition $w(-\pi) = w(\pi)$ is equivalent to

$$v(-\pi) + c_1\pi - c_2\pi^2 = v(\pi) - c_1\pi - c_2\pi^2,$$

and thus to

$$c_1 = \frac{v(\pi) - v(-\pi)}{2\pi}.$$

The condition $w'(-\pi) = w'(\pi)$ is equivalent to $v'(-\pi) + 2c_2\pi = v'(\pi) - 2c_2\pi$, and thus to

$$c_2 = \frac{v'(\pi) - v'(-\pi)}{4\pi}.$$

This concludes the proof of the lemma. \square

As a corollary, each function $f \in C^1[-\pi, \pi]$ has a unique representation

$$(4.5) \quad f(t) = c_1 t + c_2 t^2 + a_0 + \sum_{k=1}^{\infty} (a_k \cos(kt) + b_k \sin(kt)),$$

where

$$c_1 = \frac{f(\pi) - f(-\pi)}{2\pi}, \quad c_2 = \frac{f'(\pi) - f'(-\pi)}{4\pi}.$$

Theorem 4.2. *Let $f \in C[-\pi, \pi]$ be given by (4.5). If $f \notin C^1[-\pi, \pi]$, then we take $c_2 = 0$. Then the nonperiodic Tikhonov regularization ϕ_λ of f corresponding to λ is given by*

$$\phi_\lambda(t) = \gamma_1 t + \gamma_2 t^2 + \alpha_0 + \sum_{k=1}^{\infty} (\alpha_k \cos(kt) + \beta_k \sin(kt))$$

with

$$(4.6) \quad \gamma_1 = c_1 - \frac{1}{2B(\lambda)} \sum_{k=1}^{\infty} \frac{(-1)^k k^3 b_k}{1 + \lambda k^4},$$

$$(4.7) \quad \gamma_2 = \frac{1}{2 + 4A(\lambda)} \left(4A(\lambda)c_2 + \sum_{k=1}^{\infty} \frac{(-1)^k k^2 a_k}{1 + \lambda k^4} \right),$$

$$(4.8) \quad \alpha_0 = a_0 + \frac{(c_2 - \gamma_2)\pi^2}{3},$$

$$(4.9) \quad \alpha_k = \frac{1}{1 + \lambda k^4} \left(a_k + \frac{4(-1)^k (c_2 - \gamma_2)}{k^2} \right), \quad k \geq 1,$$

$$(4.10) \quad \beta_k = \frac{1}{1 + \lambda k^4} \left(b_k + \frac{2(-1)^{k-1} (c_1 - \gamma_1)}{k} \right), \quad k \geq 1,$$

where

$$A(\lambda) := \sum_{k=1}^{\infty} \frac{1}{1 + \lambda k^4}, \quad B(\lambda) := \sum_{k=1}^{\infty} \frac{k^2}{1 + \lambda k^4}.$$

Proof. The Fourier series of t and t^2 on $(-\pi, \pi)$ are

$$t = \sum_{k=1}^{\infty} \frac{2(-1)^{k-1} \sin(kt)}{k}, \quad -\pi < t < \pi,$$

$$t^2 = \frac{\pi^2}{3} + \sum_{k=1}^{\infty} \frac{4(-1)^k \cos(kt)}{k^2}, \quad -\pi \leq t \leq \pi,$$

with pointwise convergence. From Theorem 3.2, we know that $f = \phi_\lambda + \lambda \phi_\lambda^{(4)}$. Comparing the Fourier coefficients of f and $\phi_\lambda + \lambda \phi_\lambda^{(4)}$ in the expansions

$$f(t) = c_1 \sum_{k=1}^{\infty} \frac{2(-1)^{k-1} \sin(kt)}{k} + c_2 \left(\frac{\pi^2}{3} + \sum_{k=1}^{\infty} \frac{4(-1)^k \cos(kt)}{k^2} \right) \\ + a_0 + \sum_{k=1}^{\infty} (a_k \cos(kt) + b_k \sin(kt))$$

and

$$\begin{aligned} \phi_\lambda(t) + \lambda\phi_\lambda^{(4)}(t) &= \gamma_1 \sum_{k=1}^{\infty} \frac{2(-1)^{k-1} \sin(kt)}{k} + \gamma_2 \left(\frac{\pi^2}{3} + \sum_{k=1}^{\infty} \frac{4(-1)^k \cos(kt)}{k^2} \right) \\ &\quad + \alpha_0 + \sum_{k=1}^{\infty} (1 + \lambda k^4)(\alpha_k \cos(kt) + \beta_k \sin(kt)), \end{aligned}$$

we obtain the equalities

$$\begin{aligned} \frac{c_2\pi^2}{3} + a_0 &= \frac{\gamma_2\pi^2}{3} + \alpha_0 \\ \frac{4c_2(-1)^k}{k^2} + a_k &= \frac{4\gamma_2(-1)^k}{k^2} + (1 + \lambda k^4)\alpha_k, \quad k \geq 1, \\ \frac{2c_1(-1)^{k-1}}{k} + b_k &= \frac{2\gamma_1(-1)^{k-1}}{k} + (1 + \lambda k^4)\beta_k, \quad k \geq 1, \end{aligned}$$

which are equivalent to (4.8)–(4.10). From Theorem 3.2 we know that $(\phi_\lambda)''(\pi) = 0$ and $\phi_\lambda^{(3)}(\pi) = 0$, and thus,

$$\begin{aligned} (\phi_\lambda)''(\pi) &= 2\gamma_2 - \sum_{k=1}^{\infty} (-1)^k k^2 \alpha_k = 0, \\ \phi_\lambda^{(3)}(\pi) &= - \sum_{k=1}^{\infty} (-1)^k k^3 \beta_k = 0. \end{aligned} \tag{4.11}$$

Inserting (4.8)–(4.10) into (4.11) yields

$$2\gamma_2 = \sum_{k=1}^{\infty} \frac{(-1)^k k^2 a_k}{1 + \lambda k^4} + 4(c_2 - \gamma_2) \sum_{k=1}^{\infty} \frac{1}{1 + \lambda k^4}$$

and

$$0 = \sum_{k=1}^{\infty} (-1)^k k^3 \beta_k = \sum_{k=1}^{\infty} \frac{(-1)^k k^3 b_k}{1 + \lambda k^4} - 2(c_1 - \gamma_1) \sum_{k=1}^{\infty} \frac{k^2}{1 + \lambda k^4}.$$

These two conditions are equivalent to

$$2\gamma_2 = \sum_{k=1}^{\infty} \frac{(-1)^k k^2 a_k}{1 + \lambda k^4} + 4(c_2 - \gamma_2)A(\lambda) \tag{4.12}$$

and

$$2(c_1 - \gamma_1)B(\lambda) = \sum_{k=1}^{\infty} \frac{(-1)^k k^3 b_k}{1 + \lambda k^4}. \tag{4.13}$$

Now (4.6) follows from (4.12), and (4.7) follows from (4.13). \square

5. L_2 -error bounds for penalized least squares fitting. In this section we consider the inner product spaces $X = L_2(a, b)$ and $Y = W_2^2[a, b]$ of Section 4 with the same inner products $\langle f, g \rangle = \int_a^b fg$ and $[f, g] = \int_a^b f''g''$, but we now take S to be a proper subspace of Y . For $f \in L_2[a, b]$ and $\lambda > 0$, we compare the penalized least squares fit $\phi_\lambda \in Y$ of Section 3 (with weight function $w = 1$) satisfying

$$\int_a^b (f - \phi_\lambda)^2 + \lambda \int_a^b (\phi_\lambda'')^2 = \min_{u \in Y} \left\{ \int_a^b (f - u)^2 + \lambda \int_a^b (u'')^2 \right\}$$

with the penalized least squares fit $s_\lambda = Q_\lambda(f)$ from S satisfying

$$\int_a^b (f - s_\lambda)^2 + \lambda \int_a^b (s_\lambda'')^2 = \min_{u \in S} \left\{ \int_a^b (f - u)^2 + \lambda \int_a^b (u'')^2 \right\}.$$

Theorem 5.1. *Let $f \in W_2^2[a, b]$ and $\lambda > 0$. Then s_λ minimizes the expression*

$$(5.1) \quad \|\phi_\lambda - u\|_2^2 + \lambda \|(\phi_\lambda)'' - u''\|_2^2$$

with respect to all $u \in S$.

Proof. s_λ is characterized by the orthogonality relations

$$(5.2) \quad \int_a^b (f - s_\lambda)u - \lambda \int_a^b (s_\lambda)''u'' = 0, \quad \text{for all } u \in S.$$

Since $f = \phi_\lambda + \lambda\phi_\lambda^{(4)}$ by Theorem 3.2, (5.2) is equivalent to

$$\int_a^b (\phi_\lambda - s_\lambda)u + \lambda \int_a^b (\phi_\lambda^{(4)}u - (s_\lambda)''u'') = 0, \quad \text{for all } u \in S,$$

and since $\phi_\lambda''(a) = \phi_\lambda''(b) = \phi_\lambda^{(3)}(a) = \phi_\lambda^{(3)}(b) = 0$ by Theorem 3.2, integration by parts shows that (5.2) is equivalent to

$$(5.3) \quad \int_a^b (\phi_\lambda - s_\lambda)u + \lambda \int_a^b (\phi_\lambda'' - (s_\lambda)'')u'' = 0, \quad \text{for all } u \in S.$$

But (5.3) are the orthogonality relations for the minimum problem (5.1). \square

Corollary 5.2. *Let $\mathcal{S} \subset W_2^2[a, b]$ be a linear space. Suppose that for each $g \in W_2^2[a, b]$ there exists $u \in \mathcal{S}$ such that*

$$(5.4) \quad \begin{aligned} \|g - u\|_2 &\leq Ch^2\|g''\|_2, \\ \|g'' - u''\|_2 &\leq C\|g''\|_2, \end{aligned}$$

for positive numbers C and h independent of g . Then the penalized least squares fit s_λ of f in \mathcal{S} corresponding to λ satisfies

$$(5.5) \quad \|f - s_\lambda\|_2 \leq \lambda^{1/2} \left(1 + C\sqrt{h^4\lambda^{-1} + 1}\right) \|f''\|_2$$

and

$$(5.6) \quad \|(s_\lambda)''\|_2 \leq \left(1 + C\sqrt{h^4\lambda^{-1} + 1}\right) \|f''\|_2.$$

Proof. By Theorem 3.2, the Tikhonov regularization ϕ_λ of f satisfies

$$\|f - \phi_\lambda\|_2 \leq \lambda^{1/2} \|f''\|_2.$$

Let M_0 be the minimum in (5.1). Applying (5.4) for $g := \phi_\lambda$, it follows that

$$M_0 \leq C^2 \left(h^4 + \lambda\right) \|(\phi_\lambda)''\|_2^2.$$

Then, since $\|(\phi_\lambda)''\|_2 \leq \|f''\|_2$ by Theorem 3.2,

$$\begin{aligned} \|f - s_\lambda\|_2 &\leq \|f - \phi_\lambda\|_2 + \|\phi_\lambda - s_\lambda\|_2 \\ &\leq \lambda^{1/2} \|f''\|_2 + M_0^{1/2} \\ &\leq \left(\lambda^{1/2} + C\sqrt{h^4 + \lambda}\right) \|f''\|_2, \end{aligned}$$

which yields (5.5). Since $\|(\phi_\lambda)''\|_2 \leq \|f''\|_2$ by Theorem 3.2, while

$$(5.7) \quad \lambda \|(\phi_\lambda)'' - (s_\lambda)''\|_2^2 \leq M_0,$$

it follows that

$$\begin{aligned} \|(s_\lambda)''\|_2 &\leq \|(\phi_\lambda)''\|_2 + \lambda^{-1/2} M_0^{1/2} \\ &\leq \left(1 + C \left[h^4\lambda^{-1} + 1\right]^{1/2}\right) \|f''\|_2, \end{aligned}$$

which proves (5.6). \square

Corollary 5.3. *Let $\mathcal{S} \subset W_2^2[a, b]$ be a linear space. Suppose that for each $g \in W_2^4[a, b]$ there exists $u \in \mathcal{S}$ such that*

$$(5.8) \quad \begin{aligned} \|g - u\|_2 &\leq Ch^4 \|g^{(4)}\|_2, \\ \|g'' - u''\|_2 &\leq Ch^2 \|g^{(4)}\|_2, \end{aligned}$$

for positive numbers C and h independent of g . Let $f \in W_2^4[a, b]$ and $f''(a) = f''(b) = f^{(3)}(a) = f^{(3)}(b) = 0$. Then the penalized least squares fit s_λ of f in \mathcal{S} corresponding to λ satisfies

$$(5.9) \quad \|f - s_\lambda\|_2 \leq \left(\lambda + C[\lambda + h^4] \right) \|f^{(4)}\|_2,$$

$$(5.10) \quad \|f'' - (s_\lambda)''\|_2 \leq \left(\lambda^{1/2} + Ch^2[h^4\lambda^{-1} + 1]^{1/2} \right) \|f^{(4)}\|_2,$$

Proof. By Theorem 3.3, the Tikhonov regularization ϕ_λ of f satisfies

$$\|f - \phi_\lambda\|_2 \leq \lambda \|f^{(4)}\|_2.$$

Let M_0 be the minimum in (5.1). Applying (5.8) for $g := \phi_\lambda$, it follows that

$$M_0 \leq C^2 \left(h^8 + \lambda h^4 \right) \|\phi_\lambda^{(4)}\|_2^2.$$

Then, since $\|\phi_\lambda^{(4)}\|_2 \leq \|f^{(4)}\|_2$ by Theorem 3.3,

$$\begin{aligned} \|f - s_\lambda\|_2 &\leq \|f - \phi_\lambda\|_2 + \|\phi_\lambda - s_\lambda\|_2 \\ &\leq \lambda \|f^{(4)}\|_2 + M_0^{1/2} \\ &\leq \left(\lambda + Ch^2 \sqrt{h^4 + \lambda} \right) \|f^{(4)}\|_2, \end{aligned}$$

which implies (5.9).

Since $\|f'' - (\phi_\lambda)''\|_2 \leq \lambda^{1/2} \|f^{(4)}\|_2$ by Theorem 3.3, using (5.7) gives

$$\|f'' - (s_\lambda)''\|_2 \leq \lambda^{1/2} \|f^{(4)}\|_2 + \lambda^{-1/2} M_0^{1/2},$$

and (5.10) follows. \square

6. A general penalized least squares problem. Keeping in mind the problems and results of Sections 2–5, we now investigate the more general

setting described in the introduction. Let X, Y and S be linear spaces on \mathbb{R} , where $S \subseteq Y \subseteq X$; $\|\cdot\|_X : X \rightarrow \mathbb{R}$ and $\|\cdot\|_Y : Y \rightarrow \mathbb{R}$ are semi-norms induced by semi-definite inner products $\langle \cdot, \cdot \rangle$ on X and $[\cdot, \cdot]$ on Y , respectively. For simplicity, we assume throughout the rest of this paper that S has dimension n and that

the restriction $\|\cdot\|_X$ onto S is a norm on S .

This implies that for any $f \in X$ and $\lambda > 0$, there exists a unique $s_\lambda := s_\lambda(f, S) := Q_\lambda f$ in S with the property

$$\|f - s_\lambda\|_X^2 + \lambda \|s_\lambda\|_Y^2 = \inf_{u \in S} \left\{ \|f - u\|_X^2 + \lambda \|u\|_Y^2 \right\}.$$

We denote the nonpenalized least squares fit in S by s_0 , i.e.,

$$\|f - s_0\|_X^2 = \min_{u \in S} \|f - u\|_X^2.$$

The penalized least squares approximation s_λ of f is characterized by the orthogonality relations

$$(6.1) \quad \langle f - s_\lambda, u \rangle = \lambda [s_\lambda, u], \quad \text{for all } u \in S,$$

while s_0 is characterized by

$$(6.2) \quad \langle f - s_0, u \rangle = 0, \quad \text{for all } u \in S.$$

By (6.1)–(6.2), we have

$$(6.3) \quad \langle s_0 - s_\lambda, u \rangle = \lambda [s_\lambda, u], \quad \text{for all } u \in S,$$

which proves that $s_\lambda = Q_\lambda s_0$, that is

$$\|s_0 - s_\lambda\|_X^2 + \lambda \|s_\lambda\|_Y^2 = \min_{u \in S} \left\{ \|s_0 - u\|_X^2 + \lambda \|u\|_Y^2 \right\}.$$

In what follows, we will discuss the error function $s_0 - s_\lambda$ which indicates the consequences of the penalty, and is easier to analyse than the error function $f - s_\lambda$ itself. This suffices since

$$\|f - s_0\|_X \leq \|f - s_\lambda\|_X \leq \|f - s_0\|_X + \|s_0 - s_\lambda\|_X.$$

In analogy with Theorem 3.2, we have

Theorem 6.1. For each $f \in X$ and $\lambda > 0$, $s_\lambda := s_\lambda(f, S)$ satisfies

$$(6.4) \quad \|s_\lambda\|_Y \leq \|s_0\|_Y$$

and

$$(6.5) \quad \|s_0 - s_\lambda\|_X \leq \sqrt{\lambda} \|s_0\|_Y.$$

Proof. Because of (6.3),

$$\begin{aligned} M(s_0, \lambda) &:= \|s_0 - s_\lambda\|_X^2 + \lambda \|s_\lambda\|_Y^2 \\ &= \langle s_0 - s_\lambda, s_0 - u \rangle + \lambda [s_\lambda, u], \quad \text{for all } u \in S. \end{aligned}$$

Inserting $u = s_0$ yields

$$M(s_0, \lambda) = \lambda [s_\lambda, s_0] \leq \lambda \|s_\lambda\|_Y \|s_0\|_Y,$$

and inequalities (6.4) and (6.5) follow. \square

Our next result provides an improvement of (6.5) for small λ which implies that

$$\|s_0 - s_\lambda\|_X = \mathcal{O}(\lambda) \quad \text{as } \lambda \rightarrow 0.$$

Theorem 6.2. For each $f \in X$ and $\lambda > 0$, $s_\lambda := s_\lambda(f, S)$ satisfies

$$(6.6) \quad \|s_0 - s_\lambda\|_X \leq K_S \lambda \|s_0\|_Y.$$

where

$$K_S := \sup \left\{ \frac{\|u\|_Y}{\|u\|_X} : u \in S, u \neq 0 \right\}.$$

Proof. Coupling the orthogonality relation (6.3) for $u = s_0 - s_\lambda$ with the Cauchy-Schwarz inequality and (6.4), we have

$$\begin{aligned} \|s_0 - s_\lambda\|_X^2 &= \langle s_0 - s_\lambda, s_0 - s_\lambda \rangle = \lambda [s_\lambda, s_0 - s_\lambda] \\ &\leq \lambda \|s_\lambda\|_Y \|s_0 - s_\lambda\|_Y \leq \lambda \|s_0\|_Y K_S \|s_0 - s_\lambda\|_X, \end{aligned}$$

which implies (6.6). \square

We conclude this section by examining the special case where X, Y, S are function spaces on some set $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$. Our next result gives a bound on $\|s_0 - s_\lambda\|_{L_\infty(\Omega)}$ which implies that

$$\|s_0 - s_\lambda\|_{L_\infty(\Omega)} = \mathcal{O}(\lambda) \quad \text{as } \lambda \rightarrow 0.$$

Theorem 6.3. *Suppose $X \subseteq L_\infty(\Omega)$, and let*

$$\kappa_S := \sup \left\{ \frac{\|u\|_{L_\infty(\Omega)}}{\|u\|_X} : u \in S, u \neq 0 \right\} < \infty.$$

Then

$$\|s_0 - s_\lambda\|_{L_\infty(\Omega)} \leq \kappa_S \sqrt{\lambda} \|s_0\|_Y \min \left\{ 1, K_S \sqrt{\lambda} \right\}.$$

Proof. The assertion follows directly from (6.5), Theorem 6.2, and the definition of κ_S . \square

7. Applications.

7.1. Cubic splines on an interval. Let $X = C[a, b]$, $Y = W_2^2[a, b]$, and suppose $S := S_4(\Delta) \subset C^2[a, b]$ is the linear space of cubic splines with simple knots $\Delta : a = x_0 < x_1 < \dots < x_n = b$. Suppose that $D := \{t_i\}_{i=1}^m$ are distinct points in $[a, b]$ so that the restriction of $S_4(\Delta)$ onto D has full dimension $n + 3$. We can now apply the results of Section 6 for the inner products

$$\langle f, g \rangle = \frac{1}{m} \sum_{i=1}^m f(t_i)g(t_i), \quad f, g \in C[a, b]$$

and

$$[f, g] = \int_a^b f'' g'', \quad f, g \in W_2^2[a, b].$$

In the notation of Sect. 6,

$$\|u\|_X = \left(\frac{1}{m} \sum_{i=1}^m u(t_i)^2 \right)^{1/2}, \quad u \in S.$$

We write $\|\cdot\|_2$ for the L_2 -norm on $[a, b]$ and suppose that D and Δ are such that

$$\delta := \sup \left\{ \frac{\|u\|_2}{\|u\|_X} : u \in S, u \neq 0 \right\}$$

is finite. Let $h_0 := h_0(\Delta) := \min\{x_{j+1} - x_j : j = 0, \dots, n - 1\}$. Since $\|u\|_2 \leq \delta \|u\|_X$, $u \in S$, it follows that for some absolute constant C_0 ,

$$\|u\|_Y = \|u''\|_2 \leq C_0 h_0^{-2} \|u\|_2 \leq C_0 \delta h_0^{-2} \|u\|_X,$$

and thus

$$K_S \leq C_0 \delta h_0^{-2}.$$

Moreover, for some other absolute constant C_1 ,

$$\kappa_S \leq C_1 \delta h_0^{-1/2}.$$

For given real numbers $\{y_i\}_{i=1}^m$, let $s_0 \in S$ be such that

$$\frac{1}{m} \sum_{i=1}^m (y_i - s_0(t_i))^2 = \min_{u \in S} \frac{1}{m} \sum_{i=1}^m (y_i - u(t_i))^2.$$

Recall that the penalized least squares spline s_λ minimizes the expression

$$\frac{1}{m} \sum_{i=1}^m (y_i - u(t_i))^2 + \lambda \int_a^b (u''(t))^2 dt.$$

The following result follows immediately from Theorems 6.1–6.3.

Theorem 7.1. *For all $\lambda > 0$,*

$$\|s_0 - s_\lambda\|_2 \leq \delta \sqrt{\lambda} \min\{1, K_S \sqrt{\lambda}\} \|s_0''\|_2.$$

Moreover,

$$\|s_0 - s_\lambda\|_{L_\infty[a,b]} \leq C_1 \delta h_0^{-1/2} \sqrt{\lambda} \min\{1, C_0 \delta h_0^{-2} \sqrt{\lambda}\} \|s_0''\|_2.$$

7.2. Bivariate spline spaces with stable local bases. Let Ω be a bounded set in \mathbb{R}^2 with polygonal boundary, and suppose that $X \subseteq L_\infty(\Omega)$, $Y = W_2^2(\Omega)$. Let $S \subset X$ be a linear space of polynomial splines defined on a regular triangulation Δ of Ω . For results on the approximation properties of these types of spaces, see [1, 2, 4].

Suppose that $D = \{t_i\}_{i=1}^m$ are distinct points in Ω so that the restriction of S onto D has the same dimension as S . We apply the results of Section 6 for the inner products

$$\langle f, g \rangle = \frac{1}{m} \sum_{i=1}^m f(t_i)g(t_i), \quad f, g \in X,$$

and

$$[f, g] := \int_{\Omega} (f_{xx}g_{xx} + 2f_{xy}g_{xy} + f_{yy}g_{yy}) \, dx \, dy, \quad f, g \in Y.$$

For given real numbers $(z_i)_{i=1}^m$, $s_0 \in S$ is the unique function in S which satisfies

$$\frac{1}{m} \sum_{i=1}^m (z_i - s_0(t_i))^2 = \min_{u \in S} \frac{1}{m} \sum_{i=1}^m (z_i - u(t_i))^2.$$

In the notation of Sect. 6,

$$\|u\|_X = \left(\frac{1}{m} \sum_{i=1}^m u(t_i)^2 \right)^{1/2}, \quad u \in S.$$

We have investigated properties of $s_0 \in S$ in [3].

Let $\|\cdot\|_2$ denote the L_2 -norm on Ω , and suppose that D and Δ are such that

$$\delta := \sup \left\{ \frac{\|u\|_2}{\|u\|_X} : u \in S, u \neq 0 \right\}$$

is finite. Let $h_0 := h_0(\Delta)$ be the minimum side length of the triangles in Δ . Since $\|u\|_2 \leq \delta \|u\|_X$, $u \in S$, it follows that for some absolute constant C_0

$$\|u\|_Y \leq C_0 h_0^{-2} \|u\|_2 \leq C_0 \delta h_0^{-2} \|u\|_X$$

so that

$$K_S \leq C_0 \delta h_0^{-2}.$$

Moreover, for some other absolute constant C_1 ,

$$\kappa_S \leq C_1 \delta h_0^{-1}.$$

As for univariate splines, we can now apply Theorems 6.1–6.3 to get bounds on $s_0 - s_\lambda$ in both the L_2 and uniform norms on Ω .

REFERENCES

- [1] C. K. CHUI, D. HONG, R.-Q. JIA. Stability of optimal order approximation by bivariate splines over arbitrary triangulations. *Trans. Amer. Math. Soc.* **347** (1995), 3301–3318.

- [2] O. DAVYDOV, L. L. SCHUMAKER. On stable local bases for bivariate polynomial spline spaces. *Constr. Approx.* **18** (2001), 87–116.
- [3] MANFRED V. GOLITSCHKEK, LARRY L. SCHUMAKER. Bounds on projections onto bivariate polynomial spline spaces with stable local basis. *Constr. Approx.* **18** (2002), 241–254.
- [4] M. J. LAI, L. L. SCHUMAKER. On the approximation power of bivariate splines. *Advances in Comp. Math.* **9** (1998), 251–279.
- [5] DAVID L. RAGOZIN. Error bounds for derivative estimates based on spline smoothing for exact or noisy data. *J. Approx. Theory* **37** (1983) 335–355.
- [6] JOHN RICE, MURRAY ROSENBLATT. Integrated mean squared error of a smoothing spline; *J. Approx. Theory* **33** (1981), 353–369.

Manfred von Golitschek

*Institut für Angewandte Mathematik und Statistik
der Universität Würzburg, 97074 Würzburg, Germany
e-mail: goli@mathematik.uni-wuerzburg.de*

Larry L. Schumaker

*Department of Mathematics
Vanderbilt University, Nashville
TN 37240, USA
e-mail: s@mars.cas.vanderbilt.edu*

Received September 19, 2002