

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

PLISKA
STUDIA MATHEMATICA
BULGARICA

ПЛИСКА
БЪЛГАРСКИ
МАТЕМАТИЧЕСКИ
СТУДИИ

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Pliska Studia Mathematica Bulgarica
visit the website of the journal <http://www.math.bas.bg/~pliska/>
or contact: Editorial Office

Pliska Studia Mathematica Bulgarica
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: pliska@math.bas.bg

CONCERNING TRIVIAL MAXIMAL ABELIAN SUBALGEBRAS OF $B(X)$

W. ŻELAZKO

To the memory of Y. A. Tagamlitzki

We call a complex or real Banach algebra trivial, if it is either a Banach space with trivial (zero) multiplication or it is the unitization of such an algebra. Thus a trivial algebra is always commutative and in the case of an algebra with unit element it is a local ring, i. e. it has exactly one maximal ideal equal to its radical. In this paper we prove that for any real or complex Banach space X the algebra $B(X)$ of all its continuous endomorphisms has always a trivial maximal Abelian subalgebra and we give description of all such subalgebras.

Let X be a real or complex Banach space. For a non-void subset S of $B(X)$ denote by S' its commutant, i. e. the set

$$S' = \{T \in B(X) : TA = AT \text{ for all } A \text{ in } S\}.$$

It is a closed subalgebra of $B(X)$ containing its unity I , and in case when S consists of mutually commuting operators we have

$$S' = \bigcup \{ \mathcal{A} : \mathcal{A} \text{ is a maximal Abelian subalgebra of } B(X) \text{ with } S \subset \mathcal{A} \}.$$

This implies that S' is a maximal Abelian subalgebra of $B(X)$, provided it is commutative. This simple remark will be used in the proof of our theorem.

In the sequel we denote by X^* the conjugate space of a Banach space X and by T^* the conjugate operator of an element T in $B(X)$. We put also $\text{rad } \mathcal{A}$ for the radical of a commutative Banach algebra \mathcal{A} . Thus in case of a trivial algebra \mathcal{A} with unit element we have $\mathcal{A} = \text{rad } \mathcal{A} \oplus KI$, where K is the field of scalars ($K = \mathbb{C}$ or $K = \mathbb{R}$) and KI is the one-dimensional subspace of \mathcal{A} spanned by the unit element I .

Since for $\dim X \leq 1$ the whole algebra $B(X)$ is commutative and trivial, we assume in our result that $\dim X > 1$. In this case we say that a closed linear subspace X_0 of X is proper, if $\{0\} \neq X_0 \neq X$. For an operator A in $B(X)$ denote by $\ker A$ its kernel and by $\text{im } A$ its range, i. e. the sets $\ker A = \{x \in X : Ax = 0\}$ and $\text{im } A = \{Ax : x \in X\}$. Our result reads as follows

Theorem. *Let X be a real or complex Banach space with $\dim X > 1$ and let X_0 be a proper closed linear subspace of X . Then the set*

$$(1) \quad \{A \in B(X) : \text{im } A \subset X_0 \text{ and } X_0 \subset \ker A\}$$

is a trivial Abelian subalgebra of $B(X)$ and its unitization \mathcal{A} is a trivial maximal Abelian subalgebra of $B(X)$.

Conversely, if \mathcal{A} is a trivial maximal Abelian subalgebra of $B(X)$, then its radical $\text{rad } \mathcal{A}$ is of the form (1), where

$$(2) \quad X_0 = \bigcap \{ \ker A : A \in \text{rad } \mathcal{A} \}.$$

Proof. Denote by M the set (1). Obviously, it is a trivial subalgebra of $B(X)$. Let T be an operator in the commutant M' . For a functional f in X^*

with $X_0 \subset \ker f$ and for an element z in X_0 denote by $A(f, z)$ the one-dimensional operator given by $A(f, z)x = f(x)z$, this operator is clearly in M and so it commutes with T . Thus, for all x in X , all z in X_0 and all f in X^* with $X_0 \subset \ker f$ we have

$$(3) \quad f(Tx)z = f(x)Tz.$$

Choosing $f_0 \neq 0$ with $X_0 \subset \ker f_0$ and substituting for x in (3) an element x_0 in X with $f_0(x_0) = 1$ we obtain

$$Tz = \alpha_T z$$

for all z in X_0 , where α_T is the scalar given by $\alpha_T = f_0(Tx_0)$. Put $T_1 = T - \alpha_T I$. We have $T_1 \in M'$ and $X_0 \subset \ker T_1$. We shall show that the operator T_1 is in M , i. e. $\text{im } T_1 \subset X_0$. If not, then there is an element u_0 in X with $T_1 u_0 \notin X_0$ and we can find an element A in M with $AT_1 u_0 \neq 0$ (A can be chosen to be of the form $A(f, z)$). But this is impossible, since $AT_1 u_0 = T_1 A u_0$ and $A u_0 \in X_0 \subset \ker T_1$. Thus, T_1 is in M and so T is in its unitization \mathcal{A} which is a commutative algebra and thus a maximal Abelian subalgebra of $B(X)$ since it equals M' .

Conversely, suppose that \mathcal{A} is a trivial maximal Abelian subalgebra of $B(X)$ and put $M = \text{rad } \mathcal{A}$. For any two operators T_1 and T_2 in M we have $\text{im } T_1 \subset \ker T_2$, and so $\text{im } T_1 \subset X_0$, where X_0 is given by (2). Since $X_0 \subset \ker T_1$ and T_1 is an arbitrary element of M , it follows that M is contained in the set (1). By the maximality of \mathcal{A} M equals to this set, and so $\text{rad } \mathcal{A}$ is of the form (1). The conclusion follows.

Corollary 1. Any subset S of $B(X)$ consisting of mutually annihilating operators (i. e. $T_1 T_2 = 0$ for all T_i in S , $i = 1, 2$) is contained in some trivial maximal Abelian subalgebra of $B(X)$. In particular, any trivial subalgebra of $B(X)$ is contained in a trivial maximal Abelian subalgebra of $B(X)$.

Denote by $\mathcal{A}(X_0)$ the trivial maximal Abelian subalgebra of $B(X)$ whose radical is (1).

Corollary 2. The algebra $\mathcal{A}(X_0)$ is isomorphic as a Banach space to the space $B(X/X_0, X_0) \oplus K$, where $B(U, V)$ denotes the Banach space of all continuous linear operators from a Banach space U to a Banach space V and K is the field of scalars (the one-dimensional Banach space).

Examples. Taking as X_0 any subspace of X of codimension one, we obtain a trivial maximal Abelian subalgebra $\mathcal{A}(X_0)$ isomorphic as a Banach space to the space X . Its radical consists of one-dimensional operators of the form $A(f_0, z)$, where f_0 is a fixed functional in X with $\ker f_0 = X_0$ and $z \in X_0$. The isomorphism between $\mathcal{A}(X_0)$ and X is given by

$$A(f_0, z) + \lambda I \leftrightarrow z + \lambda e_0,$$

where e_0 is a fixed element in X with $f_0(e_0) = 1$.

Similarly, taking as X_0 a linear subspace of X of dimension one $X_0 = Kx_0$ with $x_0 \in X$ and $\|x_0\| = 1$, we obtain an algebra $\mathcal{A}(X_0)$ isomorphic as a Banach space to the conjugate space X^* . It consists of all operators of the form $A(f, x_0) + \lambda I$, where $f \in X^*$ with $x_0 \in \ker f$ and $\lambda \in K$. The Banach space isomorphism between $\mathcal{A}(x_0)$ and X^* is given by

$$A(f, x_0) + \lambda I \leftrightarrow f + \lambda f_0,$$

where f_0 is a fixed element in X^* with $f_0(x_0) = 1$.

In case when the space X has a direct sum decomposition $X = X_0 \oplus X_1$, where X_1 is isomorphic to X_0 , then the algebra $\mathcal{A}(X_0)$ is isomorphic as a Banach space to the space $B(X_0)$. In particular when $X = H$ — an infinite-

dimensional Hilbert space, then H can be orthogonally decomposed as $H = H_0 \oplus H_1$, where H_0 and H_1 are isometrically isomorphic to H . In this case the algebra $\mathcal{A}(H_0)$ is isomorphic as a Banach space to the space $B(H)$. It can be proved that in this case the operators in $\mathcal{A}(H_0)$ are of the following form. Let R be a partial isometry on H , which maps H_1 isometrically onto H_0 and maps H_0 onto $\{0\}$. Then

$$\mathcal{A}(H_0) = \{RA + AR : A \in B(H) \text{ and } RAR = \alpha(A)R\},$$

where $\alpha(A)$ is a scalar depending upon A . It can be shown that if $RA + AR = RA_1 + A_1R$, then $\alpha(A) = \alpha(A_1)$ and so it defines on $\mathcal{A}(H_0)$ a functional f given by $f(RA + AR) = \alpha(A)$. It is a multiplicative linear functional on $\mathcal{A}(H_0)$ and its kernel equals to $\text{rad } \mathcal{A}(H_0)$ (we have $\alpha(R^*) = 1$ and $R^*R + RR^* = I$).

If $\dim X = n < \infty$, then by Corollary 2 the possible dimensions of algebras $\mathcal{A}(X_0)$ are $(n-k) \cdot k + 1$, $k = 1, 2, \dots, n-1$, and so there are $[\frac{n}{2}]$ non-isomorphic trivial maximal Abelian subalgebras of $B(X)$, where $[r]$ is the integral part of a number r . The largest possible dimension of $\mathcal{A}(X_0)$ is in this case $[\frac{n^2}{4}] + 1$ and the smallest dimension is n . All these results in the case of finite dimensional spaces are well known even for more general scalars (cf. [2, Chapt. 2, § 3]), however, in the case of real or complex scalars our reasoning seems to be shorter. In case when X is a Hilbert space the maximal Abelian subalgebras of $B(X)$ which are local rings are known in the literature (cf. [1], or [3, p. 81, proposition 4.4]), however, the existence of such trivial algebras seems to be new and somewhat surprising. We finish this paper with some simple results on invariant subspaces for algebras $\mathcal{A}(X_0)$. In the sequel we denote by $\text{lin}(X)$ the family of all closed linear subspaces of a Banach space X , and for a subset S of $B(X)$ we denote by $\text{lat}(S)$ the set (it has a structure of a lattice) of all subspaces in $\text{lin}(X)$ which are invariant with respect to all operators in S . In case when S consists of a single operator T we simply write $\text{lat}(T)$.

Proposition 1. *Let X be a Banach space with $\dim X > 1$. Then*

$$(4) \quad \text{lat}(\mathcal{A}(X_0)) = \{Y \in \text{lin}(X) : \text{either } X_0 \subset Y, \text{ or } Y \subset X_0\},$$

where X_0 is a proper linear subspace of X .

Proof. It is clear that all subspaces in the family (4) are invariant with respect to all operators in $\mathcal{A}(X_0)$. On the other hand, if Y is a closed linear subspace of X which contains some element $x_0 \notin X_0$ and does not contain some element $z_0 \in X_0$, then it cannot be invariant with respect to all elements in $\mathcal{A}(X_0)$, since there always exists an operator of the form $A(f, z_0)$ which sends x_0 to z_0 . The conclusion follows.

A subalgebra \mathcal{A} of $B(X)$ is said to be reflexive (sf. [3]), if the condition $\text{lat}(\mathcal{A}) \subset \text{lat}(T)$ implies $T \in \mathcal{A}$.

Proposition 2. *Let H be a Hilbert space, $\dim H > 1$, then no trivial maximal Abelian subalgebra of $B(H)$ is reflexive.*

Proof. For a closed proper linear subspace H_0 of H denote by $P(H_0)$ the orthogonal projection of H onto H_0 . Clearly, we have $\text{lat}(\mathcal{A}(H_0)) \subset \text{lat}(P(H_0))$ and $P(H_0) \notin \mathcal{A}(H_0)$. The conclusion follows.

REFERENCES

1. E. Albrecht. Maximal Abelian quasinilpotent subalgebras of $B(H)$. *Mathematisches Forschungsinstitut Oberwolfach, Tagungsbericht*, 10, 1985.
2. D. A. Suprunenko, R. I. Tyshkevich. *Commutative Matrices*. New York, 1968.
3. H. Radjavi, P. Rosenthal. *Invariant Subspaces*. Berlin — Heidelberg — New York, 1973.

*Mathematical Institute,
Polish Academy of Sciences
00-950 Warszawa Sniadeckich 8*

Received 10. 7. 1986