

Provided for non-commercial research and educational use.  
Not for reproduction, distribution or commercial use.

# Serdica

## Mathematical Journal

# Сердика

## Математическо списание

---

The attached copy is furnished for non-commercial research and education use only.  
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.  
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on  
Serdica Mathematical Journal  
which is the new series of  
Serdica Bulgaricae Mathematicae Publicationes  
visit the website of the journal <http://www.math.bas.bg/~serdica>  
or contact: Editorial Office  
Serdica Mathematical Journal  
Institute of Mathematics and Informatics  
Bulgarian Academy of Sciences  
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49  
e-mail: [serdica@math.bas.bg](mailto:serdica@math.bas.bg)

## WEAKLY COMPACT GENERATING AND SHRINKING MARKUŠEVIČ BASES

M. Fabian\*, P. Hájek†, V. Montesinos‡, V. Zizler§

*Communicated by G. Godefroy*

ABSTRACT. It is shown that most of the well known classes of nonseparable Banach spaces related to the weakly compact generating can be characterized by elementary properties of the closure of the coefficient space of Markuševič bases for such spaces. In some cases, such property is then shared by all Markuševič bases in the space.

Let  $X$  be a Banach space and let  $\langle \cdot, \cdot \rangle$  denote the canonical duality pairing between  $X$  and its dual space  $X^*$ . A system  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$ , where  $x_\gamma \in X$ ,  $x_\gamma^* \in$

---

2000 *Mathematics Subject Classification*: 46B30, 46B03.

*Key words*: Weakly compactly generated spaces, shrinking Markuševič bases, Eberlein compacta.

\*Supported by Institutional Research Plan of the Academy of Sciences of Czech Republic AVOZ 10190503 and grant IAA100190610 (Czech Republic).

†Supported in part by Institutional Research Plan of the Academy of Sciences of Czech Republic AVOZ 10190503 and grant GAČR 201/04/0090 (Czech Republic).

‡Supported in part by Project MTM2005-08210, the Universidad Politécnica de Valencia and the Generalitat Valenciana (Spain).

§Supported by grant NSERC 7926 (Canada), Institutional Research Plan of the Academy of Sciences of Czech Republic AVOZ 10190503 and GAČR A1019205 (Czech Republic).

$X^*$ ,  $\gamma \in \Gamma$ , is called a *Markuševič basis* for  $X$  if  $\langle x_\gamma, x_\gamma^* \rangle = 1$  for every  $\gamma \in \Gamma$ ,  $\langle x_\gamma, x_{\gamma'}^* \rangle = 0$  whenever  $\gamma, \gamma' \in \Gamma$  and  $\gamma \neq \gamma'$ , the linear span  $\text{sp}\{x_\gamma; \gamma \in \Gamma\}$  is dense in  $X$  and  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  is weak\* dense in  $X^*$ .

A compact space  $K$  is a *Corson compact* if for some set  $\Gamma$ ,  $K$  is homeomorphic to a subset  $T$  of a product  $[-1, 1]^\Gamma$  taken in its pointwise topology, such that for every  $t \in T$ ,  $\{\gamma \in \Gamma; t(\gamma) \neq 0\}$  is at most countable. A compact space  $K$  is called an *Eberlein compact* respectively a *uniform Eberlein compact* if for some set  $\Gamma$ ,  $K$  is homeomorphic to a compact set in  $c_0(\Gamma)$  (respectively  $\ell_2(\Gamma)$ ) endowed with the weak topology.

A Banach space  $X$  is called *weakly compactly generated* (WCG) if it contains a weakly compact subset whose linear span is dense in  $X$ . We will call a set  $S \subset X$  total if its linear span is dense in  $X$ . A Banach space  $X$  is called a *Vašák space* (i.e., *weakly countably determined Banach space*) if there exists a countable family  $\mathcal{K}$  of weak\* compact subsets of  $X^{**}$  such that, given any  $x \in X$  and  $x^{**} \in X^{**} \setminus X$ , there is  $K \in \mathcal{K}$  such that  $x \in K$  and  $x^{**} \notin K$ . A Banach space  $X$  is called *weakly Lindelöf determined* (WLD) if  $B_{X^*}$  in its weak star topology is a Corson compact which is the same as if there is a set  $\Delta \subset X$ , with  $\text{sp}\Delta$  dense in  $X$ , such that  $\Delta$  countably supports  $X^*$ , that is, for every  $x^* \in X^*$  the set  $\{\delta \in \Delta; \langle \delta, x^* \rangle \neq 0\}$  is at most countable (cf. e.g. [8]). It is well known that every WLD space admits a Markuševič basis, see, e.g., [8, Propositions 8.3.1, 6.1.7, 6.2.4], [12, Theorem 12.50] and the proof of Theorem 4 below. A Banach space is called an *Asplund space* if every of its separable subspaces has separable dual. We refer to [6], [8] and [12] for more on these concepts.

A result in [7] has the following known consequence, see, e.g., [8, page 112 and Theorem 8.3.3]:

**Theorem 0.** *A Banach space  $X$  is simultaneously WCG and Asplund if and only if  $X$  admits a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  which is shrinking, that is,  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  is norm dense in  $X^*$ .*

Note that not every Markuševič basis on a WCG Asplund space is shrinking. Indeed, in any nonreflexive separable Asplund space there exists a Schauder basis which is not shrinking, [22].

The purpose of this note is to show that the property of biorthogonality, together with known techniques of projectional resolutions of the identity operator make it possible to dualize and extend some results in [10], [9] and [13] to the case of Markushevich bases in the spirit of Theorem 0. More precisely, we show that most of the classes of Banach spaces related to the weakly compact generating, like WCG spaces, subspaces of WCG spaces, Vašák spaces can be characterized by replacing the norm topology in Theorem 0 by the topology of uniform convergence on an appropriate family of subsets of the Markuševič basis

for  $X$ . Note that the classes of spaces involved are known not to coincide (cf. e.g [8] and references therein).

The approach to these spaces that uses Markušević bases provides a good insight into these spaces (compare the statement in Theorems 2 and 3 for instance).

This is useful in questions in the renormings by smooth norms, smooth approximations, weak Asplund spaces, Asplund generated spaces and in the area of topology of special compacta (Eberlein, Corson, Gul'ko, Talagrand compacta), cf.e.g. [10], [13], [16]).

We made an effort to systematically present the results of this paper together with some folklore results in the area in order to provide the reader with a compact information on this subject.

**Theorem 1.** *A Banach space  $X$  is WCG if and only if it admits a Markušević basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  such that  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  is dense in  $X^*$  in the topology of uniform convergence on the set  $\{x_\gamma; \gamma \in \Gamma\}$ .*

It follows that the Markušević basis from Theorem 1 is weakly compact, that is, the set  $\{x_\gamma; \gamma \in \Gamma\} \cup \{0\}$  is weakly compact. Indeed, any sequence of distinct elements in it converges to 0 in the topology of the pointwise convergence on  $\{x_\gamma^*\}$  (by the orthogonality) and thus converges to 0 in the weak topology by the uniformity in the condition in Theorem 1. Note that it follows from [19] and from known results on Markušević bases that there exists a WCG space  $X$  with a Markušević basis such that the set  $\{x_\gamma; \gamma \in \Gamma\} \cup \{0\}$  cannot be written as the union of countably many weakly compact sets [2], [13].

On the other hand, W. Johnson proved that any unconditional basis in WCG space can be so decomposed (see Proposition 1.3 in [19]).

Note also that  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  from the Markušević basis in Theorem 1 may not be weak\* sequentially dense in  $X^*$ , i.e not every element of  $X^*$  can be reached as the weak star limit of a sequence from  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$ . Indeed, according to S. Banach, [5, Theorem 1, Annexe] there exists a weak\* dense subspace  $Y \subset c_0^*$  which is not weak\* sequentially dense in  $c_0^*$ . Then, [12, Theorem 6.41] yields a Markušević basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  such that  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\} \subset Y$ . See also Godun [15] who showed such a phenomenon in every separable quasireflexive Banach space.

Given a system of vectors  $x_\gamma \in X$ ,  $\gamma \in \Gamma$  and a set  $\Delta \subset \Gamma$ , we define a semimetric

$$\rho_\Delta(x_1^*, x_2^*) = \sup_{\gamma \in \Delta} |\langle x_\gamma, x_1^* - x_2^* \rangle|, \quad x_1^*, x_2^* \in X^*.$$

For  $x^* \in X^*$  and a set  $M \subset X^*$  we define

$$\rho_\Delta(x^*, M) = \inf\{\rho_\Delta(x^*, y^*); y^* \in M\}.$$

**Theorem 2.** *A Banach space  $X$  is a subspace of a WCG space if and only if  $X$  admits a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  with the following property: there are sets  $\Gamma_n \subset \Gamma$ ,  $n \in \mathbb{N}$ , such that*

$$\forall \epsilon > 0 \forall \gamma \in \Gamma \exists n \in \mathbb{N} \text{ so that } \gamma \in \Gamma_n \text{ and}$$

$$\forall x^* \in B_{X^*} \rho_{\Gamma_n}(x^*, \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}) < \epsilon.$$

*In this case every Markuševič basis in  $X$  has this property.*

It should be noted that the above sets  $\Gamma_n$ 's usually overlap, see [10, Theorem 2] for how to get a “non-overlapping” version of the above theorem.

**Theorem 3.** *A Banach space  $X$  is a Vašák space if and only if  $X$  admits a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  with the following property: there are sets  $\Gamma_n \subset \Gamma$ ,  $n \in \mathbb{N}$ , such that*

$$\forall \epsilon > 0 \forall \gamma \in \Gamma \forall x^* \in B_{X^*} \exists n \in \mathbb{N} \text{ so that } \gamma \in \Gamma_n \text{ and}$$

$$\rho_{\Gamma_n}(x^*, \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}) < \epsilon.$$

*In this case every Markuševič basis in  $X$  has this property.*

**Theorem 4.** *A Banach space  $X$  is WLD if and only if it has a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  with one of the equivalent properties listed in Proposition 1 below.*

*If this is the case, all Markuševič bases in  $X$  share the same properties.*

**Proposition 1.** *Given a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  in a Banach space  $X$ , then the following are equivalent.*

- (i) *For every  $x^* \in X^*$  the set  $\{\gamma \in \Gamma; \langle x_\gamma, x^* \rangle \neq 0\}$  is at most countable.*
- (ii)  *$\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  is sequentially dense in  $X^*$  endowed with the topology of the pointwise convergence on the set  $\{x_\gamma; \gamma \in \Gamma\}$ .*
- (iii)  *$\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  is countably dense in  $X^*$  in the weak\* topology, i.e. every element of  $X^*$  is in the closure in this topology of a countable set in  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$ .*

It is known that a Banach space  $X$  admits an equivalent uniformly Gâteaux smooth norm if and only if its dual unit ball  $B_{X^*}$  in the weak\* topology is a uniform Eberlein compact, see [11], [10] for definitions and proofs. For  $M \subset X^*$  and  $\kappa \in \mathbb{N}$  we define

$$\text{sp}_\kappa M = \left\{ \sum_{i=1}^{\kappa} a_i x_i^*; a_i \in \mathbb{R}, x_i^* \in M, i = 1, \dots, \kappa \right\}.$$

**Theorem 5.** *The dual ball  $(B_{X^*}, w^*)$  of a Banach space  $X$  is a uniform Eberlein compact if and only if  $X$  admits a Markuševič basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  with the*

following property: there are numbers  $\kappa(n) \in \mathbb{N}$  and sets  $\Gamma_n \subset \Gamma$ ,  $n \in \mathbb{N}$ , such that

$$\forall \epsilon > 0 \forall \gamma \in \Gamma \exists n \in \mathbb{N} \text{ so that } \gamma \in \Gamma_n \text{ and}$$

$$\forall x^* \in B_{X^*} \rho_{\Gamma_n}(x^*, \text{sp}_{\kappa(n)}\{x_\gamma^*; \gamma \in \Gamma_n\}) < \epsilon.$$

In this case every Markušević basis in  $X$  has this property.

The next theorem characterizes a subclass (called in [9] *strongly uniformly Gâteaux smooth* if  $\text{dens } X \leq \omega_1$ ) of the uniformly Gâteaux smooth Banach spaces –spaces admitting an equivalent uniformly  $M$ -smooth norm, with  $M \subset B_X$  a total set, see [10, Theorem 8].

**Theorem 6.** *A Banach space  $(X, \|\cdot\|)$  admits a total set  $\Gamma \subset B_X$  such that*

$$(1) \quad \forall \epsilon > 0 \exists \kappa \in \mathbb{N} \forall x^* \in B_{X^*} \#\{\gamma \in \Gamma; |\langle \gamma, x^* \rangle| \geq \epsilon\} \leq \kappa.$$

*if and only if  $X$  admits a Markušević basis  $\{x_\lambda; x_\lambda^*\}_{\lambda \in \Lambda}$  such that  $x_\lambda \in \text{sp}\Gamma$  for every  $\lambda \in \Lambda$ , and denoting  $\Delta = \{x_\lambda; \lambda \in \Lambda\}$ , we have*

$$\forall \epsilon > 0 \exists \kappa \in \mathbb{N} \forall x^* \in B_{X^*} \rho_\Delta(x^*, \text{sp}_\kappa\{x_\lambda^*; \lambda \in \Lambda\}) < \epsilon.$$

**Theorem 7.** *Let  $1 < p \leq 2$ , let  $\Gamma$  be an uncountable set, and let  $e_\gamma$ ,  $\gamma \in \Gamma$ , denote the canonical basis vectors in  $\ell_p(\Gamma)$ . A Banach space  $X$  admits a linear bounded mapping from  $\ell_p(\Gamma)$  onto a dense subset of  $X$  if and only if  $X$  admits a Markušević basis  $\{x_\lambda; x_\lambda^*\}_{\lambda \in \Lambda}$ , and a bounded linear operator  $T : \ell_p(\Lambda) \rightarrow X$  such that  $x_\lambda = Te_\lambda$  for every  $\lambda \in \Lambda$ .*

Note that the property exhibited in Theorem 7 characterizes, in case that  $\text{dens } X \leq \omega_1$ , the superreflexive generated spaces (see [9]).

**Proofs.** The following simple statement will be of frequent use.

**Lemma 1.** *Let  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  be a biorthogonal system in a Banach space  $X$ . Let  $x^* \in X^*$ ,  $\epsilon > 0$ , and  $\kappa \in \mathbb{N}$  be given. Then*

$$(i) \#\{\gamma \in \Gamma; |\langle x_\gamma, x^* \rangle| \geq \epsilon\} \leq \kappa \text{ if and only if } \rho_\Gamma(x^*, \text{sp}_\kappa\{x_\gamma^*; \gamma \in \Gamma\}) < \epsilon.$$

$$(ii) \#\{\gamma \in \Gamma; |\langle x_\gamma, x^* \rangle| \geq \epsilon\} < \omega \text{ if and only if } \rho_\Gamma(x^*, \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}) < \epsilon.$$

**Proof.** (i) Necessity. Denote  $F = \{\gamma \in \Gamma; |\langle x_\gamma, x^* \rangle| \geq \epsilon\}$ . If  $\gamma_0 \in \Gamma \setminus F$  then

$$\left| \left\langle x_{\gamma_0}, x^* - \sum_{\gamma \in F} \langle x_\gamma, x^* \rangle x_\gamma^* \right\rangle \right| = |\langle x_{\gamma_0}, x^* \rangle| < \epsilon.$$

If  $\gamma_0 \in F$ , then even

$$\left| \left\langle x_{\gamma_0}, x^* - \sum_{\gamma \in F} \langle x_\gamma, x^* \rangle x_\gamma^* \right\rangle \right| = 0.$$

It follows that

$$\rho_\Gamma(x^*, \text{sp}_\kappa\{x_\gamma^*; \gamma \in \Gamma\}) \leq \rho_\Gamma(x^*, \text{sp}\{x_\gamma^*; \gamma \in F\}) < \epsilon.$$

Conversely, find  $F \subset \Gamma$ , with  $\#F \leq \kappa$ , such that  $\rho_\Gamma(x^*, \text{sp}\{x_\gamma^*; \gamma \in F\}) < \epsilon$ . Assume that there exists  $\gamma_0 \in \Gamma \setminus F$  such that  $|\langle x_{\gamma_0}, x^* \rangle| \geq \epsilon$ . Then, for every  $y^* \in \text{sp}\{x_\gamma^*; \gamma \in F\}$  we get  $\rho_\Gamma(x^*, y^*) \geq |\langle x_{\gamma_0}, x^* - y^* \rangle| = |\langle x_{\gamma_0}, x^* \rangle| \geq \epsilon$ . Hence  $\rho_\Gamma(x^*, \text{sp}\{x_\gamma^*; \gamma \in F\}) \geq |\langle x_{\gamma_0}, x^* \rangle| \geq \epsilon$ , a contradiction. Therefore  $\{\gamma \in \Gamma; |\langle x_\gamma, x^* \rangle| \geq \epsilon\} \subset F$  and we are done.

(ii) follows immediately from (i).  $\square$

Let  $(X, \|\cdot\|)$  be a non-separable Banach space. Let  $\mu$  be the first ordinal whose cardinality is equal to the density  $\text{dens } X$  of  $X$ . A transfinite sequence of linear projections  $(P_\alpha)_{\omega \leq \alpha \leq \mu}$  on  $X$  is called a *projectional resolution of identity* (PRI) if  $P_\omega \equiv 0$ ,  $P_\mu \equiv I_X$  (the identity on  $X$ ) and for all  $\alpha, \beta \leq \mu$  we have  $\|P_\alpha\| = 1$ ,  $\text{dens } P_\alpha(X) \leq \#\alpha$  (the cardinality of  $\alpha$ ),  $P_\alpha P_\beta = P_\beta P_\alpha = P_{\min(\alpha, \beta)}$ , and for every  $x \in X$  the mapping  $\alpha \mapsto P_\alpha(x)$  from the ordinal segment  $[\omega, \mu]$  in its standard topology into  $X$  is continuous. A *separable projectional resolution of the identity* (separable PRI) on  $X$  is a transfinite sequence  $\{Q_\alpha : \omega \leq \alpha \leq \mu\}$  of linear projections on  $X$  such that  $Q_\omega \equiv 0$ ,  $Q_\mu \equiv I_X$ ,  $(Q_{\alpha+1} - Q_\alpha)X$  is separable for  $\omega \leq \alpha < \mu$ , for all  $\alpha, \beta \leq \mu$  we have  $\|Q_\alpha\| < +\infty$ ,  $Q_\alpha Q_\beta = Q_\beta Q_\alpha = Q_{\min(\alpha, \beta)}$ , and for every  $x \in X$ ,  $x \in \overline{\text{sp}}\{(Q_{\alpha+1} - Q_\alpha)(x); \alpha < \mu\}$ . If  $\Gamma$  is a subset of  $X$ , a PRI (a separable PRI)  $(P_\alpha)_{\omega \leq \alpha \leq \mu}$  on  $X$  is said to be *subordinated to*  $\Gamma$  if  $P_\alpha(\gamma) \in \{0, \gamma\}$  for all  $\gamma \in \Gamma$  and all  $\alpha \in [\omega, \mu]$ .

**Proposition 2.** *Let  $X$  be a Banach space with a total subset  $\Gamma$  which countably supports  $X^*$ . Then  $X$  has a separable PRI subordinated to  $\Gamma$ .*

**Proof.** If  $X$  is separable there is nothing to prove. Assume now that the lemma holds for every Banach space with density character less than a certain uncountable cardinal  $\aleph$ . Let  $X$  be a Banach space with density character  $\aleph$  and with a total subset  $\Gamma$  which countably supports  $X$ . By [10, Proposition 1],  $X$  has a PRI  $(P_\alpha)_{\omega \leq \alpha \leq \mu}$  subordinated to  $\Gamma$ . Now, for  $\omega \leq \alpha < \mu$ , the set  $(P_{\alpha+1} - P_\alpha)\Gamma (\subset \Gamma \cup \{0\})$  is total in  $(P_{\alpha+1} - P_\alpha)X$  and countably supports the dual  $((P_{\alpha+1} - P_\alpha)X)^*$ . Moreover,  $\text{dens}(P_{\alpha+1} - P_\alpha)X$  is less than  $\aleph$ . Then, by the induction hypothesis,  $(P_{\alpha+1} - P_\alpha)X$  has a separable PRI subordinated to  $(P_{\alpha+1} - P_\alpha)\Gamma$ .

Now, it is enough to use [8, Proposition 6.2.7].  $\square$

**Proof of Theorem 1. Necessity.** A well known result of Amir and Lindenstrauss [1] yields a weakly compact Markušević basis  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  in  $X$ , i.e. an Markušević basis  $\{x_\gamma, x_\gamma^*\}$  such that  $\{x_\gamma\} \cup 0$  is a weakly compact set in  $X$ . This means that for every  $\epsilon > 0$  and every  $x^* \in X^*$  the set  $\{\gamma \in \Gamma; |\langle x_\gamma, x_\gamma^* \rangle| \geq \epsilon\}$  is finite. Now it is enough to apply Lemma 1. The sufficiency is obvious since then the set  $\{x_\gamma : \gamma \in \Gamma\} \cup \{0\}$  must be weakly compact. Alternatively, one can use the Mackey-Arens theorem in this context.  $\square$

**Proof of Theorem 2 resp. 5.** The sufficiency follows from [10, Thm. 2 resp. 6] and our Lemma 1. As regards the necessity, let  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  be any Markušević basis in  $X$  with all  $x_\gamma$ 's in  $B_X$ . Write  $\gamma$  instead of  $x_\gamma$ . Thus we have that  $\Gamma \subset B_X$ . For this  $\Gamma$  find sets  $\Gamma_n^\epsilon$ ,  $n \in \mathbb{N}$ ,  $\epsilon > 0$ , as in (ii) of [10, Theorem 2 resp. 6]. Then the (countable) family of sets  $\Gamma_n^{1/i}$ ,  $n, i \in \mathbb{N}$ , satisfies, according to Lemma 1, the condition of Theorem 2, resp. Theorem 5.  $\square$

**Proof of Theorem 3.** Combine [10, Theorem 3] with Lemma 1 as it was done in the the previous proof.  $\square$

**Proof of Theorem 4.** The condition (i) in Proposition 1 implies that  $X$  is WLD. Conversely, assume that  $X$  is WLD, that is, there exists a total set  $\Delta \subset B_X$  which countably supports  $X^*$ . By Proposition 2, we find in  $X$  a separable PRI  $(P_\alpha)_{\omega \leq \mu}$  subordinated to the set  $\Delta$ . Fix an arbitrary  $\alpha \in [\omega, \mu)$ . We note that  $\Delta_\alpha := (P_{\alpha+1} - P_\alpha)\Delta \subset \Delta \cup \{0\}$  and that this set is total in the (separable) subspace  $(P_{\alpha+1} - P_\alpha)X$ . By the classical Markušević Theorem (see, for example, [12, Theorem 6.41]), in  $(P_{\alpha+1} - P_\alpha)X$ , there exists a Markušević basis  $\{x_{\alpha,n}; x_{\alpha,n}^*\}_{n \in \mathbb{N}}$  such that  $x_{\alpha,n} \in \text{sp}\Delta$  for every  $n \in \mathbb{N}$ . Define  $Q_\alpha : X \rightarrow (P_{\alpha+1} - P_\alpha)X$  by  $Q_\alpha x = (P_{\alpha+1} - P_\alpha)x$ ,  $x \in X$ . Then, replacing PRI by separable PRI in the proof of [8, Proposition 6.2.4], we can conclude that the system  $\{x_{\alpha,n}, Q_\alpha^* x_{\alpha,n}^*\}_{n \in \mathbb{N}, \omega \leq \alpha < \mu}$  forms a Markušević basis in  $X$ . It remains to check the cardinality condition in (i) of Proposition 1. Consider any  $x^* \in X^*$  and any  $n \in \mathbb{N}$ . If  $\alpha \in [\omega, \mu)$  satisfies  $\langle x_{\alpha,n}, x^* \rangle \neq 0$ , then  $\langle \delta, x^* \rangle \neq 0$  for some  $\delta \in \Delta_\alpha \subset \Delta \cup \{0\}$ . Thus

$$\#\{(\alpha, n) \in [\omega, \mu) \times \mathbb{N}; \langle x_{\alpha,n}, x^* \rangle \neq 0\} \leq \#\{\delta \in \Delta; \langle x^*, \delta \rangle \neq 0\} \cdot \omega = \omega,$$

and (i) in Proposition 1 is verified.

Finally, assume that  $X$  is WLD and let  $\{x_\gamma; x_\gamma^*\}_{\gamma \in \Gamma}$  be any Markušević basis in  $X$ . Put  $Y = \{x^* \in X^*; \#\{\gamma \in \Gamma; \langle x_\gamma, x^* \rangle \neq 0\} \leq \omega\}$ ; this is a linear set. Take any  $\xi$  in the weak\* closure of the intersection  $Y \cap B_{X^*}$ . Since  $(B_{X^*}, w^*)$  is a Corson compact, there exists a sequence in  $Y \cap B_{X^*}$  which weak\* converges to  $\xi$  ([12, Exercise 12.35]). Therefore  $\xi \in Y \cap B_{X^*}$ . We have thus proved that the latter set is weak\* closed. Then Banach-Dieudonné Theorem ensures that  $Y$  is weak\* closed. Moreover it contains  $\{x_\gamma^*; \gamma \in \Gamma\}$ , so  $Y = X^*$  and (i) is verified.  $\square$



**Proof of Proposition 1.** (i) $\Rightarrow$ (ii). Fix any  $x^* \in X^*$ . Enumerate the set  $\{\gamma \in \Gamma; \langle x_\gamma, x^* \rangle \neq 0\}$  as  $\{\gamma_1^o, \gamma_2^o, \dots\}$ . Find  $x_1^* \in \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  so that  $|\langle x_{\gamma_1^o}, x^* - x_1^* \rangle| < 1$ . Enumerate  $\{\gamma \in \Gamma; \langle x_\gamma, x_1^* \rangle \neq 0\}$  by  $\{\gamma_1^1, \gamma_2^1, \dots\}$ . Find  $x_2^* \in \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  so that  $|\langle x_{\gamma_1^o}, x^* - x_2^* \rangle| < \frac{1}{2}$ ,  $|\langle x_{\gamma_2^o}, x^* - x_2^* \rangle| < \frac{1}{2}$ ,  $|\langle x_{\gamma_1^1}, x^* - x_2^* \rangle| < \frac{1}{2}$ , and  $|\langle x_{\gamma_2^1}, x^* - x_2^* \rangle| < \frac{1}{2}$ . Assume that for some  $i \in \mathbb{N}$  we found  $x_i^*$  with “support” on  $\Gamma$  given by  $\{\gamma_1^j, \gamma_2^j, \dots\}$ ,  $j = 1, 2, \dots, i$ . Find then  $x_{i+1}^* \in \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  so that  $|\langle x_{\gamma_l^j}, x^* - x_{i+1}^* \rangle| < \frac{1}{i+1}$  for all  $j = 0, 1, \dots, i$  and  $l = 1, 2, \dots, i$ . Then we can easily see that  $\langle x_\gamma, x^* - x_i^* \rangle \rightarrow 0$  as  $i \rightarrow \infty$  for every  $\gamma \in \Gamma$ , and (ii) is proved.

(ii) $\Rightarrow$ (i). Take any  $x^* \in X^*$ . Let  $x_i^* \in \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$ ,  $i \in \mathbb{N}$ , be such that  $\langle x_\gamma, x^* - x_i^* \rangle \rightarrow 0$  as  $i \rightarrow \infty$  for every  $\gamma \in \Gamma$ . Now if  $\gamma \in \Gamma$  satisfies  $\langle x_\gamma, x^* \rangle \neq 0$ , then necessarily  $\langle x_\gamma, x_i^* \rangle \neq 0$  for some  $i \in \mathbb{N}$ . Hence

$$\{\gamma \in \Gamma; \langle x_\gamma, x^* \rangle \neq 0\} \subset \bigcup_{i=1}^{\infty} \{\gamma \in \Gamma; \langle x_\gamma, x_i^* \rangle \neq 0\}$$

and the set on the right hand side is countable.

(i) $\Rightarrow$ (iii). Let  $Y$  denote the set of all  $x^* \in X^*$  which lie in the weak\* closure of a countable subset of  $\text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$ . We want to show that  $Y = X^*$ . Clearly,  $Y$  is linear. Let  $\xi$  be any element of the weak\* closure of  $B_Y$ . (i) guarantees that  $(B_{X^*}, w^*)$  is a Corson compact, hence  $\xi$  can be reached as the weak\* limit of a sequence  $(x_i^*)_{i=1}^\infty$  in  $B_Y$ . Now, for every  $i \in \mathbb{N}$  we can find a suitable at most countable set  $C_i \subset \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  so that  $x_i^*$  lies in the weak\* closure of  $C_i$ . Then  $\xi$  lies in the (at most countable) set  $\bigcup_{i=1}^\infty C_i$ , and so  $\xi \in Y$ . Now, the Banach-Dieudonné Theorem guarantees that  $Y$  is weak\* closed. But  $Y$  contains  $\{x_\gamma^*; \gamma \in \Gamma\}$ . Therefore  $Y = X^*$ .

(iii) $\Rightarrow$ (i). Fix any  $x^* \in X^*$ . Find an at most countable set  $C \subset \text{sp}\{x_\gamma^*; \gamma \in \Gamma\}$  so that  $x^*$  belongs to the weak\* closure of  $C$ . Then

$$\{\gamma \in \Gamma; \langle x_\gamma, x^* \rangle \neq 0\} \subset \bigcup_{y^* \in C} \{\gamma \in \Gamma; \langle x_\gamma, y^* \rangle \neq 0\},$$

and the latter set is countable.  $\square$

**Proof of Theorem 6.** The sufficiency is trivial.

The necessity. Assume first that  $X$  is separable. By [12, Theorem 6.41], there exists a Markuševič basis  $\{x_n; x_n^*\}_{n \in \mathbb{N}}$  in  $X$  such that  $x_n \in \text{sp}\Gamma$  and  $\|x_n\| < \frac{1}{n}$  for every  $n \in \mathbb{N}$ . Then for every  $\epsilon > 0$  and for every  $x^* \in B_{X^*}$  we have  $\#\{n \in \mathbb{N}, |\langle x_n, x^* \rangle| \geq \frac{1}{\epsilon}\} < \frac{1}{\epsilon}$  and Lemma 1 finishes the proof.

Second, assume that  $X$  is non-separable. Let  $\Gamma \subset B_X$  be a total set satisfying (1). This set countably supports  $X^*$ . Thus, by Proposition 2, there exists a separable PRI  $(P_\alpha)_{\omega \leq \alpha < \mu}$  in  $X$  subordinated to  $\Gamma$ . Fix any  $\alpha \in [\omega, \mu)$  and denote  $\Gamma_\alpha = (P_{\alpha+1} - P_\alpha)\Gamma \subset (\Gamma \cup \{0\})$ . Fix any  $\alpha \in [\omega, \mu)$ . Let  $\{x_{\alpha,n}; x_{\alpha,n}^*\}_{n \in \mathbb{N}}$  be a Markušević basis in the (separable) subspace  $(P_{\alpha+1} - P_\alpha)X$  such that  $x_{\alpha,n} \in \text{conv}(\Gamma_\alpha \cup -\Gamma_\alpha)$  and  $\|x_{\alpha,n}\| < \frac{1}{n}$  for every  $n \in \mathbb{N}$ ; this can be done owing to [12, Theorem 6.41] and by an eventual rescaling. Define  $Q_\alpha : X \rightarrow (P_{\alpha+1} - P_\alpha)X$  by  $Q_\alpha x = (P_{\alpha+1} - P_\alpha)x$ ,  $x \in X$ . Then, replacing PRI by separable PRI in the proof of [8, Proposition 6.2.4], we can conclude that the system  $\{x_{\alpha,n}, Q_\alpha^* x_{\alpha,n}^*\}_{n \in \mathbb{N}, \omega \leq \alpha < \mu}$  forms a Markušević basis in  $X$ . Now fix any  $x^* \in B_X^*$  and consider any  $\omega \leq \alpha < \mu$  and any  $n \in \mathbb{N}$  such that  $|\langle x_{\alpha,n}, x^* \rangle| \geq \epsilon$ . Then  $n < \frac{1}{\epsilon}$  and there is  $\gamma \in \Gamma_\alpha$  so that  $|\langle \gamma, x^* \rangle| \geq \epsilon$ . Now, if  $\kappa \in \mathbb{N}$  was found for our  $\epsilon$  and our  $x^*$  by (1), we can estimate that

$$\#\{(\alpha, n) \in [\omega, \mu) \times \mathbb{N}; |\langle x_{\alpha,n}, x^* \rangle| \geq \epsilon\} < \frac{1}{\epsilon} \cdot \kappa.$$

Finally, Lemma 1 completes the proof.  $\square$

**Proof of Theorem 7.** The sufficiency part is trivial. Let us prove the necessity. To achieve this, assume for simplicity that  $\ell_p(\Gamma)$  is a dense subset of  $X$  and that  $\|f\| \leq \|f\|_{\ell_p}$  for every  $f \in \ell_p(\Gamma)$ . Fix any  $x^* \in X^*$ . Then the restriction  $x^*|_{\ell_p(\Gamma)}$  lies in  $\ell_p(\Gamma)^*$  ( $\equiv \ell_q(\Gamma)$ ) where  $q = \frac{p}{p-1}$ . Thus the set  $\{\gamma \in \Gamma; \langle e_\gamma, x^* \rangle \neq 0\}$  is at most countable which means that the set  $\{e_\gamma; \gamma \in \Gamma\}$  countably supports all elements of  $X^*$ . Then we can apply Proposition 2 and get a separable PRI  $(P_\alpha)_{\omega \leq \alpha < \mu}$  on  $X$  subordinated to the set  $\tilde{\Gamma} := \{e_\gamma; \gamma \in \Gamma\}$ . Fix any  $\alpha \in [\omega, \mu)$ . Put  $\tilde{\Gamma}_\alpha = (P_{\alpha+1} - P_\alpha)\tilde{\Gamma}$ . Note that  $\tilde{\Gamma}_\alpha \subset \tilde{\Gamma} \cup \{0\}$  and that  $\tilde{\Gamma}_\alpha$  is linearly dense in the (separable) subspace  $(P_{\alpha+1} - P_\alpha)X$ . By [12, Theorem 6.41], we find a Markušević basis  $\{x_{\alpha,n}; x_{\alpha,n}^*\}_{n \in \mathbb{N}}$  in the subspace  $(P_{\alpha+1} - P_\alpha)X$  such that  $x_n^\alpha \in \text{sp}\tilde{\Gamma}_\alpha \subset \ell_p(\Gamma)$  and  $\|x_n^\alpha\|_{\ell_p} = 1$  for every  $n \in \mathbb{N}$ . Define  $Q_\alpha : X \rightarrow (P_{\alpha+1} - P_\alpha)X$  by  $Q_\alpha x = (P_{\alpha+1} - P_\alpha)x$ ,  $x \in X$ . Performing this for every  $\omega \leq \alpha < \mu$ , we get the system  $\{\frac{1}{n}x_{\alpha,n}; nQ_\alpha^* x_{\alpha,n}^*\}_{n \in \mathbb{N}, \omega \leq \alpha < \mu}$ , which will be a Markušević basis in  $X$ , see, e.g, the proof of [8, Proposition 6.2.4]. For every element  $(a_{\alpha,m}; \omega \leq \alpha < \mu, m \in \mathbb{N})$  of  $\ell_p([\omega, \mu) \times \mathbb{N})$ , with finite support, we define

$$T(a_{\alpha,m}) = \sum_{m=1}^{\infty} \sum_{\omega \leq \alpha < \mu} a_{\alpha,m} \frac{1}{m} x_{\alpha,m}.$$

This is a linear mapping from a dense subset of  $\ell_p([\omega, \mu) \times \mathbb{N})$  into  $X$ . Now, using Hölder inequality and a disjoint support argument in the last of the following

inequalities, we can estimate

$$\begin{aligned} \|T(a_{\alpha,m})\| &\leq \sum_{m=1}^{\infty} \frac{1}{m} \left\| \sum_{\omega \leq \alpha < \mu} a_{\alpha,m} x_{\alpha,m} \right\| \\ &\leq \left( \sum_{m=1}^{\infty} \frac{1}{m^q} \right)^{\frac{1}{q}} \left( \sum_{m=1}^{\infty} \left\| \sum_{\omega \leq \alpha < \mu} a_{\alpha,m} x_{\alpha,m} \right\|^p \right)^{\frac{1}{p}} \\ &\leq C \left( \sum_{m=1}^{\infty} \left\| \sum_{\omega \leq \alpha < \mu} a_{\alpha,m} x_{\alpha,m} \right\|_{\ell_p}^p \right)^{\frac{1}{p}} \leq C \left( \sum_{m=1}^{\infty} \sum_{\omega \leq \alpha < \mu} |a_{\alpha,m}|^p \right)^{\frac{1}{p}} = C \|(a_{\alpha,m})\|_{\ell_p}; \end{aligned}$$

here we put  $\left( \sum_{m=1}^{\infty} \frac{1}{m^q} \right)^{\frac{1}{q}} = C$ . Therefore the mapping  $T$  can be extended to the whole space  $\ell_p([\omega, \mu) \times \mathbb{N})$ . Now, every canonical basic vector from this space is mapped by  $T$  to  $\frac{1}{m}x_{\alpha,m}$  with a suitable  $m \in \mathbb{N}$  and  $\omega \leq \alpha < \mu$ . Therefore the range of  $T$  is dense in  $X$  and the proof is finished.  $\square$

**Open problem.** Characterize Banach spaces  $X$  such that every subspace of  $X$  is WCG.

**Remark.** It is likely that in this problem additional axioms of set theory may play a role (see, for example, the use of Martin's axiom in [4] and [18]).

**Acknowledgements.** This paper originated when the first named author visited the School of Mathematical and Physical Sciences, University of New Castle, NSW, Australia in the fall of 2002. He would like to thank this Department and especially John Giles for their hospitality. We thank also the referee for his/hers remarks which helped to improve the final version of this paper.

## REFERENCES

- [1] D. AMIR, J. LINDENSTRAUSS. The structure of weakly compact sets in Banach spaces. *Ann. Math.* **88** (1968), 35–46.
- [2] S. ARGYROS, S. MERCOURAKIS. Examples concerning heredity problems of WCG Banach spaces. *Proc. Amer. Math. Soc.* (to appear).

- [3] S. ARGYROS, S. MERCOURAKIS. On weakly Lindelöf Banach spaces. *Rocky Mountain J. Math.* **23** (1993), 395–446.
- [4] A. AVILÉS. Radon-Nikodym compact spaces of low weight and Banach spaces. *STUDIA MATH.* **166**, 1 (2005), 71–82.
- [5] S. BANACH. *Théorie de Opérations Linéaires*. Chelsea Pub. Co., New York, 1932.
- [6] R. DEVILLE, G. GODEFROY, V. ZIZLER. *Smoothness and Renormings in Banach Spaces*. Pitman Monographs vol. **64**, London, Longman 1993.
- [7] M. FABIAN. Each weakly countably determined Asplund space admits a Fréchet differentiable norm. *Bull. Australian Math. Soc.* **36** (1987), 367–374.
- [8] M. FABIAN. *Gâteaux Differentiability of Convex Functions and Topology. Weak Asplund Spaces*. John Wiley & Sons, New York 1997.
- [9] M. FABIAN, G. GODEFROY, P. HÁJEK, V. ZIZLER. *Hilbert-generated spaces*, *J. Functional Analysis* **200** (2003), 301–323.
- [10] M. FABIAN, G. GODEFROY, V. MONTESINOS, V. ZIZLER. Inner characterizations of weakly compactly generated Banach spaces and their relatives. *J. Math. Anal. and Appl.* **297** (2004), 419–455.
- [11] M. FABIAN, G. GODEFROY, V. ZIZLER. The structure of uniformly Gâteaux smooth Banach spaces. *Israel Math. J.* **124** (2001), 243–252.
- [12] M. FABIAN, P. HABALA, P. HÁJEK, J. PELANT, V. MONTESINOS, V. ZIZLER. *Functional analysis and infinite dimensional geometry*. *Canad. Math. Soc. Books in Mathematics*, vol. **8**, Springer-Verlag, New York 2001.
- [13] M. FABIAN, V. MONTESINOS, V. ZIZLER. Biorthogonal systems in weakly Lindelöf spaces. *Canadian Math. Bull.* **48** (2005), 69–79.
- [14] V. FARMAKI. The structure of Eberlein, uniformly Eberlein and Talagrand compact spaces in  $\Sigma(\mathbb{R}^\Gamma)$ . *Fundamenta Math.* **128** (1987), 15–28.
- [15] B. V. GODUN. On weak\* derivatives of sets of linear operators. *Matematičeskije zametki* **23** (1978), 607–616 (in Russian).
- [16] P. HÁJEK, V. MONTESINOS, J. VANDERWERFF, V. ZIZLER. *Biorthogonal Systems in Banach spaces* (a book in preparation).
- [17] K. JOHN, V. ZIZLER. Smoothness and its equivalents in weakly compactly generated Banach spaces. *J. Funct. Anal.* **15** (1974), 161–166.

- [18] S. MERCOURAKIS, E. STAMATI. A new class of weakly  $\mathcal{K}$  analytic Banach spaces. (to appear).
- [19] H. P. ROSENTHAL. The heredity problem for weakly compactly generated Banach spaces. *Compositio Math.* **28** (1974), 83–111.
- [20] G. A. SOKOLOV. On some classes of compact spaces lying in  $\Sigma$ -products. *Comment. Math. Univ. Carolinae* **25** (1984), 219–231.
- [21] M. VALDIVIA. Biorthogonal systems in certain Banach spaces. *Revista Matemática* **9** (1996), 191–220.
- [22] M. ZIPPIN. A remark on bases and reflexivity in Banach spaces. *Israel J. Math.* **6** (1968), 74–79.
- [23] V. ZIZLER. Nonseparable Banach spaces. In: Handbook of the Geometry of Banach spaces (Eds W. B. Johnson, J. Lindenstrauss), Vol. 2, 2003, 1743–1816.

M. Fabian  
Mathematical Institute  
Czech Academy of Sciences  
Žitná 25  
11567 Prague 1, Czech Republic  
e-mail: fabian@math.cas.cz

V. Montesinos  
Departamento de Matemática Aplicada  
E.T.S.I. Telecomunicación  
Universidad Politécnica de Valencia  
Instituto de Matemática  
Pura y Aplicada  
C/Vera, s/n. 46071 Valencia, Spain  
e-mail: vmontesinos@mat.upv.es

P. Hájek  
Mathematical Institute  
Czech Academy of Sciences  
Žitná 25  
11567 Prague 1, Czech Republic  
e-mail: hajek@math.cas.cz

V. Zizler  
Mathematical Institute  
Czech Academy of Sciences  
Žitná 25  
11567 Prague 1, Czech Republic  
e-mail: zizler@math.cas.cz

Received September 23, 2005  
Revised September 29, 2006