

Provided for non-commercial research and educational use.  
Not for reproduction, distribution or commercial use.

# Serdica

## Mathematical Journal

# Сердика

## Математическо списание

---

The attached copy is furnished for non-commercial research and education use only.  
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.  
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on  
Serdica Mathematical Journal  
which is the new series of  
Serdica Bulgaricae Mathematicae Publicationes  
visit the website of the journal <http://www.math.bas.bg/~serdica>  
or contact: Editorial Office  
Serdica Mathematical Journal  
Institute of Mathematics and Informatics  
Bulgarian Academy of Sciences  
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49  
e-mail: [serdica@math.bas.bg](mailto:serdica@math.bas.bg)

## ON AN ODE RELEVANT FOR THE GENERAL THEORY OF THE HYPERBOLIC CAUCHY PROBLEM

Enrico Bernardi, Antonio Bove

*Communicated by L. Stoyanov*

*Dedicated to Professor Vesselin Petkov on the occasion of his 65th birthday*

ABSTRACT. In this paper we study an ODE in the complex plane. This is a key step in the search of new necessary conditions for the well posedness of the Cauchy Problem for hyperbolic operators with double characteristics.

**1. Introduction and statements.** The purpose of this paper is to study the existence of solutions, bounded on the real axis, for an ordinary differential equation (ode). It is a well known fact that many problems in the theory of PDEs reduce to the study of an ode; the problem that motivates us here is the well posedness of the Cauchy problem for a class of hyperbolic operators with double characteristics.

---

2000 *Mathematics Subject Classification*: 34E20, 35L80, 35L15.

*Key words*: Cauchy problem, zeroes of entire functions, asymptotic expansions, Hamilton systems.

More specifically it is known, see e.g. [3] and [1], that the Cauchy problem in the presence of double characteristics is not well posed if there are bicharacteristic curves issuing from a simple characteristic point and tending to some double characteristic point. We say that in this case the Hamilton field gives rise to an unstable dynamical system.

In order to show that the Cauchy problem in such a case is not well posed in  $C^\infty$  one has to construct an asymptotic solution violating the a priori estimates implied by the well posedness of the Cauchy problem. This is done by showing that a certain ode has a solution which is bounded on the real axis (actually exponentially decreasing at infinity) if the complex parameter appearing in the equation is suitably chosen. Sometimes this parameter is called a nonlinear eigenvalue.

From a geometrical point of view we know that the Hamilton system is unstable if the principal symbol of the operator is transversally non degenerate and there is a Jordan block of size 4 in the canonical form of the Hamilton matrix, which can roughly be thought of as  $dH_p$ , where  $p$  is the principal symbol and  $H_p$  its Hamilton field. There are though other more complex situations where one may have an unstable Hamilton vector field, exhibiting higher degeneracies or a non constant rank of the symplectic form.

In this paper we study the equation

$$(1.1) \quad u''(x) = (x^{2k+1} + zx^k)u(x),$$

where  $z \in \mathbb{C}$  and  $k = 1, 2$ . The case  $k = 1$  corresponds to a neat transversally non degenerate situation, while  $k > 1$  corresponds to higher order degeneracies.

It is a striking phenomenon that the proof for  $k > 1$ , actually  $k = 2$ , is much more difficult than that where  $k = 1$ . The whole proof boils down to showing that the Wronskian of two linearly independent solutions—an entire function of  $z$ —vanishes. In the  $k = 1$  case some simple complex analysis is enough, while if  $k = 2$  we must estimate the order of the entire function and use some supplementary arguments. This is done following an idea of Christ [6] (see also [8]).

The statement of our result is given in Theorem 3.2.2 at the end of the paper. We have no doubts that an analogous result should hold for arbitrary  $k$ , but as of now the proof seems to be much longer and difficult.

**2. An upper bound for the order of an entire function.** Aim of this section is to study the exponential order of growth of the Wronskian of

equation (1.1), which can be written as

$$(2.1) \quad P_z(u) = \left( -\frac{d^2}{dx^2} + p_z(x) \right) u(x),$$

with  $p_z(x) = x^{2k+1} + zx^k$ . We define:

$$(2.2) \quad \phi_z(x) = \frac{2}{2k+3} x^{\frac{2k+3}{2}} + zx^{\frac{1}{2}}.$$

Then  $\phi'_z(x) = x^{\frac{2k+1}{2}} + \frac{1}{2}zx^{-\frac{1}{2}}$ . To keep the notation simple we omit the  $z$  parameter and write just  $u, P, p, \phi$ . We see that  $p(x) = \phi'^2(x) - \frac{1}{4}z^2x^{-1}$ . Let

$$G(x) = \frac{1}{\phi'^{1/2}(x)} e^{-\phi(x)}$$

and

$$A = A(z) = |z|^{\frac{2}{1-\delta}},$$

where  $\delta < 1$  and will be chosen suitably later. Let  $\psi = \log G = -\phi - \frac{1}{2} \log \phi'$  and  $D = \frac{d}{dx} + \psi'$ ,  $\tilde{D} = -\frac{d}{dx} + \psi'$ , where we choose the principal branch of the complex logarithm. Then we have:

$$(2.3) \quad D\tilde{D} = \left( \frac{d}{dx} + \psi' \right) \left( -\frac{d}{dx} + \psi' \right) = -\frac{d^2}{dx^2} + \psi'' + \psi'^2,$$

where  $\psi' = -\phi' - \frac{1}{2} \frac{\phi''}{\phi'}$ ,  $\psi'' = -\phi'' - \frac{1}{2} \left( \frac{\phi''}{\phi'} \right)'$ .

So that:

$$(2.4) \quad \begin{aligned} \psi'' + \psi'^2 &= -\phi'' - \frac{1}{2} \left( \frac{\phi''}{\phi'} \right)' + \left( \phi' + \frac{1}{2} \frac{\phi''}{\phi'} \right)^2 \\ &= E + \phi'^2 = E + p(x) + \frac{1}{4}z^2x^{-1}, \end{aligned}$$

where we put  $E = -\frac{1}{2} \left( \frac{\phi''}{\phi'} \right)' + \frac{1}{4} \left( \frac{\phi''}{\phi'} \right)^2$ .

Thus

$$(2.5) \quad D\tilde{D} = -\frac{d^2}{dx^2} + p(x) + E + \frac{1}{4}z^2x^{-1} = P + E + \frac{1}{4}z^2x^{-1}.$$

Let us compute

$$PG = \left( -\frac{d^2}{dx^2} + p \right) \left( \frac{1}{\phi^{1/2}(x)} e^{-\phi(x)} \right).$$

We have:

$$\begin{aligned} -\frac{d^2}{dx^2} \left( \frac{1}{\phi^{1/2}(x)} e^{-\phi(x)} \right) \\ = - \left( \frac{1}{(\phi')^{1/2}} \right)'' e^{-\phi} + 2\phi' \left( \frac{1}{(\phi')^{1/2}} \right)' e^{-\phi} - \frac{e^{-\phi}}{(\phi')^{1/2}} (\phi'^2 - \phi'') \end{aligned}$$

And thus

$$\begin{aligned} (2.6) \quad & \left( -\frac{d^2}{dx^2} + p(x) \right) G \\ & = - \left( \frac{1}{(\phi')^{1/2}} \right)'' e^{-\phi} + 2\phi' \left( \frac{1}{(\phi')^{1/2}} \right)' e^{-\phi} - \frac{e^{-\phi}}{(\phi')^{1/2}} (\phi'^2 - \phi'') \\ & \quad + \phi'^2 \frac{1}{(\phi')^{1/2}} e^{-\phi} - \frac{1}{4} z^2 x^{-1} \frac{1}{(\phi')^{1/2}} e^{-\phi} \\ & = - \left( \frac{1}{(\phi')^{1/2}} \right)'' e^{-\phi} - \frac{1}{4} z^2 x^{-1} \frac{1}{(\phi')^{1/2}} e^{-\phi} \\ & \quad = -(\phi')^{1/2} \left( \frac{1}{(\phi')^{1/2}} \right)'' G - \frac{1}{4} z^2 x^{-1} G \end{aligned}$$

Now

$$\frac{\left( \frac{1}{(\phi')^{1/2}} \right)''}{\left( \frac{1}{(\phi')^{1/2}} \right)} = (\phi')^{1/2} \left( \frac{1}{(\phi')^{1/2}} \right) = \mathcal{O}(x^{-2}).$$

So that

$$|PG| \leq C|x|^{-2}|G(x)| + \frac{1}{4}|z|^2|x|^{-1}|G(x)|.$$

Let  $A < x < y$  and denote by  $u_y(x)(= u(x))$  the solution of:

$$(2.7) \quad \begin{cases} Pu = PG, & A < x < y \\ u(y) = u'(y) = 0. \end{cases}$$

Set

$$\begin{aligned} v &= \tilde{D}u \\ e^{-\psi} \frac{d}{dx} e^{\psi} v &= Dv = D\tilde{D}u \\ &= Pu + Eu + \frac{1}{4}z^2x^{-1}u, \end{aligned}$$

then, because  $v(y) = 0$ , for  $A \leq s \leq y$  we may write:

$$(2.8) \quad e^{\psi(s)}v(s) = \int_y^s e^{\psi(t)}(PG(t) + Eu + \frac{1}{4}z^2t^{-1}u)dt.$$

From (2.8) we then deduce that

$$|v(s)| \leq |e^{-\psi(s)}| \int_y^s |e^{\psi(t)}| [ |PG(t)| + |Eu| + \frac{1}{4}|z^2|t^{-1}|u| ] dt.$$

**Lemma 2.1.** *If  $x \geq |z|^{\frac{2}{1-\delta}}$  we have*

$$|\phi'(x)| \sim x^{\frac{2k+1}{2}}.$$

The proof is straightforward and is left to the reader. Thus, due to Lemma 2.1 we have that:

$$\begin{aligned} |e^{-\psi}| &= \left| \frac{1}{G} \right| = |(\phi')^{1/2}| |e^{\phi}| \lesssim x^{\frac{2k+1}{4}} |e^{\phi}| \\ |e^{\psi}| &= |G| \lesssim x^{-\frac{2k+1}{4}} |e^{-\phi}|, \end{aligned}$$

with suitable constants.

The function  $v$  can be then estimated as follows:

$$(2.9) \quad |v(s)| \leq s^{\frac{2k+1}{4}} |e^{\phi}| \int_s^y t^{-\frac{2k+1}{4}} |e^{-\phi(t)}| [ Ct^{-2}|G(t)| + \frac{1}{4}|z|^2t^{-1}|G(t)| + Ct^{-2}|u(t)| + \frac{1}{4}|z|^2t^{-1}|u(t)| ] dt = \sum_1^4 I_j,$$

and

$$\begin{aligned}
I_1 &= C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{4}} |e^{-\phi(t)}| C t^{-2} |G(t)| dt \\
&\leq C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{4}} |e^{-\phi(t)}| t^{-\frac{2k+1}{4}-2} |e^{-\phi(t)}| dt \\
&\leq C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{2}-2} |e^{-2\phi(t)}| dt \\
&\leq C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{2}-2} \frac{1}{(-2 \operatorname{Re} \phi)'} (-2 \operatorname{Re} \phi(t))' e^{-2 \operatorname{Re} \phi(t)} dt.
\end{aligned}$$

As above  $(\operatorname{Re} \phi)' \sim \operatorname{Re} \phi' \sim x^{\frac{2k+1}{2}}$ , so that

$$\begin{aligned}
I_1 &\leq C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-(2k+1)-2} (-2 \operatorname{Re} \phi(t))' e^{-2 \operatorname{Re} \phi(t)} dt \\
&\leq C s^{\frac{2k+1}{4}} |e^\phi(s)| s^{-(2k+1)-2} |e^{-2\phi(s)}| \\
&= s^{-\frac{3(2k+1)}{4}-2} |e^{-\phi(s)}|.
\end{aligned}$$

$$\begin{aligned}
I_2 &= C |z|^2 s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{4}} |e^{-\phi(t)}| t^{-1} |G(t)| dt \\
&\leq C |z|^2 s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{2}-1} |e^{-2\phi(t)}| dt \\
&\leq C |z|^2 s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{2}-1} \frac{1}{(-2 \operatorname{Re} \phi)'} (-2 \operatorname{Re} \phi)' e^{-2 \operatorname{Re} \phi(t)} dt \\
&\leq C |z|^2 s^{\frac{2k+1}{4}} |e^\phi(s)| s^{-(2k+1)-1} |e^{-2\phi(s)}| \\
&= C |z|^2 s^{-\frac{3(2k+1)}{4}-1} |e^{-\phi(s)}|.
\end{aligned}$$

But  $|z|^2 \leq s^{1-\delta}$ , so that

$$I_2 \leq C s^{-\frac{3(2k+1)}{4}-\delta} |e^{-\phi(s)}|.$$

Let us now estimate  $I_3$ :

$$\begin{aligned}
I_3 &= C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{4}} |e^{-\phi(t)}| t^{-2} |u(t)| dt \\
&= C s^{\frac{2k+1}{4}} |e^\phi(s)| \int_s^y t^{-\frac{2k+1}{4}-2} |e^{-\phi(t)}| u(t) dt.
\end{aligned}$$

Finally:

$$I_4 = C s^{\frac{2k+1}{4}} |e^{\phi(s)}| \int_s^y |z|^2 t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| dt.$$

As above  $|z|^2 \leq s^{1-\delta}$  so that:

$$I_4 \leq C s^{\frac{2k+1}{4}+1-\delta} |e^{\phi(s)}| \int_s^y t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| dt.$$

Altogether:

$$\begin{aligned} \sum_1^4 I_j &\leq C s^{-\frac{3(2k+1)}{4}-\delta} |e^{-\phi(s)}| \\ &+ C s^{\frac{2k+1}{4}} |e^{\phi(s)}| \int_s^y t^{-\frac{2k+1}{4}-2} |e^{-\phi(t)} u(t)| dt \\ &+ C s^{\frac{2k+1}{4}+1-\delta} |e^{\phi(s)}| \int_s^y t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| dt. \end{aligned}$$

Since  $|z| \geq 1$  and hence  $s \geq 1$  the third quantity dominates the second:

$$(2.10) \quad |v(s)| \leq \sum_1^4 I_j \leq C \left\{ s^{-\frac{3(2k+1)}{4}-\delta} |e^{-\phi(s)}| + s^{\frac{2k+1}{4}+1-\delta} |e^{\phi(s)}| \int_s^y t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| dt \right\}.$$

Let us estimate the function  $u$  for  $A \leq x \leq y$ :

$$\begin{aligned} |u(x)| &\leq |e^{\psi(x)}| \int_x^y |e^{-\psi(s)} v(s)| ds \\ &\leq x^{-\frac{2k+1}{4}} |e^{-\phi(x)}| \int_x^y s^{\frac{2k+1}{4}} |e^{\phi(s)}| |v(s)| ds \\ &\leq C x^{-\frac{2k+1}{4}} |e^{-\phi(x)}| \int_x^y s^{\frac{2k+1}{4} - \frac{3(2k+1)}{4} - \delta} ds \\ &+ C x^{-\frac{2k+1}{4}} |e^{-\phi(x)}| \int_x^y s^{\frac{2k+1}{4} + \frac{2k+1}{4} + 1 - \delta} |e^{2\phi(s)}| ds \\ &\quad \cdot \int_s^y t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| dt ds \\ &= J_1 + J_2. \end{aligned}$$



$$\begin{aligned} J_1 &\leq C \left| \frac{1}{(\phi')^{1/2}} e^{-\phi} \right| \int_x^y s^{-\frac{2k+1}{2}-\delta} ds \\ &\leq C x^{-\frac{2k+1}{2}+1-\delta} |G(x)|. \end{aligned}$$

$$\begin{aligned} J_2 &= C x^{-\frac{2k+1}{4}} |e^{-\phi(x)}| \int_x^y ds s^{\frac{2k+1}{2}+1-\delta} |e^{2\phi(s)}| \\ &\quad \cdot \int_s^y t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} u(t)| \\ &\leq C |G(x)| \int_x^y dt \left( \int_x^y ds s^{\frac{2k+1}{2}+1-\delta} |e^{2\phi(s)}| \right) \\ &\quad \cdot t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)}|. \end{aligned}$$

Since  $\phi'(s) \sim s^{\frac{2k+1}{2}}$  the inner integral can be estimated by

$$t^{1-\delta} |e^{2\phi(t)}|,$$

so that

$$J_2 \leq C |G(x)| \int_x^y t^{-\frac{2k+1}{4}-1+1-\delta} |e^{\phi(t)}| |u(t)| dt.$$

Hence, defining

$$B_\alpha = \sup_{[A,y]} \frac{|u(x)|}{x^{-\alpha} |G(x)|},$$

for a suitable  $\alpha > 0$ , we have

$$\begin{aligned} |u(x)| \leq J_1 + J_2 &\leq C \left\{ x^{-\frac{2k+1}{4}-\delta} |G(x)| \right. \\ &\quad \left. + |G(x)| \int_x^y t^{-\frac{2k+1}{4}-\delta} |e^{\phi(t)}| |u(t)| dt \right\} \\ &= C |G(x)| \left\{ x^{-\frac{2k+1}{2}+1-\delta} \right. \\ &\quad \left. + \int_x^y t^{-\frac{2k+1}{4}-\delta} \frac{|u(t)|}{t^{-\alpha} \left| \frac{1}{(\phi')^{1/2}} e^{-\phi(t)} \right|} t^{-\alpha} t^{-\frac{2k+1}{4}} dt \right\} \end{aligned}$$

$$\begin{aligned} &\leq C|G(x)| \left\{ x^{-\frac{2k+1}{2}+1-\delta} + \int_x^y t^{-\frac{2k+1}{2}-\alpha-\delta} B dt \right\} \\ &\leq C|G(x)| \left\{ x^{-\frac{2k+1}{2}+1-\delta} + B_\alpha x^{-\frac{2k+1}{2}-\alpha-\delta+1} \right\} \\ &= C|G(x)| x^{-\frac{2k+1}{2}+1-\delta} [1 + B_\alpha x^{-\alpha}]. \end{aligned}$$

By choosing

$$\alpha = \frac{2k+1}{2} + \delta - 1 > 0$$

we obtained that  $B_\alpha \leq C \frac{1}{1-cx^{-\alpha}}$ . Thus we proved the following result:

**Proposition 2.1.** *Let  $\delta > -\frac{2k+1}{2} + 1$ . For  $|z|^{\frac{2}{1-\delta}} \leq x \leq y$  we have the estimate:*

$$(2.11) \quad |u_y(x)| \leq Cx^{-\alpha}|G(x)|,$$

with a constant  $C$  independent of  $z$  and  $y$ .

Define  $u_y$  to be such that:

$$\begin{cases} P_z u_y(x) = P_z G(x) \\ u_y(y) = u'_y(y) = 0 \end{cases}$$

As before define  $\alpha = \frac{2k+1}{2} + \delta - 1 > 0, k = 1, 2, \dots$

For  $A \leq x \leq y_1 \leq y_2$  we want to find an upper bound for

$$|u_{y_1}(x) - u_{y_2}(x)|$$

It has already been proved in Proposition 2.1 that

$$|u_{y_2}(y_1)| \leq Cy_1^{-\alpha}|G(y_1)|.$$

Furthermore, from the estimate (2.10) we have:

$$\begin{aligned}
 |\tilde{D}u_{y_2}(s)| &\leq Cs^{-\frac{3(2k+1)}{4}-\delta}|e^{-\phi(s)}| + Cs^{\frac{2k+1}{4}+1-\delta}|e^{\phi(s)}| \\
 &\quad \cdot \int_s^{y_2} t^{-\frac{2k+1}{4}-1}|e^{-\phi(t)}u_{y_2}(t)|dt \\
 &\leq Cs^{-\frac{2k+1}{2}-\delta}|G(s)| + Cs^{\frac{2k+1}{4}+1-\delta}|e^{\phi(s)}| \\
 &\quad \cdot \int_s^{y_2} t^{-(2k+1)-\delta}|e^{-2\phi(t)}|dt \\
 &\leq Cs^{-\frac{2k+1}{2}-\delta}|G(s)| + Cs^{-\frac{5(2k+1)}{4}+1-2\delta}|e^{-\phi(s)}| \\
 &\leq Cs^{-\frac{2k+1}{2}-\delta}|G(s)|,
 \end{aligned}$$

since  $0 \geq -\frac{3(2k+1)}{4} + 1 - \delta$ , for  $k = 1, 2, \dots$

Set

$$w_1 = u_{y_1} - u_{y_2} \quad v = \tilde{D}w_1.$$

Since  $u_{y_1}(y_1) = u'_{y_1}(y_1) = 0$ , we have:

$$\begin{aligned}
 |w_1(y_1)| = |u_2(y_1)| &\leq Cy_1^{-\alpha}|G(y_1)| \\
 &\leq Cy^{-\alpha-\frac{2k+1}{4}}|e^{-\phi(y_1)}| \\
 |v(y_1)| = |\tilde{D}u_{y_2}(y_1)| &\leq Cy_1^{-\frac{2k+1}{2}-\delta}|G(y_1)| \\
 &\leq y_1^{-\frac{3(2k+1)}{4}-\delta}|e^{-\phi(y_1)}|.
 \end{aligned}$$

Now:

$$\begin{aligned}
 e^{-\psi} \frac{d}{ds} e^{\psi} v &= Dv = D\tilde{D}w_1 \\
 &= Pw_1 + Ew_1 + \frac{1}{4}z^2x^{-1}w_1 \\
 &= Ew_1 + \frac{1}{4}z^2x^{-1}w_1,
 \end{aligned}$$

so that, for  $A \leq s \leq y_1$ ,

$$\begin{aligned}
 |e^{\psi(s)}v(s)| &\leq |e^{\psi(y_1)}v(y_1)| + \int_s^{y_1} |e^{\psi(t)}(Ew_1 + \frac{1}{4}z^2t^{-1}w_1)|dt \\
 &\leq Cy_1^{-(2k+1)-\delta}|e^{-2\phi(y_1)}| + \int_s^{y_1} t^{-\frac{2k+1}{4}-2}|e^{-\phi(t)}||w_1|dt \\
 &\quad + \frac{1}{4}z^2 \int_s^{y_1} t^{-\frac{2k+1}{4}-1}|e^{-\phi(t)}||w_1|dt
 \end{aligned}$$

On the other hand, since  $-e^\psi d_x e^{-\psi} w_1 = v$ , we have:

$$\begin{aligned} |e^{-\psi(x)} w_1(x)| &\leq |e^{-\psi(y_1)} w_1(y_1)| \\ &\quad + \int_x^{y_1} |e^{-\psi(s)} v(s)| ds \\ &= |e^{-\psi(y_1)} w_1(y_1)| + \int_x^{y_1} |e^{-2\psi(s)}| |e^{\psi(s)} v(s)| ds. \end{aligned}$$

We estimate the two pieces of the above expression:

$$|e^{-\psi(y_1)} w_1(y_1)| \leq \frac{1}{|G(y_1)|} C y_1^{-\alpha} |G(y_1)| = C y_1^{-\alpha}.$$

$$\begin{aligned} \int_x^{y_1} |e^{-2\psi(s)}| |e^{\psi(s)} v(s)| ds &\leq C \int_x^{y_1} |e^{-2\psi(s)}| y_1^{-(2k+1)-\delta} |e^{-2\phi(y_1)}| ds \\ &\quad + C \int_x^{y_1} |e^{-2\psi(s)}| \int_s^{y_1} t^{-\frac{2k+1}{4}-2} |e^{-\phi(t)} w_1(t)| dt ds \\ &\quad + C |z|^2 \int_x^{y_1} |e^{-2\psi(s)}| \int_s^{y_1} t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} w_1(t)| dt ds = \sum_1^3 H_j. \end{aligned}$$

Let us consider the different terms  $H_j$ :

$$\begin{aligned} H_1 &\leq C y_1^{-(2k+1)-\delta} |e^{-2\phi(y_1)}| \int_x^{y_1} s^{\frac{2k+1}{2}} |e^{2\phi(s)}| ds \\ &\leq C y_1^{-(2k+1)-\delta} |e^{-2\phi(y_1)}| |e^{2\phi(y_1)}| \\ &= C y_1^{-(2k+1)-\delta}. \end{aligned}$$

$$\begin{aligned} H_2 &= C \int_x^{y_1} t^{-\frac{2k+1}{4}-2} |e^{-\phi(t)} w_1(t)| \int_x^t |e^{-2\psi(s)}| ds dt \\ &\leq C \int_x^{y_1} t^{-\frac{2k+1}{4}-2} |e^{\phi(t)} w_1(t)| dt, \end{aligned}$$

arguing as above for the inner integral.

$$\begin{aligned} H_3 &\leq C |z|^2 C \int_x^{y_1} t^{-\frac{2k+1}{4}-1} |e^{-\phi(t)} w_1(t)| \int_x^t |e^{-2\psi(s)}| ds dt \\ &\leq C |z|^2 \int_x^{y_1} t^{-\frac{2k+1}{4}-1} |e^{\phi(t)} w_1(t)| dt. \end{aligned}$$

Since  $|z|^2 \leq x^{1-\delta} \leq t^{1-\delta}$  we see that  $H_2 \leq H_1$ . Thus, since  $\alpha = \frac{2k+1}{2} - 1 + \delta$ ,

$$\begin{aligned} |e^{-\psi(x)}w_1(x)| &\leq Cy_1^{-\alpha} + C \int_x^{y_1} t^{-\frac{2k+1}{4}-\delta}|e^{\phi(t)}w_1(t)|dt \\ &\leq Cy_1^{-\alpha} + C \int_x^{y_1} t^{-\frac{2k+1}{2}-\delta}|e^{-\psi(t)}w_1(t)|dt. \end{aligned}$$

Set

$$B_1 = \sup_{A \leq x \leq y_1} \frac{|w_1(x)|}{|G(x)|}.$$

Then we have:

$$B_1 \leq Cy_1^{-\alpha} + C \int_x^{y_1} B_1 t^{-\alpha-1} dt \leq Cy_1^{-\alpha} + \frac{B_1}{2},$$

which implies

$$B_1 \leq Cy_1^{-\alpha}.$$

We have thus proved the following

**Proposition 2.2.** *Let  $\delta > -\frac{2k+1}{2} + 1$ . Then for  $A(z) \leq x \leq y$  we have the inequality*

$$(2.12) \quad |u_{y_1}(x) - u_{y_2}(x)| \leq Cy_1^{-\alpha}|G(x)|,$$

where the positive constant  $C$  is independent of  $y_1, y_2$  and  $z$ .

Using both Proposition 2.1 and Proposition 2.2 we now proceed to prove an upper bound for the exponential order of the Stokes coefficient.

Denote by  $u^+$  a solution of  $Pu^+ = 0$  subdominant e.g. in a sector containing the positive real axis. Then we know that

$$(2.13) \quad u^+(x, z) = x^{-\frac{2k+1}{4}} e^{-\phi_z(x)}(1 + \mathcal{O}(x^{-1}))$$

**Proposition 2.3.** *Let  $A(z) = C_0|z|^{\frac{2}{1-\delta}}$ , where  $C_0 > 0$  and  $k \geq 2$ . Then there exists a constant  $C \in ]0, 1[$  and independent of  $z$ , such that*

$$(2.14) \quad |u^+(x, z) - x^{-\frac{2k+1}{4}} e^{-\phi_z(x)}| \leq C|x|^{-\frac{2k+1}{4}}|e^{-\phi_z(x)}|,$$

and

$$(2.15) \quad |u^{+'}(x, z) + x^{\frac{2k+1}{4}} e^{-\phi_z(x)}| \leq C|x|^{\frac{2k+1}{4}}|e^{-\phi_z(x)}|$$

for  $x \geq A(z)$ .

**Proof.** We prove only the second estimate, since the first is trivial. We write  $u$  instead of  $u^+$  for the sake of simplicity. Let  $w(x) = G(x) - u(x)$ . Then  $u' = (G - W)' = G' + \tilde{D}w - \psi'w$ . Thus

$$(2.16) \quad u' + x^{\frac{2k+1}{4}} e^{-\phi(x)} = G' + x^{\frac{2k+1}{4}} e^{-\phi(x)} + \tilde{D}w - \psi'w.$$

It is straightforward to verify that

$$|G' + x^{\frac{2k+1}{4}} e^{-\phi(x)}| \leq \theta |x|^{\frac{2k+1}{4}} |e^{-\phi(x)}|.$$

Let us consider the other two terms of the expression above. Due to (2.10) proved above and Proposition 2.1, we have that

$$\begin{aligned} |\tilde{D}w(x)| &\leq C|x|^{-\frac{3}{4}(2k+1)-\delta} |e^{-\phi(x)}| \\ &\quad + C|x|^{\frac{1}{4}(2k+1)+1-\delta} |e^{\phi(x)}| \int_x^y t^{-\frac{1}{4}(2k+1)-1} |e^{-\phi(t)} u(t)| dt \\ &\leq C|x|^{-\frac{3}{4}(2k+1)-\delta} |e^{-\phi(x)}| \\ &\quad + C|x|^{\frac{1}{4}(2k+1)+1-\delta} |e^{\phi(x)}| \int_x^y t^{-\frac{3}{2}(2k+1)-\delta} |t^{\frac{1}{2}(2k+1)} e^{-2\phi(t)}| dt \\ &\leq C|x|^{-\frac{3}{4}(2k+1)-\delta} |e^{-\phi(x)}| + C|x|^{-\frac{5}{4}(2k+1)+1-2\delta} |e^{-\phi(x)}| \\ &\leq C|x|^{-\frac{3}{4}(2k+1)-\delta} |e^{-\phi(x)}|, \end{aligned}$$

because  $\alpha > 0$ . Now the latter quantity above is

$$|x|^{-(2k+1)-\delta} |x|^{\frac{1}{4}(2k+1)}$$

and the first factor is small if  $k \geq 2$  for every  $\delta$  for which  $\alpha = \alpha(\delta) > 0$ .

Let us consider the third term in (2.16):  $\psi'w$ . Because of Proposition 2.1, we may write

$$|\psi'w(x)| \leq C|x|^{-\alpha} |x|^{\frac{1}{4}(2k+1)} |e^{-\phi(x)}|,$$

and this ends the proof of the proposition.  $\square$

An analogous estimate can be derived for any subdominant solution in its Stokes sector (see next section for more details).

We may now state the main result of this section.

**Theorem 2.1.** *Let  $k \geq 2$ . Let  $A(z) = C_0|z|^{\frac{2}{1-\delta}}$  be as above. Denote by  $W$  the Wronskian of  $u_0$  and  $u_k$ , where  $u_j$  denotes the subdominant solution in the sector of amplitude  $2/(2k+3)\pi$  and centered at the line of argument  $2j/(2k+3)\pi$ .*

Then

$$|W(z)| \leq C e^{C_1 |z|^{\frac{2k+3}{1-\delta}}}$$

for suitable positive constants  $C$  and  $C_1$ .

Proof. We argue on  $u = u_0$ . The same argument applied to the other solution allows us to achieve the proof.

First notice that

$$\begin{cases} |u(A)| \leq C A^{-\frac{2k+1}{4}} |e^{-\phi(A)}| \\ |u'(A)| \leq C A^{\frac{2k+1}{4}} |e^{-\phi(A)}| \end{cases}$$

where  $C > 0$  does not depend on  $z$ . Now define

$$E(x, z) = |z|^{2\frac{2k+1}{1-\delta}} |u(x)|^2 + |u'(x)|^2.$$

Due to the above properties we easily see that

$$\frac{d}{dx} E(x) \leq C |z|^{\frac{2k+1}{1-\delta}} E(x),$$

which implies that

$$(2.17) \quad E(0, z) \leq C E(A, z) e^{A|z|^{\frac{2k+1}{1-\delta}}}.$$

This allows us to conclude that

$$|W(z)| \leq C e^{C_1 |z|^{\frac{2k+3}{1-\delta}}}.$$

This ends the proof of the theorem.  $\square$

Since  $\alpha > 0$  implies that  $1 - \delta < \frac{2k+1}{2}$ . Thus we may always assume that  $W(z)$  is an entire function of order  $< 3$ .

**3. Proof of the Main Result.** We recall from [9] the following facts relative to equation (1.1):

- (a) the complex plane can be divided into  $2k+3$  sectors  $\mathcal{S}_j$ ,  $j = 0, \dots, 2k+2$  of amplitude  $\frac{2\pi}{2k+3}$ : to each sector one can associate a subdominant solution  $\mathcal{Y}_j(x; z)$  of the equation (i.e. a solution exponentially decreasing at  $\infty$ ).

(b)  $\forall j = 0, \dots, 2k + 3$  we have the Stokes connection formula:

$$(3.1) \quad \mathcal{Y}_j(x; z) = c_j(z)\mathcal{Y}_{j+1}(x; z) - \omega\mathcal{Y}_{j+2}(x; z),$$

where it is understood that  $\mathcal{Y}_{2k+3} = \mathcal{Y}_0$  and  $\mathcal{Y}_{2k+4} = \mathcal{Y}_1$ .

(c) Here we put  $c_j(z) = c_0(G^j(z))$ , where

$$c_0(z) = \frac{W_{0,2}(z)}{W_{1,2}(z)}$$

and  $G^j(z) = \omega^{-(k+1)j}z$ , with  $\omega = \exp \frac{2i\pi}{2k+3}$ . Moreover we recall that  $c_0(z)$  is not a constant function, since  $c'_j(0) \neq 0$ .

(d) In the above expression  $W_{0,2}(z)$  denotes the Wronskian determinant of  $\mathcal{Y}_0, \mathcal{Y}_2$  and  $W_{0,1}(z)$  that of  $\mathcal{Y}_0, \mathcal{Y}_1$ . Furthermore it is easily checked that  $W_{1,2}(z) = \omega^{\frac{2k-3}{4}}$ . On the other hand  $W_{0,2}(z)$  coincides modulo a constant with that estimated in Theorem 2.1.

(e) Define

$$S_j(z) = \begin{bmatrix} c_j(z) & 1 \\ -\omega & 0 \end{bmatrix} \quad j = 0, \dots, 2k + 2,$$

then

$$S_{2k+2}(z)S_{2k+1}(z) \cdots S_1(z)S_0(z) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

(f) The solution  $\mathcal{Y}_0(x; z)$  has in the sector  $\mathcal{S} = \left\{ x \in \mathbb{C} \mid \arg x < \frac{3}{2k+5} \right\}$  the following asymptotic development:

$$\mathcal{Y}_0(x; z) \sim x^{-\frac{2k+1}{4}} (1 + \mathcal{O}(x^{-1/2}))e^{-E(x;z)},$$

where  $E(x; z) = \frac{2}{2k+3}x^{\frac{2k+3}{2}} + zx^{\frac{1}{2}}$ .

From (3.1) we see that:

$$\mathcal{Y}_0 = A_j(z)\mathcal{Y}_j - \omega A_{j-1}(z)\mathcal{Y}_{j+1},$$

where the  $A_j(z)$  denote entire functions made up from the Stokes coefficients  $c_j$ , whose explicit expression can be given as follows:

$$(3.2) \quad A_l(z) = \sum_{j=0}^{\lfloor \frac{l}{2} \rfloor} (-\omega)^j \sum_{k_1, \dots, k_j=0}^{l-1} c_0 c_1 \cdots \hat{c}_{k_1} \hat{c}_{k_1+1} \cdots \hat{c}_{k_j} \hat{c}_{k_j+1} \cdots c_{l-2} c_{l-1},$$



where by  $\hat{c}_{k_j}$  we mean that the corresponding term has been omitted. (3.2) now yields in the special case of  $k = 1, 2$  that:

$$(3.3) \quad A_1(z) = c_0(z),$$

and that:

$$(3.4) \quad A_2(z) = c_0(z)c_1(z) - \omega.$$

The aim of this section is to prove that  $A_1(z)$  has at least one zero in  $\mathbb{C}$  when  $k = 1$  and that if  $k = 2$  the same holds for  $A_2(z)$ .

**3.1. The case  $k = 1$ .** This case has been essentially already treated in [1] and for the sake of completeness we briefly recall the argument.

From the holonomy equation in (d) we obtain the functional equation for  $A_1(z)$

$$A_1(z) + \omega^2 A_1(\omega z) A_1(\omega^4 z) - \omega^3 = 0, \quad \forall z \in \mathbb{C}.$$

Suppose that  $A_1(z) \neq 0$  for all  $z \in \mathbb{C}$ . Then from the above equation that it follows that  $A_1(z) \neq \omega^3$  for all  $z \in \mathbb{C}$ . Since  $A_1(z)$  is an entire function Picard's Little Theorem implies that  $A_1(z)$  would be constant because  $A_1(z)$  avoids two distinct values 0 and  $\omega^3$ . But this contradicts what has been stated in item (c) above.

**3.2. The case  $k = 2$ .** Here we prove that (3.4) has at least one zero in  $\mathbb{C}$ .

In fact if we assume that  $A_2(z) \neq 0 \forall z \in \mathbb{C}$  we may conclude that  $c_0(z)$  has in infinite number of zeros in  $\mathbb{C}$  due to the Borel-Hadamard lemma, see e.g. [5]. Recalling that  $c_0(0) \neq 0$  and the Hadamard representation theorem ( see e.g. [10] Theorem 5.1) we have:

$$(3.2.1) \quad c_0(z) = e^{g(z)} \prod_1^{\infty} \left( 1 - \frac{z}{a_n} \right) e^{\frac{z}{a_n} + \frac{1}{2} \left( \frac{z}{a_n} \right)^2}.$$

Here  $g(z)$  is a polynomial with degree less or equal than 2 due to Theorem 2.1. On the other hand, because of our assumption

$$A_2(z) = e^{p(z)}$$

where  $p(z)$  is another polynomial whose degree is less or equal than 2. Let us now study the equation  $A_2(z) = e^{p(z)} = -\omega$ . Taking the logarithm of both sides we get for the zeros:

$$p(z) = i\pi \frac{9}{7} + 2i\pi h, \quad h \in \mathbb{Z}$$

We argue in the case when  $p$  has degree 2. The case when the degree is 1 is similar but simpler. Let  $p(z) = \alpha z^2 + \beta z + \gamma$ , with  $\alpha \neq 0$ : from the preceding equation and the fact that, due to item c above,  $c_1(z) = c_0(\omega^4 z)$ , we obtain these two asymptotic representations for the complex numbers  $a_h, h \in \mathbb{Z}$ :

$$(3.2.2) \quad \begin{aligned} \alpha a_h^2 &\sim 2i\pi h'(h) \\ \alpha \omega^6 a_h^2 &\sim 2i\pi h''(h), \end{aligned}$$

where  $h'(h), h''(h) \in \mathbb{Z}$  and  $h'(h), h''(h) \rightarrow \infty$  if  $h$  tends to infinity. Since  $\text{Im } \omega^6 \neq 0$  it is clear from (3.2.2) that we have a contradiction. Thus we have proved the following Theorem:

**Theorem 3.2.2.** *The equation (1.1) has a bounded rapidly decreasing solution defined on the whole real line for suitable values of the complex number  $z$ .*

*Proof.* It is a straightforward consequence of the previous argument and of the fact that, due to the  $P$ - $T$  invariance properties of equation (1.1), it is always possible to choose the zero of  $A_2(z)$  in a such a way that its imaginary part is strictly negative. (See e.g. [4]).

## REFERENCES

- [1] E. BERNARDI, A. BOVE. On the Cauchy problem for some hyperbolic operator with double characteristics. In: *Phase Space Analysis of Partial Differential Equations* (Eds A. Bove et al), Birkhäuser, 2006, 29–44.
- [2] E. BERNARDI, A. BOVE. Geometric Results for a Class of Hyperbolic Operators with Double Characteristics. *Comm. Partial Differential Equations* **13** (1988), 61–86.
- [3] E. BERNARDI, A. BOVE. Remark on the Cauchy Problem for a Model Hyperbolic Operator. In: *Hyperbolic Differential Operators and Related Problems* (Eds V. Ancona, J. Gaveau) Marcel Dekker, New York, 2002, 41–52.
- [4] E. BERNARDI, A. BOVE. The Cauchy Problem is not well posed if the Hamilton field is unstable, in preparation.
- [5] E. BOREL. *Leçons sur les Fonctions Entières*, Paris, Gauthier-Villars, 1921.

- [6] M. CHRIST. Certain sums of squares of vector fields fail to be analytic hypoelliptic. *Comm. Partial Differential Equations* **16** (1991), 1695–1707.
- [7] M. CHRIST. A family of degenerate differential operators. *J. Geom. Anal.* **3** (1993), 579–597.
- [8] T. HOSHIRO. Failure of analytic hypoellipticity for some operators of  $X^2 + Y^2$  type. *J. Math. Kyoto Univ.* **35** (1993), 569–581.
- [9] Y. SIBUYA. Global Theory of a Second Order Linear Ordinary Equation with a Polynomial Coefficient. North-Holland Mathematical Studies vol. **18**, 1975.
- [10] E. M. STEIN, R. SHAKARCHI. Complex Analysis. Princeton Lectures in Analysis vol. **2**, Princeton University Press, 2003.

*Enrico Bernardi*

*Dipartimento di Matematica per le Scienze Economiche e Sociali*

*Università di Bologna*

*Viale Filopanti 5*

*40126 Bologna, Italia*

*e-mail: enrico.bernardi@unibo.it*

*Antonio Bove*

*Dipartimento di Matematica*

*Università di Bologna*

*Piazza di Porta S. Donato 5*

*40127 Bologna, Italia*

*e-mail: bove@bo.infn.it*

*Received November 19, 2007*