

Int J Clin Exp Pathol 2013;6(9):1874-1879
www.ijcep.com /ISSN:1936-2625/IJCEP1307020

Original Article

Immunohistochemical evaluation of cardiac connexin43 in rats exposed to low-frequency noise

Eduardo Antunes¹, Gonçalo Borrecho¹, Pedro Oliveira¹, José Brito¹, Artur Águas², José Martins dos Santos¹

¹Center for Interdisciplinary Research Egas Moniz, Health Sciences Institute, Monte de Caparica, Portugal;

²Department of Anatomy and UMIB of ICBAS, Abel Salazar Institute for Biomedical Sciences, University of Porto, Porto, Portugal

Received July 11, 2013; Accepted July 26, 2013; Epub August 15, 2013; Published September 1, 2013

Abstract: Introduction: Low-frequency noise (LFN) leads to an abnormal proliferation of collagen and development of tissue fibrosis. It has been shown that myocardial fibrosis in association with gap junction remodeling occurs in several cardiac diseases and can be implicated in the development of ventricular tachyarrhythmias. We previously reported a strong development of myocardial fibrosis induced by LFN in rats but it is not known whether LFN induces any modification on cardiac connexin43 (Cx43). Objectives: The aim of this study was to evaluate modifications on cardiac Cx43 induced by LFN in Wistar rats. Methods: Two groups of rats were considered: A LFN-exposed group with 10 rats submitted continuously to LFN during 3 months and a control group with 8 rats. The hearts were sectioned from the ventricular apex to the atria and the mid-ventricular fragment was selected. The immunohistochemical evaluation of Cx43 was performed using the polyclonal antibody connexin-43m diluted 1:1000 overnight at 4 °C. Quantifications of Cx43 and muscle were performed with the *image J software* and the ratio Cx43/muscle was analyzed in the left ventricle, interventricular septum and right ventricle. Results: The ratio Cx43/muscle was significantly reduced in LFN-exposed rats ($p=0.001$). The mean value decreased 46.2%, 22.2% and 55.6% respectively in the left ventricle ($p=0.008$), interventricular septum ($p=0.301$) and right ventricle ($p=0.004$). Conclusions: LFN induces modifications on cardiac Cx43 in rats. The Cx43 reduction observed in our study suggests that LFN may induce an arrhythmogenic substrate and opens a new investigational area concerning the effects of LFN on the heart.

Keywords: Low-frequency noise, connexin43, gap junction, intercalated disks, ventricular myocardium

Introduction

Low-frequency noise (LFN) leads to pathological changes in the extracellular matrix, characterized by an abnormal proliferation of collagen and the development of tissue fibrosis, in the absence of an inflammatory process [1-7]. We previously reported a significant fibrotic development in ventricular myocardium of rats exposed to LFN [8] and an increase of perivascular fibrosis in the arterial coronary vessels after exposure to industrial noise which is rich in LFN components [9], but it is not known whether LFN induces modifications on the electrophysiological *milieu*.

Gap junctions are composed by proteins known as connexins, form the intercellular pathway for electrical impulse transmission and are deter-

minants in the genesis of cardiac arrhythmias. Changes on gap junctional connexin43 (Cx43) have been implicated in ventricular remodeling and development of arrhythmias in several cardiac diseases [10-16]. Additionally, experimental studies provided evidence that a reduction of Cx43 expression is critical to increase the propensity for ventricular tachyarrhythmias [17-22].

As LFN induces the development of myocardial fibrosis [8] and perivascular fibrosis [9] and taking into account that the presence of interstitial fibrosis in association with a decrease of Cx43 seems to have arrhythmogenic consequences [23-25], the importance of quantifying this protein is crucial to establish the occurrence of a morphological arrhythmogenic substrate induced by LFN. Thus, the aim of this study was to




Figure 1. Immunostained connexin43 observed at the intercalated disks in a section taken from the left ventricle of a LFN-exposed rat (A) and control rat (B) (x 400).

evaluate modifications on cardiac Cx43 induced by LFN in Wistar rats.

Materials and methods

Eighteen adult Wistar rats from a Spanish breeder (Charles River Laboratories España, SA, Spain) were studied. The animals were treated in accordance with the EU Commission on Animal Protection for Experimental and Scientific Purposes and with the Portuguese legislation for the same purpose. Ten rats were continuously exposed to LFN for a period of three months. The control group of 8 rats was kept in a silent environment. All the animals were kept in cages, fed standard rat food and had free access to water.

The sound signal was emitted by an analog noise generator and the noise level was the same as previously reported [26].

The hearts were fixed in 10% buffered formalin, transversely sectioned from the ventricular apex to the atria and the mid-ventricular fragment was selected for the study. The samples were incubated with polyclonal antibody connexin-43m (GJA1) diluted 1:1000 overnight at 4°C for immunohistochemical analysis.

The histological images were acquired with an optical microscope using 400 x magnifications. In each section the optical fields were selected from the left ventricle, the interventricular septum and the right ventricle. Criteria used to select each field were defined by the myocardium portions containing the highest visualiza-

tion of Cx43 immunostained intercalated disks. A total of 146 optical fields were selected from all the anatomical components, by three observers, under blinded assessment, and analyzed using the *Image J* software that gives a quantification based on the image color analysis. The signal intensity threshold value of 140 on the 0 – 255 scale was identified to distinguish Cx43 from other structures. All areas with signal intensity between 0 and 140 were considered gap junctions at the intercalated disks and the following parameters were measured: 1- Cx43, 2- muscle. Then a ratio of Cx43/muscle was calculated.

Data are presented as mean \pm SD. Comparisons among groups simultaneously for the three anatomical regions were performed by One-Way MANOVA, while comparisons *in totum* were performed using a t-test for independent samples. Statistical tests were applied at the 5% level of significance.

Results

The histological observation showed immunostained Cx43 at the intercalated disks and examples of sections from LFN-exposed and control rats are shown in **Figure 1**. In general, less immunoreactive particles were observed among the samples of LFN exposed rats.

The ratio Cx43/muscle in each anatomical region is shown in **Table 1** and is graphically depicted in **Figure 2**. The ratio Cx43/muscle *in totum* is also shown in **Table 1**.

Cardiac connexin43 in rats exposed to low-frequency noise

Table 1. Ratio Cx43/muscle in each anatomical region and *in totum* in LFN-exposed (n=10) and control (n=8) animals

Anatomical region	Group	Mean	Std. Deviation	Percent decrease with LFN exposure	p
LV	exposed	0.014	0.006	46.2%	0.008
	control	0.026	0.008		
IVS	exposed	0.014	0.006	22.2%	0.301
	control	0.018	0.008		
RV	exposed	0.016	0.012	55.6%	0.004
	control	0.036	0.012		
<i>In totum</i>	exposed	0.015	0.007	43.3%	0.009
	control	0.026	0.009		

LV = Left Ventricle; IVS = Interventricular Septum; RV = Right Ventricle; LFN = Low-Frequency Noise.




Figure 2. Ratio Cx43/muscle in the left ventricle (LV), interventricular septum (IVS) and right ventricle (RV) in LFN exposed and control animals. A significant reduction was observed in exposed animals in the left ventricle ($p=0.008$) and in the right ventricle ($p=0.004$) but not in the interventricular septum ($p=0.301$).

The mean values of cardiac muscle were not significantly different between the exposed and control animals in any of the anatomical regions considered in this study ($p \geq 0.664$).

The total amount of Cx43 was significantly reduced in the left ventricle ($p=0.011$) and in the right ventricle ($p=0.009$) of LFN-exposed

animals when compared to controls. No differences were detected between the two groups concerning the total amount of Cx43 in the interventricular septum ($p=0.237$).

The ratio Cx43/muscle *in totum* decreased 43.3% among the LFN-exposed rats ($p=0.009$). Multivariate comparisons over the three anatomical regions showed significant differences between groups ($p=0.001$) with a decrease of 46.2%, 22.2% and 55.6% respectively in the left ventricle ($p=0.008$), in the interventricular septum ($p=0.301$) and in the right ventricle ($p=0.004$).

Discussion

As far as we know this is the first study concerning the evaluation of cardiac Cx43 in rats submitted to LFN.

In humans, gap junction remodeling has been studied in several pathologies and arrhythmias [10-16, 27, 28]. The loss of Cx43 expression has been shown to be a key for the development of an arrhythmic anatomic substrate in chronically hypertrophied myocardium [15]. Modifications in gap junction organization and on Cx43 expression contribute to conduction disturbances and development of arrhythmias in myocardial infarction [27], non-ischemic dilated cardiomyopathy [12, 13], cardiac heart failure [11] and valvular heart disease [14].

In several studies, a reduction of Cx43 from 30 to 50% occurs in ventricular remodeling [14, 23, 29], but changes in gap junction expression alone are presumably not sufficient for conduction slowing and enhanced arrhythmogenicity, apparently because there is a large conduction reserve [30].

Meanwhile, it is known that an increase of the intercellular collagen deposits may lead to

anisotropic reentry [23, 24], and a strong enhancement of arrhythmogenic vulnerability can be attained by the association of increased fibrosis with a 50% reduction of Cx43 expression [31].

As LFN induces the development of interstitial myocardial fibrosis [8] and perivascular fibrosis [9] we hypothesized that the finding of significant modifications on gap junctions after LFN exposure could lead to a morphological arrhythmogenic substrate.

In our study the measurement of Cx43 was performed in equivalent tissue mass among exposed and control animals and showed a significant reduction in rats exposed to LFN. The modification was evident at the free ventricular wall but not in the interventricular septum suggesting that this anatomical region could be more protected against the effects of LFN. However, this does not discard the possibility of a strong alteration of the electrophysiological *milieu*. In fact, the observed statistical powers of 99% for the multiple comparisons between groups and in excess of 80% for the comparisons between groups regarding the left and right ventricles are noteworthy, in view of the sample dimension, suggesting the marked effects of LFN exposure.

Taking into consideration the universal existence of LFN in modern societies and having in mind the difficulties to explain some ventricular tachyarrhythmias without structural heart disease, the hypothesis of idiopathic ventricular fibrillation as being also a consequence of gap junction remodeling mediated through the effects of LFN, should not be despicable. On the other hand, in patients with already known specific cardiac disorders, a reduction of gap junctions by the exposure to LFN makes the development of arrhythmic events possible, carrying out an adverse prognosis.

We still do not know the mechanisms underlying the fibrotic development we previously reported in rats exposed to LFN [8] or to industrial noise [9] as well as the mechanism of the Cx43 alteration observed in this study. Nevertheless, the Cx43 remodeling has been linked to intrinsic factors occurring on biosynthesis at transcriptional or posttranscriptional phase [32, 33]. As connexin43 is degraded through lysosomal or proteasomal pathway

[34, 35] one can also speculate that the Cx43 reduction observed in this study can be related to an activation of these pathways. Theoretically, LFN acting as an external mechanical force could also lead to activation of protein kinases which might modify the level of Cx43 phosphorylation [36, 37].

Regardless of the mechanism how LFN induces loss of Cx43, our results suggest that the significant reduction of this protein can lead to electrophysiological modifications. The occurrence of a significant myocardial and periarterial fibrosis induced respectively by LFN and industrial noise reported earlier by our group [8, 9], together with a possible gap junction remodeling observed in this study, makes the development of a morphological arrhythmogenic substrate by LFN possible. Further experimental and clinical studies are needed to evaluate the functional and arrhythmic consequences. With this study we put forward the hypothesis of a link between LFN and ventricular tachyarrhythmias, opening a new investigational area concerning the effects of LFN on the heart.

In conclusion, we can state that low frequency noise induces modifications on cardiac connexin43 in rats. The connexin43 reduction observed in our study can contribute to a morphological arrhythmogenic substrate.

Disclosure of conflict of interest

None.

Address correspondence to: Eduardo Antunes, CIEM, Centro de Investigação Interdisciplinar Egas Moniz, Instituto Superior das Ciências da Saúde Egas Moniz, ISCSEM, Quinta da Granja, Monte de Caparica, 2829-511 Caparica, Portugal. E-mail: ejpantunes@sapo.pt

References

- [1] Alves-Pereira M, Castelo Branco NAA. Vibroacoustic disease: biological effects of infrasound and low-frequency noise explained by mechanotransduction cellular signalling. *Prog Bioph Mol Biol* 2007; 93: 256-279.
- [2] da Fonseca J, dos Santos JM, Branco NC, Alves-Pereira M, Grande NR, Oliveira P, Martins AP. Noise-induced gastric lesions: a light and scanning electron microscopy study of the alterations of the rat gastric mucosa induced by

Cardiac connexin43 in rats exposed to low-frequency noise

- low frequency noise. *Cent Eur J Public Health* 2006; 14: 35-38.
- [3] Grande NR, Águas AP, Sousa Pereira A, Monteiro E, Castelo Branco NAA. Morphological changes in the rat lung parenchyma exposed to low frequency noise. *Aviat Space Environ Med* 1999; 70: A70-A77.
- [4] de Sousa Pereira A, Águas AP, Grande NR, Mirones J, Monteiro E, Castelo Branco NAA. The effect of chronic exposure to low frequency noise on rat tracheal epithelia. *Aviat Space Environ Med* 1999; 70: A186-A190.
- [5] Martins dos Santos J, Grande NR, Castelo Branco NAA, Zagalo C, Oliveira P, Alves-Pereira M. Lymphatic lesions and vibroacoustic disease. *Eur J Lymphol* 2004; 12: 17-20.
- [6] Oliveira PM, Pereira da Mata AD, Martins dos Santos JA, da Silva Marques DN, Branco NC, Silveira JM, Correia da Fonseca JC. Low-frequency noise effects on the parotid gland of the Wistar rat. *Oral Dis* 2007; 13: 468-73.
- [7] Martins dos Santos J, Grande NR, Castelo Branco NA, Zagalo C, Oliveira P. Vascular lesions and vibroacoustic disease. *Eur J Anat* 2002; 6: 17-21.
- [8] Antunes E, Borrecho G, Oliveira P, Oliveira MJR, Brito J, Águas A, Martins dos Santos J. Myocardial Fibrosis in Rats Exposed to Low Frequency Noise. *Acta Cardiol* 2013; 68: 241-245.
- [9] Antunes E, Oliveira P, Oliveira MJR, Brito J, Águas A, Martins dos Santos J. Histomorphometric evaluation of the coronary artery vessels in rats submitted to industrial noise. *Acta Cardiol* 2013; 68: 285-289.
- [10] Smith JH, Green CR, Peters NS, Rothery S, Severs NJ. Altered Patterns of gap junction distribution in ischemic heart disease. An immunohistochemical study of human myocardium using laser scanning confocal microscopy. *Am J Pathol* 1991; 139: 871-878.
- [11] Kostin S, Rieger M, Dammer S, Hein S, Richter M, Klovekorn WP, Bauer EP, Schaper J. Gap junction remodeling and altered connexin43 expression in the failing human heart. *Mol Cell Biochem* 2003; 242: 135-144.
- [12] Kitamura H, Ohnishi Y, Yoshida A, Okajima K, Azumi H, Ishida A, Galeano EJ, Kubo S, Hayashi Y, Itoh H, Yokoyama M. Heterogeneous loss of connexin43 protein in nonischemic dilated cardiomyopathy with ventricular tachycardia. *J Cardiovasc Electrophysiol* 2002; 13: 865-887.
- [13] Kitamura H, Ohnishi Y, Yoshida A, Okajima K, Azumi H, Ishida A, Galeano EJ, Kubo S, Fukuzawa K, Takano T, Yokoyama M. Correlation of connexin43 expression and late ventricular potentials in nonischemic dilated cardiomyopathy. *Circ J* 2003; 67: 1017-1021.
- [14] Kostin S, Dammer S, Hein S, Klovekorn WP, Bauer EP, Schaper J. Connexin 43 expression and distribution in compensated and decompensated cardiac hypertrophy in patients with aortic stenosis. *Cardiovasc Res* 2004; 62: 426-436.
- [15] Peters NS, Green CR, Poole-Wilson PA, Severs NJ. Reduced content of connexin43 gap junctions in ventricular myocardium from hypertrophied and ischemic human hearts. *Circulation* 1993; 88: 864-875.
- [16] Takeuchi S, Akita T, Takagishi Y, Watanabe E, Sasano C, Honjo H, Kodama I. Disorganization of gap junction distribution in dilated atria of patients with chronic atrial fibrillation. *Circ J* 2006; 70: 575-582.
- [17] Gutstein DE, Morley GE, Tammaddon H, Vaidya D, Schneider MD, Chen J, Chien KR, Stuhlmann H, Fishman GI. Conduction slowing and sudden arrhythmic death in mice with cardiac-restricted inactivation of connexin43. *Circ Res* 2001; 88: 333-339.
- [18] Danik SB, Liu F, Zhang J, Suk HJ, Morley GE, Fishman GI, Gutstein DE. Modulation of cardiac gap junction expression and arrhythmic susceptibility. *Circ Res* 2004; 95: 1035-1041.
- [19] van Rijen HVM, Eckardt D, Degen J, Theis M, Ott T, Willecke K, Jonsma HJ, Opthof T, Bakker JMT. Slow conduction and enhanced anisotropy increase the propensity for ventricular tachyarrhythmias in adult mice with induced deletion of connexin43. *Circulation* 2004; 109: 1048-1055.
- [20] Uzzaman M, Honjo H, Tagagishi Y, Emdad L, Magee AI, Severs NJ, Kodama I. Remodeling of gap junctional coupling in hypertrophied right ventricles of rats with monocrotaline-induced pulmonary hypertension. *Circ Res* 2000; 86: 871-879.
- [21] Mayama T, Matsumura K, Lin H, Ogawa K, Imanaga I. Remodeling of cardiac gap junction connexin 43 and arrhythmogenesis. *Exp Clin Cardiol* 2007; 12: 67-76.
- [22] Lin H, Ogawa K, Imanaga I, Tribulova N. Remodeling of connexin 43 in the diabetic rat heart. *Mol Cell Biochem* 2006; 290: 69-78.
- [23] Peters NS, Coromilas J, Severs NJ, Wit AL. Disturbed connexin43 gap junction distribution correlates with the location of reentrant circuits in the epicardial border zone of healing canine infarcts that cause ventricular tachycardia. *Circulation* 1997; 97: 988-996.
- [24] Kawara T, Derksen R, de Groot JR, Coronel R, Tasseron S, Linnenbank AC, Hauer RN, Kirkels H, Janse MJ, de Bakker JM. Activation delay after premature stimulation in chronically diseased human myocardium relates to the architecture of interstitial fibrosis. *Circulation* 2001; 104: 3069-3075.
- [25] La Vecchia L, Ometto R, Bedogni F, Finocchi G, Mosele GM, Bozzola L, Bevilacqua P, Vincenzi

Cardiac connexin43 in rats exposed to low-frequency noise

- M. Ventricular late potentials, interstitial fibrosis, and right ventricular function in patients with ventricular tachycardia and normal left ventricular function. *Am J Cardiol* 1998; 81: 790-792.
- [26] da Fonseca J, Martins dos Santos J, Castelo-Branco N, Alves-Pereira M, Grande N, Oliveira P. Noise-induced duodenal lesions. *Eur J Anat* 2005; 9: 29-33.
- [27] Kieken F, Mutsaers N, Dolmotova E, Virgil K, Wit AL, Kellezi A, Hirst-Jensen BJ, Duffy HS, Sorgen PL. Structural and molecular mechanisms of gap junction remodeling in epicardial border zone myocytes following myocardial infarction. *Circ Res* 2009; 104: 1103-1112.
- [28] Basso C, Czarnowska E, Barbera ML, Bauce B, Beffagna G, Wlodarska EK, Pilichou K, Ramondo A, Lorenzon A, Wozniak O, Corrado D, Daliento L, Danieli GA, Valente M, Nava A, Thiene G, Rampazzo A. Ultrastructural evidence of intercalated disc remodeling in arrhythmogenic right ventricular cardiomyopathy: an electron microscopy investigation on endomyocardial biopsies. *Eur Heart J* 2006; 27: 1847-1854.
- [29] Kaprielian RR, Gunning M, Dupont E, Shepard MN, Rothery SM, Underwood R, Pennell DJ, Fox K, Pepper J, Poole-Wilson PA, Severs NJ. Downregulation of immunodetectable connexin43 and decreased gap junction size in the pathogenesis of chronic hibernation in the human left ventricle. *Circulation* 1998; 97: 651-660.
- [30] van Rijen HV, de Bakker JM, van Veen TA. Hypoxia, electrical uncoupling, and conduction slowing: Role of conduction reserve. *Cardiovasc Res* 2005; 66: 9-11.
- [31] Jansen JA, van Veen AA, Bosch AA, van der Nagel R, Vos MA, Bakker JM, van Rijen HV. Arrhythmia vulnerability of aged haploinsufficient Cx43 mice is determinant by heterogeneous downregulation of Cx43 combined with increased fibrosis. *Circulation* 2008; 118: S494.
- [32] Teunissen BE, Bierhuisen MF. Transcriptional control of myocardial connexins. *Cardiovasc Res* 2004; 62: 246-255.
- [33] Yang B, Lin H, Xiao J, Lu Y, Luo X, Li BT, Zhang Y, Xu C, Bai Y, Wang H, Chen G, Wang Z. The muscle-specific micro-RNA miR-1 regulates cardiac arrhythmogenic potential by targeting GJA1 and KCNJ2. *Nat Med* 2007; 13: 486-491.
- [34] Laing JG, Tadros PN, Green K, Saffitz JE, Beyer EC. Proteolysis of connexin43-containing gap junctions in normal and heat-stressed cardiac myocytes. *Cardiovasc Res* 1998; 38: 711-718.
- [35] Berthoud VM, Minogue PJ, Laing JG, Beyer EC. Pathways for degradation of connexins and gap junctions. *Cardiovasc Res* 2004; 62: 256-267.
- [36] Saffitz JE, Kleber AG. Effects of mechanical forces and mediators of hypertrophy on remodeling of gap junction in the heart. *Circ Res* 2004; 94: 585-591.
- [37] Yamada K, Green KG, Samarel AM, Saffitz JE. Distinct pathways regulate expression of cardiac electrical and mechanical junction proteins in response to stretch. *Circ Res* 2005; 97: 346-353.