

CLINICAL INQUIRIES

Evidence-based answers from the
Family Physicians Inquiries Network

Geetha Chilakamarri, MD
Sparrow/Michigan State
University Family Medicine
Residency Program, Lansing

**David Weismantel,
MD, MS**
Department of Family
Medicine, Michigan State
University, East Lansing

Arlene Weismantel, MLIS
Michigan State University
Libraries, East Lansing



**In terms of
augmenting
antidepressants,
T₃ may have a
slight advantage
over lithium
because it has
fewer side
effects.**

Q/ Can nonantidepressants help treat depression?

EVIDENCE-BASED ANSWER

A/ YES, LITHIUM, TRIIODOTHYRONINE (T₃), AND ATYPICAL ANTIPSYCHOTICS are all effective adjuncts. Lithium (serum levels >0.5 mEq/L) can produce clinical improvement when added to ineffective antidepressant treatment (strength of recommendation [SOR]: **A**, meta-analysis of randomized controlled trials [RCTs]).

Thyroid supplementation using T₃ at doses no higher than 50 mcg per day also increases the effectiveness of antidepressant therapy (SOR: **B**, meta-analysis of RCT and cohort studies).

Atypical antipsychotic agents are less effective adjuncts for patients with treatment-resistant major depressive disorder (SOR: **B**, meta-analysis of RCT and cohort studies).

1.80-5.37), yielding a number needed to treat (NNT) of 5.³ The mean response rate was 41.2% in the lithium group and 14.4% in the placebo group ($P < .001$). The meta-analysis included only RCTs that enrolled subjects with unipolar or bipolar disorder (depressive phase) who were treated with any antidepressant plus lithium in any dose compared with placebo.

A previous meta-analysis published in 1999 concluded that a lithium

Evidence summary

As many as 30% of patients with major depression fail to respond to treatment with a single antidepressant drug given in adequate dosage for an appropriate period.¹ Pharmacologic approaches such as switching antidepressant classes are often attempted first, followed by augmentation with another agent if needed.

The most widely studied medications used for augmentation are lithium and T₃.² An important limitation to their use is that most of the supporting evidence comes from studies of patients who didn't respond initially to tricyclic antidepressants.

Lithium boosts response to antidepressants

A 2007 meta-analysis of 10 augmentation studies reported that adding lithium to various antidepressant agents increased the chances of clinical response 3-fold relative to placebo (odds ratio [OR]=3.11; 95% confidence interval [CI],

dose sufficient to produce serum levels of at least 0.5 mEq/L and a minimum treatment duration of 2 weeks resulted in a pooled OR of response to lithium augmentation compared with placebo of 3.31 (95% CI, 1.46-7.53).⁴ Lithium augmentation is a reasonable alternative for depressed patients who don't respond to conventional antidepressants.

T₃ works well with tricyclics, but what about newer antidepressants?

A 1996 meta-analysis of 8 studies with a total of 292 patients found that patients who received T₃ augmentation of tricyclic antidepressant therapy were twice as likely to respond as controls (relative response=2.09; 95% CI, 1.31-3.32).⁵ The corresponding pooled absolute difference in response rate was 23.2% with a corresponding NNT of 4.3.

T₃ dosage in the studies ranged from 25 to 50 mcg/day and duration varied from 7 to 35 days. Analysis of data from RCTs alone produced a lower pooled relative response of 1.53

(95% CI, 0.70-3.35) and an NNT of 12.5. A major drawback of T₃ augmentation is that little information is available about its efficacy in combination with newer antidepressant agents.

T₃ may have a slight edge over lithium in side effects

As a part of the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study, lithium and T₃ augmentation were compared directly in patients who had failed 2 medication treatments for depression. A total of 142 adult outpatients with nonpsychotic major depressive disorder who had not achieved remission with (or who were intolerant to) citalopram, and who similarly had no luck with either a therapeutic switch or an augmentation trial, were randomly assigned to lithium or T₃ augmentation for as long as 14 weeks.

After a mean of 9.6 weeks of treatment, remission rates were 15.9% with lithium and 24.7% with T₃, although the difference between the 2 drugs wasn't statistically significant. Because T₃ has fewer side effects and is easier to use, the researchers suggested that it may have a slight advantage over lithium.

Atypical antipsychotic agents are another alternative

A 2007 meta-analysis of 10 clinical trials involving 1500 outpatients studied the efficacy of augmenting standard antidepressants with atypical antipsychotic agents to treat resistant major depressive disorder.⁶ Across the trials, the pooled risk ratio (RR) for remission was 1.75 (95% CI, 1.36-2.24) and for response rates was 1.35 (95% CI, 1.13-1.63). Pooled rates for

remission and response were 47.4% vs 22.3% (NNT=4) and 57.2% vs 35.4% (NNT=4.6), respectively.

Although the meta-analysis found no difference in overall discontinuation rates ($P=.929$), the rate of discontinuation because of adverse events was lower among placebo-treated patients (RR=3.38, $P<.0001$). These results suggest a role for atypical antipsychotic agents in augmenting standard antidepressants for treatment-resistant major depressive disorder.

Recommendations

Both the American Psychiatric Association⁷ and the Institute for Clinical Systems Improvement⁸ recommend considering a trial of lithium or thyroid augmentation for patients who respond only partially to initial antidepressant therapy. Many experienced clinicians consider lithium to be the most effective adjunct. The APA cautions, however, that its Major Depressive Disorder Practice Guideline, published in 2000, is no longer current.⁹ An update is expected by the end of the year.

Other options include maximizing the initial treatment, switching to another agent, or augmenting initial treatment with another antidepressant agent or psychotherapy. Before prescribing any additional treatments for patients who fail to respond to initial antidepressant therapy, however, primary care physicians need to be mindful of the fact that empirical data regarding the relative effectiveness of these strategies are limited—and that they should consider whether the patients they're treating have bipolar—rather than unipolar—depression. **JFP**

References

1. Cowen PJ. New drugs, old problems. Revisiting pharmacological management of treatment resistant-depression. *Adv Psychiatr Treat.* 2005;11:19-27.
2. Nierenberg AA, Fava M, Trivedi MH, et al. A comparison of lithium and T₃ augmentation following two failed medication treatments for depression: a STAR*D report. *Am J Psychiatry.* 2006;163:1519-1530.
3. Crossley NA, Bauer M. Acceleration and augmentation of antidepressants with lithium for depressive disorders: two meta-analyses of randomized, placebo-controlled trials. *J Clin Psychiatry.* 2007;68:935-940.
4. Bauer M, Dopfmer S. Lithium augmentation in treatment-resistant depression: meta-analysis of placebo-controlled studies. *J Clin Psychopharmacol.* 1999;19:427-434.
5. Aronson R, Offman HJ, Joffe RT, et al. Triiodothyronine augmentation in the treatment of refractory depression. A meta-analysis. *Arch Gen Psychiatry.* 1996;53:842-848.
6. Papakostas GI, Shelton RC, Smith J, et al. Augmentation of antidepressants with atypical antipsychotic medications for treatment-resistant major depressive disorder: a meta-analysis. *J Clin Psychiatry.* 2007;68:826-831.
7. Practice guideline for the treatment of patients with major depressive disorder (revision). American Psychiatric Association. *Am J Psychiatry.* 2000;157(4 suppl):S1-S45.
8. Institute for Clinical Systems Improvement. Depression, Major, in Adults in Primary Care. Bloomington, Minn: Institute for Clinical Systems Improvement; 2009. Available at: www.icsi.org/guidelines_and_more/gl_os_prot/behavioral_health/depression_5/depression_major_in_adults_in_primary_care_4.html. Accessed July 14, 2009.
9. American Psychiatric Association. APA Practice Guidelines. Major Depressive Disorder. Arlington, Va: American Psychiatric Publishing, Inc. Available at: http://www.psychiatryonline.com/pracGuide/pracGuideChapTop_7.aspx. Accessed September 16, 2009.

 **JOIN THE DISCUSSION**

What agent have you had the most success with when antidepressant therapy required augmentation?

Tell us @ journaloffamilypractice.blogspot.com