

FUNCTIONAL VOMITING PATTERNS AND DISORDERS IN INFANTS, CHILDREN AND ADOLESCENTS

In infants: “*Innocent Vomiting*” (a.k.a. “Functional GER”)

“*Nervous Vomiting*”

“*Infant Rumination Syndrome*”

Rumination in the developmentally disabled

Adult-type Rumination

Gag-induced vomiting: *Tussive Vomiting*

Vomiting due to aberrant oral phase of swallowing

Anticipatory Nausea and Vomiting

Vomiting associated with rage: (Overt) *Vomiting during tantrums*

(less overt) *Hill’s “Psychogenic Vomiting”*

“*Contentious Vomiting*” (self-induced for manipulative purpose during an altercation)

Vomiting as an act of malingering

Vomiting triggered by disgust or revulsion

Oral-Defensive Vomiting

Vomiting associated with panic attacks

Chronic Nausea with little or no vomiting

Cyclic Vomiting Syndrome

Vomiting as a symptom of Conversion Disorder

Vomiting associated with Somatization Disorder

Self-induced vomiting other than “Contentious Vomiting” and Malingered Vomiting:

Bulimia, Bulimarexia

Factitious Vomiting

David R. Fleisher, MD
Associate Professor, Department of Child Health
University of Missouri
School of Medicine
Columbia, Missouri, USA

FleisherD@health.missouri.edu