Public Abstract First Name:Sarah Middle Name:Kathleen Last Name:Mobley Adviser's First Name:Pam Adviser's Last Name:hinton Co-Adviser's First Name: Co-Adviser's Last Name: Graduation Term:SS 2010 Department:Exercise Physiology Degree:MA

Title:Effects of Isometric Exercise on the Relationship Between Muscle Pain and Resting Blood Pressure and Resting Heart Rate

Silent myocardial ischemia is a common phenomenon in patients with coronary heart disease (CHD). This may be related to findings that high BP is associated with suppression of chest pain during episodes of myocardial ischemia. Purpose: The primary objective of this study was to examine the relationship between isometric exercise-induced muscle pain and resting blood pressure (BP) and heart rate (HR). Five exploratory objectives related to the effects of eccentric exercise pre- and two days post-exercise were also examined. Methods: Thirty-two participated this randomized controlled trial, which included two exercise sessions. In the first session baseline measurements of blood pressure, HR, cortisol and pain ratings were taken. Tonic and phasic isometric muscle contraction tests were performed following all of the baseline measurements. After the isometric muscle contraction tests, participants (n=16) performed three sets of 12 eccentric muscle contractions. All of the measurements and isometric muscle contraction tests were completed again, two days post-eccentric exercise. Results: Isometric exercise-induced pain and resting BP and HR was not found to be significantly or negatively correlated. None of the relationships examined in the exploratory objectives were significant from pre- to post-eccentric exercise. Salivary cortisol concentration did not significantly change pre- to post-eccentric exercise. Conclusion: A negative relationship between exercise-induced pain and resting BP and HR was not detected, which may be due to several factors, such as the type of pain induced during exercise or the timing of the BP and HR measurement.