

Psychosis in the Elderly

Background

- Definition:
 - Psychosis is a common psychiatric symptom characterized by hallucinations and delusions²
- 2. General information:
 - Associated with aggressive or disruptive behavior
 - Source of great distress to caregivers
 - Can result in neglect and abuse of elderly patients
 - Persistent symptoms may lead to institutionalization

Pathophysiology

1. Causes of psychosis can include:
 - Pharmacological
 - Amphetamines, cocaine, PCP, methylphenidate
 - Psychiatric
 - Mania, major depression, schizophrenia
 - Infectious disease
 - Sepsis
 - Behavioral/environmental circumstances
 - Substance abuse
 - Degenerative diseases
 - Alzheimer's, Parkinson's dz
 - Organic disease
 - Brain damage
 - Vitamin deficiencies
 - B3 and B12
 - Alcohol withdrawal
 - Neurological carcinomas
 - Any space occupying mass/lesion
2. Incidence, prevalence:
 - Prevalence
 - 2-4.75% in community dwelling elderly
 - 10-65% in nursing home populations
 - Prevalence of common causes of psychosis
 - Schizophrenia
 - 0.6% of US population ages 45-65
 - 0.2% of 65+ population
 - Women at greater risk of late onset schizophrenia
 - Delirium^{3,4}
 - Up to 60% in >70 yo nursing home residents
 - 14-56% of elderly, hospitalized patients
 - Polypharmacy is a major cause of psychosis in elderly⁵
 - Incidence 5.7 per 100
 - Prevalence 6.1 per 100⁶
 - Alzheimer's disease
 - Prevalence 33% >85 yo

- Major depression with psychotic features
 - 0.4% of elderly
 - Bipolar disorder with psychotic features
 - 0.1-0.4% of adults >65 yo
3. Risk factors:
- Schizophrenia
 - Alzheimer's dz
 - Parkinson's dz
 - AIDS
 - Cognitive decline/Dementia not otherwise specified
 - Diabetes (hypoglycemic episodes of psychosis)
 - ICU admission is an independent risk factor for delirium in older population⁷
 - Polypharmacy
 - Age-related pharmacokinetic and pharmacodynamic changes
 - Biopsychosocial risk factors:
 - Female
 - Low socioeconomic status
 - Presence of sensory or perceptual deficits
 - Immigrant status
 - Social isolation
4. Morbidity / mortality:
- Psychiatric illness has significant negative impact on survival of elders⁸

Diagnostics

1. History:
- Gather history from patient, caretakers, medical staff
 - Cognitive impairment may decrease accuracy
 - Review medication list (including OTC, herbals)
 - Substance use history
 - Elicit previous psychiatric history
 - Assess suicidality
 - Assess homicidality
2. Physical examination:
- Vital signs
 - Fever could be a sign of delirium
 - Thorough neurological exam and psychological exams
 - Speech: rate, rhythm, volume
 - Type and quality of hallucinations/illusions (may be concealed by patient)
 - Thought processes:
 - Goal directed vs. derailed
 - Thought content:
 - Delusions, overvalued ideas, paranoia
 - Insight/judgment
3. Screening tests
- Folstein Mini-Mental State Examination
 - Reliably differentiates between dementia/delirium (MMSE <23) and psychiatric disorders (MMSE >24)
 - Delirium Rating Scale

- Confusion Assessment Method
 - Sensitivity of 93-100% for diagnosis of delirium
 - Specificity of 90-95% (delirium)
 - Diagnosis of delirium requires presence of both features 1 and 2, plus either feature 3 or 4:
 - Acute onset and fluctuating course
 - Inattention, distractibility
 - Disorganized thinking, illogical or unclear ideas
 - Alteration in consciousness
- 4. Diagnostic testing
 - Laboratory evaluation
 - Chemistry panel
 - CBC
 - UA
 - Chest film
 - Toxicology screen
 - HIV and VDRL in high risk cases
- 5. Diagnostic imaging
 - Radiologic imaging:
 - Only if organic neurologic disorder or stroke is suspected
 - Head CT:
 - When no other cause can be found
- 6. Other studies
 - Lumbar puncture:
 - If patient has meningeal signs
- 7. Diagnostic criteria
 - Symptoms of psychosis vary by cause, but generally include some of the following:
 - Hallucinations (in any modality)
 - Delusions or diminished sense of reality
 - Disorganized speech
 - Disorganized and/or catatonic behavior

Differential Diagnosis

1. Key DDx: see Table

- Dementia (37% of psychosis in elderly)
- Affective disorder (major depression or bipolar disorder; 20% of psychosis in elderly)
 - Manic episode of Bipolar Disorder
 - Major Depression, severe
 - Mood congruent psychotic features
 - Mood incongruent psychotic features
- Delirium (12% of psychosis in elderly)
- Schizophrenia (0.1-0.5% of elderly population)
 - Early and late onset
 - 4% of schizophrenics very late onset (>age 65)

2. Extensive DDx

- Neurological
 - Alzheimer's disease
 - Carcinomas
 - Neurological damage
 - Temporal Lobe Epilepsy
 - Multiple Sclerosis
 - Stroke
 - Toxins/Intoxications (marijuana)
- Psychological
 - Delusional Disorders
 - Post Traumatic Stress Disorder
 - Substance-abuse Disorders
- Infectious
 - Lyme disease
 - Neurosyphilis
 - AIDS Dementia
 - Leprosy
 - Malaria
- Medication use/withdrawal¹⁵
 - Benzodiazepines
 - Antihistamines
 - Anti-Parkinsonian drugs
 - Anti-arrhythmics
 - Anti-inflammatory drugs
 - Anticonvulsants
 - Steroids
 - Anticancer agents
 - Opiates
 - Digoxin
 - Calcium channel blockers
 - SSRI (serotonin syndrome)
- Other
 - Sleep deprivation
 - Vitamin B12 deficiency
 - Sarcoidosis
 - Parkinson's
 - Lupus
 - Electrolyte abnormalities
 - Dehydration
 - Hypoglycemia
 - Hypoxemia
 - Poisons
 - Hypertensive encephalopathy

Therapeutics

1. Acute treatment

- Standard treatment for acute psychotic agitation
 - IM lorazepam and haloperidol¹⁶
 - Elderly have increased sensitivity to antipsychotics
 - Start low, titrate slowly
 - Scheduled dosing preferred over PRN
 - Efficacy: conventional antipsychotics = atypical antipsychotics¹⁷
 - Atypical antipsychotics < adverse effects
 - Now considered to be 1st line treatment in elderly
 - But Black box warning
 - "Elderly patients with dementia-related psychosis treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo"¹⁸
 - Possible weight gain, hyperglycemia, diabetes
 - Antipsychotics associated with increased risk of:
 - Venous thromboembolism
 - Sudden cardiac death
 - Neuroleptic malignant syndrome
 - Dystonias, akathisia
 - Agranulocytosis
- Major depression
 - Antidepressant plus antipsychotic
 - Consider ECT
- Bipolar disorder
 - Lithium
 - Lower dose if impaired renal clearance, cardiac disease, or diuretic treatment
 - Divalproex

2. Further management (24 hrs):

- Depends on etiology
- Seizures-secondary to alcohol withdrawal
 - Benzodiazepines to prevent delirium tremens
- Hypoglycemia, especially in diabetic patients
- Monitor electrolytes
- Monitor temperature
- Volume status
- In case of delirium
 - Keep pts oriented to date, time and place
 - Frequent monitoring
 - Rooms with windows help to keep sleep-wake cycle on track^{19,20}
 - Stable staffing helps

3. Long-term care

- Varies by cause of psychosis:
- Psychotic depression:
 - Maintain antipsychotic 6 months²¹
- Schizophrenia:
 - Referral to psychiatrist²²

- Hypoglycemia:
 - Improve glucose management
- Stroke:
 - Physical and occupational therapy
- Intoxication:
 - Substance abuse counseling
- Alzheimer's and Parkinson's:
 - Antipsychotic drugs associated with higher mortality in patients with dementia
 - Mortality: Conventional antipsychotics > atypical antipsychotics in elderly

Follow-Up

1. Return to office
 - Time frame for return visit
 - After acute episode of psychosis, patient should have follow-up with their PCP within a week, then regular visits Q monthly until stable on medication
 - Recommendations for earlier follow-up- this depends on the cause of episode
2. Refer to specialist
 - Recommendations / urgency
 - If episode was due to psychiatric problem, psychiatric evaluation is urgent
 - Assessment by neurologist may be necessary if patient had stroke
3. Admit to hospital
 - Recommendations / urgency
 - Depending on situation, acute episode of psychosis may warrant hospital admission and workup
 - Patients in nursing home or hospice setting may not need hospitalization

Prognosis

1. Early onset schizophrenia²³
 - Positive symptoms tend to decrease in severity with aging
 - Negative symptoms may become more severe with advancing age
 - Social and functional decline positively correlate with negative symptoms
 - Greater risk of cognitive deficits
 - Treatment resistance common
 - Positive symptoms respond to antipsychotics, sometimes partially
2. Late-onset schizophrenia
 - Positive symptoms prominent
 - Negative symptoms & formal thought disorder rare
 - Usually respond well to antipsychotic medications
3. Major Depression with psychotic features²⁴
 - Only 10% "good outcome" at 1 year
 - Likely to require readmission within 5 years
 - 5x greater risk of suicide than non-psychotic depressed elders

- 2x greater risk of mortality than non-psychotic depressed elders (after controlling for health status)
4. Bipolar disorder with psychotic features
- Greater suicide risk
 - High treatment resistance
 - Constant risk of manic episodes throughout lifespan

Prevention

1. It is crucial to obtain a thorough psychiatric history on every new patient
2. Evaluating your patient's medication list will help prevent complications due to polypharmacy

Patient Education

1. Psychosis explained (from government of Australia)
http://www.betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Psychosis_explained?OpenDocument
2. Antipsychotic medications: (from government of Australia)
http://www.betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Antipsychotic_medications_explained?open
3. Delirium and Dementia (Merk Home Manual)
<http://www.merck.com/mmhe/sec06/ch083/ch083a.html?qt=delirium%20&alt=s>
h

References

1. McGlade N, Behan C, Hayden J, O'Donoghue T, Peel R, Haq F, Gill M, Corvin A, O'Callaghan E, Donovan G: Mental state decoding v. mental state reasoning as a mediator between cognitive and social function in psychosis. *British Journal of Psychiatry* 2008;193:77-78. <http://bjp.rcpsych.org/cgi/content/full/193/1/77>
2. Broadway J, Mintzer J: The many faces of psychosis in the elderly. *Current Opinion in Psychiatry* 2007;20(6):551-558. <http://journals.lww.com/co-psychiatry/pages/articleviewer.aspx?year=2007&issue=11000&article=00006&type=abstract>
3. Fann JR: The epidemiology of delirium: a review of studies and methodological issues. *Semin Clin Neuropsychiatry* 2000;5(2):64-74.
http://www.ncbi.nlm.nih.gov/pubmed/10837095?ordinalpos=29&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum
4. Inouye S, Viscoli C, Horwitz R, Hurst L, Tinetti M: A Predictive Model for Delirium in Hospitalized Elderly Medical Patients Based on Admission Characteristics. *Annals of Internal Medicine* Sept 1993;119(6):474-481.
<http://www.annals.org/cgi/content/full/119/6/474>
5. Veehof L, Stewart B, Meyboom-de Jong B, Haaijer-Ruskamp F: Adverse drug reactions and polypharmacy in the elderly in general practice. *European Journal of Clinical Pharmacology* 1999;55(7):533-536.
http://www.ncbi.nlm.nih.gov/pubmed/10501824?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum
6. Voaklander D, Rowe B, Dryden D, Pahal J, Saar P, Kelly K. Medical illness, medication use and suicide in seniors: a population-based case control study. *J*

- Epidemiology and Community Health 2008; 62:138-146.
<http://jech.bmj.com/cgi/content/full/62/2/138>
7. McNicoll, L, Pisani, MA, Zhang, Y, et al. Delirium in the intensive care unit: occurrence and clinical course in older patients. *J Am Geriatr Soc* 2003;51(5):591-598.
<http://www3.interscience.wiley.com/journal/118861915/abstract?CRETRY=1&;SRETRY=0>
 8. Zubenko G, Mulsant B, Sweet R, Pasternak R, Tu X: Mortality of elderly patients with psychiatric disorders. *Am J Psychiatry* 1997; 154:1360-1368.
<http://ajp.psychiatryonline.org/cgi/content/abstract/154/10/1360>
 9. Bunney EB. Altered mental status in the elderly: Neurologic nightmares. American College of Emergency Physicians Scientific Assembly, Chicago, IL. October, 2008.
 10. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth edition - Text Revision. 2000. Washington, DC: American Psychiatric Association. <http://allpsych.com/disorders/psychotic/index.html>
 11. Sharma V, Mazmanian D. Sleep loss and postpartum psychosis. *Bipolar Disorders* 2003; 5(2):98-105.
<http://www3.interscience.wiley.com/journal/118852722/abstract>
 12. Bona JR, Fackler SM, Fendley MJ, Nemeroff CB: Neurosarcoidosis as a cause of refractory psychosis: A complicated case report. *American Journal of Psychiatry* August 1998; 155:1106-1108.
<http://ajp.psychiatryonline.org/cgi/content/full/155/8/1106>
 13. Fallon B, Niels J. Lyme disease: a neuropsychiatric illness. *American Journal of Psychiatry* 1994; 151(11):1571-1583.
<http://ajp.psychiatryonline.org/cgi/reprint/151/11/1571>
 14. Kararizou E, Mitsonis C, Dimopoulos N, Gkiatas K, et al. Psychosis or simply a new manifestation of neurosyphilis? *J. Int. Med. Res*;2006, 34(3): 335-337.
<http://www.ingentaconnect.com/content/field/jimr/2006/00000034/00000003/art00014?token=004e1b416d5f264e0dc7e2a46762c47655d23703b706a2356457c673f7b2f27375f2a72752d7>
 15. Karim S, Byrne E: Treatment of psychosis in elderly people. *Advances in Psychiatric Treatment* 2005; 11(4):286-296.
<http://apt.rcpsych.org/cgi/reprint/11/4/286?maxtoshow=&;HITS=10&hits=10&RESULTFORMAT=&fulltext=karim+s&searchid=1&FIRSTINDEX=0&resourcetype=HWCIT>
 16. Currier G, Chou J, Feifel D, Bossie C, Turkoz I, Mahmoud R, Gharabawi G. Acute treatment of psychotic agitation: a randomized comparison of oral treatment with haloperidol and lorazepam. *J Clin Psychiatry* 2004; 65(3):386-394.
http://www.ncbi.nlm.nih.gov/pubmed/15096079?ordinalpos=20&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum
 17. Kindermann SS, Dolder CR, Bailey A, Katz IR, Jeste DV. Pharmacological treatment of psychosis and agitation in elderly patients with dementia: four decades of experience. *Drugs and Aging* 2002, 19(4): 257-276.
http://www.ncbi.nlm.nih.gov/pubmed/12038878?ordinalpos=7&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum

18. Cascade E, Kalali AH, Cummings JL. Use of atypical antipsychotics in the elderly. *Psychiatry* 2008;5(7):28-31. <http://www.psychiatrymmc.com/use-of-atypical-antipsychotics-in-the-elderly/>
19. Swigart S, Yasuhiro K, Thurber S, Kathol R, Meller W. Misdiagnosed delirium in patient referrals to a university based hospital psychiatry department. *Psychosomatics* 2008; 49:104-108. <http://psy.psychiatryonline.org/cgi/content/full/49/2/104>
20. Inouye S, Bogardus S, Charpentier P, Leo-Summers D, et al. A multicomponent intervention to prevent delirium in hospitalized older patients. *N Engl J Med* 1999; 340(9):669-676. <http://content.nejm.org/cgi/content/abstract/340/9/669>
21. Alexopoulos GS, Katz IR, Reynolds CF 3rd, Carpenter D, Docherty JP, Ross RW. Pharmacotherapy of depression in older patients: a summary of the expert consensus guidelines. *J Psychiatr Pract.* 2001 Nov;7(6):361-76. http://www.ncbi.nlm.nih.gov/pubmed/15990550?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum
22. Cohen C, Vahia I, Reyes P, Diwan S, Bankole A, Palekar N, Kehn M, Ramirez P: Focus on geriatric psychiatry: Schizophrenia in Later Life. *Clinical Symptoms and Social Well-being.* *Psychiatr Serv* 2008; 59:232-234. <http://ps.psychiatryonline.org/cgi/content/full/59/3/232>
23. Hasset, A. Defining psychotic disorders in an aging population. In *Psychosis in the Elderly, 2005* (pp. 11-22). Eds.:Hasset A, Ames D, Chiu E. Taylor and Frances: London. <http://www.informaworld.com/smpp/title~content=t733000386~db=all>
24. Ames, D. Psychosis in affective disorders-depression. In: *Psychosis in the Elderly, 2005* (pp. 150-161). Eds.Hasset A, Ames D, Chiu E. Taylor and Frances: London. <http://www.informaworld.com/smpp/title~content=t733000386~db=all>

Authors: Radha Raman, MD, & Michele Larzelere, MD, *LSU FMRP-Kenner, LA*

Editor: Melissa Stiles, MD, *Fox Valley FMR, University of Wisconsin*