Brief Pain Inventory (Elderly) - Short Form

Background

- 1. Brief measuring tool for pain and its effect on function of daily living
- 2. Based on the Wisconsin Brief Pain Questionnaire developed by the Pain Research Group Self-administered survey
- 3. Produces similar data in many languages; multi-lingual forms are available
- 4. Can be used to determine the effectiveness of pain intervention

Validity

- 1. In a 2008 RCT, the BPI-SF did not lead to more information about osteoarthritis pain in the elderly than open-ended questions
- 2. In a 2004 study of the BPI-SF in arthritic and low back pain patients, results supported the validity of the BPI-SF
- 3. BPI-SF
 - $\circ \quad http://www3.mdanderson.org/depts/prg/bpisf.pdf \\$

References

- Keller, S. Validity of the Brief Pain Inventory for Use in Documenting the Outcomes of Patients with Noncancer Pain. Clin J Pain. 20(5) Sept-Oct 2004: 309-18.
- 2. Mcdonald, D. Older Adult Pain Communication and the Brief Pain Inventory Short Form. Pain Management Nursing. 9, December 2008:154-159.

Author: Isabel Garcia, MD, Baylor COM-Kesley Seybold, TX

Editor: Melissa Stiles, MD, Fox Valley FMR, University of WI