

## **Galanin N-Terminal fragment (1-15) induces a stronger effect than Galanin in depression- and anxiety- like related behavioural tests in rats.**

C. Millón<sup>1</sup>, M. Narváez<sup>1</sup>, A. Flores-Burgess<sup>1</sup>, C. Parrado<sup>1</sup>, A. Puigcerver<sup>2</sup>, R. Coveñas<sup>4</sup>, L. Santín<sup>2</sup>, DO. Borroto-Escuela<sup>3</sup>, K. Fuxe<sup>3</sup>, J. A. Narváez<sup>1</sup>, Z. Díaz-Cabiale<sup>1</sup>.

<sup>1</sup>. Faculty of Medicine, University of Málaga, Málaga

<sup>2</sup>. Faculty of Psychology, University of Málaga, Málaga

<sup>3</sup>. Department of Neuroscience, Karolinska Institute, Stockholm

<sup>4</sup>. Institute of Neuroscience of Castilla y León, Salamanca.

Galanin (GAL) and GAL Receptors are implicated in anxiety- and depression-related behaviours. The GAL N-terminal fragment (1-15) [GAL(1-15)] also induces depressant- and anxiogenic- like actions. In this work, we have compared the effects of GAL and GAL(1-15) in depression- and anxiety- related behaviours using the Forced Swimming Test (FST) and the Open Field Test (OFT).

In the first set of experiments, rats (n=8-10) were injected with GAL 3nmol, GAL(1-15) 3nmol or vehicle 15min before the FST. Two sessions were conducted: a 15min. pre-test followed 24h later by 5min. test. The duration of immobility and climbing were recorded during the test. In the second set, rats (n=6-10) were injected with GAL 3nmol, GAL(1-15) 3nmol or vehicle 15min before the OFT. Time and entries in the central square during 300 Sec. were scored.

In the FST, the increase in the immobility induced by GAL(1-15) was significantly higher ( $p<0.01$ ) than the one induced by GAL. In the climbing response GAL(1-15) also induced a significantly stronger decrease ( $p<0.05$ ) in climbing compared with GAL.

In the OFT, GAL(1-15) significantly decrease the number of entries ( $p<0.01$ ) and time ( $p<0.05$ ) in centre while GAL lacks of effect.

These results indicate that GAL(1-15) has more powerful effects than GAL in depression-like behaviour in the FST and anxiogenic-like effects in the OFT. These findings may give the basis for the development of novel therapeutic drugs targeting GAL(1-15) system for treatment of depression and anxiety disorders.

This study was supported by Spanish CVI6476 and TV3-Marató 090130/31/32.