

Interferometric Versus Projective Measurement of Anyons

Claire Levaillant¹ and Michael Freedman²

1 Department of Mathematics, South Hall Room 6607, University of California, Santa Barbara, CA 93106, USA

2 Station Q, Microsoft Research, Santa Barbara, CA 93106-6105, USA

Abstract

Two distinct methods for measuring topological charge in a nonabelian anyonic system have been discussed in the literature: projective measurement of a single point-like quasiparticle and interferometric measurement of the total topological charge of a group of quasiparticles. Projective measurement by definition is only applied near a point and will project to a topological charge sector near that point. Thus, if it is to be applied to a *group* of anyons to project to a *total* charge, then the anyons must first be fused one by one to obtain a single anyon carrying the collective charge. We show that interferometric measurement is strictly stronger: Any protocol involving projective measurement can be simulated at low overhead by another protocol involving only interferometric measurement.

1998 ACM Subject Classification J.2 Physical Sciences and Engineering

Keywords and phrases anyons, measurement, interferometry, physics

Digital Object Identifier 10.4230/LIPIcs.TQC.2015.245

1 Introduction

We clarify a foundational issue regarding the relative power of two computational schemes which have been long proposed for nonabelian anyons. The two are projective measurement [8] and interferometric measurement; abstractly [11, 10, 5, 4, 1] and in a physical system, [7].

The discussion is restricted to the basic case: unitary modular tensor categories (UMTC) [13]. This is a mathematical idealization of a $(2 + 1)$ -dimension topological quantum field theory (TQFT). The unitary requirement is necessary for the system to be a legitimate model of the non-dissipative physics of the low energy states of a gapped quantum mechanical system. The modularity assumption is a nonsingularity requirement. Abstractly this condition assures us that the quantum system can be consistently defined on a physical torus (and this implies all surfaces [12]) – the system is *not* tied to the plane. Concretely it tells us that every topologically nontrivial excitation (or equivalently “quasiparticle” or “anyon”) of the theory a is detected by its braiding with some other anyon b (see page 29 of [1]). For example, when anyons a and b are mutually abelian, i.e., have a unique fusion channel c , braiding assumes the form:

$$\begin{array}{c} \text{Diagram: A loop with two crossings, labeled } b \text{ and } a \text{ at the bottom.} \\ = M_{b,a} \begin{array}{c} \text{Diagram: Two parallel vertical lines, labeled } b \text{ and } a \text{ at the bottom.} \end{array}, M_{b,a} \text{ a nontrivial phase.} \end{array}$$

Modularity implies that for each nontrivial particle a , there is a (possibly composite) b so that not only is $M_{b,a} \neq 1$ but $M_{b,a} \neq M_{b,a'}$, $a' = a$, a' an anyon of the theory. Even for *non-abelian* theories, modularity implies that particles are distinguished via braiding.



© Claire Levaillant and Michael Freedman;
licensed under Creative Commons License CC-BY


10th Conference on the Theory of Quantum Computation, Communication and Cryptography (TQC 2015).

Editors: Salman Beigi and Robert König; pp. 245–249

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



■ **Figure 1**

This distinguishability was exploited in [1, 4] to give a full analysis of anyonic interferometers of Mach-Zehnder and Fabry-Perot designs. We note that much of the applied and even experimental literature on measuring topological charge, e.g. [14], has focused on Fabry-Perot geometries. In the low tunnelling (single pass) limit, the Fabry-Perot interferometer determines the same evolution of the system density matrix ρ as the simpler Mach-Zehnder interferometer. Even so, the evolution of ρ under operation of the interferometer is quite complicated. Fortunately, and this is the main conclusion of [1, 4], there is an easy *topological* interpretation of the asymptotic action of an interferometer, Int_a , which has converged to a measurement “a”, meaning the total topological charge within the interferometric loop has been measured to be that of an anyon of type a . See Figure 1. Furthermore, convergence to this limit is efficient – exponentially fast.

In contrast, a projective measurement Proj_a to particle type a is the Hermitian orthogonal projection to that particle sector. It is visualized as occurring by bringing some external prob, such as an STM tip, up to an isolated point-like anyon and directly detecting some, perhaps non-universal, signature of that particular particle type in that particular system. For example, even for an electrically neutral ψ , in $\nu = \frac{5}{2}$ fractional quantum Hall effect (FQHE), higher moments of the electric field might provide a signature. In any case, it is this hope which has led to the projective measurement model.

We now show that within the UMTC formalism, any protocol using projective measurement can be efficiently simulated by a protocol which instead uses interferometric measurement.

To do this we first need to define, through a density matrix diagram, the asymptotic topological action of Int_a . The diagrams, Figure 2b and on, have a $|\text{ket}\rangle\langle\text{bra}|$ aspect when read from top to bottom. The diagrams in $|\text{ket}\rangle\langle\text{bra}|$ format, of course, represent operators (density matrices), and Figure 2a represents a state vector ($|\text{ket}\rangle$). It should be noted that such representations of operators and states obey topological rules [13] and may or may not correspond bit by bit to a physical process.

We start with a vacuum and create a, perhaps complex, system of anyons which we divide into two halves “inside” and “outside” the interferometer. In Figure 2 these two halves at any given time are depicted simply as points, but they may represent composite anyons, in dual groups, drawn out of the vacuum.




Figure 2




Figure 3

From [1, 4], Int_a asymptotically transforms from Figure 2b to Figure 3, assuming that the measurement outcome a , indeed, has nonzero probability. In Figure 3 and below, we drop overall nonzero scalars from the diagrams.


Notation: ω_a is the a^{th} row of the normalized S -matrix which operates as a projector onto the a particle:

$$\begin{array}{c} b \\ | \\ \omega_a = \delta_{a,b} \\ | \\ b \end{array}$$

The presence of ω_a is expected – it projects onto the a -particle type sector. The ω_0 loop has long been regarded as an unfortunate but unavoidable consequence of running the interferometer. ω_0 encodes a kind of decoherence between inside and outside caused by the intervening stream of prob particles b . In [4], this severing of charge lines k running from the inside to the outside of the interferometer was called *anyonic charge line decoherence*. Topologically, ω_0 surgers the a -lines (Figure 3c) so that a and \bar{a} have forgotten that they came out of the vacuum together – they are no longer correlated.

The presence of this ω_0 -loop leads to a general supposition in the community that Int_a and Proj_a were incomparable: Int_a permits non-demolition measurement, as the internal correlations within the two groups of quasiparticles, inside and outside, are *not* disturbed, whereas in order to obtain a localized particle on which to apply Proj_a , the internal structure of the quasiparticle group to be measured would first need to be destroyed by a series of fusions (which produces decoherence even if the fusion outcomes are presumed not to be observed). In contrast, Proj_a does *not* cause anyonic charge line decoherence between the measured subsystem from its complement. Each seemed to have its own peculiar advantages and disadvantages with respect to the preservation of quantum information.

However, we will now show that the anyonic charge line decoherence of a and \bar{a} can be reversed (oddly, this oximoron *is* possible in a topological system) by additional interferometric measurements. The key observation is that if after Int_a we interferometrically measure the collective state of $\bar{a} \cup a$, there is still a nonzero probability of observing the outcome 0, the trivial particle. This is because if the two lines in Figure 3c are recoupled, the F -symbol,



■ Figure 4

$F_{\bar{a},a;0,0}^{\bar{a},a} \neq 0$, reflecting the fact that a particle and its uncorrelated antiparticle may fuse into the vacuum. (This, in fact, is the definition of an antiparticle.) Suppose we measure $\bar{a} \cup a$ and observe 0, so that Int_0 is applied to the system. The result is given in Figure 4.

We see that Int_0 has restored us to the situation we would have been in if the ω_0 loop associated to the initial Int_a had *not* been present. The decoherence has been reversed.

Of course it is not certain that the second measurement will result in Int_0 ; other outcomes Int_k are also possible. But now there is an easy “ping-pong” strategy (referred to as “forced measurement” in [3]): Bounce back and forth between measuring the initial inside – which will always return outcome a , Int_a will be applied on the odd steps of this cycle, and measuring the entire system $\bar{a} \cup a$. Each of the odd steps decoheres the inside and outside of the initial interferometer and returns the density matrix to that shown in Figure 3c. Actually, since on the odd steps 3, 5, 7, . . . , there is no doubt that topological charge a will be measured, it is *not* necessary to *read* the output of the interferometer, but merely to run the probe particles around the initial loop, thus producing anyonic charge line decoherence. Each of these even step measurements on the entire system constitute an *independent* chance to apply Int_0 . Because of independence, the tail event that after $2t$ measurements Int_0 has not been applied decays exponentially in t . The exact exponential rate is easily calculated from the data of any particular UMTC. Since $F_{\bar{a},a;0,0}^{\bar{a},a} = \frac{1}{d_a}$, charge 0 is observed at each step $2t$ with probability $p = \left(\frac{1}{d_a}\right)^2$, so the exponential rate of decay in t is $2 \log_e \left(1 - \frac{1}{d_a^2}\right)$.

2 Conclusion and outlook

We have shown how to “projectively”¹ measure the total topological charge with repeated interferometric measurements of groups of anyons without decohering the group from its complement. In contrast, projective measurement of anyonic charge is in the usual model [8] limited to projecting to the topological charge of a single anyon. A priori, this looks like a strictly *weaker* operation.

It should be remarked, though, that as with all issues of complexity, at this point in history there are no “lower bounds.”

Any proof that it is impossible to simulate interferometric measurement by projective measurement would necessarily rely on complexity assumptions. We regard this as an area for future work.

However, evidence of the enhanced strength of interferometric measurement is presented in a series of papers [2, 9] on universal gate systems for qubit and qutrit systems within $SU(2)_4$ and its Jones-Kauffman partner. Previously a universal protocol for a certain qutrit

¹ We place “projectively” in quotes because unlike the usual usage in the arena of anyonic systems, this measurement is both nonlocal and nondemolitional. It is projective in the usual quantum mechanical sense of Hermitian orthogonal projection onto an eigenbasis, in this case the eigenbasis of total topological charge.

within $SU(2)_4$ was found [6] using projective measurement but the argument appears quite special and not applicable to qubits.

We call attention to a shortcut for graphically exploring interferometry protocols. Because of the iterative process we have just described for eliminating the decohering ω_0 -loops associated to interferometric measurement, one may proceed – in the manner of a person writing computer code in a higher order language – only to manipulate the $|\text{ket}\rangle$ which describes the current state of the system of anyons. The $|\text{ket}\rangle$ is used at any given time to describe the state that has been pulled out of the vacuum. It is not necessary to double the diagram by adding the dual bra (and the linking $|\text{ket}\rangle\langle\text{bra}|$ by ω_0 -loops). Any ω_0 -loop will eventually be removed by some even numbered step of our protocol. Thus it is not really necessary to draw the ω_0 -loops, or even the $\langle\text{bra}|$, but merely to keep track of the $|\text{ket}\rangle$. In the end, if a density matrix ρ is desired, one may obtain ρ as the outer product of the final $|\text{ket}\rangle_{\text{final}}$ with its dual $\langle\text{bra}|_{\text{final}}$, $\rho = |\text{ket}\rangle_{\text{final}}\langle\text{bra}|_{\text{final}}$.

References

- 1 Bonderson, P. *Non-abelian anyons and interferometry*. PhD thesis, Caltech, 2007. Physics.
- 2 Bonderson, P., Freedman, M., Lutchyn, R., Nayak, C., and Wang, Z. Unpublished.
- 3 Bonderson, P., Freedman, M., and Nayak, C. Measurement-only topological quantum computation via anyonic interferometry. *Annals of Physics* 324 (2009), 787.
- 4 Bonderson, P., Shtengel, K., and Slingerland, J.K. Decoherence of anyonic charge in interferometry measurements. *Phys. Rev. Lett.* 98 (2007), 070401.
- 5 Bonderson, P., Shtengel, K., and Slingerland, J.K. Interferometry of non-Abelian Anyons. *Annals of Physics* 323 (2008), 2709.
- 6 Cui, S.X., and Wang, Z. Universal quantum computation with metaplectic anyons. arXiv:1405.7778, April 2014.
- 7 Das Sarma, S., Freedman, M., and Nayak, C. Topologically-protected qubits from a possible non-abelian fractional quantum hall state. *Phys. Rev. Lett.* 94 (2005), 166802.
- 8 Kitaev, A. Fault-tolerant computation by anyons. *Ann. Phys.* 303 (2003), 2–111.
- 9 Levaillant, C. 2014, to appear.
- 10 Overbosch, B., and Bais, F. Inequivalent classes of interference experiments with non-abelian anyons. *Phys. Rev. A* 64 (2001), 062107.
- 11 Preskill, J. Fault-tolerant quantum computation. In *Introduction to Quantum Computation*, H.-K. Lo, S. Popescu, and T. P. Spiller, Eds. World Scientific, 1998.
- 12 Turaev, V.G. *Quantum invariants of knots and 3-manifolds*, vol. 18 of *Studies in Mathematics*. Walter de Gruyter, 1994.
- 13 Wang, Z. *Topological quantum computation*, vol. 112 of *Regional Conference Series in Mathematics*. Conference Board of the Mathematical Sciences, 2010.
- 14 Willet, R., Pfeiffer, L., West, K., and Manfra, M. Aharonov-bohm effect and coherence length of charge $\frac{e}{4}$ quasiparticles at $\frac{5}{2}$ filling factor measured in multiple small fabry-perot interferometers. arXiv:1301.2594, January 2013.