

Total Space in Resolution Is at Least Width Squared*

Ilario Bonacina[†]

KTH Royal Institute of Technology, School of Computer Science and
Communication, Stockholm, Sweden
ilario@kth.se

Abstract

Given an unsatisfiable k -CNF formula φ we consider two complexity measures in Resolution: width and total space. The width is the minimal W such that there exists a Resolution refutation of φ with clauses of at most W literals. The total space is the minimal size T of a memory used to write down a Resolution refutation of φ , where the size of the memory is measured as the total number of literals it can contain. We prove that $T = \Omega((W - k)^2)$.

1998 ACM Subject Classification F.2.2 Nonnumerical Algorithms and Problems – Complexity of proof procedures

Keywords and phrases Resolution, width, total space

Digital Object Identifier 10.4230/LIPIcs.ICALP.2016.56

1 Introduction

Resolution is a well known propositional proof system introduced by Blake in [16] and proposed by Robinson in [38] for automated theorem proving. Since then this proof system became the most studied proof system in the sub-area of complexity theory that is *Proof Complexity*. Given a set of clauses φ , that is a set of disjunctions of literals or, equivalently, given a formula in Conjunctive Normal Form, *Resolution* is a method to infer new clauses according to the following inference rule:

$$\frac{C \vee x \quad D \vee \neg x}{C \vee D}, \quad (1)$$

where C, D are clauses and x is a variable. Resolution is sound and complete, that is it is possible to derive the empty clause \perp if and only if φ is unsatisfiable. A Resolution *refutation* of φ is then just a sequence of clauses C_1, \dots, C_ℓ with $C_\ell = \perp$ and each clause of the sequence is either a clause from φ or it is inferred by previous clauses in the sequence according to the inference rule in equation (1).

Nowadays, the main reason for the interest in Resolution comes from a practical perspective: it is at the core of most of the state-of-the-art SAT solvers since the introduction of the DPLL algorithm [22, 23] and its improvements, the so called *Conflict Driven Clause Learning* (CDCL) algorithms [4, 32, 40]. The track of the running of such algorithms on unsatisfiable instances produces a (particular form of) Resolution proofs. Hence Resolution is a valuable tool to study their performances and limitations. In this work we are interested in more theoretical questions about the Resolution proof system and the reader interested

* A full version of this paper is available online on ECCC [18].

[†] The author was partially funded by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013) / ERC grant agreement no. 279611.



© Ilario Bonacina;

licensed under Creative Commons License CC-BY

43rd International Colloquium on Automata, Languages, and Programming (ICALP 2016).

Editors: Ioannis Chatzigiannakis, Michael Mitzenmacher, Yuval Rabani, and Davide Sangiorgi;

Article No. 56; pp. 56:1–56:13



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



in more details on the connections between Resolution and SAT solvers could look at the recent survey [35].

Given an unsatisfiable φ we are interested in measuring how complex a Resolution proof of φ must be. Certainly there are many ways of measuring the complexity of proofs and in this work we are interested in connecting two of such measures. The main complexity measure we can associate to φ in Resolution, and by far the most important, is the minimal length of a Resolution refutation of φ . This measure is denoted with $\text{size}(\varphi \vdash \perp)$ and, since a long time now, we know that there are certain formulas φ_n requiring exponentially long proofs, e.g. the encodings of the Pigeonhole Principle [28] or Tseitin formulas [39, 42]. Another, easier to study, complexity measure is the *width*. Suppose that we focus on Resolution refutations of a formula φ with clauses up to a certain length w . The minimal w such that we have a refutation of φ with clauses of length at most w is the *width*, $\text{width}(\varphi \vdash \perp)$. We have a trivial upper bound connecting size and width, that is for every set of clauses φ in n variables

$$\text{size}(\varphi \vdash \perp) \leq n^{O(\text{width}(\varphi \vdash \perp))},$$

and indeed this trivial bound could be asymptotically tight, cf. [3]. Another, more useful, connection between width and size is the following result from the seminal paper by Ben-Sasson and Wigderson [12]:

$$\log_2 \text{size}(\varphi \vdash \perp) \geq \frac{(\text{width}(\varphi \vdash \perp) - k)^2}{16n}, \quad (2)$$

where φ is a collection of clauses over n variables and each of them has at most k literals. Hence, if $\text{width}(\varphi \vdash \perp) = \omega(\sqrt{n \log n})$ then, immediately by the previous inequality, $\text{size}(\varphi \vdash \perp)$ is super-polynomial. Moreover this size-width inequality is essentially optimal [21].

Regarding the space complexity of proofs, its investigation was proposed in 1998 by Armin Haken as a natural analogue of the space complexity in the context of Turing machines and the first definitions of space measures in Resolution were given in [25, 1]. When talking about space, Resolution proofs are seen as a sequence of memory configurations $\mathfrak{M}_0, \dots, \mathfrak{M}_\ell$, where each \mathfrak{M}_i is a set of clauses, $\perp \in \mathfrak{M}_\ell$ and each \mathfrak{M}_{i+1} derive from \mathfrak{M}_i in one of the two following ways:

Axiom download: $\mathfrak{M}_{i+1} \subseteq \mathfrak{M}_i \cup \{C\}$, where $C \in \varphi$;

Inference: $\mathfrak{M}_{i+1} = \mathfrak{M}_i \cup \{D \vee E\}$, where both $D \vee x$ and $E \vee \neg x$ belong to \mathfrak{M}_i , for some variable x .

We then have some notions of how “spacious” a memory configuration can be. The most natural space measure for a memory configuration is of course the number of bits needed to write down it. Unfortunately it turns out that this notion of space is quite hard to study and hence some alternative notions of space were introduced [25, 1]. For example, the *clause space* of a memory configuration is the number of distinct clauses it can contain. The *total space*¹ of a memory configuration instead is the total number of literals it can contain. The minimal s such that we have a refutation of φ with memory configurations with total space at most s is the *Total Space* (needed to refute φ), $\text{TSpace}(\varphi \vdash \perp)$. Similarly for the clause space we obtain $\text{CSpace}(\varphi \vdash \perp)$. A more formal definition of $\text{TSpace}(\varphi \vdash \perp)$, to avoid misunderstandings, is provided in Section 2.

¹ In [1] this space complexity measure is called *variable space*, but we follow [10, 9, 33, 11, 34] in calling it *total space*. This is due to distinguish it from a different space complexity measure in which different occurrences of the same variable are not counted, the *variable space*, investigated for instance in [43].

We now recall some known results about space complexity measures. Given any unsatisfiable set of clauses φ in n variables, in [25] it was proven that

$$\text{CSpace}(\varphi \vdash \perp) \leq n + 1,$$

and, as a trivial consequence, we have that

$$\text{TSpace}(\varphi \vdash \perp) \leq n(n + 1).$$

Both upper bounds are asymptotically tight, for example for random k -CNF formulas [8, 20]. Regarding lower bounds, in [2] it is proved that

$$\text{CSpace}(\varphi \vdash \perp) \geq \text{width}(\varphi \vdash \perp) - k + 1, \quad (3)$$

where φ consists of clauses of at most k literals. Clearly $\text{TSpace}(\varphi \vdash \perp) \geq \text{CSpace}(\varphi \vdash \perp)$ and whenever $\text{TSpace}(\varphi \vdash \perp) = \omega(\text{CSpace}(\varphi \vdash \perp))$ we say to have a *non-trivial* total space lower bound.

The total space measure was introduced in [1] and there the first non-trivial lower bounds were proven, for two particular class of formulas the Complete Tree formulas and the Pigeonhole Principle formulas. After that, in [20] it was introduced a technique to prove total space lower bounds in Resolution. That technique was sufficiently strong to prove asymptotically optimal total space lower bounds for instance for random k -CNFs [20, 13] but the proofs given there are quite long and involved. This paper, as a corollary, deeply simplify such proofs. Space complexity measures are also studied concerning trade-offs with other complexity measures, see for example [9, 36, 11, 34, 7, 5].

1.1 Contributions

This work is about proving an analogue of the inequality in (3) for the total space. This will add a nice bit to our knowledge of the lattice of relations between complexity measures in Resolution; it will simplify the proofs of existing total space lower bounds and it will imply new non-trivial total space lower bounds.

► **Theorem 1.** *Let φ be a k -CNF formula, then*

$$\text{TSpace}(\varphi \vdash \perp) \geq \frac{1}{16} (\text{width}(\varphi \vdash \perp) - k - 4)^2.$$

The general idea of the proof is the following: given a Resolution refutation, we identify a memory configuration where some small clause appear and then show that before that moment there must have been some memory configuration with a lot of clauses (and hence with large total space). This idea was originally used in [1] in some particular cases and in more generality in [20]. Indeed the proof we show has some close structural similarities with the total space lower bound from [20] and essentially it is a simplification of the proof of Theorem 2.5 from the author's PhD. Thesis [17]. The proof we give is not direct since it involves another, less studied, complexity measure: the *asymmetric width*, $\text{awidth}(\varphi \vdash \perp)$, and families of assignments closely related to it. The asymmetric width was introduced in [29, 30] and the definition is quite technical so we defer it to Section 2 where we collect all the preliminary definitions and notations. The proof of Theorem 1 is purely combinatorial and implicitly uses some properties from a characterization of the asymmetric width from [15], cf. Section 3 for more details, together with a result tightly connecting the width and the asymmetric width, Theorem 2 (Lemma 8.5 from [29]).

Although defined quite differently, asymmetric width and width indeed share many properties. For instance, an analogue of the size-width inequality by Ben-Sasson and Wigderson [12]: given an unsatisfiable CNF formula φ in n variables

$$\ln(\text{size}(\varphi \vdash \perp)) \geq \frac{\text{awidth}(\varphi \vdash \perp)^2}{8n},$$

cf. Theorem 6.12 of [31]. For more information and history on the asymmetric width we refer to [15].

1.2 Examples of applications (and limitations) of Theorem 1

Since the seminal work of Ben-Sasson and Wigderson [12], the width measure has become one of the main tool to study Resolution proofs and their complexity. Hence we already have many relevant width lower bounds for many interesting class of formulas and then the range of applications of Theorem 1 is quite large. Below we recall some relevant examples.

Tseitin formulas. Given a d -regular graph G over n vertices, the Tseitin formula over G , $\text{Tseitin}(G)$, is a CNF formula over $dn/2$ variables based on a propositional encoding of the fact that the total degree in any graph is even, see for example [12] for a formal definition. Such formulas were used by Tseitin to prove the first super-polynomial size lower bound for Resolution size [41]. Since then, Tseitin formulas became one of the standard tools in proof complexity to prove lower bounds and trade-offs, see for example [39, 43, 12, 25, 7]. In particular given a connected 3-regular graph G over n vertices which is an expander, we have that $\text{width}(\text{Tseitin}(G) \vdash \perp) \geq \Omega(n)$, cf. [12]. Hence, by Theorem 1, we have an asymptotically optimal total space lower bound: $\text{TSpace}(\text{Tseitin}(G) \vdash \perp) = \Theta(n^2)$. This answers the open question 4 from [1] in the case of Resolution.

Random k -CNFs. A random k -CNF with n variables and clause density Δ is a CNF formula picked as follows: choose independently uniformly at random Δn clauses from the set of all possible clauses in the variables $\{x_1, \dots, x_n\}$ containing exactly k literals. If $\Delta = o(n^{1/4})$, Beame et al. [6] showed that random k -CNFs require exponential size Resolution proofs. Such result was simplified in [12] by showing a lower bound on width: if φ is a random k -CNF ($k \geq 3$) in n variables and Δn clauses, and Δ is a constant for simplicity, then with high probability $\text{width}(\varphi \vdash \perp) \geq \Omega(n)$. Hence, by Theorem 1, with high probability $\text{TSpace}(\varphi \vdash \perp) \geq \Omega(n^2)$. That is, almost every k -CNF require asymptotically optimal total space to be refuted in Resolution. This result was proven in [20] for $k \geq 4$ and for $k = 3$ in [13] with some explicit but quite involved constructions. Instead, as we saw, an asymptotically optimal total space lower bound for such formulas follows immediately from Theorem 1.

Formulas with short proofs. Bonet and Galesi [21] showed that the size-width inequality by Ben-Sasson and Wigderson [12] is essentially optimal. That is they showed that there are arbitrarily large 3-CNF formulas φ_n with $\Theta(n^3)$ clauses, $\Theta(n^2)$ variables and such that

- $\text{width}(\varphi_n \vdash \perp) = \Theta(n)$,
- $\text{CSpace}(\varphi_n \vdash \perp) = \Theta(n)$,

but φ_n has some Resolution proof of size $O(n^3)$, width $O(n)$ and clause space $O(n)$. Theorem 1 in this case tells us that $\text{TSpace}(\varphi_n \vdash \perp) = \Omega(n^2)$, which is just a linear lower bound in the number of variables of φ_n . On the other hand this is a non-trivial total space lower bound since $\text{TSpace}(\varphi_n \vdash \perp) = \omega(\text{CSpace}(\varphi_n \vdash \perp))$.

Regarding the limitations, Theorem 1 suffers from the same kind of limitations of size-width inequality, equation (2), and the clause space-width inequality, equation (3). That is it became trivially vacuous for CNF formulas in n variables with clauses with many literals. For example we see such phenomenon when considering encodings of the negation of the Pigeonhole Principle as CNFs having clauses of n literals, the PHP_n^{n+1} formulas. For such formulas $\text{width}(\text{PHP}_n^{n+1} \vdash \perp) = \Theta(n)$ and hence no size lower bound or clause space lower bound could be implied directly from equations (2) - (3). The same applies for Theorem 1. On the other hand, by different techniques, we still have size lower bounds [42, 37], $\text{size}(\text{PHP}_n^{n+1} \vdash \perp) \geq 2^{\Omega(n)}$, clause space lower bounds [1], $\text{CSpace}(\text{PHP}_n^{n+1} \vdash \perp) \geq n$, and total space lower bounds [1, 20], $\text{TSpace}(\text{PHP}_n^{n+1} \vdash \perp) \geq \frac{1}{4}n^2$.

1.3 Organization of the paper

Section 2 contains all the preliminary definitions and notations needed for the proof of Theorem 1. In Section 3 we prove Theorem 1 and we give some more detailed comments on the proof. Section 4 contains some open questions about total space.

2 Preliminaries

We consider fixed a set of variables X and, given a natural number n , we denote as $[n]$ the set $\{1, \dots, n\}$. Given a set A , $\binom{A}{\leq 2}$ is the subset of the power set of A consisting of all the subsets of size at most 2.

Partial assignments. Given a set of variables X , a *partial (Boolean) assignment over X* is a function $\alpha : X \rightarrow \{0, 1\} \cup \{\star\}$. The *domain* of α is $\text{dom}(\alpha) = \alpha^{-1}(\{0, 1\})$ and we say that α assigns a value to x if $x \in \text{dom}(\alpha)$. Given two partial assignments over X , α and β we say that α extends β , $\beta \subseteq \alpha$, if for all $x \in X$, $\beta(x) \in \{\alpha(x), \star\}$. We denote by $\{x \mapsto b\}$ the partial assignment with domain the variable x mapped to $b \in \{0, 1\}$. Given two partial assignments α and β with disjoint domains, with $\alpha \cup \beta$ we denote the partial assignment with domain $\text{dom}(\alpha) \cup \text{dom}(\beta)$ such that for each $x \in \text{dom}(\alpha) \cup \text{dom}(\beta)$

$$\alpha \cup \beta(x) = \begin{cases} \alpha(x) & \text{if } x \in \text{dom}(\alpha), \\ \beta(x) & \text{if } x \in \text{dom}(\beta). \end{cases}$$

CNF formulas. A *literal* is a variable in X or the negation of a variable in X . A clause C is a formula of the form $\ell_1 \vee \dots \vee \ell_k$, where the ℓ_i are literals and m is the *width* of the clause C , denoted as $|C|$. A formula in *Conjunctive Normal Form* (CNF) is a formula φ with variables in X of the form $C_1 \wedge \dots \wedge C_m$, where the C_j s are clauses. A k -CNF formula is a CNF formula where each clause has at most k distinct literals. With $\text{var}(\varphi)$ we denote the set of variables occurring in the formula φ .

Given a CNF formula φ over a set of variables X and a partial assignment α over X , we can apply α to φ , obtaining a new CNF formula, denoted as $\varphi \upharpoonright_\alpha$ or $\alpha(\varphi)$, in the following way: for each variable $x \in \text{dom}(\alpha)$ substitute each occurrence of x in φ with $\alpha(x)$. Then simplify the resulting CNF according to the following rules: $\neg 0 \equiv 1$, $\neg 1 \equiv 0$, $0 \vee A \equiv A$, $1 \vee A \equiv 1$, $1 \wedge A \equiv A$, $0 \wedge A \equiv 0$. We say that α *satisfies* φ if $\alpha(\varphi) = 1$ and we say that α *falsifies* φ if $\alpha(\varphi) = 0$. Similarly, we can apply a partial assignment α to set of formulas $A = \{C_1, \dots, C_\ell\}$ component-wise: $A \upharpoonright_\alpha = \{C_1 \upharpoonright_\alpha, \dots, C_\ell \upharpoonright_\alpha\}$. Given a set of formulas F and a partial assignment α we say that α satisfies F , $\alpha \models F$, if and only if for every formula $\varphi \in F$, $\alpha(\varphi) = 1$.

Resolution proofs. A *Resolution derivation* of a clause C from a CNF formula φ is a sequence of clauses $\pi = (C_1, \dots, C_\ell)$ such that $C_\ell = C$ and each C_i is either a clause from φ or it is inferred from C_j, C_k with $j, k < i$ and such that $\frac{C_j \neg C_k}{C_i}$ is a valid instance of the Resolution rule:

$$\frac{C \vee x \quad D \vee \neg x}{C \vee D}$$

where C, D are clauses and x is a variable; or, C_i is inferred from a C_j with $j < i$ and such that $\frac{C_j}{C_i}$ is a valid instance of the *weakening* inference rule²

$$\frac{C}{C \vee D}$$

where C, D are clauses. A *Resolution refutation* of a CNF formula φ is a Resolution derivation of the empty clause \perp from φ . Resolution is sound and complete, that is it is possible to infer the empty clause \perp from φ if and only if φ is unsatisfiable.

Width. Given a sequence of clauses $\pi = (C_1, \dots, C_\ell)$ we recall that

$$\text{width}(\pi) = \max_{C_j \in \pi} |C_j|$$

and the minimal width needed to refute φ in Resolution is

$$\text{width}(\varphi \vdash \perp) = \min_{\pi} \text{width}(\pi),$$

where the min is taken over all refutations of φ in Resolution³.

Asymmetric width. The notion of *asymmetric width* was introduced in [30, 31]. Let φ be a CNF formula and $\pi = (C_1, \dots, C_\ell)$ be a Resolution derivation from φ . To define the asymmetric width of π , $\text{awidth}(\pi)$ we preliminary need the notion of *witness function*. A *witness function for $\pi = (C_1, \dots, C_\ell)$* is a function $\sigma : [\ell] \rightarrow \binom{[\ell]}{\leq 2} \cup \{\star\}$ witnessing the fact that π is a derivation from φ , that is such that

- $\sigma(i) = \{j, k\}$ implies that $j, k < i$ and $\frac{C_j \neg C_k}{C_i}$ is a valid instance of the inference rule of Resolution and if $j = k$ we require $\frac{C_j}{C_i}$ to be a valid instance of the weakening rule; and
- $\sigma(i) = \star$ implies that C_i is a clause from φ .

Given $\pi = (C_1, \dots, C_\ell)$ a Resolution derivation from φ and a witness function σ for π , the *asymmetric width of C_i with respect to π and σ* , $\text{aw}_{\pi, \sigma}(C_i)$, is defined as follows

$$\text{aw}_{\pi, \sigma}(C_i) = \begin{cases} 0 & \text{if } \sigma(i) = \star, \text{ that is } C_i \in \varphi, \\ \min_{j \in \sigma(i)} |C_j| & \text{otherwise.} \end{cases}$$

Then $\text{awidth}(\pi)$ is the minimum over all the possible functions σ witnessing the validity of π of the maximum over i of $\text{aw}_{\pi, \sigma}(C_i)$, that is

$$\text{awidth}(\pi) = \min_{\sigma} \max_{C_i \in \pi} \text{aw}_{\pi, \sigma}(C_i).$$

² Notice that the weakening rule is not really needed but it will make simpler the exposition when dealing with restrictions of Resolution proofs.

³ If φ is a satisfiable CNF formula then is customary to define $\text{width}(\varphi \vdash \perp) = \infty$.

Finally, the asymmetric width needed to refute φ , $\text{awidth}(\varphi \vdash \perp)$, is the minimum of $\text{awidth}(\pi)$ over all possible sequence of clauses $\pi = (C_1, \dots, C_\ell)$ that are Resolution refutations of φ .

Clearly it holds that $\text{awidth}(\varphi \vdash \perp) \leq \text{width}(\varphi \vdash \perp)$. Interestingly, the width cannot be much bigger than the asymmetric width.

► **Theorem 2** (Lemma 8.5 of [29]). *Let φ be an unsatisfiable k -CNF formula, then*

$$\text{width}(\varphi \vdash \perp) \leq \text{awidth}(\varphi \vdash \perp) + \max\{\text{awidth}(\varphi \vdash \perp), k\}.$$

A self-contained proof of this result, essentially based on [14], is proven in the full version of this paper [18].

Total Space. As we saw in the introduction, a Resolution refutation of a CNF formula φ can be seen as a sequence of memory configurations $\pi = (\mathfrak{M}_0, \dots, \mathfrak{M}_\ell)$, where each \mathfrak{M}_i is a set of clauses, $\perp \in \mathfrak{M}_\ell$ and each \mathfrak{M}_{i+1} derive from \mathfrak{M}_i in one of the two following ways:

Axiom download: $\mathfrak{M}_{i+1} \subseteq \mathfrak{M}_i \cup \{C\}$, where $C \in \varphi$;

Inference: $\mathfrak{M}_{i+1} = \mathfrak{M}_i \cup \{D \vee E\}$, where both $D \vee x$ and $E \vee \neg x$ belong to \mathfrak{M}_i , for some variable x .

Given π as above, the *total space* of π is

$$\text{TSpace}(\pi) = \max_{i \in [\ell]} \sum_{C \in \mathfrak{M}_i} |C|$$

and given an unsatisfiable CNF formula φ , the total space needed to refute φ in Resolution is

$$\text{TSpace}(\varphi \vdash \perp) = \min_{\pi} \text{TSpace}(\pi),$$

where the min is taken over all the possible Resolution refutations of φ given as a sequence of memory configurations⁴.

3 Proof of Theorem 1

First let's prove the main result of this work, Theorem 1, for convenience of the reader restated below. We postpone more detailed comments on the proof after the proof itself.

► **Restated Theorem 1.** *Let φ be a k -CNF formula, then*

$$\text{TSpace}(\varphi \vdash \perp) \geq \frac{1}{16} (\text{width}(\varphi \vdash \perp) - k - 4)^2.$$

Proof. Let $\text{awidth}(\varphi \vdash \perp) = r + 1$. We prove that

$$\text{TSpace}(\varphi \vdash \perp) \geq \frac{1}{4} (r - 1)^2,$$

or, more precisely, we prove that every Resolution refutation of φ must pass through a memory configuration of at least $(r - 1)/2$ clauses each of width at least $(r - 1)/2$. Once we prove this, the desired lower bound between total space and width follows:

$$\text{TSpace}(\varphi \vdash \perp) \geq \frac{1}{4} (r - 1)^2 \geq \frac{1}{16} (\text{width}(\varphi \vdash \perp) - k - 4)^2,$$

⁴ If φ is a satisfiable CNF formula then is customary to define $\text{TSpace}(\varphi \vdash \perp) = \infty$.

where the last inequality uses that $\text{width}(\varphi \vdash \perp) \leq 2(r+1) + k$, a consequence of Theorem 2.

Let Ξ and Ψ be two functions respectively mapping subsets of clauses into subsets of partial assignments and viceversa. Given a set of clauses A ,

$$\Xi(A) = \{\alpha \text{ partial assignment} : \forall C \in A, \alpha(C) \neq 0\},$$

and given a set of partial assignments F ,

$$\Psi(F) = \{C \text{ clause} : \exists \alpha \in F, \alpha(C) = 0\}.$$

Notice that, by construction, for every set of clauses A , $A \cap \Psi \circ \Xi(A) = \emptyset$ and $\perp \in \Psi(F)$ whenever F is non-empty. We consider the following special set:

$$W_r = \{C \text{ clause} : \text{awidth}(\varphi \vdash C) \leq r\},$$

and its images $\Xi(W_r)$ and $\mathcal{S} = \Psi \circ \Xi(W_r)$. The main reason to consider the set $\Xi(W_r)$ is the following property:

► **Claim 3** (Extension Property of $\Xi(W_r)$). *Let α be a \subseteq -maximal partial assignment in $\Xi(W_r)$ and x a variable not in $\text{dom}(\alpha)$, then for every $\beta \subseteq \alpha$ such that $|\text{dom}(\beta)| < r$ both $\beta \cup \{x \mapsto 0\}$ and $\beta \cup \{x \mapsto 1\}$ are in $\Xi(W_r)$.*

Proof. By contradiction let $\beta \subseteq \alpha$ such that $|\text{dom}(\beta)| < r$ and $b \in \{0, 1\}$ such that $\beta_b = \beta \cup \{x \mapsto b\} \notin \Xi(W_r)$. Without loss of generality we can restrict to consider $b = 0$. Since $\beta_0 \notin \Xi(W_r)$ it means that there exists a clause D in W_r such that $\beta_0(D) = 0$ but $\alpha(D) \neq 0$. This means that $D = D' \vee x$, $|D| \leq r$ and $\beta(D') = \alpha(D') = 0$. By maximality of α then both $\alpha_0 = \alpha \cup \{x \mapsto 0\} \notin \Xi(W_r)$ and $\alpha_1 = \alpha \cup \{x \mapsto 1\} \notin \Xi(W_r)$. In particular there exists a clause $E \in W_r$ such that $\alpha_1(E) = 0$, so, as before, we must have that $E = E' \vee \neg x$ and $\alpha(E') = 0$. But now

$$\frac{D' \vee x \quad E' \vee \neg x}{D' \vee E'}$$

is a valid instance of the Resolution rule. Hence, by definition of asymmetric width,

$$\text{awidth}(\varphi \vdash D' \vee E') \leq \max\{\text{awidth}(\varphi \vdash D), \text{awidth}(\varphi \vdash E), r\} \leq r,$$

since both D and E belong to W_r and $|D| \leq r$. So $D' \vee E' \in W_r$ and $\alpha(D' \vee E') = 0$ which is a contradiction. ◀

Let $\pi = (\mathfrak{M}_0, \dots, \mathfrak{M}_\ell)$ be a Resolution refutation of φ given as a sequence of memory configurations. By definition of W_r , $\perp \notin W_r$ and hence the empty partial assignment is in $\Xi(W_r)$, so, in particular $\perp \in \mathcal{S}$. Hence the following set is non-empty:

$$\mathcal{A} = \{i \in [\ell] : \exists C \in \mathfrak{M}_i \cap \mathcal{S}, |C| < (r-1)/2\}.$$

Let $t = \min \mathcal{A}$ and let $C \in \mathfrak{M}_t \cap \mathcal{S}$ be a clause of width less than $(r-1)/2$. Since $C \in \mathcal{S}$ there must exist a partial assignment $\alpha \in \Xi(W_r)$ that falsifies C and let α_C be the minimal partial assignment contained in α falsifying C . Notice that $|\text{dom}(\alpha_C)| = |C| < (r-1)/2$. Our goal now is to show that there exists some $i < t$ such that $|\mathfrak{M}_i \cap \mathcal{S}| \geq (r-1)/2$. Since for every $i < t$ every clause in $\mathfrak{M}_i \cap \mathcal{S}$ has width at least $(r-1)/2$, this will give the desired result.

For sake of contradiction, suppose that for each $i < t$, $|\mathfrak{M}_i \cap \mathcal{S}| < (r-1)/2$. We inductively construct a sequence of assignments β_0, \dots, β_t in $\Xi(W_r)$ such that for each $i \leq t$ we have

that $\alpha_C \subseteq \beta_i$ and that $\beta_i \models \mathfrak{M}_i \cap \mathcal{S}$. This immediately give a contradiction when we reach β_t , since α_C falsifies the clause $C \in \mathfrak{M}_t \cap \mathcal{S}$ and $\beta_t \supseteq \alpha_C$.

The first memory configuration \mathfrak{M}_0 is empty, so we can put $\beta_0 = \alpha$. Supposing that $0 \leq i < t$ and that we already have a suitable β_i , we construct β_{i+1} distinguishing between two cases.

Axiom download case. $\mathfrak{M}_{i+1} \subseteq \mathfrak{M}_i \cup \{D\}$, where D is a clause from φ . Since each clause D from φ belongs to W_r and we have that $W_r \cap \mathcal{S} = \emptyset$, then $\mathfrak{M}_i \cap \mathcal{S} = \mathfrak{M}_{i+1} \cap \mathcal{S}$ and hence we can simply put $\beta_{i+1} = \beta_i$.

Inference case. $\mathfrak{M}_{i+1} \subseteq \mathfrak{M}_i \cup \{D \vee E\}$ where $D \vee E$ follows by Resolution on some variable x from two clauses $D \vee x$ and $E \vee \neg x$ in \mathfrak{M}_i . Then, by the inductive hypothesis, there exists $\beta_i \in \Xi(W_r)$ such that $\beta_i \models \mathfrak{M}_i \cap \mathcal{S}$, let $\bar{\beta}_i \in \Xi(W_r)$ be a \subseteq -maximal partial assignment containing β_i and let β be an assignment contained in $\bar{\beta}_i$ \subseteq -minimal such that $\alpha_C \subseteq \beta$ and $\beta \models \mathfrak{M}_i \cap \mathcal{S}$. We have that

$$|\text{dom}(\beta)| \leq |\text{dom}(\alpha_C)| + |\mathfrak{M}_i \cap \mathcal{S}| < (r-1)/2 + (r-1)/2 = r-1,$$

where the first inequality follows easily from the fact that to satisfy a clause $F \in \mathfrak{M}_i \cap \mathcal{S}$ an assignment just have to satisfy a single literal in F . Notice that since $|\text{dom}(\beta)| \leq r-2$ the extension property from Claim 3 can be applied twice and we will use this later. The main property of β that we now use is the following:

► **Claim 4.** *Let $\gamma \in \Xi(W_r)$ and F be any clause in \mathfrak{M}_i , if $\text{var}(F) \subseteq \text{dom}(\gamma)$ and $\beta \subseteq \gamma$, then $\gamma \models F$.*

Proof. Since $\text{var}(F) \subseteq \text{dom}(\gamma)$, then $\gamma(F) \in \{0, 1\}$. If by contradiction $\gamma(F) = 0$, then, by construction $F \in \mathcal{S}$ and, again by construction, $\beta \models \mathfrak{M}_i \cap \mathcal{S}$. So $\beta \models F$, which is a contradiction since $\beta \subseteq \gamma$. ◀

The remaining part of the proof is just case analysis. If there is some variable y in $D \vee E$ unassigned by $\bar{\beta}_i$ then we can use the extension property (Claim 3) extending β to some $\beta' \in \Xi(W_r)$ setting y and satisfying $D \vee E$.

If $\text{var}(D \vee E) \subseteq \text{dom}(\beta)$ then we can extend β to some assignment $\beta' \in \Xi(W_r)$ setting x to some value (either by choosing $\bar{\beta}_i$ if $x \in \text{dom}(\bar{\beta}_i)$ or otherwise by the extension property). Then $\text{var}(D \vee x) \subseteq \text{dom}(\beta')$, and, by the previous claim, $\beta' \models D \vee x$. The same happens for $E \vee \neg x$ and hence $\beta' \models D \vee E$ by the soundness of the Resolution rule.

The only remaining possibility is that $\text{var}(D \vee E) \not\subseteq \text{dom}(\beta)$ but $\text{var}(D \vee E) \subseteq \text{dom}(\bar{\beta}_i)$, and without loss of generality suppose that $\text{var}(D) \not\subseteq \text{dom}(\beta)$. If $x \in \text{dom}(\bar{\beta}_i)$ then, by the previous claim $\bar{\beta}_i \models (D \vee x) \wedge (E \vee \neg x)$ so $\bar{\beta}_i \models D \vee E$. Suppose then that $x \notin \text{dom}(\bar{\beta}_i)$. By Claim 3 we have that $\beta' = \beta \cup \{x \mapsto 0\} \in \Xi(W_r)$. Take a \subseteq -maximal assignment in $\Xi(W_r)$ containing β' , let $\bar{\beta}'$ be such assignment. If $\text{var}(D \vee x) \subseteq \text{dom}(\bar{\beta}')$ then, by the previous claim, $\bar{\beta}_i \models D \vee x$, but $\beta'(x) = 0$ so $\bar{\beta}' \models D$ and hence $\bar{\beta}' \models D \vee E$. If $\text{var}(D \vee x) \not\subseteq \text{dom}(\bar{\beta}')$ then there is some variable y in $D \vee x$ not assigned by $\bar{\beta}'$ and since $|\text{dom}(\beta')| = |\text{dom}(\beta)| + 1 < r$ we can apply the extension property to β' extending it setting y and satisfying D . ◀

First of all notice that we proved something actually stronger, that is we proved that given an unsatisfiable CNF formula φ , every Resolution refutation of φ must pass through a memory configuration of at least $\frac{1}{2}(\text{awidth}(\varphi \vdash \perp) - 2)$ clauses each of width at least $\frac{1}{2}(\text{awidth}(\varphi \vdash \perp) - 2)$.

A crucial point in the proof of Theorem 1 is Claim 3. It is related with the following characterization of asymmetric width in Resolution by [15].

► **Theorem 5** (Theorem 22 from [15]). *Let φ be an unsatisfiable CNF formula, then the followings are equivalent:*

1. $\text{awidth}(\varphi \vdash \perp) > r$.

2. *There exists a non-empty set \mathcal{F} of partial assignments such that:*

Consistency: *For every $\alpha \in \mathcal{F}$ and every clause C of φ , $\alpha(C) \neq 0$;*

Extension: *If $\alpha \in \mathcal{F}$ and $\beta \subseteq \alpha$ is such that $|\text{dom}(\beta)| < r$, then for every variable $x \notin \text{dom}(\alpha)$ and for every $\epsilon \in \{0, 1\}$ there exist $\beta_\epsilon \in \mathcal{F}$ with $\beta \subseteq \beta_\epsilon$ such that $\beta_\epsilon(x) = \epsilon$.*

Claim 3 is based on the proof of the implication from 1. to 2. in the previous theorem. The other implication (easier to prove) is not needed for Theorem 1. Indeed it is easy to see that given 1. the set of \subseteq -maximal partial assignments in $\Xi(W_r)$ satisfies the properties claimed in 2., and the crucial extension property is essentially Claim 3.

4 Open questions

We conclude this work with some open questions about the behaviour of the total space measure. Most of the questions are motivated by some analogy with the behaviour of the clause space measure.

On super-linear lower bounds. Is there any family of k -CNF formulas φ_n in n variables and $n^{O(1)}$ clauses such that $\text{size}(\varphi_n \vdash \perp) = n^{O(1)}$ and $\text{TSpace}(\varphi_n \vdash \perp) = \Theta(n^2)$?

For the formulas from [21] we saw in Section 1.2 we just have a linear total space lower bound. If we could find some formulas ψ_n with polynomial size Resolution proofs⁵ but such that $\text{width}(\psi_n \vdash \perp) = \omega(\sqrt{n})$ then, by Theorem 1, we would have that $\text{TSpace}(\psi_n \vdash \perp) = \omega(n)$. This is anyway quite far from the question we are asking here and it seems that a positive answer should need some new techniques.

On simpler proofs for total space lower bounds. Is there a simpler more direct proof of a total space-width lower bound?

The clause space inequality $\text{CSpace}(\varphi \vdash \perp) \geq \text{width}(\varphi \vdash \perp) - k + 1$, where φ is a k -CNF, can be proven using some families of assignments and a characterization of Resolution width [2] or it can be proven (with some small loss in an additive constant) via some operation on Resolution proofs [26]. The proof we have of Theorem 1 is in a sense similar to the proof of the clause-width lower bound in [2] although not quite simple as that since we pass through the asymmetric width and more complicated families of assignments.

Beyond Resolution. Space measures are defined in [1] also for proof systems stronger than Resolution. For example for Polynomial Calculus, a proof system where instead of clauses we infer polynomials, or $\text{Res}(k)$, a (stronger) version of Resolution where instead of clauses k -DNF can be inferred, or Frege systems. In all such systems very little is known about space especially when it comes to total space. Regarding other space measures something is known for example for $\text{Res}(k)$ [10, 24] and for Polynomial Calculus [1, 19, 27]. In [1] it is

⁵ Equation 2 in this case implies that $\text{width}(\psi_n \vdash \perp) = O(\sqrt{n \log n})$.

proven that in Frege system the total space is always at most linear and regarding Polynomial Calculus, the only lower bounds known for total space are from [1] and those are for the PHP_n^{n+1} formulas and the Complete Tree formulas. Is there any family of k -CNF formulas in n variables φ_n with $n^{O(1)}$ clauses requiring $\omega(n)$ total space to be refuted say in Polynomial Calculus?

Acknowledgements. The main question about width and total space is the result of some discussions had at the Dagstuhl seminar 15171: we want to thank Schloss Dagstuhl for the inspiring environment and the kind hospitality. We want to thank Jakob Nordström and Nicola Galesi for several long discussions on space complexity.

References

- 1 Michael Alekhovich, Eli Ben-Sasson, Alexander A. Razborov, and Avi Wigderson. Space complexity in propositional calculus. *SIAM J. Comput.*, 31(4):1184–1211, 2002. doi:10.1137/S0097539700366735.
- 2 Albert Atserias and Víctor Dalmau. A combinatorial characterization of resolution width. *J. Comput. Syst. Sci.*, 74(3):323–334, 2008. doi:10.1016/j.jcss.2007.06.025.
- 3 Albert Atserias, Massimo Lauria, and Jakob Nordström. Narrow proofs may be maximally long. In *IEEE 29th Conference on Computational Complexity, CCC 2014, Vancouver, BC, Canada, June 11-13, 2014*, pages 286–297. IEEE, 2014. doi:10.1109/CCC.2014.36.
- 4 Roberto J. Bayardo Jr. and Robert Schrag. Using CSP look-back techniques to solve real-world SAT instances. In Benjamin Kuipers and Bonnie L. Webber, editors, *Proceedings of the Fourteenth National Conference on Artificial Intelligence and Ninth Innovative Applications of Artificial Intelligence Conference, AAAI 97, IAAI 97, July 27-31, 1997, Providence, Rhode Island.*, pages 203–208. AAAI Press / The MIT Press, 1997. URL: <http://www.aaai.org/Library/AAAI/1997/aaai97-032.php>.
- 5 Paul Beame, Christopher Beck, and Russell Impagliazzo. Time-space tradeoffs in resolution: superpolynomial lower bounds for superlinear space. In Howard J. Karloff and Toniann Pitassi, editors, *Proceedings of the 44th Symposium on Theory of Computing Conference, STOC 2012, New York, NY, USA, May 19-22, 2012*, pages 213–232. ACM, 2012. doi:10.1145/2213977.2213999.
- 6 Paul Beame, Richard M. Karp, Toniann Pitassi, and Michael E. Saks. On the complexity of unsatisfiability proofs for random k -CNF formulas. In Jeffrey Scott Vitter, editor, *Proceedings of the Thirtieth Annual ACM Symposium on the Theory of Computing, Dallas, Texas, USA, May 23-26, 1998*, pages 561–571. ACM, 1998. doi:10.1145/276698.276870.
- 7 Chris Beck, Jakob Nordström, and Bangsheng Tang. Some trade-off results for polynomial calculus: extended abstract. In Dan Boneh, Tim Roughgarden, and Joan Feigenbaum, editors, *Symposium on Theory of Computing Conference, STOC'13, Palo Alto, CA, USA, June 1-4, 2013*, pages 813–822. ACM, 2013. doi:10.1145/2488608.2488711.
- 8 Eli Ben-Sasson and Nicola Galesi. Space complexity of random formulae in resolution. *Random Struct. Algorithms*, 23(1):92–109, 2003. doi:10.1002/rsa.10089.
- 9 Eli Ben-Sasson and Jakob Nordström. Understanding space in resolution: optimal lower bounds and exponential trade-offs. In Peter Bro Miltersen, Rüdiger Reischuk, Georg Schnitger, and Dieter van Melkebeek, editors, *Computational Complexity of Discrete Problems, 14.09.–19.09.2008*, volume 08381 of *Dagstuhl Seminar Proceedings*. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Germany, 2008. URL: <http://drops.dagstuhl.de/opus/volltexte/2008/1781/>.

- 10 Eli Ben-Sasson and Jakob Nordström. A space hierarchy for k -DNF resolution. *Electronic Colloquium on Computational Complexity (ECCC)*, 16:47, 2009. URL: <http://eccc.hpi-web.de/report/2009/047>.
- 11 Eli Ben-Sasson and Jakob Nordström. Understanding space in proof complexity: Separations and trade-offs via substitutions. In Bernard Chazelle, editor, *Innovations in Computer Science – ICS 2010, Tsinghua University, Beijing, China, January 7-9, 2011. Proceedings*, pages 401–416. Tsinghua University Press, 2011. URL: <http://conference.itcs.tsinghua.edu.cn/ICS2011/content/papers/3.html>.
- 12 Eli Ben-Sasson and Avi Wigderson. Short proofs are narrow – resolution made simple. *J. ACM*, 48(2):149–169, 2001. doi:10.1145/375827.375835.
- 13 Patrick Bennett, Ilario Bonacina, Nicola Galesi, Tony Huynh, Mike Molloy, and Paul Wollan. Space proof complexity for random 3-CNFs. *CoRR*, abs/1503.01613, 2015. URL: <http://arxiv.org/abs/1503.01613>.
- 14 Olaf Beyersdorff and Oliver Kullmann. Hardness measures and resolution lower bounds. *CoRR*, abs/1310.7627, 2013. URL: <http://arxiv.org/abs/1310.7627>.
- 15 Olaf Beyersdorff and Oliver Kullmann. Unified characterisations of resolution hardness measures. In Carsten Sinz and Uwe Egly, editors, *Theory and Applications of Satisfiability Testing – SAT 2014 – 17th International Conference, Held as Part of the Vienna Summer of Logic, VSL 2014, Vienna, Austria, July 14-17, 2014. Proceedings*, volume 8561 of *Lecture Notes in Computer Science*, pages 170–187. Springer, 2014.
- 16 Archie Blake. *Canonical Expressions in Boolean Algebra*. PhD thesis, University of Chicago, 1937. University of Chicago.
- 17 Ilario Bonacina. *Space in weak propositional proof systems*. PhD thesis, Department of Computer Science, Sapienza University of Rome, 12 2015.
- 18 Ilario Bonacina. Total space in resolution is at least width squared. *Electronic Colloquium on Computational Complexity (ECCC)*, 2016. <http://eccc.hpi-web.de/report/2016/057/>.
- 19 Ilario Bonacina and Nicola Galesi. A framework for space complexity in algebraic proof systems. *J. ACM*, 62(3):23, 2015. doi:10.1145/2699438.
- 20 Ilario Bonacina, Nicola Galesi, and Neil Thapen. Total space in resolution. In *55th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2014, Philadelphia, PA, USA, October 18-21, 2014*, pages 641–650. IEEE Computer Society, 2014. doi:10.1109/FOCS.2014.74.
- 21 Maria Luisa Bonet and Nicola Galesi. Optimality of size-width tradeoffs for resolution. *Computational Complexity*, 10(4):261–276, 2001. doi:10.1007/s000370100000.
- 22 Martin Davis, George Logemann, and Donald W. Loveland. A machine program for theorem-proving. *Commun. ACM*, 5(7):394–397, 1962. doi:10.1145/368273.368557.
- 23 Martin Davis and Hilary Putnam. A computing procedure for quantification theory. *J. ACM*, 7(3):201–215, 1960. doi:10.1145/321033.321034.
- 24 Juan Luis Esteban, Nicola Galesi, and Jochen Messner. On the complexity of resolution with bounded conjunctions. *Theor. Comput. Sci.*, 321(2-3):347–370, 2004. doi:10.1016/j.tcs.2004.04.004.
- 25 Juan Luis Esteban and Jacobo Torán. Space bounds for resolution. *Inf. Comput.*, 171(1):84–97, 2001. doi:10.1006/inco.2001.2921.
- 26 Yuval Filmus, Massimo Lauria, Mladen Mikša, Jakob Nordström, and Marc Vinyals. From small space to small width in resolution. In Ernst W. Mayr and Natacha Portier, editors, *31st International Symposium on Theoretical Aspects of Computer Science (STACS 2014), STACS 2014, March 5-8, 2014, Lyon, France*, volume 25 of *LIPIcs*, pages 300–311. Schloss Dagstuhl – Leibniz-Zentrum fuer Informatik, 2014. doi:10.4230/LIPIcs.STACS.2014.300.

- 27 Yuval Filmus, Massimo Lauria, Jakob Nordström, Noga Ron-Zewi, and Neil Thapen. Space complexity in polynomial calculus. *SIAM J. Comput.*, 44(4):1119–1153, 2015. doi:10.1137/120895950.
- 28 Armin Haken. The intractability of resolution. *Theor. Comput. Sci.*, 39:297–308, 1985. doi:10.1016/0304-3975(85)90144-6.
- 29 Oliver Kullmann. Investigating a general hierarchy of polynomially decidable classes of CNF's based on short tree-like resolution proofs. *Electronic Colloquium on Computational Complexity (ECCC)*, (41), 1999. URL: <http://eccc.hpi-web.de/eccc-reports/1999/TR99-041/index.html>.
- 30 Oliver Kullmann. An improved version of width restricted resolution. In *AMAI*, 2000. URL: <http://rutcor.rutgers.edu/~amai/aimath00/regular/kullmann.ps>.
- 31 Oliver Kullmann. Upper and lower bounds on the complexity of generalised resolution and generalised constraint satisfaction problems. *Ann. Math. Artif. Intell.*, 40(3-4):303–352, 2004. doi:10.1023/B:AMAI.0000012871.08577.0b.
- 32 Matthew W. Moskewicz, Conor F. Madigan, Ying Zhao, Lintao Zhang, and Sharad Malik. Chaff: Engineering an efficient SAT solver. In *Proceedings of the 38th Design Automation Conference, DAC 2001, Las Vegas, NV, USA, June 18-22, 2001*, pages 530–535. ACM, 2001. doi:10.1145/378239.379017.
- 33 Jakob Nordström. Narrow proofs may be spacious: Separating space and width in resolution. *SIAM J. Comput.*, 39(1):59–121, 2009. doi:10.1137/060668250.
- 34 Jakob Nordström. Pebble games, proof complexity, and time-space trade-offs. *Logical Methods in Computer Science*, 9(3), 2013. doi:10.2168/LMCS-9(3:15)2013.
- 35 Jakob Nordström. On the interplay between proof complexity and SAT solving. *ACM SIGLOG News*, 2(3):19–44, August 2015.
- 36 Jakob Nordström and Johan Håstad. Towards an optimal separation of space and length in resolution. *Theory of Computing*, 9:471–557, 2013. doi:10.4086/toc.2013.v009a014.
- 37 Alexander A. Razborov. Proof complexity of pigeonhole principles. In Werner Kuich, Grzegorz Rozenberg, and Arto Salomaa, editors, *Developments in Language Theory, 5th International Conference, DLT 2001, Vienna, Austria, July 16-21, 2001, Revised Papers*, volume 2295 of *Lecture Notes in Computer Science*, pages 100–116. Springer, 2001.
- 38 John Alan Robinson. A machine-oriented logic based on the resolution principle. *J. ACM*, 12(1):23–41, 1965. doi:10.1145/321250.321253.
- 39 Uwe Schöning. Resolution proofs, exponential bounds, and Kolmogorov complexity. In Igor Prívara and Peter Ruzicka, editors, *Mathematical Foundations of Computer Science 1997, 22nd International Symposium, MFCS'97, Bratislava, Slovakia, August 25-29, 1997, Proceedings*, volume 1295 of *Lecture Notes in Computer Science*, pages 110–116. Springer, 1997. doi:10.1007/BFb0029954.
- 40 João P. Marques Silva and Karem A. Sakallah. GRASP: A search algorithm for propositional satisfiability. *IEEE Trans. Computers*, 48(5):506–521, 1999. doi:10.1109/12.769433.
- 41 G.S. Tseitin. On the complexity of derivation in propositional calculus. In Jörg H. Siekmann and Graham Wrightson, editors, *Automation of Reasoning, Symbolic Computation*, pages 466–483. Springer Berlin Heidelberg, 1983. URL: http://dx.doi.org/10.1007/978-3-642-81955-1_28, doi:10.1007/978-3-642-81955-1_28.
- 42 Alasdair Urquhart. Hard examples for resolution. *J. ACM*, 34(1):209–219, 1987. doi:10.1145/7531.8928.
- 43 Alasdair Urquhart. The depth of resolution proofs. *Studia Logica*, 99(1-3):349–364, 2011. doi:10.1007/s11225-011-9356-9.