

contrasting grasslands by time averaged gradients (TAG)

Daniela Famulari^{1,2}, Robert L. Storeton-West¹, Eiko Nemitz¹, Y. Sim Tang¹, Kenneth J. Hargreaves¹, Keith Weston², Mark A. Sutton¹, David Fowler¹


¹ = Centre for Ecology and Hydrology Edinburgh, Bush Estate, Penicuik, EH26 0QB, Scotland, UK. ² = Institute of Atmospheric and Environmental Science, University of Edinburgh, West Mains Road, EH9 3JJ, Edinburgh, UK.

Corresponding author e-mail: danf@ceh.ac.uk

INTRODUCTION & METHOD

The TAG (Time Averaged Gradient) is a low-cost system to measure long-term (weekly to monthly) deposition fluxes of trace gases between the surface and the atmosphere using the flux-gradient method by conditional sampling. Over daily periods, atmospheric conditions can range from high stability, where the vertical gradients of ambient concentration are enhanced due to very small diffusivity, to highly unstable conditions, in which concentration gradients are small due to the intense turbulent activity of the surface layer. The large vertical gradients generated by high stability would bias the estimate of the actual flux: to avoid this, the TAG system samples conditionally, within a carefully defined range of stability (see Famulari et al., 2007). In Fig. 1 is shown a diagram of the TAG system applied to measure long-term dry net exchange fluxes of NH₃ and SO₂ to three contrasting temperate grasslands in the UK described below (see Fig. 2).

Fig. 1. Diagram of TAG: CA: cup anemometer T: temperature sensor WV: wind vane GM: gas flow meters WG: wind power generator SP: solar panels B: battery P: air pump CL: Campbell 21X data logger D1,2,3: denuder trains RM: rotameters R: relay.



THE SITES





Fig. 2 Map of the UK, showing the locations of the 3 sites.

Auchencorth Moss: blanket bog over a surface of 1000 ha, located in Midlothian (NT221562, lat. 55°47'30", long. 3°14'20", alt. 270m ASL). The moorland vegetation comprises a succession of hummocks and hollows (vertical range 60 cm) including grasses, sedges, rushes, Sphagnum, and other bryophyte species. The site is used for sheep grazing at a very low density of about 0.1 to 1 sheep/ha. Low intensity agricultural activities in the vicinity. Flux monitoring of NH₃ and SO₂ during 1999-2003.

Plynlimon located in the Cambrian Mountains at SN 806853 (lat. 52°27'26", long. 3°47'46", alt. 500m). The site is quite exposed, windy and rainy. The land cover is a mixture of dwarf-shrub heath, blanket bog and unimproved acid grassland: it is used for sheep grazing almost all year round. Surrounding the area monitored by the TAG system there are farming activities at a distance of 1 to 1.5 miles. Flux monitoring of NH₃ and SO₂ during 2000-2003.

Easter Bush intensively managed grassland located south of Edinburgh (NT245641, lat. 55°52', long. 3°2', alt. 190 m ASL). The field is used for silage production, cattle and sheep grazing covered by more than 90% with Lolium Perenne, with a minor presence of *Phleum Pratense*, *Ranunculus repens*, *Poa Annua*, *Trifolium Repens*. Around the field site is a mixture of woodland, roads, rough grazing land and farms. Flux monitoring of NH₃ during 2001-2003.

CHARACTERISATION OF THE DIFFERENT SITES: METEOROLOGY AND STABILITY




CHEMICAL CLIMATE FOR NH₃ AND SO₂ AT THE THREE SITES AND EXCHANGE FLUXES

AUCHENCORTH MOSS

PLYNLIMON

EASTER BUSH



CONCLUSIONS

Several years of net deposition fluxes of NH₃ and SO₂ have been measured at three contrasting grasslands in the UK using Time Averaged Gradient technique (TAG): the low-cost system provides long term estimates of S and N dry deposition for comparison between different environments. The data show the meteorological differences between the sites, and through the seasonal cycles. This information, coupled with the measurement of the chemical climate and the land cover description, provide a very useful tool in the description of long-term trends, for validation of models and development of their parameterisation, although the data do not permit accurate description of short term fluxes.

The description of the sites shows their suitability for the application of the TAG system: in general, for similar fetches, windier sites are better suited for atmospheric neutrality, and allow better data coverage: all sites in this work showed sufficient data coverage to describe reliably the deposition fluxes, but the scenario might change radically in more continental (more stable) conditions.

For ammonia the dry deposition fluxes were on average -7.6, -5.98, and -12.2 ng m⁻² s⁻¹ for Auchencorth Moss, Plynlimon and Easter Bush respectively, indicating differences in NH₃ concentrations and exchange processes, reflecting the interaction of temperature, rain and different grassland managing regimes. The measurements of SO₂ fluxes have suffered from the low background concentrations at the sites (close to detection limit): the overall results show deposition is dominating, but values are very often in the noise-around-zero region, with average values at -7.04 ng m⁻² s⁻¹ and -3.4 ng m⁻² s⁻¹ at Plynlimon and Auchencorth Moss, respectively.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the contribution of the EU under the ACCENT BIAFLUX project and the NitroEurope IP, as well as funding for this work from Department for Environment, Food and Rural Affairs (DEFRA) through the "Acid Deposition Processes"

REFERENCES

D. Famulari, D. Fowler, R.L. Storeton-West, E. Nemitz, K. Hargreaves, G. Rutherford, Y.S. Tang, M.A. Sutton, K.J. Weston. "Development of a low-cost system for measuring SO₂ and NH₃ deposition fluxes on the field scale by micrometeorology" Submitted to Environmental Pollution, 2007.

