



Univerza v Mariboru

Fakulteta za kemijo
in kemijsko tehnologijo

Diploma thesis

Extraction of curcuminoids with subcritical water

June, 2015

Ignacio Macicior Borregón



Univerza v Mariboru

Fakulteta za kemijo
in kemijsko tehnologijo

Ignacio Macicior Borregón

Extraction of curcuminoids with subcritical water

Diploma thesis

Maribor, 2015

ACKNOWLEDGMENTS

I want render thanks to my tutor, prof. Mojca Skerget, for offering me the opportunity of being part of this project. I am grateful also for Matej Ravber and Tina Perko, because they were my guides, preaching patient and advices. For their help, I thank Marko, Neven and Igor.

Finally, to my parents: you made everything possible, thanks for all.

Abstract

The purpose of this diploma thesis is to determine the best conditions for the extraction of antioxidants in curcuma, mainly curcuminoids, with pressurized hot water, while the currently industrial way to obtain these antioxidants is the ethanol extraction. This can be a more environmental friendly way, without organic solvent.

The extractions were performed for different times between 5 minutes and two hours, and in the temperature range from 100 to 200°C, while the was high enough to assure liquid water at this conditions. The influence of process parameters of the yield of extraction was determined. The extracts were analyzed of the content of curcuminoids by HPLC and their antioxidant activity by DPPH.

The results obtained are a maximum extraction yield at 175°C, being practically constant in the range of temperatures performed. The highest concentration of curcuminoids in the extract has been obtained at 100°C and 120 minutes of extraction. The extraction with the greatest amount of curcuminoids extracted per gram of curcuma has been obtained at 125°C, after 30 min of extraction.

At temperatures of 175 and 200°C all the curcuminoids are degraded to hydroxyl-cinnamic acids as paracumaric and ferulic acid. These substances have also a good antioxidant character; extractions with the highest antioxidant capability were those at 200°C and 60 or 120 minutes. Higher antioxidant capability can be expected in extractions with longer times or higher temperatures.

Key words: subcritical water extraction, curcuma, antioxidants.



Univerza v Mariboru

Fakulteta za kemijo
in kemijsko tehnologijo

Datum: 2. julij 2015

Na osnovi 330. člena Statuta Univerze v Mariboru (Ur. l. RS, št. 44/2015)

izdajam

SKLEP O DIPLOMSKEM DELU

Ignacio Macicior Borregón, študent/ka študijskega programa 1. stopnje Kemijska tehnologija, izpolnjuje pogoje, zato se mu/ji dovoljuje izdelati diplomsko delo.

MENTORICA: red. prof. dr. Mojca Škerget
SOMENTOR: red. prof. dr. Željko Knez

Naslov diplomskega dela:

EKSTRAKCIJA KURKUMINOIDOV S SUBKRITIČNO VODO

Naslov diplomskega dela v angleškem jeziku:

EXTRACTION OF CURCUMINOIDS WITH SUBCRITICAL WATER

Diplomsko delo je potrebno izdelati v skladu z "Navodili za kakovostno izvajanje diplomskega dela" in ga oddati v enem izvodu do 10. 7. 2015 v referat za študentske zadeve Fakultete za kemijo in kemijsko tehnologijo.

Pravni pouk: zoper ta sklep je možna pritožba na senat članice v roku 3 delovnih dni.

Obvestiti:

kandidata-ko
mentorja
somentorja
odložiti v arhiv



Dekan
Red. prof. dr. Željko KNEZ



Univerza v Mariboru

Fakulteta za kemijo
in kemijsko tehnologijo

Date: July, 2nd 2015

Based on the Article 330 of the Statute of the University of Maribor (Official Gazette of the Republic of Slovenia, No. 44/2015)

I issue a

DECREE ON BACHELOR'S THESIS

Ignacio Macicior Borregón, graduate student of Bachelor's study programme Chemical Technology is allowed to prepare his Bachelor's thesis.

MENTOR: Full Prof. Dr. Mojca Škerget

COMENTOR: Full Prof. Dr. Željko Knez

Bachelor's thesis title:

EXTRACTION OF CURCUMINOIDS WITH SUBCRITICAL WATER

Bachelor's thesis contents

Guidelines by mentor.

Bachelor's thesis is to be prepared in accordance to "Navodila za kakovostno izvajanje diplomskega dela (Guidelines for quality Bachelor's thesis preparation)" and delivered in one copy due 10. 7. 2015.

Legal caution: Notice of appeal against this decree must be filed with the Senate of Faculty of Chemistry and Chemical Engineering within 3 working days.

Forwarded to:

- The candidate
- The mentor
- The co-mentor
- Record depository



Dean:
Prof. Dr. Željko KNEZ

Index

| | |
|---|----|
| Abstract | 2 |
| Index | 3 |
| Index of figures and illustrations:..... | 5 |
| Index of tables..... | 7 |
| Index of equations..... | 8 |
| Symbols | 9 |
| Abbreviations | 10 |
| 1. Introduction..... | 11 |
| 2. Theory and literature review | 13 |
| 2.1 Curcumin, curcuminoids and curcuma | 13 |
| 2.2 Solid liquid extraction..... | 15 |
| 2.3 Extraction of natural products | 16 |
| 2.3.1 History: | 16 |
| 2.3.2 Methods of extraction:..... | 16 |
| 2.3.3 Solvent:..... | 19 |
| 2.4 High pressure water extraction..... | 23 |
| 2.4.1 Pressure:..... | 24 |
| 2.4.2 Density:..... | 25 |
| 2.4.3 Viscosity:..... | 26 |
| 2.4.4 Surface tension:..... | 27 |
| 2.4.5 Dielectric constant: | 28 |
| 2.5 Chemical reactions in high pressure water | 29 |
| 3. Procedure | 34 |
| 3.1 Material and equipment | 34 |
| 3.2 Experimental Procedure..... | 35 |
| 3.3 Analytical procedures..... | 36 |
| 3.3.1 HPLC analysis of curcuminoids..... | 36 |
| 3.3.2 DPPH analysis | 36 |
| 4. Results | 38 |

| | |
|---|----|
| 4.1 Extract yield | 38 |
| 4.2 Curcuminoids in extract | 40 |
| 4.3 Antioxidant capability | 42 |
| Color and bright | 43 |
| 4.5 Results of extractions at 100°C | 44 |
| 4.6 Results of extractions at 125°C | 45 |
| 4.5 results of extraction at 150°C | 46 |
| 4.5 results of extraction at 175°C | 48 |
| 4.9 Results of extractions at 200°C | 49 |
| 5. Conclusions: | 51 |
| 6. Literature | 52 |
| 7. Appendix | 55 |

Index of figures and illustrations:

| | |
|--|----|
| figure 1: water at critical point | 23 |
| Figure 2: Location of triple and critical points of water..... | 23 |
| figure 1: water at critical point | 23 |
| figure 3: temperature/pressure relation of saturated water as fluids definition | 24 |
| Figure 4: density of saturated liquid and gas water as fluid definition | 25 |
| Figure 5: viscosity of saturated liquid and gas water as fluids definition | 26 |
| Figure 6: surface tension of saturated liquid water. | 27 |
| Figure 7: dielectric constant of saturated liquid and gas water | 28 |
| figure 8: diketo-curcumin isomer | 29 |
| figure 9: dieno-cucumin isomer | 29 |
| figure 10: keto-eno-curcumin isome | 29 |
| Figure 11: comparison of diketo and keto-enol curcumin isomers behavior with water molecules in methanol solution..... | 30 |
| figure 12: ferulic acid..... | 33 |
| figure 13: vanilin | 33 |
| figure 14: p-coumaric acid | 33 |
| figure 15: 4-hidroxy-benzaldehyde | 33 |
| figure 16: autoclave used in the experiments | 35 |
| figure 17: extraction yield at different times and temperatures..... | 38 |
| figure 18: total curcuminuids extracted for each time and temperature and time. | 40 |
| figure 19: results of HPLC for extracions at rising temperatures..... | 41 |
| figure 20: Antioxidant capability, % inhibition for all temperatures and times..... | 42 |
| figure 21: color for each sample..... | 43 |
| figure 22: Extraction yield at 100°C..... | 44 |
| figure 23: HPLC & DPPH results for 100°C extractions | 44 |
| figure 24: Extraction yield at 125°C..... | 45 |
| figure 25: HPLC & DPPH results for 125°C extractions | 46 |
| figure 26:Extraction yield at 150°C..... | 46 |
| figure 27: HPLC & DPPH results for 150°C extractions | 47 |
| figure 28: stratification of the solid residue after centrifugation..... | 47 |
| figure 29:Extraction yield at 175°C..... | 48 |
| figure 30: HPLC & DPPH results for 175°C extractions | 48 |

| | |
|---|----|
| figure 32:Extraction yield at 200°C..... | 49 |
| figure 31: comparison of color of samples at 175°C, 10 and 60 minutes | 49 |
| figure 33: HPLC & DPPH results for 200°C extractions | 50 |

Index of tables

| | |
|---|----|
| Table 1: Properties of curcumin | 14 |
| Table 2: Properties of organic solvents | 20 |
| Table 3: Properties of organic solvents (continuation) | 21 |
| Table 4: physical properties of products of curcuminoids decomposition..... | 32 |
| Table 5: extraction yield of all samples | 55 |
| Table 6: HPLC results: mass of curcuminoids..... | 56 |
| Table 7: DPPH results. antioxidant capability. | 57 |

Index of equations

| | |
|---|----|
| equation 1: surface tension of saturated water | 27 |
| equation 2: O ₂ concentration of pressurized inert atmosphere..... | 35 |
| equation 3: % inhibition, antioxidant activity | 37 |
| equation 4: extract yield | 38 |

Symbols

| Symbol | Description | Units |
|---------------|--|-----------|
| MW | Molecular weight | g/mol |
| σ | Surface tension | N/m |
| B | Constant (surface tension) | N/m |
| b | Constant (surface tension) | [] |
| μ | Constant (surface tension) | [] |
| T_c | Critical temperature of water | K |
| T | Temperature | K |
| C_{O_2} | O ₂ concentration of inert atmosphere | %(volume) |
| $C_{O_2 atm}$ | O ₂ concentration of air | %(volume) |
| P_{atm} | Atmospheric pressure | bar |
| P_b | Blanketing pressure | Bar |
| P_{set} | Extraction pressure | Bar |
| n | number of blanketing | [] |
| % inhibition | Inhibition of UVA-Vis (antioxidant activity) | % |
| A_A | Blank absorbance | % |
| A_B | Sample absorbance | % |
| E | Extract yield | % |
| m_e | Extract and flask mass | g |
| m_f | Empty flask mass | g |
| m_c | Curcuma mass | g |

Abbreviations

| | |
|-------|--|
| HPSCW | High pressure subcritical water |
| C | Curcumin |
| DMC | Demethoxy curcumin |
| BDMC | Bidemethoxy curcumin |
| HPLC | High performance liquid chromatography |
| DPPH | 2,2-diphenyl-1-picrylhydrazyl |
| DME | Dimethoxy ethane |
| DMF | Dimethyl formamide |
| DMSO | Dymetyl sulfoxide |
| HMPA | Hexamethyl phosphoramide |
| HMPT | Hexamethyl phosphortriamide |
| MTBE | Methyl tert-butyl ether |
| NMP | N-Methyl-2-pyrrolidone |
| THF | Tetrahydrofuran |

1. Introduction

Curcuma longa is a common plant that grows in rainy and warm regions, and its rhizomes have been used the last 4000 years as food colorant, spicy and dye for clothes and skin [1]. The most interesting kind of compounds of this rhizome, and the reason of its properties, are curcuminoids, which represent the 3-5% of the rhizome mass. These compounds are curcumin (50-60%), demethoxy-curcumin (20-30%) and bis-demethoxy-curcumin (7-20%) [2], all of them with an antioxidant, anti microbial and anti protozoan, anti proliferative, anti-angiogenic, and anti-inflammatory powerful capability. The dry and minced rhizome is used as ingredient for traditional homemade medicine in India and China [2].

Nowadays, curcuminoids are being extracted with organic solvents, mainly ethanol, due to the low solubility of curcuminoids in neutral and acid water solutions, and due to the fact that ethanol is the harmless organic solvent for human being [3].

This thesis forms part of the study of alternative ways to obtain curcuminoids, as supercritical fluid extraction, Soxhlet extraction, and ultrasonic maceration, performed by Tina Perko, Matej Ravber, Zeljko Knez, Mojca Skerget, in the Faculty of Chemistry and Chemical Engineering, Laboratory for Separation Processes and Product Design, University of Maribor.

Extraction with pressurized water is a harmless and more environmentally friendly way to obtain curcuminoids, following the steps of the green chemistry of the XXI century. This green chemistry means the obtainment and use of natural products instead of synthetic ones, by eco and respectful ways, with the environment, society, producers, users and handlers.

Extraction of curcuminoids with pressurized water meets all the targets above mentioned. Curcuminoids are natural compounds with interesting properties for health and welfare, and extraction with pressurized water is a non organic solvent extraction, without toxic wastes for people and environment, harmless solvent for users and handlers and the high pressure industry is enough developed to be considered as safe.

The use of high pressure subcritical water, HPSCW, as solvent in extraction is a feasible candidate to obtain natural curcuminoids from turmeric due to the excellent properties of pressurized water as solvent and totally innocuous at room conditions.

The purpose of this study is to determinate the feasibility of this technique and to find the best conditions for the extraction of curcuminoids with HPSCW in order to define the boundaries and optimize the work conditions of real plants in the future.

2. Theory and literature review

2.1 Curcumin, curcuminoids and curcuma

Curcumin, a bis- α,β -unsaturated β -diketone, known also as diferuloylmethane [4] [2], is a natural polyphenolic compound, the main curcuminoid one, obtained in the turmeric (*curcuma longa*) rhizomes. Curcumin has interesting pharmacological properties, being anti proliferative (prevent the production of new cancerous cells), anti-angiogenic (hinder the production of new blood vessels in tumors), anti-inflammatory and antioxidant [2]; due his high yellow color, his main purpose along history has been as colorant. His code is E-100, in the group III, by the European Commission of Food and Feed safety [5].

Curcuma longa is a rhizomatous herbaceous plant, from the family of the zingiberaceae. It demands a temperature between 20-30°C and often rainfall. It grows in the southwest of India, China and Indonesia, where has been used as medicine due the curcuminoids properties. Turmeric contains moisture (>9%), curcumin (5–6.6%), extraneous matter (<0.5% by weight), mould (<3%), and volatile oils (<3.5%). The components responsible for the aroma of turmeric are turmerone, arturmerone, and zingiberene [1].

Curcuminoids, C, are a kind of di-phenyl-heptanoids, secondary metabolites produced by plants. As his name means, di-phenyl-heptanoids, are a heptane chain with two phenyl rings in both ends, with a keto or alcoholic functional group in 3 and 7 carbons [6]. Their properties are the same as the curcuma's ones, being this just the most common of them. These curcuminoids are demethoxy-curcumin, DMC, and bis-demethoxy-curcumin, BDMC.

The main difficulty in the curcuminoids application is the difficulty of ingest due his very limited solubility in water and acid solutions as the gastric acid [7]. By this way, his bioavailability is scarce, not just in order to supply it as medicament, but allocate it in the specific tissue in effectives amounts. As additional fact, curcumin in particular has also a rapid metabolism, making this bioavailability even lower [2] [8] [9].

Nowadays new methods to improve the bioavailability are been developed, as complex with nanoparticles, liposomes, micelles and phospholipids [7]. This new methods allow a slower metabolism, increasing the permeability against the intestinal conditions, like pH and enzymes. Other way for this purpose is encapsulation [2].

Table 1: Properties of curcumin, obtained from toxnet.nlm.nih.gov database [9]

| Curcumin | | | |
|--|--|---|--|
| Molecular Formula: | C ₂₁ H ₂₀ O ₆ | [10] | Molecular Weight: 368.39 [10] |
| Color/Form: | | | |
| Orange-yellow, crystal powder; gives brownish-red color with alkali; light-yellow color with acids | | | [10] |
| the crystal powder is Orange yellow prisms, rhombic prisms from methanol | | | [11] |
| Melting Point: | 183°C | [10] | Vapor Pressure(25°C): 3.08 · 10 ⁻¹² mmHg (est) [12] |
| Spectral Properties: | | | |
| MAX ABSORPTION (DIOXANE): | | 265 NM (LOG E= 4.18) | |
| | | 420 NM (LOG E= 4.77) [13] | |
| MASS: | 75938 | [11] | IR: 13460 [11] UV: 6-924 [11] |
| Solubility: | | | |
| In water(25°C): | 3.12 mg/L(est) | [12] | Insoluble in ether [10] |
| Soluble in glac. acetic acid | | [10] | Soluble in alcohol [10] |
| Octanol/Water Partition Coefficient: | | log Kow = 3.29 (est) [14] | |
| Henry's Law constant (25°C): | | 7.04X10 ⁻²² atm m ³ /mol (est) [12] | |
| Other Chemical/Physical Properties: | | | |
| Specific gravity (15°C): | 0.982-1.01 | [15] | refractive index (20°C): 1.5023-1.5088 [15] |
| optical rotation (90% ethanol, 10% turmerone): | | | +8° - +17° [15] |
| Slightly fluorescent | | [16] | acid value: 0.3-2.4 [15] |
| ester value: | 16-22.4 | [15] | ester value (after acetylation): 56-73.4 [15] |

2.2 Solid liquid extraction

Solid liquid extraction, or leaching, is the separation process of a substance originally in a solid matrix by another, based on the capability of dilute the first one, called solute, in the second one, called solvent. This solvent must be selective, to dissolve just the substances target. After the extraction, there are two phases, one is the solution of solute in the solvent, this is called extract; the other phase is the solid phase, this one often is not dry, being wet by the extract, the name of this wet solid phase is waste.

To perform the extraction, it's required:

Good contact between the solid and solvent phase,

Two different and separable phases.

The factors that determine the amount of solute extracted can be about steady or transitory process. Being the first ones:

- the solubility equilibrium between solute and solvent,
- the liquid kept by the solid in the phases separation process.

The transitory factors are all the mass transfer process:

- diffusion of solvent in solid,
- dissolution of the solute in the solvent,
- diffusion of solute in solvent through solid, and finally,
- diffusion of solute in solvent out the solid particle.

In order to increase mass transfer, the particle's size must be as small as it's possible, to enable short times of diffusion throw the solid; and the extraction process under agitation conditions, to improve the diffusion of the solute in the solvent. The solution process is just function of chemical parameters that cannot be improved changing mechanic properties.

With enough agitation and particles smallness, the extraction process can be considered stationary, being the yield ratio just function of temperature, that changes dissolution properties as transitory speed process and saturation capacity of the solvent. In this experience, this affirmation is not true due the reaction process, as it will be explained later.

2.3 Extraction of natural products

2.3.1 History:

Along history, different methods have been developed to obtain natural products from natural raw materials. Extraction of natural compounds started in Egypt and Mesopotamia during the ancient history, looking for compounds to produce high value products as pharmaceutical oils, waxes and perfumes, being the oldest evidence obtained dates 3500BC. Lately, several Sumerian texts (2100BC) show technological knowledge in extraction. This develops use water as solvent and inorganic additives, in order to change the water properties [17].

The first evidence of extraction with organic substances dates 1600BC, being used beer and wine as solvent. But the real development of this field started in 900AC, once ethanol was obtained in enough high concentrations [17].

2.3.2 Methods of extraction:

There are different ways of extraction, in order to the properties of the substance to be extracted.

Maceration:

The easiest extraction is supply contact between solvent and the particles where the solute is placed, keeping it together the needed time. This process can be with or without agitation. The particles must be pulverized, in order to increase the surface per mass ratio, being transferred the solute to the solvent.

The container can be emptied and filled with fresh solvent, to supply more capacity, if the previous one was saturated, or just to reduce the concentration of solute in the solvent kept by the waste, being a crosscurrent process. This process can be done too using the extract of this step as solvent in the previous one, being now countercurrent.

As last step it's needed a way to separate the waste and the extract, for this purpose the industrial ways are filtration, mechanic press or centrifuge.

Due his easy mechanism, maceration was the first extraction method executed.

According to the experiments performed by T. Perko et Al. [3] the extraction yield of 6 hours maceration of 10g curcuma with 250 mL ethanol at room temperature is 10.40%, being the amount of curcuminoids 166.45 mg curcumin (C), 84.35 mg demethoxy curcumin (DMC), 100.51 mg bis-demethoxy curcumin (BDMC), 351.32 mg curcuminoids per extract gram, and its antioxidant capability is 34%.

Extraction with solvent reflux:

The main feature of extraction is that remove the solute from one substance to another so, obviously, this new solution must have an easier way to be separated in his compounds. If boiling is the way, the solution can be evaporated in order to obtain fresh solvent, to be used again after being condensed. This way of work allows a small use of solvent per solute extracted. Extraction with soxhlet can be taken as example of extraction with solvent reflux.

The results obtained with soxhlet [3] in the same conditions (6h at room temperature, 10g curcuma in 250 mL ethanol) shows a raising of extraction, composition and antioxidant capability: 161.27 mg C, 111.04 mg DMC and 125.24 mg BDMC: 397.55 mg curcuminoids. Being the inhibition value 52%.

Percolation:

The process of fluids moving through porous solids, also called percolation, is a way to produce the contact between the solvent, as fluid, and the solute in the solid. As industrial method, the solid is placed on a filter and sustained by gravity, or fixed between two. The feed of solvent use to be fresh, in the inlet of the filter and it gets solute as it passes through the solid, may be in equilibrium in part of the thickness. As the fluid flow, the inlet gets empty of solute, so the equilibrium is done with the next particles, until all the solvent has been removed from the entire solid. The easiest example for this process is a coffee machine.

Percolation has not been studied as method to extract curcuminoids.

Ultrasound maceration:

As method for small extractions, as essential oils extraction, ultrasound can be a support in maceration. High resonance mechanic waves produce cavitation, due the fast change of

pressure. This cavitation produces the rupture of the cell wall. The cavitation produces bubbles just in contact with the solid, being an implicit mixer that increases the mass transfer in the place of maximum concentration. The ultrasonic produces also an increasing of the temperature, fact that raises the solubility of the solute in the solvent [18].

Maceration of curcuma supported with ultrasounds, has been performed by Perko et Al. [3], being used 1 gram of curcuma and 25 mL of Ethanol. The maceration takes 30 minutes, at 20°C, being the ultrasound fixed at 400W. The results obtained are 13.44 mg C, 69.44 mg DMC and 14.63 mg BDMC: 97.51 mg curcuminoids per gram of extract, and the absorption, around 20%.

Microwave maceration:

The use of microwave energy gives to the maceration the chance to allocate the energy in specific places, the organic compound and the cell wall. This allocated heat produces a dilatation of the cell, and his wall rupture. Once the cell is broken, the solvent gets inside, being heated by the cell itself in a mass and heat diffusion mechanism. This bigger temperature of the solvent raises the solubility of the solute in the solvent [18].

There are not studies of microwave maceration of curcuma due the good solubility of this one in ethanol at room temperature.

Chromatography:

Based in a multi infinitesimal stage distribution due the different affinity of each extract substance with the static and mobile phases, withhold and dragged respectively, chromatography reaches the total separation of the solute after enough time or length. Chromatography is used as analytic tool, qualitative and quantitative, not as separation process due his low mass capacity of operation. By this way chromatography has not been studied for curcuminoids procurement, but has been used for analysis study after all the experiments.

Supercritical fluids extraction:

Supercritical fluid is a fluid with a pressure and temperature higher than his critical point. In these conditions the different liquid and vapor phases doesn't exist. This fluid has particular properties, having his diffusion coefficient the same orders as his liquid phase, and his viscosity

the same order as his gas phase. But the most interesting property of supercritical fluids is the high reliance of solubility with density and this one with pressure and temperature [19] [20]. The fact of dispose solubility as controllable variable has especial interest in extraction process. The most common fluids for this kind of extraction are CO_2 and water.

Several experiments, using CO_2 as solvent, have taken place [3], being the best results obtained at 60°C and 200 bars: 18.88 mg C, 2.70 mg DMC and 5.03 mg BDMC: 26.60 mg of curcuminoids per gram of extract. This result reveals this method is not the best for curcumin extraction. The absorption is around 24%.

2.3.3 Solvent:

Speaking about natural products, there are a lot of restrictions, due the final purpose of the extract, being this often nutritional. The most common because his abundance, nonexistent hazard and usability, has been water, perfect to extract polarity compounds. To extract non polarity compounds can be used foodstuff desirably, or other organic solvents like propene, butane, acetylene, CO_2 , ethanol, etc. The rule “like extracts like” must be used as first key to choose the solvent, according to Van Laar theory [21], but solution solubility can be improved using mixtures [22].

Water or foodstuff residue in final extract is assumed to be harmless, for the rest of organic solvents their amount is limited in order not just to guarantee quality but safety.

The most common organic solvents and their solubility properties are shown in the table 2 [23]:

Table 2: Properties of organic solvents, compiled by Steve Murov [23]

| Solvent | formula | MW | boiling point (°C) | melting point (°C) | density (g/mL) | solubility in water (g/100g) | Dielectric Constant | flash point (°C) |
|-------------------------|---|--------|-----------------------|-----------------------|-------------------|------------------------------------|------------------------|---------------------|
| acetic acid | C ₂ H ₄ O ₂ | 60.052 | 118 | 16.6 | 10.446 | Miscible | 6.20 | 39 |
| acetone | C ₃ H ₆ O | 58.079 | 56.05 | -94.7 | 0.7845 | Miscible | 21.01 | -20 |
| acetonitrile | C ₂ H ₃ N | 41.052 | 81.65 | -43.8 | 0.7857 | Miscible | 36.64 | 6 |
| benzene | C ₆ H ₆ | 78.11 | 80.1 | 5.5 | 0.8765 | 0.18 | 2.28 | -11 |
| 1-butanol | C ₄ H ₁₀ O | 74.12 | 117.7 | -88.6 | 0.8095 | 6.3 | 17.8 | 37 |
| 2-butanol | C ₄ H ₁₀ O | 74.12 | 99.5 | -88.5 | 0.8063 | 15 | 17.26 | 24 |
| 2-butanone | C ₄ H ₈ O | 72.11 | 79.6 | -86.6 | 0.7999 | 25.6 | 18.6 | -9 |
| <i>t</i> -butyl alcohol | C ₄ H ₁₀ O | 74.12 | 82.4 | 25.7 | 0.7887 | Miscible | 12.5 | 11 |
| carbon tetrachloride | CCl ₄ | 153.82 | 76.8 | -22.6 | 1.594 | 0.08 | 2.24 | -- |
| chlorobenzene | C ₆ H ₅ Cl | 112.56 | 131.7 | -45.3 | 11.058 | 0.05 | 5.69 | 28 |
| chloroform | CHCl ₃ | 119.38 | 61.2 | -63.4 | 14.788 | 0.795 | 4.81 | -- |
| cyclohexane | C ₆ H ₁₂ | 84.16 | 80.7 | 6.6 | 0.7739 | <0.1 | 2.02 | -20 |
| 1,2-dichloroethane | C ₂ H ₄ Cl ₂ | 98.96 | 83.5 | -35.7 | 1.245 | 0.861 | 10.42 | 13 |
| diethylene glycol | C ₄ H ₁₀ O ₃ | 106.12 | 246 | -10 | 11.197 | 10 | 31.8 | 124 |
| diethyl ether | C ₄ H ₁₀ O | 74.12 | 34.5 | -116.2 | 0.713 | 7.5 | 4.267 | -45 |
| diglyme | C ₆ H ₁₄ O ₃ | 134.17 | 162 | -68 | 0.943 | Miscible | 7.23 | 67 |
| (glyme, DME) | C ₄ H ₁₀ O ₂ | 90.12 | 84.5 | -69.2 | 0.8637 | Miscible | 7.3 | -2 |
| (DMF) | C ₃ H ₇ NO | 73.09 | 153 | -60.48 | 0.9445 | Miscible | 38.25 | 58 |
| (DMSO) | C ₂ H ₆ OS | 78.13 | 189 | 18.4 | 1.092 | 25.3 | 47 | 95 |
| 1,4-dioxane | C ₄ H ₈ O ₂ | 88.11 | 101.1 | 11.8 | 1.033 | Miscible | 2.21(25) | 12 |
| ethanol | C ₂ H ₆ O | 46.07 | 78.5 | -114.1 | 0.789 | Miscible | 24.6 | 13 |
| ethyl acetate | C ₄ H ₈ O ₂ | 88.11 | 77 | -83.6 | 0.895 | 8.7 | 6(25) | -4 |
| ethylene glycol | C ₂ H ₆ O ₂ | 62.07 | 195 | -13 | 1.115 | Miscible | 37.7 | 111 |
| glycerin | C ₃ H ₈ O ₃ | 92.09 | 290 | 17.8 | 1.261 | Miscible | 42.5 | 160 |

Table 3: Properties of organic solvents (continuation)

| Solvent | formula | MW | boiling point (°C) | melting point (°C) | density (g/mL) | solubility in water (g/100g) | Dielectric Constant | flash point (°C) |
|--------------------|--|--------|-----------------------|-----------------------|-------------------|------------------------------------|------------------------|---------------------|
| heptane | C ₇ H ₁₆ | 100.20 | 98 | -90.6 | 0.684 | 0.01 | 1.92 | -4 |
| (HMPA) | C ₆ H ₁₈ N ₃ OP | 179.20 | 232.5 | 7.2 | 1.03 | Miscible | 31.3 | 105 |
| (HMPT) | C ₆ H ₁₈ N ₃ P | 163.20 | 150 | -44 | 0.898 | Miscible | ?? | 26 |
| hexane | C ₆ H ₁₄ | 86.18 | 69 | -95 | 0.659 | 0.014 | 1.89 | -22 |
| methanol | CH ₄ O | 32.04 | 64.6 | -98 | 0.791 | Miscible | 32.6(25) | 12 |
| (MTBE) | C ₅ H ₁₂ O | 88.15 | 55.2 | -109 | 0.741 | 5.1 | ?? | -28 |
| methylene chloride | CH ₂ Cl ₂ | 84.93 | 39.8 | -96.7 | 1.326 | 1.32 | 9.08 | 1.6 |
| (NMP) | CH ₃ H ₉ NO | 99.13 | 202 | -24 | 1.033 | 10 | 32 | 91 |
| nitromethane | CH ₃ NO ₂ | 61.04 | 101.2 | -29 | 1.382 | 9.50 | 35.9 | 35 |
| pentane | C ₅ H ₁₂ | 72.15 | 36.1 | -129.7 | 0.626 | 0.04 | 1.84 | -49 |
| ligroine | -- | -- | 30-60 | -40 | 0.656 | -- | -- | -30 |
| 1-propanol | C ₃ H ₈ O | 88.15 | 97 | -126 | 0.803 | Miscible | 20.1(25) | 15 |
| 2-propanol | C ₃ H ₈ O | 88.15 | 82.4 | -88.5 | 0.785 | Miscible | 18.3(25) | 12 |
| pyridine | C ₅ H ₅ N | 79.10 | 115.2 | -41.6 | 0.982 | Miscible | 12.3(25) | 17 |
| (THF) | C ₄ H ₈ O | 72.106 | 65 | -108.4 | 0.8833 | 30 | 7.52 | -14 |
| toluene | C ₇ H ₈ | 92.14 | 110.6 | -93 | 0.867 | 0.05 | 2.38(25) | 4 |
| triethyl amine | C ₆ H ₁₅ N | 101.19 | 88.9 | -114.7 | 0.728 | 0.02 | 2.4 | -11 |
| water | H ₂ O | 18.02 | 100.00 | 0.00 | 0.998 | -- | 78.54 | -- |
| water, heavy | D ₂ O | 20.03 | 101.3 | 4 | 1.107 | Miscible | ?? | -- |
| <i>o</i> -xylene | C ₈ H ₁₀ | 106.17 | 144 | -25.2 | 0.897 | Insoluble | 2.57 | 32 |
| <i>m</i> -xylene | C ₈ H ₁₀ | 106.17 | 139.1 | -47.8 | 0.868 | Insoluble | 2.37 | 27 |
| <i>p</i> -xylene | C ₈ H ₁₀ | 106.17 | 138.4 | 13.3 | 0.861 | Insoluble | 2.27 | 27 |

In order to choose the best solvent, there are basic rules:

-selectivity: a high selectivity for the compound target is desired, by this way it's not necessary more process after or before the extraction.

-capacity: the amount of solute in solvent is a important factor mainly due to reduce the amount of solvent needed to extract the solute, but in order to allow a fast extraction too, increasing the difference between the solvent an solid phase, and by this way, increasing the speed also.

-chemically inert: the reaction between the solvent and solute is not desired, because reduce the amount of both of them and produce undesired byproducts. The reaction between solvent and waste is not desired too, because new molecules produce undesired changes properties of the solution.

-highly volatile: in separation process the target is to remove determined compounds from a current, and have it in other. The extraction process just produce the first step, to make the second one it is needed a way to separate solvent and solute definitely. For this purpose evaporation is desired, being demanded for this a high difference between the volatility of both compounds.

-harmless.

-environment friendly.

-as cheap as possible: where price doesn't means just the price of the solvent but too all the features mentioned before: cost of the extraction process, cost of the separation, insurances, patents, licenses, fines...

2.4 High pressure water extraction



figure 1: water at critical point. Udo Van Hes [23]

Pressurized water is water with a pressure between 1 and 218 bars being the critical point at this pressure, and 374 °C temperatures [25]. A higher pressure and temperature, water stays as supercritical fluid, this doesn't fits in the target of the study.

Water as solvent has a very particular properties. At room temperature, water has a disproportionately high boiling pressure, high dielectric constant and polarity according to his own mass. The cause of this is the hydrogen bonded structure has a relative very high strength (5-6 Kcal/mol) due the high polarity density of the hydrogen atom. This strength is of course smaller than molecular bonding (being around 5% of the weakest bond), but it's much bigger than the Van der Waals ones [26].




Figure 2: Location of triple and critical points of water. [27]

These properties that define the solvent, and others, change along the temperature goes up, as follows.

2.4.1 Pressure:

First of all, as it's shown in [25], for biological extractions, pressure can supply mechanic strength to broke cells, but, being enough to guarantee the liquid phase of the solvent, pressure is not a factor in the chemical properties of the solvent.

In an isochoric saturated system, the pressure is function of temperature, as is shown in figure 3 [28]:




figure 3: temperature/pressure relation of saturated water as fluids definition, calculated by NIST webbook [28]

The relation between pressure and temperature lacks of significance in chemical and physical properties for the extraction results, but the combination of both defines the properties of the sample and the ambient work. In an isochoric system, as the one used in this study, the control of one (Temperature) entail the change of the second one (pressure).

2.4.2 Density:

The increasing of the temperature of a liquid fluid implies the uprising of the movement of his molecules. This agitation supplies more empty space between molecules, fact that reduce the density.

The same process of the agitation because of heat that accelerate the movement of molecules, increasing the diffusion, gives also more free place to the free circulation of these particles. By this way, an increasing of density of the liquid due a higher temperature produces an increasing of mass diffusion coefficient.




Figure 4: density of saturated liquid and gas water as fluid definition, calculated by NIST webbook [28]

Density of liquid water decreases slowly, until the evaporation begins. If the pressure is fixed enough high to guarantee the liquid phase of the solvent until the critic temperature, all this range can be used, being the density at these conditions (374°C & 218 bar) 494.53 kg/m³, 48.93% density at 0°C, 218 bar [28].

The density of liquid water variation along the temperature is relevant, but stay at the same order of magnitude.

2.4.3 Viscosity:

Viscosity has very relevant properties in diffusion process. The coefficient between viscous diffusion rate and mass diffusivity is the non-dimensional Schmidt number. Between gases, this number always round the unit, this doesn't happen with liquids.

The reduction of viscosity appoint the reduction of the energy implied in the penetration of the solvent in the solid and the output of the solute, taking the diffusion process less time in be completed, and having faster mass transfer process[29].




Figure 5: viscosity of saturated liquid and gas water as fluids definition, calculated by NIST webbook [28]

Viscosity of liquid water decreases until being close to 0 at 373°C. This decreasing goes faster at colder temperatures. The change of this property is about several orders of magnitude, and produces an important improvement of the mechanic features of the fluid in diffusion scales.

Viscosity of saturated steam can be considered null.

2.4.4 Surface tension:

Surface tension means energy needed to increase the area of a fixed volume. The fact of cross this surface implies break it and by this way an increment of the surface is needed. So great values of surface tension will hinder transfer process.

The surface tension is modified also by the temperature, following, in saturated conditions, the equation 1 [30]:

$$\sigma = B \left[\frac{T_c - T}{T_c} \right]^\mu \left[1 + b \frac{T_c - T}{T_c} \right] \quad 1$$

Being $B=235.8E-3N/m$; $b=-0.625$; $\mu=1.256$. And all temperatures in Kelvin scale.

By this way, the relation between surface tension and temperature is represented as follows:




Figure 6: surface tension of saturated liquid water, calculated with equation 1.

Surface tension decreases quasi-linearly from 0.75 to zero in the critical point for saturated liquid water.

Surface tension of gasses or supercritical fluids do not exist because their total diffusivity that define them inability the occurrence of surfaces.

2.4.5 Dielectric constant:

The dielectric constant, or permittivity, is function of temperature and pressure, as was studied in [31]. In the figure 6 it's shown the dielectric constant of saturated steam and liquid water.




Figure 7: dielectric constant of saturated liquid and gas water, performed by NIST webbook [28]

The dielectric constant decreases in liquids while temperature increases because the random spin motion given by thermal energy grows, and conceals partially the dipole moment [32]. The value of the dielectric constant defines the kind of solutes that are best dissolved, being a measurement of his polarity.

In the other hand, the dielectric constant of the saturated steam grows sparingly, from 1 to 3, and can be considered constant.

2.5 Chemical reactions in high pressure water

The purpose of these experiments is to study the allowance of pressurized water as solvent in the curcuminoids extraction from curcuma. As has been shown before, one of the most important features in the solvent selection is the lack of undesired reactions, which really happen in water at these temperatures.

First of all, consider the molecular properties of curcumin, demethoxy curcumin and bis-demethoxy curcumin will be done.




figure 8: diketo-curcumin isomer. Y. Manolova [33]




figure 10: keto-eno-curcumin isomer. Y. Manolova [33]




figure 9: dieno-curcumin isomer. Y. Manolova [33]

studied by Y. Manolova et al [33] (Figure 11).

demethoxy curcumin will be done.

Curcumin are two aromatic rings, phenyl compounds, connected by two α,β -unsaturated carbonyl groups. This can be assumed as the union of two ferulic acid molecules by their acidic group. This union means a β -dihydroxycarbonyl that has the tautomeric reaction of aldol system. By this way, three compounds can exist together in equilibrium, keto-enol-curcumin, diketo-curcumin, and dienol-curcumin.

The last one really doesn't exist, and the balance between the first ones is mainly keto-enol phase, inside solid matrix and in as solution. The ratio of diketo phase can be raised in methanol/water solution as has been




Figure 11: comparison of diketo and keto-enol curcumin isomers behavior with water molecules in methanol solution. AE calculated by Y. Manolova et Al. [33]

By this way, the solubility of curcuminoids is function of his inherent tautomeric reaction.

As has been compiled by Nermin Simsek Kus [9], the reactions that can take place in organic compounds submitted to pressurized hot water embraces almost all the organic reaction, being obviously function of the substance studied. Examples of these reactions are alkylation, condensation, coupling reactions, cyclization, decomposition, decarboxylation, dies-alder reaction, disproportionation, elimination, ene reaction, isomerization, hydrogenation-dehydrogenation, hydrolysis, oxidation, organometalic reactions and rearrangement. All these reactions can take place in specific conditions in pressurized water, which have to been fixed to arrange the optimal conditions in his production.

The lowest kinetic condition are being looked for in this experience. There are reactions that can't take place due the nonexistence of some of the reactives needed, as ene-reaction, dies-alder reaction, oxidation, coupling reactions, hydrogenation, or organometalic reactions. Others reactions won't take place due the needle of catalyst, as cyclization, dehydrogenation and hydrogenation.

Reactions that are just function of physic properties in the samples performed are alkylation, condensation, decomposition, decarboxylation, disproportionation, elimination, hydrolysis and rearrangement.

As it will be show, at high temperatures there are not curcuminoids compounds, but the antioxidant capability, issued by the phenolyc ring is bigger than samples at lower temperatures, emerging ferulic acid, paracumaric acid, 4-hydroxybenzaldehyde and vanillin. is also observable that the amount of demethoxy and bis-demethoxy compounds is bigger than curcumin along total amount of curcuminoids increases. It is reasonable understand that there are taking place reactions of hydrolysis, producing the disruption of the aldol group and the methoxy substituent, between others.

Extraction yield of specific compounds is function of the solubility of each compound in the solvent and additional facts, as common-ion effect and chelation in coordination complex, but the amount of solute is bounded by solubility. The reaction of the obtained compounds in others, obtained indirectly, with close properties can mean an increasing of these properties due the overshoot of the solubility, which still notes the limit of each component. If the solubility of the solvent fixes the limit, an increasing of the antioxidant properties of the solution can be obtained hydrolyzing the curcuminoids of a first extraction and performing a second one with a new sample. The interest of the studies is find the biggest amount of curcuminoids and

antioxidant capability per mass of solid, so the limits are really fixed by the solid, the mass transfer and the kinetic, being the solid-solvent ratio reduced if the saturation is accomplished, so this second addition of solid doesn't have interest.

Properties of interesting compounds existing in the samples after extraction are compiled in Table 4 and defined in figures 12-15:

Table 4: physical properties of products of curcuminoids decomposition. Compiled from TOXNET [9] and PUBCHEM [34] databases.

| Physical Property | Units | vanillin | p-coumaric | ferulic acid | 4-Hydroxy-benzaldehyde |
|-----------------------------------|-------------------------------|----------|------------|--------------|------------------------|
| molecular weigh | g/mol | 152.1473 | 164.15802 | 194.184 | 124.38 |
| molecular formula | (none) | C8H8O3 | C9H8O3 | C10H10O4 | C7H6O2 |
| Melting Point (760 mmHg) | °C | 81.5 | 211.5 | --- | 117 |
| Boiling Point (760 mmHg) | °C | 285 | --- | --- | 310 |
| pKa Dissociation Constant (25°C) | (none) | 7 | 4.64 | 4.58 | --- |
| log P (octanol-water) | (none) | 1,21 | 1.79 | --- | 1.35 |
| Water Solubility (25°C) | mg/L | 1.10E+03 | 1.83E+03 | 5970 | 8450 |
| Vapor Pressure (25°C) | mm Hg | 1.18E-05 | 1.61E-05 | 2.69E-06 | --- |
| Henry's Law Constant (25°) | atm-m ³ /mole | 2.15E-09 | 1.35E-12 | 7.96E-14 | --- |
| Atmospheric OH Rate Constant (°C) | cm ³ /molecule-sec | 2.73E-11 | 5.17E-11 | 4.83E-11 | --- |




figure 13: vanilin [9]




figure 12: ferulic acid [9]




figure 14: p-coumaric acid [9]




figure 15: 4-hidroxy-benzaldehyde [9]

3. Procedure

The aim of the work was to find the best time and temperature conditions for the extraction of curcuminoids compounds from curcuma. The ratio has been fixed for all experiments to 1:20 (g:mL). The autoclave was loaded with 1.5g of curcuma and 30mL of milli-Q water.

25 experiments have been performed. The pressure was set initially to 20 bars at room temperature with pressurized nitrogen, and then raised along temperature raises. Experiments have been performed changing the temperature, at 100, 125, 150, 175 and 200°C and time of extraction, 5, 10, 30, 60 and 120 minutes. The measurements obtained during the extraction procedure have been the solid mass, the empty flask mass, the flask with the solute mass. With these values the extraction yield can be obtained. The analyses carried out have been HPLC and DPPH antioxidant activity.

3.1 Material and equipment

The equipment used has been a high pressure (400 bar) high temperature (400°C) reactor and valves (Autoclave Engineers), manometer (Wika), thermometer (-50°C to 1150°C) (Greisenger), HPLC (Agilent 1100 HPL), HPLC column Agilent Eclipse XDB-C18 (150 mm × 4.6 mm; 5mm particle size), UV-Vis spectrophotometer (Varian 50 Probe).

Material used in these experiments:

Curcuma comes from Hisa Zacimb, Ljubljana, being imported from India; milli-Q water has been produced in the laboratory, dymetyl sulfoxide (Merck KGaA), acetonitryl (Sigma Aldrich), methanol (Sigma Aldrich) and acetic acid (Sigma Aldrich).

3.2 Experimental Procedure

1.5 grams of curcuma are weighed on a digital scale, and loaded in the autoclave used as extractor. 30 mL of milli-Q water are also inserted in the extractor. Once the magnetic stirrer is introduced, the autoclave is ready to be closed and heated by heating jacket. In order to avoid oxidation, oxygen is removed from the extractor loading pressurized nitrogen three times until 50 bar each one and releasing the pressure. Once the atmosphere gets inert, pressure is established with nitrogen at 20 bars. The amount of oxygen inside the extractor atmosphere is:

$$C_{O_2} = C_{O_{2atm}} \times \left(\frac{P_{atm}}{P_b}\right)^n \times \frac{P_{atm}}{P_{set}} \times 100 \cong 0.00001\% \quad 2$$



figure 16: autoclave used in the experiments

Being $C_{O_{2atm}}$ the amount of atmosphere oxygen (% volume), P_{atm} the atmosphere pressure (bar), P_b the pressure reached in the blanketing process, n the times the autoclave has been loaded, and P_{set} the pressure fixed in the extractions.

At these conditions the autoclave is heated to desired temperature, where it stays the required time, and then it is cooled with water until 30°C.

Afterwards the sample is filtrated with a vacuum system, obtaining a yellow/red/brown liquid and a darker solid waste dependent on the temperature and time applied. Existence of colloids can hinder the filtration process; these colloids appear at 150 and 175°C. To avoid this, the sample is centrifugated previously and, by this way, the main amount of solids is removed from the suspension, colloid gets

precipitated and liquid gets diluted, so the filtration is easier.

The extract is introduced in a vacuum evaporator that is initially set at 50 mBar, being this pressure enough to evaporate water without bubbles that drag the extract and lately, when there is not possibility of bubbles, at 30 mBar until the extract is absolutely dry. Therefore water is absolutely removed, obtaining at the end the dry solute in the spherical flask, which has been weighed before. The difference between the empty flask and flask with solute is the amount of

solute extracted from the curcuma treated. The content of curcuminoids in extract and antioxidant activity of extract was analyzed by different methods, described in the following subchapter.

Samples are stored in freezer until the analysis.

3.3 Analytical procedures

Two kinds of analysis have been done, a DPPH antioxidant activity essay, and HPLC.

3.3.1 HPLC analysis of curcuminoids

20 mg of solute are added in 10 mL flask, and then are diluted with dimethyl sulfoxide, DMSO, due the presence of starch in the solute, insoluble in water or methanol at room temperature.

The analytical procedure has been directly taken from T. Perko et al, (2015) Isolation, characterization and formulation of curcuminoids and in vitro release study of the encapsulated particles. *The Journal of Supercritical Fluids*, 103, pp.48-54. [3]:

“The extracts were analyzed by HPLC, using the method described by Lee and Choung:

The Agilent 1100 HPLC system consisted of a binary pump, column heater, auto sampler and variable wavelength detector (VWD). The separation was achieved on chromatographic column Agilent Eclipse XDB-C18 (150 mm × 4.6 mm; 5mm particle size). The mobile phases were 2% acetic acid in water (elution A) and 2% acetic acid in acetonitrile (elution B). The solvent gradient was as follows: 0–3 min, 10% B; 8 min, 20% B; 13 min, 25% B; 18 min, 35% B; 28–33 min, 55% B and then held for 3 min before returning to initial conditions. The solvent flow rate was 1.0 mL/min and the column temperature was 30°C. The volume of injection was 10 µL and peaks were monitored at 420 nm. Quantification of single curcuminoids was done using calibration curves obtained from curcuminoid standards. All measurements were performed in triplicate and averages were calculated.”

3.3.2 DPPH analysis

10 mg of solute are added in 10 mL flask, and diluted with dimethyl sulfoxide assisted by ultrasound. The method used has been the same that described by T. Perko et al. [3].

“DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity was measured using the UV-VIS spectrophotometric method [23]. [...] To a dark flask containing 77 µL of extract solution, 3 mL of DPPH solution was added. The mixed solutions were kept in the dark for 15

min at room temperature and the absorbance was measured using UV-VIS spectrophotometer (Varian, USA) at 515 nm. Radical-scavenging activities of samples were calculated using:

$$\%inhibition = \frac{A_B - A_A}{A_B} \cdot 100 \quad 3$$

Where A_B is absorbance of blank sample ($t = 0$ min) and A_A is absorbance of extract solution ($t = 15$ min). All measurements were performed in triplicate and averages were calculated.”

Samples get frozen in the fridge between the preparation of solutions and analysis due the melting point of DMSO.

4. Results

4.1 Extract yield

The extract yield has been calculated according the equation 4.

$$E = \frac{m_e - m_f}{m_c} \times 100 \quad 4$$

Being E the extract yield, m_e the mass of the flask and the extract after the evaporation process, m_f the mass of the empty flask and m_c the mass off curcuma filled in the autoclave.

The amount of extract obtained in different times and temperatures, obey the figure 17.




figure 17: plot of extraction yield at different times and temperatures. Values of experiments done twice figure as average.

Extraction yield increases with increasing temperature until 175°C. The yield increases with time at lower temperatures, being practically constant at 175°C, and decreasing with increasing time at 200°C.

The extract yield at 175°C has three local maximums and two local minimums in a five points function. The range between global maximum and minimum is less than 10%, so the yield has been considered constant at this temperature, being 71%.

The extract yield at 150°C, 120 minutes is lower (31%) than the expected one, being this one outside of the results after 120 minutes between temperatures at 125°C (56%) and 175°C (76%). This lack of coherence may be caused by the existence of starch in the sample, that hampers the filtration, bringing about the need of various filters instead one, even though a previous centrifugation. The analysis was also hindered due the low solubility of starch in water or methanol, being used dimethyl sulfoxide as solvent to avoid this problem.

4.2 Curcuminoids in extract

The amount of curcuminoids in each sample is shown in figure 18.




figure 18: total curcuminoids extracted for each time and temperature and time.

It can be seen that the highest quantity of curcuminoids is obtained at the lowest temperatures, 100 and 125°C, being zero at temperatures higher than 150°C. A reduction of the mass of curcuminoids can be seen at 125°C and 150°C while time increase, so curcuminoids are degraded at temperatures higher than 125°C.

As has been shown before, in methanol solution, curcuminoids are almost in keto-enol isomer, being the amount diketo almost null. These substances are always in equilibrium, so during the separation process that the HPLC entail, they are together, taking place instantly the tautomeric reaction in other case. The relation between the mobile phases is monotone, so differentiable peaks for each isomer are not expected.




figure 19: from the top to the bottom, results of HPLC for extractions at rising temperatures (100, 125, 150, 175 and 200°C) after 30 minutes. Curcuminoids are the peaks at 26 to 28 min.

figure 19 shows the behavior of samples obtained at the same time of extraction (30 min) at different temperatures. Curcuminoids are compounds whose outflow takes place after 25 minutes. By increasing the temperature of extraction the decomposition of curcuminoids occurs, and new substances are formed, as described before: vanillin, p-coumaric acid, ferulic acid and 4-Hydroxy-benzaldehyde, at 21-22, 13-15 and 7-9 minutes.

4.3 Antioxidant capability

Results of antioxidant capability of extract are given in figure 20.




figure 20: Antioxidant capability, % inhibition for all temperatures and times.

The antioxidant capability is shown as % inhibition, being determined by DPPH method, where the free radicals of DPPH have to be scavenged by the antioxidants acting in the oxidative process.

The figure 20 shows a low level of antioxidant capability for all samples but the ones obtained at 200°C. The antioxidant activity of extracts obtained at 200°C increases with time of extraction until 18% after 120 minutes of extraction.

Extractions at 100°C, 30 min and 175°C, 30 min have an antioxidant activity measurement zero value, value that is not expected. These results have to be taken as mistakes.

Color and bright

A quality measurement has been taken. Figure 21 shows the color of each sample.

| Time/temp | 100 | 125 | 150 | 175 | 200 |
|-----------|----------|-----|-----|-----|-----|
| 005 | | | | | |
| 010 | XXXXXXXX | | | | |
| 030 | | | | | |
| 060 | | | | | |
| 120 | | | | | |

figure 21: color for each sample.

The yellow color is brighter at lower temperatures, specifically at 125°C and short times, this agrees with the curcuminoids analysis.

More particular explanations will be done for each temperature. A further qualitative description is also given.

4.5 Results of extractions at 100°C




figure 22: Extraction yield at 100°C, single values and averages (◆), errors (—).

At the lowest temperature, 100°C, the mixture obtained in the extractor is a bright yellow solution, good smell, and can be filtrated without problems, obtaining a dry porous solid and a yellow solution.

At this temperature, the maximum yield of extraction is obtained in short times. After one hour of extraction the yield doesn't increase any more. The extraction yield is low, never higher than 30%.




figure 23: HPLC & DPPH results for 100°C extractions at 5 min (1), 10 min (2) 30 min (3) 60 min (4) and 120 min (5)

The amount of curcuminoids increases with time of extraction, until close to 30 mg/g of extract (the amount of curcuminoids in turmeric is between 3 and 5%). Bis-demethoxy curcumin is the most abundant in all extracts. The absorbance maximum value of absorbance is 4% (10 minutes and 120 minutes). There is a direct relation between curcuminoids content and antioxidant capability.

4.6 Results of extractions at 125°C




figure 24: Extraction yield at 125°C, single values and averages (■), errors (■).

At 125°C the extracts are more or less the same in appearance and smell as 100°C extract. At 125°C and 30 min there were problems with the filtration; the flow in the filter was close to zero and was zero after few seconds. To avoid this, the solution is centrifuged previously (two minutes 11000 rpm) to remove the colloid that obstructs the filter. After centrifugation the flow decreases with time, but doesn't stop before the solid is dry.

At 125°C the extraction yield increases with time, being the maximum around 60% and almost constant after one hour. Five minutes at this temperature is enough to obtain the same extract as the optimum at 100°C, after 60 minutes of extraction.

The content of curcuminoids is the highest at 30 minutes of extraction and afterwards it decreases with time what shows that at this temperature decompositions start to take place. The composition of the sample after 30 minutes is consistent with the fact that curcumin is the most abundant component, but it loses the methoxy radical to become demethoxy-curcumin or bis-demethoxy-curcumin; with time, all of them are decomposed to easier compounds.




figure 25: HPLC & DPPH results for 125°C extractions at 5 min (1), 10 min (2) 30 min (3) 60 min (4) and 120 min (5)

It's also observable that the amount of demethoxy curcumin and bi-demethoxy curcumin remain constant during the first 30 minutes, and just the amount of curcumin increases. It can be understandable that the mass transfer of the last one is slower than the DMC and BDMC. The antioxidant activity rises until 3% after one hour of extraction.

4.5 results of extraction at 150°C




figure 26: Extraction yield at 150°C, single values and averages (▲), errors (■).

At 150°C there are again problems with filtration. To avoid this, the sample is centrifugated for 6 minutes at 11000 spins per minute. Anyway, the filtration requires a lot of time (20-30 minutes for 60 mL of solution), and news filters (up to 6 in the worst situation, what causes high losses of extract. The appearance of the mixture is yellow too, but more brown or red than at 125°C. The smell is still good.

The yield of extraction is approximately constant in time period from 5 to 60 minutes (60%) and afterwards it decreases to approximately 30% after 120 minutes of extraction. The big difference between errors of samples at 10 minutes is due the difficulties in the filtration.




figure 27: HPLC & DPPH results for 150°C extractions at 5 min (1), 10 min (2) 30 min (3) 60 min (4) and 120 min (5)

The amount of curcuminoids in extract is smaller than at lower temperatures, reaching zero values after 60 min. The amount of curcumin is almost zero or zero in all the samples, and there is just bis-demethoxy curcumin after 30 minutes of extraction, so the decomposition of the methoxy groups is also expected at 150°C. On the other hand, the antioxidant capability rises with time, while curcuminoids amount in extract decreases.



figure 28: stratification of the solid residue after centrifugation

After the centrifugation, the solid is stratified, with a big amount of brown substance and a thin layer of brighter yellow material.

4.5 results of extraction at 175°C




figure 29:Extraction yield at 175°C, single values and averages (X), errors (T).

At 175°C and short times of extraction until 10 minutes, the properties of products are close to the ones obtained at 150°C and longer times; a yellow to brown mixture is obtained that is bad filterable also after centrifugation. If the time is 30 minutes or higher, the mixture starts to be brown, and it's easy filterable, so the sample is not centrifugated. The smell is lightly unpleasant, and softer than of the ones samples obtained at lower temperatures.

Extraction yield at 175°C is almost constant with time (around 70%), and is also the biggest.




figure 30: HPLC & DPPH results for 175°C extractions at 5 min (1), 10 min (2) 30 min (3) 60 min (4) and 120 min (5)

The amount of curcuminoids is zero regardless of extraction time, what indicates that total decomposition of curcuminoids has taken place even in the shortest times. Antioxidant activity

reaches low values of 5% after 1 or 2 hours. The antioxidant activity result is zero if time is 30 min, this unexpected result is probably an experimental error.



10 min 175°C

60 min 175°C

figure 31: comparison of color of samples at 175°C, 10 and 60 minutes

The color of the solid waste after filtration is absolutely different after 10 to 60 minutes of extraction due to the Maillard reactions that occur at these temperatures with organic compounds. The waste obtained after 10 minutes 175 °C samples is yellow, but doesn't have the bright as samples at lower temperatures had. The Maillard's reaction has started after 10 minutes, but his effects are not so visual as after 60.

4.9 Results of extractions at 200°C




figure 32:Extraction yield at 200°C, single values and averages (■), errors (■).

The properties of the product obtained at 200°C are close to the ones at 175°C, The color strength obtained after the filtration is lower than of the product obtained at 175°C. The smell is unpleasant too, more deep than of the previous one, but still softer than of the samples obtained at 150°C and lower temperatures, and has a toasted connotation. The extract after evaporation is not a solid, but an oily substance.

The extraction yield is high, not so high as at 175°C, but close to 70%, at short times of extraction. Extraction yield decreases after 30 and reaches being 40% after two hours.




figure 33: HPLC & DPPH results for 200°C extractions at 5 min (1), 10 min (2) 30 min (3) 60 min (4) and 120 min (5)

Similar as at 175°C, all the curcuminoids have been degraded to simpler compounds, while the antioxidant activity increases with time. At this temperature the antioxidant activity increases until 18%, what is more than three times higher as at lower temperatures. This value is still lower than the values obtained with other extraction procedures (24% for the best supercritical CO₂ extraction, 35% for traditional extraction with ethanol) [3].

5. Conclusions:

175°C is high enough temperature to degrade the total amount of curcuminoids already in 5 minutes. At 150°C the same result is obtained after 30 minutes of extraction.

The degradation of curcuminoids can be considered zero at 100°C in pressurized water.

The best conditions for curcuma extraction are 125°C and 60 min, in order to obtain the maximum amounts of curcuminoids; or 125°C, 30 min to have the most concentrated extract. These maximums are 10.2mg/g curcuma (125°, 60 min) and 19.53mg/g extract (125, 30 min). The best values of antioxidant activity are obtained at the highest temperature, after 1 and 2 hours, being 18%. All these values are lower than those obtained by the traditional extraction with ethanol. This fact reveals this method as not the best one for this process.

6.Literature

- [1] Benzie, I. and Wachtel-Galor, S. (2011). *Herbal medicine*. Boca Raton: CRC Press.
- [2] Anand, P., Kunnumakkara, A., Newman, R. and Aggarwal, B. (2007). Bioavailability of Curcumin: Problems and Promises. *Molecular Pharmaceutics*, 4(6), pp.807-818.
- [3] Perko, T., Ravber, M., Knez, Ž. and Škerget, M. (2015). Isolation, characterization and formulation of curcuminoids and in vitro release study of the encapsulated particles. *The Journal of Supercritical Fluids*, 103, pp.48-54.
- [4] Tapal, A. and Tiku, P. (2012). Complexation of curcumin with soy protein isolate and its implications on solubility and stability of curcumin. *Food Chemistry*, 130(4), pp.960-965.
- [5] on colours for use in foodstuff. (1994). 1st ed. [ebook] Oficial Journal of European Activities, pp.L237\13. Available at: http://ec.europa.eu/food/fs/sfp/addit_flavor/flav08_en.pdf [Accessed 12 Jun. 2015].
- [6] Brahmachari, G. (n.d.). *Chemistry and pharmacology of naturally occurring bioactive compounds*.
- [7] Prasad, S., Tyagi, A. and Aggarwal, B. (2014). Recent Developments in Delivery, Bioavailability, Absorption and Metabolism of Curcumin: the Golden Pigment from Golden Spice. *Cancer Research and Treatment*, 46(1), pp.2-18.
- [8] Sharma, R., Gescher, A. and Steward, W. (2005). Curcumin: The story so far. *European Journal of Cancer*, 41(13), pp.1955-1968.
- [9] Toxnet.nlm.nih.gov, (2015). *TOXNET*. [online] Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search2/f?/.temp/~IVlcxR:1> [Accessed 12 Jun. 2015].
- [10] O'Neil, M.J. (ed.). *The Merck Index - An Encyclopedia of Chemicals, Drugs, and Biologicals*. Cambridge, UK: Royal Society of Chemistry, 2013., p. 474
- [11] CRC Handbook of Chemistry and Physics: A Ready-Reference of Chemical and Physical Data, 85th ed Edited by David R. Lide (National Institute of Standards and Technology). CRC Press LLC: Boca Raton, FL. 2004. 2712 ISBN 0-8493-0485-7. (2005). *J. Am. Chem. Soc.*, 127(12), pp.4542-4542.
- [12] US EPA; Estimation Program Interface (EPI) Suite. Ver. 4.1. Nov, 2012
- [13] Book Listing of CRC Handbook of Chemistry and Physics, 92nd ed., 2011–2012 CRC Handbook of Chemistry and Physics, 92nd ed., 2011–2012 . Editor-in-Chief: W. M.Haynes (National Institute of Standards and Technology). CRC Press (an imprint of Taylor & Francis Group) : Boca Raton, FL . 2011 . 2656 ISBN 978-1-4398-5511-9 . (2011). *J. Am. Chem. Soc.*, 133(34), pp.13766-13766.
- [14] Socaciu, C. (2007). *Food colorants*. Boca Raton: Taylor & Francis.

- [15] Merory, J. Food Flavorings: Composition, Manufacture, and Use
- [16] Fenaroli, G. and Burdock, G. (1994). *Fenaroli's handbook of flavor ingredients*. Boca Raton: CRC Press.
- [17] Jörg Bart, H. (2015). *Extraction of Natural Products from Plants – An Introduction*. 1st ed. [ebook] pp.1-5. Available at: http://www.wiley-vch.de/books/sample/3527325042_c01.pdf [Accessed 28 Apr. 2015].
- [18] Alupupui A. Calinescu I. Lavric V. Ultrasonic vs. Microwave extraction intensification of active principles from medicinal plants.
- [19] Mesa J.L. capitulo 2, fluidos supercríticos.
- [20] SUPERCRITICAL FLUID EXTRACTION. (2010). 1st ed. [ebook] Osmadabad, pp.729-732. Available at: [http://www.sadgurupublications.com/ContentPaper/2010/3_8\(2\)2010\).pdf](http://www.sadgurupublications.com/ContentPaper/2010/3_8(2)2010).pdf) [Accessed 12 Jun. 2015].
- [21] Van Laar , J.J. (1990) Z. Phys. Chem. 723
- [22] Hildebrand , J.M. and Scott , R.L. (1950) The Solubility of Nonelectrolytes , 3rd edn , Reinhold , New York, USA .
- [23] Common Organic Solvents: Table of Properties This table was orginally from: Prof. Murov's Orgsoltab, which was edited and reposted by Erowid
- [24] Critical point of water Udo van Hes universty of amsterdamm
- [25] Extraction with superheat water, Roger M. Smith
- [26] A. D. McNaught and A. Wilkinson IUPAC golden book hydrogen bond *in theoretical organic chemistry*
- [27] Chemguide.co.uk, (2015). *phase diagrams of pure substances*. [online] Available at: <http://www.chemguide.co.uk/physical/phaseeqia/phasediags.html> [Accessed 8 Jun. 2015].
- [28] Webbook.nist.gov, (n.d.). *Thermophysical Properties of Fluid Systems*. [online] Available at: <http://webbook.nist.gov/chemistry/fluid/> [Accessed 26 Apr. 2015].
- [29] Teo, C., Tan, S., Yong, J., Hew, C. and Ong, E. (2010). Pressurized hot water extraction (PHWE). *Journal of Chromatography A*, 1217(16), pp.2484-2494.

- [30] NIST, (2015). *international tables of surface tension of water*. [online] Available at: <http://www.nist.gov/data/PDFfiles/jpcrd231.pdf> [Accessed 2 May 2015].
- [31] static dielectric constant of water and steam. (2015). 1st ed. [ebook] NIST. Available at: <http://www.nist.gov/data/PDFfiles/jpcrd172.pdf> [Accessed 2 May 2015].
- [32] Doitpoms.ac.uk, (2015). *DoITPoMS - TLP Library Dielectric materials - Effect of temperature on the dielectric constant*. [online] Available at: <http://www.doitpoms.ac.uk/tlplib/dielectrics/temperature.php> [Accessed 1 May 2015].
- [33] Manolova, Y., Deneva, V., Antonov, L., Drakalska, E., Momekova, D. and Lambov, N. (2014). The effect of the water on the curcumin tautomerism: A quantitative approach. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 132, pp.815-820.
- [34] Pubchem.ncbi.nlm.nih.gov, (2015). *The PubChem Project*. [online] Available at: <http://pubchem.ncbi.nlm.nih.gov/> [Accessed 11 Jun. 2015].

7. Appendix

Table 5: extraction yield of all samples

| sample | curcuma mass | empty flask | extract and flask | extract | extract yield |
|--------------|--------------|-------------|-------------------|---------|---------------|
| 100°C;5min | 1.5006 | 167.5599 | 167.6616 | 0.1017 | 6.78 |
| 100°C;10min | 1.5038 | 181.7532 | 182.047 | 0.2938 | 19.54 |
| 100°C;30min | 1.5002 | 167.3875 | 167.7069 | 0.3194 | 21.29 |
| 100°C;60min | 1.5016 | 181.7532 | 182.1467 | 0.3935 | 26.21 |
| 100°C;60min | 1.4999 | --- | --- | 0.3937 | 26.25 |
| 100°C;120min | 1.5007 | 167.3871 | 167.855 | 0.4679 | 31.18 |
| 100°C;120min | 1.5007 | --- | --- | 0.3072 | 20.47 |
| 125°C;5min | 1.5007 | --- | --- | 0.3072 | 20.47 |
| 125°C;5min | 1.5001 | --- | --- | 0.4828 | 32.18 |
| 125°C;10min | 1.5002 | 167.2853 | 167.91 | 0.6247 | 41.64 |
| 125°C;30min | 1.5009 | 167.3895 | 168.0762 | 0.6867 | 45.75 |
| 125°C;60min | 1.5044 | 167.2853 | 168.224 | 0.9387 | 62.40 |
| 125°C;120min | 1.4997 | 172.7764 | 173.6096 | 0.8332 | 55.56 |
| 150°C;5min | 1.499 | 167.3854 | 168.2195 | 0.8341 | 55.64 |
| 150°C;10min | 1.4991 | 167.3854 | 167.9308 | 0.5454 | 36.38 |
| 150°C;10min | 1.5006 | 225.56 | 226.67 | 1.11 | 73.97 |
| 150°C;30min | 1.5013 | 167.3897 | 168.2216 | 0.8319 | 55.41 |
| 150°C;60min | 1.5013 | 167.3897 | 168.3164 | 0.9267 | 61.73 |
| 150°C;120min | 1.5003 | 173.5291 | 173.9912 | 0.4621 | 30.80 |
| 175°C;5min | 1.5 | 167.3858 | 168.5022 | 1.1164 | 74.43 |
| 175°C;10min | 1.4994 | 167.3886 | 168.3854 | 0.9968 | 66.48 |
| 175°C;30min | 1.4992 | 167.3886 | 168.3443 | 0.9557 | 63.75 |
| 175°C;30min | 1.5001 | 167.389 | 168.5378 | 1.1488 | 76.58 |
| 175°C;30min | 1.5001 | 167.389 | 168.5378 | 1.1488 | 76.58 |
| 175°C;60min | 1.4997 | 172.6738 | 173.6885 | 1.0147 | 67.66 |
| 175°C;120min | 1.5007 | 167.3862 | 168.5289 | 1.1427 | 76.14 |
| 200°C;5min | 1.4982 | 167.3886 | 168.321 | 0.9324 | 62.23 |
| 200°C;10min | 1.4991 | 167.3886 | 168.3675 | 0.9789 | 65.30 |
| 200°C;30min | 1.5003 | 167.3871 | 168.3126 | 0.9255 | 61.69 |
| 200°C;30min | 1.5 | 167.3882 | 168.3776 | 0.9894 | 65.96 |
| 200°C;60min | 1.5024 | 169.3673 | 170.0636 | 0.6963 | 46.35 |
| 200°C;120min | 1.5002 | 167.3886 | 168.1408 | 0.7522 | 50.14 |
| 200°C;120min | 1.5034 | 167.3896 | 167.9331 | 0.5435 | 36.15 |

Table 6: HPLC results: mass of curcuminoids

| sample | HPL mass | BDMC area | DMC area | C area | BDMC mg/ g ext. | BMC mg/g ext. | C mg/ g ext. | total mg/g ext. |
|--------------|----------|-----------|----------|--------|-----------------|---------------|--------------|-----------------|
| 100°C;5min | 20.44 | 114.05 | 115.45 | 243.90 | 3.53 | 3.54 | 1.19 | 8.27 |
| 100°C;10min | 20.13 | 275.30 | 266.90 | 615.35 | 5.68 | 5.64 | 5.43 | 16.75 |
| 100°C;30min | 20.09 | 107.95 | 96.85 | 180.20 | 3.45 | 3.29 | 0.47 | 7.20 |
| 100°C;60min | 19.86 | 321.30 | 297.15 | 580.90 | 6.30 | 6.06 | 5.04 | 17.39 |
| 100°C;60min | 17.88 | 216.30 | 192.05 | 365.30 | 4.90 | 4.60 | 2.58 | 12.08 |
| 100°C;120min | --- | --- | --- | --- | --- | --- | --- | --- |
| 100°C;120min | 21.35 | 642.20 | 554.75 | 923.55 | 10.58 | 9.62 | 8.95 | 29.15 |
| 125°C;5min | 23.43 | 164.20 | 174.90 | 480.15 | 4.20 | 4.36 | 3.89 | 12.46 |
| 125°C;5min | --- | --- | --- | --- | --- | --- | --- | --- |
| 125°C;10min | 26.1 | 188.00 | 206.50 | 593.70 | 4.52 | 4.80 | 5.19 | 14.51 |
| 125°C;30min | 22.07 | 267.50 | 300.60 | 827.35 | 5.58 | 6.10 | 7.85 | 19.54 |
| 125°C;60min | 24.03 | 237.10 | 257.25 | 641.20 | 5.17 | 5.50 | 5.73 | 16.41 |
| 125°C;120min | 22.86 | 174.10 | 161.40 | 291.80 | 4.33 | 4.18 | 1.74 | 10.25 |
| 150°C;5min | 21.44 | 92.65 | 79.95 | 177.95 | 3.25 | 3.05 | 0.44 | 6.74 |
| 150°C;10min | 10.8 | 35.75 | 32.55 | 65.75 | 2.49 | 2.40 | 0.00 | 4.88 |
| 150°C;10min | 23.84 | 185.75 | 169.10 | 335.75 | 4.49 | 4.28 | 2.24 | 11.01 |
| 150°C;30min | 22.12 | 20.50 | 0.00 | 0.00 | 2.28 | 0.00 | 0.00 | 2.28 |
| 150°C;60min | 21.73 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 150°C;120min | 22.09 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;5min | 21.07 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;10min | 12.16 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;30min | --- | --- | --- | --- | --- | --- | --- | --- |
| 175°C;30min | 26.27 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;30min | 23.69 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;60min | 26.44 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 175°C;120min | 28.62 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;5min | 20.55 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;10min | 23.14 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;30min | 27.62 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;30min | 26.85 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;60min | 22.29 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;120min | 24.5 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 200°C;120min | 23.63 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |

Table 7: DPPH results. antioxidant capability.

| | anti-oxidative mass | valor | % inhibición |
|--------------|---------------------|--------|--------------|
| 100°C;5min | 10.1 | 0.8335 | 2.95 |
| 100°C;10min | 9.31 | 0.824 | 4.05 |
| 100°C;30min | 11.25 | 0.8575 | 0.15 |
| 100°C;60min | 9.98 | 0.8339 | 2.90 |
| 100°C;60min | 10.44 | 0.8367 | 2.57 |
| 100°C;120min | --- | --- | --- |
| 100°C;120min | 10.88 | 0.8261 | 3.81 |
| 125°C;5min | 9.91 | 0.8565 | 0.27 |
| 125°C;5min | --- | --- | --- |
| 125°C;10min | 11.56 | 0.8445 | 1.67 |
| 125°C;30min | 10.17 | 0.841 | 2.07 |
| 125°C;60min | 11.29 | 0.8332 | 2.98 |
| 125°C;120min | 11.4 | 0.8382 | 2.40 |
| 150°C;5min | 9.59 | 0.8471 | 1.36 |
| 150°C;10min | 10.02 | 0.8448 | 1.63 |
| 150°C;10min | 10.48 | 0.8236 | 4.10 |
| 150°C;30min | 12.13 | 0.8352 | 2.75 |
| 150°C;60min | 10.73 | 0.821 | 4.40 |
| 150°C;120min | 13.58 | 0.8197 | 4.55 |
| 175°C;5min | 11.82 | 0.8471 | 1.36 |
| 175°C;10min | 9.09 | 0.8329 | 3.02 |
| 175°C;30min | --- | --- | --- |
| 175°C;30min | 10.1 | 0.8771 | -2.13 |
| 175°C;30min | --- | --- | --- |
| 175°C;60min | 10.69 | 0.8205 | 4.46 |
| 175°C;120min | 10.16 | 0.8173 | 4.83 |
| 200°C;5min | 9.78 | 0.8243 | 4.02 |
| 200°C;10min | 10.23 | 0.8265 | 3.76 |
| 200°C;30min | 9.76 | 0.7542 | 12.18 |
| 200°C;30min | 10.11 | 0.7659 | 10.82 |
| 200°C;60min | 10.1 | 0.7105 | 17.27 |
| 200°C;120min | 9.88 | 0.7237 | 15.73 |
| 200°C;120min | 12.78 | 0.6731 | 21.62 |