

Marquette University
e-Publications@Marquette

College of Professional Studies Faculty Research
and Publications

Professional Studies, College of

1-1-2012

Child Health Providers' Precautionary Discussion of Emotions During Communication about Results of Newborn Genetic Screening

Michael Farrell
Medical College of Wisconsin

Jodi Speiser
Loyola University Chicago

Lindsay Deuster
Medical College of Wisconsin

Stephanie Christopher
Marquette University, stephanie.christopher@marquette.edu

Archives of Pediatrics & Adolescent Medicine, Volume 166, No.1, pp 62-67 (January, 2012). DOI. ©
2012 American Medical Association. Used with permission.

Marquette University

e-Publications@Marquette

Professional Studies Faculty Research and Publications/School of Professional Studies

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Archives of Pediatrics & Adolescent Medicine, Vol. 166, No. 1 (January, 2012): 62-67. [DOI](#). This article is © American Medical Association and permission has been granted for this version to appear in [e-Publications@Marquette](#). American Medical Association does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from American Medical Association.

Child Health Providers' Precautionary Discussion of Emotions During Communication About Results of Newborn Genetic Screening

Michael H. Farrell

Center for Patient Care and Outcomes Research, Medical College of Wisconsin, Milwaukee, WI

Jodi Speiser

Department of Pathology, Loyola University Medical Center, Chicago, IL

Lindsay Deuster

Department of Pediatrics, University of Minnesota, Minneapolis, MN

Stephanie Christopher

Center for Patient Care and Outcomes Research, Medical College of Wisconsin, Milwaukee, WI

Abstract

Objective To demonstrate a quantitative abstraction method for Communication Quality Assurance projects to assess physicians' communication about hidden emotions after newborn genetic screening.

Design Communication quality indicator analysis.

Setting Standardized parent encounters performed in practicing physicians' clinics or during educational workshops for residents.

Participants Fifty-nine pediatrics residents, 53 pediatricians, and 31 family physicians.

Intervention Participants were asked to counsel standardized parents about a screening result; counseling was recorded, transcribed, and parsed into statements (each with 1 subject and 1 predicate). Pairs of abstractors independently compared statements with a data dictionary containing explicit-criteria definitions.

Outcome Measures Four groups of "precautionary empathy" behaviors (assessment of emotion, anticipation/validation of emotion, instruction about emotion, and caution about future emotion), with definitions developed for both "definite" and "partial" instances.

Results Only 38 of 143 transcripts (26.6%) met definite criteria for at least 1 of the precautionary empathy behaviors. When partial criteria were counted, this number increased to 80 of 143 transcripts (55.9%). The most common type of precautionary empathy was the "instruction about emotion" behavior (eg, "don't be worried"), which may sometimes be leading or premature.

Conclusions Precautionary empathy behaviors were rare in this analysis. Further study is needed, but this study should raise concerns about the quality of communication services after newborn screening.

The quality of parent-physician communication after newborn screening is an important area of concern. Communication influences patients' understanding and psychological outcomes¹ and may be crucial when screening reveals that an infant is a heterozygous "carrier" for cystic fibrosis (CF) or sickle cell hemoglobinopathy (SCH). Carrier infants are healthy, but some say that such infants have a "nondisease"² because of the possibility for misconceptions and psychosocial complications.³⁻¹² Such problems have been cited in arguments against the routine use of genetic and molecular screening technologies.¹³⁻¹⁵ One of us (M.H.F.) has counter-argued that newborn screening could result in "more good than harm" if psychosocial risks were managed over the same statewide populations served by the screening programs.¹⁶ This article demonstrates the next step in our long-term effort to develop Communication Quality Assurance for population-scale use; it uses the new tool to assess a pilot sample for communication about the potential for emotions after newborn screening.

Emotions are an important topic for communication after newborn screening. Studies after various types of genetic screening have documented that nondisease results can be followed by clinical levels of anxiety and depression, as well as other stress-related problems, such as physical symptoms, feelings of shock, sleep disturbances, reports of infant feeding problems, and uncertainty about the future.^{7-9,11,12} In other areas of health care, patients may be more likely to reveal their emotions and discuss them if the physician initiates the topic of emotions, but they may never reveal their emotions if they are not asked.¹⁷⁻¹⁹ Physicians may not notice their patients' emotions²⁰⁻²² or may fail to acknowledge them.²³ Failure to address emotions is problematic because emotions may be distressing or may impede patients' ability to process information.

For this article, we chose to examine physicians' communication about emotions that may be present but are not apparent in the patient's speech, facial expressions, or body language. Addressing potential emotions is important because patients' facial expressions do not always correlate with their emotional states for cultural reasons or because of a sense of shock.^{24,25} We developed the term *precautionary empathy* for these communication behaviors to distinguish between physician behaviors that react to obvious emotions and to avoid confusion with the broader term *empathy*. Many definitions of empathy have been proposed,²⁶⁻³⁰ and many scales³¹⁻³³ to measure empathy exist, but generally speaking, empathy can be defined as "identification with and understanding of another's situation, feelings, and motives."³⁴ We also coined the new term *precautionary empathy* to avoid confusion in the medical education field, where *empathy* has sometimes been used as a shorthand term for health care providers' responsiveness to emotions.³⁵⁻³⁷ A list of the precautionary empathy behaviors is shown in [Table 1](#) and further described in the "Types of Precautionary Empathy Behaviors" subsection in the "Methods" section.

Table 1. Precautionary Empathy Behaviors

Type	Precautionary Empathy Behavior	Example Quote
1	Assess for prior emotion	How are you feeling about the newborn screening process?
	Assessment for emotion (close-ended)	Are you feeling anxious about this?
	Assessment for emotion (open-ended)	What feelings are you experiencing right now?
2	Anticipate/validate emotion	Many people get sad when they hear this kind of news.
3	Instruction about emotion	Don't cry.
4	Caution about future emotion	When you think more about the implications of this result, you might get more worried.

A method to assess physicians' communication about the potential for emotions informs our larger efforts to improve physician communication quality. This effort, called Communication Quality Assurance, attempts to improve physicians' communication by providing quantitatively reliable assessment of communication behaviors. Our panel of communication behaviors is not intended to be exhaustively complete,³⁸⁻⁴⁶ but the theoretical framework for Communication Quality Assurance holds that each behavior's usage may incrementally improve the chance of favorable outcomes. Focusing on communication about emotion is a key component of this goal.

Method

Design

As part of a larger effort,³⁸⁻⁴⁶ we reviewed the literature to develop an explicit criteria data dictionary of communication behaviors dealing with emotions. We then used the dictionary to abstract transcripts of physicians counseling standardized parents about newborn screening results. Methods were approved by institutional review boards at Yale University and the Medical College of Wisconsin.

Types of precautionary empathy behaviors

Our literature review examined communication guidelines and other sources.^{36,37,47-60} We identified 8 behaviors consistent with our concept for precautionary empathy, but overlap between the behaviors

prompted us to group them into 4 “types” (Table 1). As with other communication quality indicators, the 4 types of communication quality indicators represent independent behaviors, each of which may also have varying degrees of usefulness for different situations.

Assessment for Emotion

The most elementary of the behaviors was for the physician to explicitly ask about emotion. Guidelines suggest that an open-ended question (“How do you feel right now?”) would be more effective at obtaining an accurate answer than a closed-ended question (“Are you worried?”) because the natural answer to the open-ended version would be more detailed than yes or no. Closed-ended questions may be particularly ineffective if a leading syntax is used (“You’re not worried, are you?”).

The Assessment for Emotion behavior may also vary by its timing relative to the emotion. For example, a physician could ask a general question (“Are you worried about all of this?”) or ask whether the patient is worried by the conversation itself (“How is this news making you feel?”). Finally, it may be wise to inquire about emotions from before the conversation (“Were you worried when my nurse asked you to come to my office to talk about this test?”). To facilitate abstraction, we developed criteria for 3 quality indicators: an Assessment of Prior Emotion behavior, an Assessment for Emotion (open-ended), and an Assessment for Emotion (closed-ended).

Anticipate or Validate Emotion

In this behavior, a physician reassures the patient that it is normal to have emotions like anxiety or fear or that emotions are nothing to be ashamed about. The patient may then be more likely to reveal emotions to the physician and to discuss them and may even self-reflect and gain a better understanding of his or her feelings. We identified 2 main variants: an “Anticipate Emotion” behavior that indicates the possibility of an emotional reaction (“What I have to say may seem frightening to you at first”) and a “Validate Emotion” behavior in which the physician comments that any current or future emotional response is understandable, common, or normal for people to feel (“Many people become sad when they hear this type of news”). We combined these 2 variants when initial tests found difficulty in discriminating between them.

Caution About Future Emotion

This behavior covers remarks about emotions arising after the conversation (“You might get more worried about this after you get home”). We separated this behavior from the Anticipate or Validate Emotion behavior because of differences from the patient perspective. For example, physicians might say, “Here is the number for a sickle cell support group to call if you get to be more worried about this screening result.”

Instruction About Emotion

Physicians sometimes tell patients how to feel (“I wouldn’t be worried about this”). The validity of some instructions may be questionable, have inappropriate timing, or impair trust or openness (eg, “Don’t be sad”). For this analysis, however, we chose to include all instructions to pave the way for future study.

Data source

The data dictionary was used to analyze 143 transcripts of conversations between child health providers and standardized parents. To include a variety of communication styles, we developed a heterogeneous sample from 2 data collection efforts. The first 64 transcripts were conversations by pediatric residents

who were participating in an educational workshop about newborn screening for CF and SCH. The other 79 transcripts came from a study of counseling about SCH carrier results by physicians who were working outside of academia.

The resident transcripts were collected as part of their curriculum; informed consent was asked for use of tapes in research. Each workshop began with a 10-minute review about newborn screening, CF, SCH, and inheritance. No teaching about communication or emotions was performed during this period. Each resident was taped in 1 SCH trait encounter and 1 likely CF carrier encounter, the order of which was randomly distributed. A handout described the screening result but did not provide any advice about how to communicate with the parent. In the SCH carrier scenarios, residents were told the infant's screening result showed hemoglobin F, A, and S, a result that had been presented in the review session as definitely indicating that an infant is an SCH carrier. In the likely CF carrier scenarios, residents were told the result showed an elevated immunoreactive trypsinogen, a single $\Delta F508$ mutation, and no multiallele follow-up screening. Our research group uses the word *likely* for this result because it is associated with a 5% to 10% chance of having the actual disease due to an undetected second mutation.⁶¹

All standardized parents were women and chosen to plausibly depict the age and ethnicity of a mother of an infant with CF or SCH. Encounters began with a simple question: "Hello, doctor. I heard you wanted to speak with me, something about my baby's screening test result?"

The standardized parents were coached to adopt a neutral facial expression and to avoid any appearance of anxiety or other emotions. These instructions helped our analysis to focus on precautionary empathy rather than on the physicians' ability to respond to visible emotions. Physicians' response to visible emotional cues (eg, crying) is a critical skill but is not the subject of this study.

For transcripts from graduated physicians, the counseling task always concerned an SCH carrier result. Names and contact information were obtained from a search of the American Medical Association Masterfile for Pediatricians in Connecticut and Wisconsin and Family Physicians in Wisconsin. Search parameters included self-report of working in direct patient care and graduation from residency at least 2 years before. Participants were offered \$20.

The standardized parent encounters followed procedures of the resident project, except that physicians were randomized to counsel either by telephone or in person at their offices. Recordings were transcribed, proofread, and stripped of identifying information. To facilitate the abstraction procedures, we used a sentence diagramming technique to parse transcripts into individual "statements," each with 1 subject and 1 predicate.

Abstraction

Our procedure for abstracting transcripts is derived from quality improvement methods used for explicit-criteria review of medical records,⁶² with a data dictionary containing explicit definitions and examples. Abstractors read transcripts statement-by-statement, comparing each statement with the data dictionary.

Abstraction used a trichotomous scale, in which the abstractors rated the statement "definite" if it met full criteria for the communication behavior described in the data dictionary or "absent" if there was no reference to emotion. As with our previous studies,^{38-42,45,46} we used "partial" ratings to represent

statements that may have been an attempt to discuss emotion, such as leading syntax, assessment questions without a pause for the patient to answer, or hinting at emotion with a vague term that fails to mention an actual emotion (“so what’s going on in your mind?”).

Each transcript was abstracted duplicatively by 2 of the abstractors, and discrepancies were automatically resolved by a spreadsheet algorithm to final results. The algorithm resulted in a final status of “definite” if the individual abstractors had entered definite or if one had entered definite and the other entered partial. The algorithm resulted in a final status of “partial” for all other combinations of abstractor entries except if both statements had been left blank or deliberately entered as “absent.”

One-third of the duplicate abstractions were discussed further to ensure quality and consistency, following the suggestion by Feinstein.⁶³

Statistical analysis

Interabstractor reliability was calculated using a weighted adaptation of the Cohen method⁶⁴ for the trichotomous (definite/partial/absent) schema. Data were analyzed using the χ^2 test for grouped categorical responses and the Wilcoxon rank sum test for nonparametric continuous responses to categorical variables.

We also ran a series of regression models to adjust for overlapping effects between hypothesized predictor variables. The rates of the individual precautionary empathy behaviors were low enough, however, that all of the individual factors lost their significance even in 2-parameter logistic models.

Results

Characteristics of the composite sample are shown in [Table 2](#). Data for the residents were similar to data for the residency program at the time of the study. Data for the graduated physicians were similar to the Masterfile data for physicians who declined participation.

Table 2. Characteristics of 143 Participants

Characteristics	No. (%)
Sex	
Male	73 (51.0)
Female	70 (49.0)
Age, y ^a	
21-40	83 (58.0)
41-60	40 (27.9)
61-80	16 (11.2)
Specialty	
Family medicine	31 (21.7)
Pediatrics	112 (78.3)
Postgraduate year	
Second	39 (27.3)
Third or fourth ^b	20 (14.0)
>2 y out of residency	84 (58.7)

^aFour participants declined to provide their age.

^bFourth-year pediatric residents were from the combined medicine-pediatrics program.

Physicians' use of precautionary empathy behaviors

Use of precautionary empathy behaviors was rare. Definite criteria were identified for 58 instances of the precautionary empathy behaviors. Partial criteria were identified for an additional 119 instances. Abstractors agreed on more than 99% of the instances, but the behaviors were rare enough in the sample that the overall Cohen κ was only 0.73.

Only 38 of 143 transcripts (26.6%) met definite criteria for at least 1 precautionary empathy behavior (Figure 1). When abstractors' partial-criteria ratings were also counted, the number of transcripts increased by 42 to 80 of 143 (55.9%) (Figure 2).




Figure 1. Number of precautionary empathy behaviors included in child health providers' counseling of standardized parents.




Figure 2. Number of precautionary empathy behaviors included when partial-criteria abstractions are counted. The bar indicating 3 or more behaviors includes 9 transcripts with 3 types, 1 with 4 types, and 1 with 5 types.

Transcripts were more likely to include at least 1 precautionary empathy behavior if the infant had likely CF carrier status (86.2% vs 48.3% for SCH carrier infants; χ^2 test; $P < .001$). Precautionary empathy was also more common if the physician were a resident (74.6% vs 42.9% for graduates; χ^2 test; $P < .001$) or were female (74.4% vs 48.0% for males; χ^2 test; $P = .004$). We tested whether effects would persist after adjustment for each other, but the numbers of instances were insufficient for 2-parameter regression.

Individual types of precautionary empathy

Assessment for Emotion

Abstractors identified definite criteria in only 1 transcript for open-ended Assessment for Emotion. No transcripts met definite criteria for closed-ended Assessment for Emotion. When partial-criteria abstractions were also counted, open-ended Assessments for Emotion were found in 8 other transcripts (6.3% total). These were more common in likely CF carrier transcripts (17.2% vs 3.6% for SCH carrier infants; χ^2 test; $P = .006$).

Twelve transcripts (8.4%) met partial criteria for at least 1 closed-ended Assessment for Emotion. Transcripts were more likely to include a closed-ended Assessment for Emotion if the physician were female (16.3% vs 5.0% for males; χ^2 test; $P = .03$) or a resident (15.3% vs 3.6% for graduates; χ^2 test; $P = .01$). Eleven transcripts (7.7%) met partial criteria for at least 1 "Assess for Prior Emotion" behavior.

Abstractors agreed on 93.5% of the open-ended assessments, 90.7% of the closed-ended assessments, and 95.1% of the Assessments of Prior Emotion. All of these were so rare that the κ coefficients were not able to exceed 0.5.

Anticipate or Validate Emotion

Definite criteria for this anticipation or validation of emotion were identified in 8 of 143 transcripts (3.1%). Partial criteria were identified in 22 transcripts (20.9% total). Transcripts were more likely to Anticipate or Validate Emotion if the physician were female (32.6% vs 16.0% for males; χ^2 test; $P = .004$), a resident (34.0% vs 11.9% for graduates; χ^2 test; $P = .004$), or younger (Wilcoxon; $P = .05$). Physicians were less likely to Anticipate or Validate Emotion if the encounter were by telephone (2.9% vs 26.9% for in-person encounters), and more likely if the infant were a likely CF carrier (51.7% vs 13.2% for SCH carrier infants). Abstractors disagreed on 10% of the Anticipate/Validate Behaviors but there were more instances, so the interabstractor reliability was better than for the other behaviors ($\kappa = 0.67$).

Caution About Future Emotion

Abstractors failed to identify any transcripts with definite criteria for this behavior. When partial criteria were counted, 8 of 143 transcripts (5.6%) were identified with this behavior. Female physicians were more likely than males to Caution About Future Emotion (11.6% vs 3.0%; χ^2 test; $P = .04$).

Instruction About Emotion was the most common type of precautionary empathy found in this analysis. The abstractors identified definite criteria in 33 of 143 transcripts (23.1%) for at least 1 instruction about emotion. Most of these instances consisted of phrases such as "don't worry" or "this is not something you need to worry about." Partial criteria were identified in 23 additional transcripts (39.2% total). Female physicians were more likely than male physicians to have included an Instruction About Emotion (48.8% vs 35.0%; χ^2 test; $P = .04$). Abstractor disagreements were more common for the Instructions About Emotion behavior than for the other behaviors (17.0% disagreement; $\kappa = 0.56$).

Comment

Communicating about the possibility for emotions requires different empathic behaviors than responding to apparent emotional cues. This article demonstrates our communication quality indicator method for measuring these precautionary empathy behaviors. This method would be an important component in a communication quality assessment toolkit because unaddressed emotions may be

distressing or impair understanding. The results raise concern about an apparent problem with communication after “nondisease” carrier newborn screening results because it would seem that many of our physicians were unaware that emotional responses may not be apparent. The sample size is small enough that it is difficult to draw firmer conclusions about associated factors, but if our findings are generalizable, then many physicians fail to recognize parents' emotional problems after newborn screening.

Our analysis has several potential limitations. We used standardized parents to evaluate physicians on a level playing field, but physicians' awareness that the “parent” is not truly harboring emotions may have reduced the usage of precautionary empathy. On the other hand, the physicians' sense of being evaluated should have led to their best behavior.

Several other factors could have accounted for the rarity of precautionary empathy. Physicians may lack experience with genetics or may not realize that parents can become distressed about nondisease results. Some physicians may not be attuned to the possibility for emotions to be hidden, having been taught to respond to feigned emotions in standardized parents. Reliability may be improved in future efforts by increasing the explicitness of the data dictionary, by increasing abstractor training, or by focusing on fewer behaviors.

Some of these questions may be addressed by our ongoing, statewide study of counseling by primary care providers of actual infants with nondisease newborn screening results.^{65,66} This effort to assess communication quality may allow us to examine both the usage of communication behaviors and the direct impact of those behaviors on parental outcomes. The fact that the community physicians did not fare any better than the residents underlines the need for Communication Quality Assurance efforts on a population scale.

Our quality indicator methods are intended to allow existing quality improvement professionals to work on physicians' communication rather than to assume that physicians will continue to use skills learned during training.^{36,37,53} Addressing communication problems in health care does not need to interfere with the biomedical importance of screening or change medical educators' traditional focus on emotions—but such problems may need to be addressed if population-scale newborn screening is to result in more good than harm.

References

1. Roter DL, Hall JA, Kern DE, Barker LR, Cole KA, Roca RP. Improving physicians' interviewing skills and reducing patients' emotional distress: a randomized clinical trial. *Arch Intern Med.* 1995;155(17):1877-18847677554
2. Hampton ML, Anderson J, Lavizzo BS, Bergmen AB. Sick cell “nondisease”: a potentially serious public health problem. *AJDC.* 1974;128(1):58-614834985
3. Billings P, Kohn M, de Cuevas M, Beckwith J, Alper J, Natowicz M. Discrimination as a consequence of genetic testing. *Am J Hum Genet.* 1992;50(3):476-4821539589
4. Kessler S, Kessler H, Ward P. Psychological aspects of genetic counseling, III: management of guilt and shame. *Am J Med Genet.* 1984;17(3):673-6976711619
5. Markel H. The stigma of disease: implications of genetic screening. *Am J Med.* 1992;93(2):209-2151497018

6. Markel H. Scientific advances and social risks: historical perspectives of genetic screening programs for sickle cell disease, Tay-Sachs disease, neural tube defects, and Down syndrome, 1970-1997. In: Holtzman NA, Watson MS, eds. *Promoting Safe and Effective Genetic Testing in the United States: Final Report of the Task Force on Genetic Testing*. Baltimore, MD: Johns Hopkins University Press; 1998:161-176
7. Mischler EH, Wilfond BS, Fost N, et al. Cystic fibrosis newborn screening: impact on reproductive behavior and implications for genetic counseling. *Pediatrics*. 1998;102(1, pt 1):44-529651412
8. Tluczek A, Kosciak RL, Farrell PM, Rock MJ. Psychosocial risk associated with newborn screening for cystic fibrosis: parents' experience while awaiting the sweat-test appointment. *Pediatrics*. 2005;115(6):1692-170315930234
9. Tluczek A, Mischler EH, Bowers B, et al. Psychological impact of false-positive results when screening for cystic fibrosis. *Pediatr Pulmonol Suppl*. 1991;7:29-371782126
10. Hewlett J, Waisbren SE. A review of the psychosocial effects of false-positive results on parents and current communication practices in newborn screening. *J Inherit Metab Dis*. 2006;29(5):677-68216917730
11. Tluczek A, Mischler EH, Farrell PM, et al. Parents' knowledge of neonatal screening and response to false-positive cystic fibrosis testing. *J Dev Behav Pediatr*. 1992;13(3):181-1861613113
12. Laird L, Dezateux C, Anionwu EN. Neonatal screening for sickle cell disorders: what about the carrier infants? *BMJ*. 1996;313(7054):407-4118761233
13. McNeil TF, Harty B, Thelin T, Aspegren-Jansson E, Sveger T. Identifying children at high somatic risk: long-term effects on mother-child interaction. *Acta Psychiatr Scand*. 1986;74(6):555-5623493618
14. Andrews L, Fullarton J, Holtzman N, Motulsky A. *Assessing Genetic Risks: Implications for Health and Social Policy*. Washington, DC: National Academy Press; 1994
15. Botkin JR, Clayton EW, Fost NC, et al. Newborn screening technology: proceed with caution. *Pediatrics*. 2006;117(5):1793-179916651338
16. Farrell MH, Farrell PM. Newborn screening for cystic fibrosis: ensuring more good than harm. *J Pediatr*. 2003;143(6):707-71214657812
17. Detmar SB, Aaronson NK, Wever LD, Muller M, Schornagel JH. How are you feeling? who wants to know? patients' and oncologists' preferences for discussing health-related quality-of-life issues. *J Clin Oncol*. 2000;18(18):3295-330110986063
18. Robinson JW, Roter DL. Psychosocial problem disclosure by primary care patients. *Soc Sci Med*. 1999;48(10):1353-136210369436
19. Maguire P, Faulkner A, Booth K, Elliott C, Hillier V. Helping cancer patients disclose their concerns. *Eur J Cancer*. 1996;32A(1):78-818695247
20. Bartlett EE, Grayson M, Barker R, Levine DM, Golden A, Libber S. The effects of physician communications skills on patient satisfaction; recall, and adherence. *J Chronic Dis*. 1984;37(9-10):755-7646501547
21. Levinson W, Gorawara-Bhat R, Lamb J. A study of patient clues and physician responses in primary care and surgical settings. *JAMA*. 2000;284(8):1021-102710944650
22. Suchman AL, Markakis K, Beckman HB, Frankel R. A model of empathic communication in the medical interview. *JAMA*. 1997;277(8):678-6829039890

23. Bylund CL, Makoul G. Examining empathy in medical encounters: an observational study using the empathic communication coding system. *Health Commun.* 2005;18(2):123-14016083407
24. Quill TE. Barriers to effective communication. In: Lipkin M, Putnam SM, Lazare A, eds. *The Medical Interview: Clinical Care, Education and Research*. New York, NY: Springer; 1995
25. Johnson TM, Hardt EJ, Kleinman A. Cultural factors in the medical interview. In: Lipkin M, Putnam SM, Lazare A, eds. *The Medical Interview: Critical Care, Education and Research*. New York, NY: Springer; 1995
26. de Waal FBM. *Good Natured: The Origins of Right and Wrong in Humans and Other Animals*. Cambridge, MA: Harvard University Press; 1996
27. Eisenberg N, Strayer J. Critical issues in the study of empathy. In: Eisenberg N, Strayer J, eds. *Empathy and Its Development*. New York, NY: Cambridge University Press; 1987:3-13
28. Krebs DL. Empathy and altruism. *J Pers Soc Psychol.* 1975;32(6):1134-11461214217
29. Mead GH. *Mind, Self and Society*. Chicago, IL: University of Chicago Press; 1934
30. Stotland E. Exploratory investigations of empathy. In: Berkowitz L, ed. *Advances in Experimental Social Psychology*. Vol 4. New York, NY: Academic; 1969:271-313
31. Kane GC, Gotto JL, Mangione S, West S, Hojat M. Jefferson Scale of Patient's Perceptions of Physician Empathy: preliminary psychometric data. *Croat Med J.* 2007;48(1):81-8617309143
32. Weaver MJ, Ow CL, Walker DJ, Degenhardt EF. A questionnaire for patients' evaluations of their physicians' humanistic behaviors. *J Gen Intern Med.* 1993;8(3):135-1398455109
33. Matthews DA, Feinstein AR. A new instrument for patients' ratings of physician performance in the hospital setting. *J Gen Intern Med.* 1989;4(1):14-222915268
34. *The American Heritage Dictionary of the English Language*. 4th ed. Boston, MA: Houghton Mifflin Co; 2006
35. Cole SA, ed, Bird J, ed. *The Medical Interview: The Three-Function Approach*. 2nd ed. St Louis, MO: Mosby; 2000
36. Coulehan JL, Block MR. *The Medical Interview: Mastering Skills for Clinical Practice*. 5th ed. Philadelphia, PA: FA Davis Co; 2006
37. Silverman J, ed, Kurtz S, ed, Draper J, ed. *Skills for Communicating With Patients*. Oxon, England: Radcliffe Medical Press; 1998
38. Christopher S, Ahmad N, Bradford L, et al. A method to assess the use of organizing behaviors in physicians' conversation. *Med Encounter.* 2010;24(3):224
39. Farrell MH, Kuruvilla P, Eskra KL, Christopher SA, Brienza RS. A method to quantify and compare clinicians' assessments of patient understanding during counseling of standardized patients. *Patient Educ Couns.* 2009;77(1):128-13519380210
40. Farrell M, Deuster L, Donovan J, Christopher S. Pediatric residents' use of jargon during counseling about newborn genetic screening results. *Pediatrics.* 2008;122(2):243-24918676539
41. Farrell MH, Kuruvilla P. Assessment of parental understanding by pediatric residents during counseling after newborn genetic screening. *Arch Pediatr Adolesc Med.* 2008;162(3):199-20418316655
42. Deuster L, Donovan J, Christopher SA, Farrell MH. Quantity of jargon used by internal medicine residents in their counseling about mammography and prostate cancer screening. Poster presented at: Society of General Internal Medicine 30th Annual Meeting; April 25-28, 2007; Toronto, Canada

43. La Pean A, Farrell MH. Initially misleading communication of carrier results after newborn genetic screening. *Pediatrics*. 2005;116(6):1499-150516322177
44. Farrell MH, La Pean A, Ladouceur L. Content of communication by pediatric residents after newborn genetic screening. *Pediatrics*. 2005;116(6):1492-149816322176
45. Deuster L, Christopher S, Donovan J, Farrell M. A method to quantify residents' jargon use during counseling of standardized patients about cancer screening. *J Gen Intern Med*. 2008;23(12):1947-195218670828
46. Farrell MH, Kuruvilla PE, Eskra KL, Christopher SA, Brienza RS. A method to quantify and compare clinicians' assessments of patient understanding during counseling of standardized patients. *Patient Educ Couns*. 2009;77(1):128-13519380210
47. Ley P. *Communicating With Patients: Improving Communication, Satisfaction and Compliance*. London, England: Croom Helm; 1988
48. Makoul G. Essential elements of communication in medical encounters: the Kalamazoo consensus statement. *Acad Med*. 2001;76(4):390-39311299158
49. Makoul G. The SEGUE Framework for teaching and assessing communication skills. *Patient Educ Couns*. 2001;45(1):23-3411602365
50. Makoul G, Clayman ML. An integrative model of shared decision making in medical encounters. *Patient Educ Couns*. 2006;60(3):301-31216051459
51. Roter D, Larson S. The Roter interaction analysis system (RIAS): utility and flexibility for analysis of medical interactions. *Patient Educ Couns*. 2002;46(4):243-25111932123
52. Roter DL, ed, Hall JA, ed. *Doctors Talking With Patients, Patients Talking With Doctors: Improving Communication in Medical Visits*. Westport, CT: Auburn House; 1992
53. Smith RC, ed. *Patient-Centered Interviewing: An Evidence-Based Method*. 2nd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2002
54. Kurtz S, Silverman J, Draper J. *Teaching and Learning Communication Skills in Medicine*. Oxon, England: Radcliffe Medical Press; 1998
55. Stewart M, ed, Belle Brown K, ed, Weston W, ed, McWhinney I, ed, McWilliam C, ed, Freeman T, ed. *Patient-Centered Medicine: Transforming the Clinical Method*. Thousand Oaks, CA: Sage; 1995
56. van Servellen G. *Communication Skills for the Health Care Professional: Concepts and Techniques*. Gaithersburg, MD: Aspen Publishers; 1997
57. Desmond J, Copeland LR. *Communicating With Today's Patient: Essentials to Save Time, Decrease Risk, and Increase Patient Compliance*. New York, NY: Jossey-Bass; 2000
58. Falvo DR, ed. *Effective Patient Education: A Guide to Increased Compliance*. 2nd ed. Gaithersburg, MD: Aspen; 1994
59. Office of Disease Prevention and Health Promotion. *Quick Guide to Health Literacy: Fact Sheets, Strategies, and Resources*. <http://www.health.gov/communication/literacy/quickguide/Quickguide.pdf>. Accessed July 10, 2007
60. Braddock CH III, Edwards KA, Hasenberg NM, Laidley TL, Levinson W. Informed decision making in outpatient practice: time to get back to basics. *JAMA*. 1999;282(24):2313-232010612318
61. Bobadilla JL, Farrell MH, Farrell PM. Applying CFTR molecular genetics to facilitate the diagnosis of cystic fibrosis through screening. *Adv Pediatr*. 2002;49:131-19012214770

62. Allison JJ, Wall TC, Spettell CM, et al. The art and science of chart review. *Jt Comm J Qual Improv.* 2000;26(3):115-13610709146
63. Feinstein A. *Clinical Epidemiology: The Architecture of Clinical Research.* Philadelphia, PA: WB Saunders; 1985
64. Cohen J. Weighted kappa: nominal scale agreement with provision for scaled disagreement or partial credit. *Psychol Bull.* 1968;70(4):213-22019673146
65. Farrell MH, Christopher SA, Tluczek A, et al. Improving communication between doctors and patients after newborn screening. *Wisc Med J.* 2011;110(5):221-227
66. La Pean A, Collins JL, Christopher SA, et al. A qualitative secondary evaluation of statewide follow-up interviews for abnormal newborn screening results for cystic fibrosis and sickle cell hemoglobinopathy [published online ahead of print]. *Genet Med*