

The Linacre Quarterly

Volume 31 | Number 4

Article 22

November 1964

Cancer and Pregnancy

Larry McGowan

Follow this and additional works at: <http://epublications.marquette.edu/lnq>

Recommended Citation

McGowan, Larry (1964) "Cancer and Pregnancy," *The Linacre Quarterly*: Vol. 31: No. 4, Article 22.
Available at: <http://epublications.marquette.edu/lnq/vol31/iss4/22>

Cancer and Pregnancy

LARRY MCGOWAN, M.D.*

Cancer associated with pregnancy arrests the attention of all physicians. A mother who has the misfortune of developing cancer is burdened by two proliferating systems, the disease and the fetus. One demands elimination or control; the other, preservation.¹¹² Cancer is becoming a more frequent cause of maternal death because many obstetrical factors that previously were responsible for maternal deaths are not as prevalent today.²⁵

The following review presents current medical practice when the more common types of cancer are associated with pregnancy. The reader is referred to a recent more extensive article on cancer and pregnancy.¹³¹

BREAST

Cancer of the breast is still the major cause of death from cancer among women.⁷⁴ One in 35 patients with carcinoma of the breast will have pregnancy as a complication, but if the cancer is during the childbearing years, the figure is 1 in 3.¹²⁰

The disease is found by the patient in approximately 90% of cases.⁷⁶ Immediate biopsy of a suggestive tumor is indicated.^{19, 33, 75, 76}

Holleb and Farrow reviewed 266 patients with carcinoma of the breast in whom the carcinoma was associated with pregnancy as a simultaneous, postpartum, or subsequent event.⁴¹ They observed 28 women in the first

trimester of pregnancy, 24 of whom were treated by radical mastectomy. Four patients were excluded because they refused treatment or received roentgen therapy only. The five-year survival rate was 25% for the radical mastectomy group in the first trimester of pregnancy. The axillary nodes were involved in 79% or 19 patients, and the five-year clinical cure rate for this group was 21%. Only 21%, 5 patients, had negative nodes, and here the clinical cure rate rose to 40%.⁴⁹ One-half of the patients were allowed to deliver normally, and one-half had therapeutic abortions.⁴⁹ In their group of 24 patients, the interruption of pregnancy did not seem to affect the clinical cure rate. The overall five-year clinical cure rate was better in the group which delivered normally compared with those patients whose pregnancy was interrupted, 33% versus 17%.⁴⁹

Peters reported 70 cases of cancer of the breast associated with pregnancy that were treated by use of radiotherapy in conjunction with mastectomy.⁹² In this group she reported 8 patients in whom prophylactic therapeutic abortion was performed for carcinoma coincident with pregnancy. None of the previously mentioned women survived over 4 years.⁹²

White recorded 12 patients operated upon for breast cancer and then became pregnant within 4 years. Eight of these patients had disease localized to the breast and 4 had axillary node involvement. Seven of the 8 patients with local disease survived 10 years or

*Dr. McGowan is associated with the Department of Obstetrics and Gynecology, Thomas M. Fitzgerald Mercy Hospital, Darby, Penna., and Hahnemann Medical College, Philadelphia, Penna.

more.¹²⁰ In Stage II cancer of the breast the fate of the patient depends more on the state of the disease than on the condition of the pregnancy.⁴⁸

If cancer is discovered during or after pregnancy, radical mastectomy and axillary lymph node dissection should be carried out promptly.^{21,48,49,73-120} Postoperatively radiation therapy can be given to the involved tissues.⁹² The pregnancy should not be interrupted nor should castration be performed.^{40,92,73} Pregnancy after therapy for cancer of the breast should be delayed for at least 3 years and longer if the axillary lymph nodes contain cancer.

CERVIX

Cervical carcinoma coexisting with pregnancy occurs once in every 2,000 to 6,000 women.^{44,57,64,72,103,115}

All women should have a cytospin of the cervix performed during pregnancy.^{4,50,55,57,105} Preinvasive carcinoma of the uterine cervix can be made on biopsies of pregnant patients and the lesions are not changes due to pregnancy.^{41,53,72,77,91} Conization of the cervix during pregnancy for the diagnosis of preinvasive or invasive cervical cancer can safely be performed.^{12,32}

Patients with carcinoma in situ can have a vaginal delivery providing there are no obstetrical contraindications.^{24,53,78} There is insufficient evidence to prove that pregnancy or delivery per se causes progression of the lesion.⁵³

Fletcher, et al. reviewed 74 cases of carcinoma of the cervix associated with pregnancy or diagnosed up to 1 year postpartum.³⁵ All patients were given radiation therapy and divided into 2 groups according to whether they were treated before supervoltage radiation therapy or with supervoltage radiation therapy. Before supervoltage

radiation therapy 30 patients were treated with 40.5% five-year survival and after supervoltage therapy was instituted, 44 patients were treated with a 55.5% five-year survival. The increased survival rate was most marked for Stage I cases with all 18 patients treated still alive.³⁵ Fletcher, et al. gave to Stage I and Stage II cases a dose of 4,000 rads to the whole pelvis.³⁵ In the first trimester of pregnancy abortion usually occurred after 2,000 to 3,000 rads were given. When external irradiation was completed at 4 weeks, the usual intracavitary radium therapy was used. In the second trimester, abortion usually occurred before 4,000 rads were delivered. When external irradiation was completed, the uterus had involuted enough to proceed with the normal radium therapy. There were 4 instances in which abortion did not occur by the time 4,000 rads had been delivered. In those cases vaginal radium only was used and a hysterectomy performed.³⁵ In the third trimester, with a viable fetus, a high cesarean section was done and whole pelvis irradiation was started 3 or 4 days afterward. The uterus was sufficiently involuted after 4,000 rads were delivered to carry the normal intracavitary radium therapy.³⁵ In the immediate postpartum period 4,000 rads were given to the whole pelvis. The uterus was fully involuted at the end of external irradiation and radium therapy was performed without difficulty.³⁵

Kinch compared 75 pregnant patients with cancer of the cervix treated by irradiation with 402 patients in a comparable age group of 35 or under and discovered an almost identical five-year survival rate of 44% as compared with 45%.⁵⁵ In 28 patients with Stage I carcinoma of the cervix in pregnancy, a five-year survival of

64% was obtained which was almost identical with that in the comparable 402 nonpregnant women. He concluded there was no general worsening of the prognosis when the lesion was complicated by pregnancy.⁵⁵

Sadugor, et al. believe women with cancer of the cervix during pregnancy should be treated without regard for the fetus up until viability was certain since any appreciable delay might decrease the mother's chance of survival. If diagnosis was made after viability they performed a cesarean section without hysterectomy and started the patient on radiation therapy in about 2 weeks. The results of their irradiation treatment for 78 women with cancer of the cervix in all stages associated with pregnancy was a 26.9% five-year survival. This was contrasted to 3,173 women with a comparable percentage in each stage of cervical cancer and a five-year survival of 27.5% was disclosed.¹⁰³

Kistner, et al. performed radical hysterectomy and pelvic node dissection alone in 13 women as the treatment for cervical cancer in pregnancy with 7 women surviving 5 years.⁵⁷

Most authors state that the clinical stage of invasive cancer during pregnancy was the single most important prognostic criterion.^{44,50,55,64,115} The predisposing factors to the rapid spread of invasive cervical cancer in pregnancy were late diagnosis and the effects of vaginal delivery.^{35,55} Cytologic follow-up postpartum of patients with preinvasive carcinoma of the cervix discovered during pregnancy with indicated cold conization of the cervix is carried out if additional children are desired.^{4,53,68,24} Patients with invasive cancer of the cervix discovered before the seventh month of pregnancy or postpartum should undergo definitive therapy.^{35,103} External irradiation can be administered, and fol-

lowed by intracavitary radium therapy after spontaneous abortion.^{35,55,103} Radical hysterectomy and pelvic node dissection in Stage I cancer of the cervix is another mode of treatment.^{57,61} Women with a pregnancy approaching fetal viability and invasive cancer can have therapy briefly delayed until viability. Patients with a viable fetus and invasive cancer of the cervix should be delivered by a classical cesarean section and radiation therapy started after operation.³⁵

HODGKIN'S DISEASE

The incidence of Hodgkin's disease is approximately 1 in 6,000 deliveries⁸³ and in women with Hodgkin's disease the incidence of pregnancy is approximately 18%.¹¹²

The cardinal symptom in the typical case was painless enlargement of the cervical lymph glands.⁴⁶ The diagnosis is based on histologic study of an excised gland.^{46,83,112} The women are permitted to go to term and delivery of healthy infants is usual.^{46,83,99,114,116}

Barry, et al.¹¹ observed 347 patients with Hodgkin's disease and compared those whose illness was associated with 1 or more pregnancies with those whose illness was not. The median survival was just under 5 years. Fertility was not impaired with Hodgkin's disease.^{11,46,54,99,110} Barry and others have discovered active disease during pregnancy to be most frequent above the diaphragm; this was treated with ionizing radiation, shielding the abdomen.^{11,46,99,112} The median survival time in Barry's study was 90 months in the group with pregnancy associated with Hodgkin's disease and it compared favorably with findings for the age-corrected nonpregnant control group (median survival 52 months). The onset of Hodgkin's disease during pregnancy does not adversely af-

fect survival.¹¹ The frequency of exacerbations of the disease was not increased by pregnancy, as the expected rates in the control subjects and pregnant patients were similar.¹¹ Pregnancy did not have a detrimental effect on survival as 54 women in the entire Barry study survived 10 years and 22 of the 54 were pregnant a total of 34 times in association with Hodgkin's disease.¹¹ Other authors have also made this observation.^{46,67,83,99,112,116}

It is desirable not to administer cytotoxic drugs in the first trimester of pregnancy because of the risk to the fetus.^{26,47,112,113,46,83}

Women desirous of pregnancy with Hodgkin's disease should be urged to wait until they have had quiescence of disease for at least 2 years.¹¹ Patients with active disease should be advised against pregnancy, not because of any adverse effect of pregnancy on the mother or danger to the fetus, but because of the poor prognosis in the mother.

LEUKEMIA

Good prenatal care is indicated in pregnant women with leukemia.³⁶ The treatment of chronic leukemia is usually external radiation therapy with shielding of the pregnant uterus.¹⁰⁸ Pregnancy does not appear to have an adverse effect on chronic leukemia.¹⁰⁸ In a disease such as acute leukemia with few survivors, the aim of therapy is to maintain the pregnant patient's general condition at least until a healthy infant is obtained.^{36,62,63,70,79,85,130} The rapidly effective steroids are the drugs of choice during all trimesters with folic acid antagonists and antimetabolites being reserved for periods during the second and third trimesters when steroid control fails or for the postpartum period.^{63,130} Most authors have observed that in-

fants born of leukemic mothers will do as well as normal controls provided that antimetabolites are not administered and radiation therapy to the uterus is not given during the first trimester of pregnancy and the fetus goes beyond the age of viability.^{30,36,62,63,79,84,87,130}

MELANOMA

George, Fortner and Pack³⁸ report a five-year survival in 27 of 54 (50%) and a ten-year survival of 8 of 19 (42%) women pregnant coincident with melanoma. Fifteen of 32 women (47%) followed 5 years survived and 9 of 23 (39%) survived 10 years in the group pregnant after melanoma. In the control or nonpregnant group in the ages of 16 to 43, 56 of 112 (50%) survived 5 years and 21 of 47 (45%) survived 10 years.

White et al.¹²⁶ compared 30 patients whose pregnancies occurred 1 year before the diagnosis of melanoma and 5 years after diagnosis with 31 patients not pregnant during this period. The five-year survival in the pregnant group was 73.3% and in the nonpregnant group 58.8%. They conclude that the five-year survival of melanoma in pregnant women is as good as, or better than, five-year survivals in nonpregnant women. They also state that the stage of disease at the time of diagnosis and treatment was of great importance in determining five-year survival, but the pregnancy did not seem to exert any influence.¹²⁶

Pack and others state that the demonstrated good prognosis for pregnant patients with melanoma establishes a basis for carrying out the same early, adequate surgical treatment for them as for their nonpregnant counterparts.^{14,38,70,80} The pregnancy is not interrupted.

OVARY

An enlarging ovary 6 cm. or more in diameter should be inspected to exclude cancer.^{17,65} The complete excision of ovarian cancer tissue is indicated at any stage of pregnancy.^{15-17,39} The use of external radiation therapy or intra-abdominal radioactive compounds should be delayed after operation until fetal viability.^{15,27} Cesarean section can then be performed or it can be combined with a primary definitive surgical operation.^{39,42,58} Interruption of pregnancy does not appear to have any beneficial effect on maternal prognosis.²⁷

THYROID

The diagnosis of thyroid cancer is made by tissue examination of an enlarged gland.^{15,43,47,81,102} The treatment is immediate thyroidectomy followed by external radiation therapy to the involved area. The pregnancy is not interrupted. Radioactive iodine to inactivate functioning thyroid metastases should not be used after 12 weeks of gestation because it traverses the placenta and may concentrate in the fetal thyroid.^{43,102} After viability of the infant cesarean section can be performed to allow earlier use of radioactive iodine.

SUMMARY

Primary cancer has been reported in most regions of the body during pregnancy. The obstetrician is in an unparalleled position to diagnose cancer. His patient is making her first visit to a physician or the first visit since the birth of her last child for a complete physical examination. The early diagnosis of cancer in a pregnant woman, as in a nonpregnant woman, is usually associated with an increase in survival. In addition to earlier diagnosis, the improvement in surgical techniques, radiation therapy, and chemotherapeutic agents have opened a new era in oncology.

The occasional case report of cancer and pregnancy is of value but precedent in therapy should not be determined until sufficient clinical material has been followed for 5 or more years. As greater numbers of group studies of pregnant women with cancer are published, the concept that pregnancy does not influence the cancer emerges. Treatment is directed at destroying the cancer if it is considered curable, preserving the fetus if the cancer patient is terminal, and compromising with both situations when palliation is indicated and the fetus needs to become more mature before it is delivered.¹⁵

BIBLIOGRAPHY

- ¹Amico, J. C.: Pregnancy complicated by primary carcinoma of the ovary, *Am. J. Obst. & Gynec.*, 74:920, 1957.
- ²Ariel, I. M.: Radioactive isotope therapy for ovarian carcinoma, *J.A.M.A.* 176:103, 1961.
- ³Arneson, A. N.: Special therapeutic problems in gynecologic cancer, *Postgrad. Med.*, 17:480, 1955.
- ⁴Ayre, J. E.: Carcinoma in situ in pregnancy, *J.A.M.A.*, 176:104, 1961.
- ⁵Bacon, H. E. and Rowe, R. J.: Abdominoperineal proctosigmoidectomy

for rectal cancer complicating pregnancy, *South. Med. J.* 40:471, 1947.

- ⁶Banner, E. A., Hunt, A. B., and Dixon, C. F.: Pregnancy associated with carcinoma of the large intestine, *Surg., Gynec. & Obst.*, 80:211, 1945.

- ⁷Barnes, J. E. and Abbott, K. H.: Cerebral complications incurred during pregnancy and the puerperium, *Am. J. Obst. & Gynec.*, 82:192, 1961.

- ⁸Barnes, J. J., Thomas, P. K. and West, O. T.: Primary carcinoma of the vagina associated with pregnancy, *Am. J. Obst. & Gynec.*, 60:1382, 1950.

- ⁹Barr, J. S.: Placental metastases from bronchial carcinoma, *J. Obst. & Gynaec. Brit. Emp.*, 61:895, 1954.
- ¹⁰Barrett, M. E.: Granulosa cell carcinoma in the third trimester of pregnancy, *J.M.A. Alabama*, 18:1, 1956.
- ¹¹Barry, R. M., Diamond, H. D., and Graver, L. E.: Influence of pregnancy on the course of Hodgkin's disease, *Am. J. Obst. & Gynec.*, 84:445, 1962.
- ¹²Beecham, C. T. and Andros, G. J.: Cervical conization in pregnancy, *Obst. & Gynec.*, 16:521, 1960.
- ¹³Bender, S.: Placental metastases in malignant disease complicated by pregnancy, *Brit. M. J.*, 1:980, 1950.
- ¹⁴Berlin, H. and Winters, H. S.: Malignant melanoma of the vulva with pregnancy, Report of a case with six-year survival. *Obst. & Gynec.*, 15:302, 1960.
- ¹⁵Betson, J. R. and Golden, M. L.: Cancer and pregnancy, *Am. J. Obst. & Gynec.*, 81:718, 1961.
- ¹⁶Betson, J. R. and Golden, M. L.: Primary carcinoma of the ovary co-existing with pregnancy, *Obst. & Gynec.*, 12:589, 1958.
- ¹⁷Bossert, L. J.: Bilateral ovarian carcinoma associated with intrauterine pregnancy, *Am. J. Obst. & Gynec.*, 44:336, 1942.
- ¹⁸Bowers, R. N. and Walters, W.: Carcinoma of the stomach complicated by pregnancy, *Minnesota Med.*, 41:30, 1958.
- ¹⁹Brown, R. N.: Carcinoma of the breast followed by pregnancy, *Surg.*, 48:862, 1960.
- ²⁰Brunschwig, A. and Barber, H. R. K.: Cesarean section immediately followed by radical hysterectomy and pelvic node excision, *Am. J. Obst. & Gynec.*, 76:199, 1958.
- ²¹Byrd, B. F., Jr., Bayer, D. S., Robertson, J. C. and Stephenson, S. E.: Therapy of breast cancer in pregnancy, *Ann. Surg.*, 155:940, 1962.
- ²²Daro, A., Carey, C. M., and Zummo, B. P.: Spontaneous rupture of a primary carcinoma of the ovary complicating pregnancy, *Am. J. Obstet. & Gynec.*, 57:1011, 1949.
- ²³Day, A. and Murray, D. J.: Malignant ovarian neoplasms complicating pregnancy, *Canad. M.A.J.* 78:941, 1958.
- ²⁴Dean, R. E., Isbell, N. P., and Woodward, D. E.: Cervical carcinoma in situ in pregnancy, *Obst. & Gynec.*, 20:633, 1962.
- ²⁵D'Esopo, D. A.: Twenty-five years of Obstetrics, *Bull. of the Sloane Hospital for Women*, 5:1, 1959.
- ²⁶Deuschle, K. W. and Wiggins, W. S.: The use of nitrogen mustard in the management of two pregnant lymphoma patients, *Blood*, 8:576, 1953.
- ²⁷Douglas, R. G. and Stromme, W. B.: *Operative Obstetrics*, New York, Appleton-Century-Crofts, Inc. 1957, 627-630.
- ²⁸Editorial: Intracranial neoplasms and pregnancy, *Brit. Med. J.*, 2:1028, 1958.
- ²⁹Einsel, I. H. and Cooks, R.: Carcinoma of the splenic flexure complicated by pregnancy, *J. Internat. Coll. Surg.*, 27:183, 1957.
- ³⁰Elizarova, N. A. and Stupnitskaia, V. M.: A case of acute leukemia in a pregnant patient. *Klin. Med.*, 40:104, 1962.
- ³¹Erf, L. A.: Leukemia (summary of 100 cases) and lymphosarcoma complicated by pregnancy, *Am. J. Clin. Path.*, 17:268, 1947.
- ³²Ferguson, J. H. and Brown, G. C.: Cervical conization during pregnancy. *Surg. Gynec. & Obstet.*, 111:603, 1960.
- ³³Finn, W. F.: Association of pregnancy with cancer of the breast and cervix, *J. M. Soc. New Jersey*, 52:61, 1955.
- ³⁴Finn, W. F. and Lord, J. W.: Carcinoma of the colon producing acute intestinal obstruction during pregnancy, *Surg., Gynec. & Obst.*, 80:545, 1945.
- ³⁵Fletcher, G. H., Rutledge, F. N., and Chau, P. M.: Cancer of the cervix uteri, *Am. J. Roentgenol.*, 87:6, 1962.
- ³⁶Frenkel, E. P. and Meyers, M. C.: Acute leukemia and pregnancy, *Ann. Int. Med.*, 53:656, 1960.
- ³⁷Gemmell, A. A. and Haines, M.: Pregnancy following radical vulvectomy for carcinoma of vulva, *J. Obst. & Gynaec. Brit. Emp.*, 67:1997, 1960.
- ³⁸George, P. A., Fortner, J. G., and Pack, G. T.: Melanoma with pregnancy, *Cancer*, 13:854, 1960.
- ³⁹Green, W. L. and Jones, E. H.: Co-existent pseudomucinous cystadenocarcinoma and pregnancy, *Obst. & Gynec.*, 14:349, 1959.
- ⁴⁰Green, G. G., Smith, A. E., and McClelland, T.: A malignant granulosa cell tumor associated with pregnancy, *Am. J. Obst. & Gynec.* 60:686, 1956.

- ⁴¹Greene, R. R., Peckham, B. M., Chung, J. T., Bayly, M. A., Benaron, H. D. W., Carrow, L. A., and Gardner, G. H.: Preinvasive carcinoma of the cervix during pregnancy, *Surg., Obst. & Gynec.*, 96:71, 1953.
- ⁴²Gustafson, G. W., Gardiner, S. and Stout, F. E.: Ovarian tumors complicating pregnancy, *Am. J. Obst. & Gynec.*, 67:1210, 1954.
- ⁴³Hamill, G. C., Jarman, J. A., and Wynne, M. D.: Fetal effects of radioactive iodine therapy in a pregnant woman with thyroid cancer, *Am. J. Obst. & Gynec.*, 81:1018, 1961.
- ⁴⁴Hayden, G. E.: Carcinoma of the cervix associated with pregnancy, *Am. J. Obst. & Gynec.*, 71:780, 1956.
- ⁴⁵Hendleman, L. and Mestel, A. L.: Multiple carcinomatosis of the colon complicating pregnancy, *Obst. & Gynec.*, 11:119, 1958.
- ⁴⁶Hennessy, J. P. and Rottino, A.: Hodgkin's disease in pregnancy with a report of twelve cases, *Am. J. Obst. & Gynec.*, 63:756, 1952.
- ⁴⁷Hesseltine, H. C. and Loth, M. F.: Malignant diseases associated with pregnancy, *West. J. Surg.*, 64:529, 1956.
- ⁴⁸Hochman, A. and Schreiber, H.: Pregnancy and cancer of the breast, *Obst. & Gynec.*, 2:268, 1953.
- ⁴⁹Holleb, A. I. and Farrow, J. H.: The relation of carcinoma of the breast and pregnancy in 283 patients, *Surg., Gynec. & Obst.*, 115:65, 1962.
- ⁵⁰Holzäepfel, J. H. and Ezell, H. E.: An evaluation of carcinoma of the cervix associated with pregnancy, *Am. J. Obst. & Gynec.*, 76:292, 1958.
- ⁵¹Horner, E. N.: Placental metastases: Case report: maternal death from ovarian cancer, *Obst. & Gynec.*, 15:566, 1960.
- ⁵²Jennings, E. M.: Multiple primary malignant tumors of rectum and breast complicating pregnancy, *J.A.M.A.*, 148:736, 1952.
- ⁵³Johnson, L. D., Hertig, A. T., Hinman, C. H., and Easterday, C. L.: Preinvasive cervical lesions in obstetric patients, *Obst. & Gynec.*, 16:133, 1960.
- ⁵⁴Kasdon, S. C.: Pregnancy and Hodgkin's disease, *Am. J. Obst. & Gynec.*, 57:282, 1949.
- ⁵⁵Kinch, R. A. H.: Factors affecting the prognosis of cancer of the cervix in pregnancy, *Am. J. Obst. & Gynec.*, 82:45, 1961.
- ⁵⁶King, A. B.: Neurologic conditions occurring as complications of pregnancy, *Arch. Neurol. & Psychiat.*, 63:611, 1950.
- ⁵⁷Kistner, R. W., Gorbach, A. C., and Smith, V.: Cervical cancer in pregnancy, *Obst. & Gynec.*, 9:554, 1957.
- ⁵⁸Klein, J.: Pregnancy complicated by teratoma of the ovary, *Obst. & Gynec.*, 3:93, 1954.
- ⁵⁹Klein, M. D., Goodfriend, M. J., and Gordimer, H.: Adenocarcinoma of the colon with bilateral ovarian metastases, *Obst. & Gynec.*, 6:410, 1955.
- ⁶⁰Lash, A. F.: Hypernephroid carcinoma of the kidney or adrenal and pregnancy, *Am. J. Obst. & Gynec.*, 50:221, 1945.
- ⁶¹Lash, A. F.: Management of carcinoma of the cervix in pregnancy, *Obst. & Gynec.*, 17:41, 1961.
- ⁶²Lee, R. A., Johnson, C. E., and Hanlon, D. G.: Leukemia during pregnancy, *Am. J. Obst. & Gynec.*, 84:455, 1962.
- ⁶³Lloyd, H. O.: Acute leukemia complicated by pregnancy, *J.A.M.A.*, 178:78, 1961.
- ⁶⁴McDuff, H. C., Carney, W. I., and Waterman, G. W.: Cancer of the cervix and pregnancy, *Obst. & Gynec.*, 8:196, 1956.
- ⁶⁵McGowan, L.: The diagnosis of ovarian cancer, *Illinois M. J.*, 120:222, 1961.
- ⁶⁶McLean, D. W., Arminski, T. C. and Bradley, G. T.: Management of primary carcinoma of the rectum diagnosed during pregnancy, *Am. J. Surg.*, 90:816, 1955.
- ⁶⁷MacDonald, I.: Hodgkin's disease in pregnancy, *J. Obst. & Gynaec. Brit. Emp.*, 63:931, 1956.
- ⁶⁸Marcus, M. D., Brandt, M. L. and Cibley, L. J.: Carcinoma of the cervix complicating pregnancy, *Obst. & Gynec.*, 10:669, 1957.
- ⁶⁹Marcus, M. D., Cibley, L. J. and Brandt, M. L.: A case of carcinoma of the rectum complicating pregnancy and a review of the literature, *Am. J. Obst. & Gynec.*, 73:1337, 1957.
- ⁷⁰Marcus, M. D. and Fagin, J.: Malignant melanoma in pregnancy, *California Med.*, 91:151, 1959.
- ⁷¹Marozzine, F.: Malignant osteoclastoma in pregnancy, *Ann. Ostet. Gynec.*, 7:559, 195.

- ⁷²Marsh, M. and Fitzgerald, P. J.: Carcinoma in situ of the human uterine cervix in pregnancy, *Cancer*, 9:1195, 1956.
- ⁷³Miller, H. K.: Cancer of the breast during pregnancy and lactation, *Am. J. Obst. & Gynec.*, 83:607, 1962.
- ⁷⁴Montgomery, T. L., Bowers, P. A., and Kittleberger, W. C.: The diagnosis and management of breast cancer, *Obst. & Gynec.*, 17:19, 1961.
- ⁷⁵Montgomery, T. L.: Cancer diagnosis in obstetrics, *South, M. J.*, 47:47, 1954.
- ⁷⁶Montgomery, T. L.: Detection and disposal of breast cancer in pregnancy, *Am. J. Obst. & Gynec.*, 81:926, 1961.
- ⁷⁷Moore, D. B. and Gusberg, S. B.: Cancer precursors in pregnancy, *Obst. & Gynec.*, 13:530, 1959.
- ⁷⁸Moore, J. G., Morton, D. G., Applegate, J. W., and Bashore, R. A.: Superficial carcinoma of the uterine cervix in pregnancy, *Surg., Gynec. & Obst.* 113:339, 1961.
- ⁷⁹Morgan, J. E. and Reyes, C. T.: The association of acute lymphatic leukemia and pregnancy, *Obst. & Gynec.*, 8:642, 1956.
- ⁸⁰Mulla, N.: Malignant melanoma in pregnancy, *Obst. & Gynec.*, 14:530, 1959.
- ⁸¹Muller, J. H. and Brunner, C.: Normal parturition after successful treatment of a malignant goitre and metastases in the mother with radioactive iodine, *Schweiz. med. Wchnschr.*, 83:54, 1953.
- ⁸²Murphy, J., Wilson, J. M., and Bickel, D. A.: Adenoid cystic carcinoma of the Bartholin gland and pregnancy, *Am. J. Obst. & Gynec.*, 83:612, 1962.
- ⁸³Myles, T. J. M.: Hodgkin's disease and pregnancy, *J. Obst. & Gynec. Brit. Emp.*, 62:884, 1955.
- ⁸⁴Neu, L. T., Jr.: Leukemia complicating pregnancy. Two additional case reports, *Missouri Med.*, 59:220, 1962.
- ⁸⁵Newson, A. A., Bruce, C. H., Tabler, J. W., and Strother, W. K.: Leukemia and pregnancy, *Am. J. Obst. & Gynec.*, 69, 892, 1955.
- ⁸⁶O'Leary, J. A. and Bepko, Jr., F. J.: Rectal carcinoma and pregnancy, *Am. J. Obst. & Gynec.*, 84:459, 1962.
- ⁸⁷Olmer, J. and Carcassonne, Y.: Leukemia in pregnancy, *Marseille Med.*, 99:31, 1962.
- ⁸⁸Osband, R. and Jones, W. N.: Carcinoma in situ in pregnancy, *Am. J. Obst. & Gynec.*, 83:559, 1962.
- ⁸⁹Pack, G. T. and Scharnagel, I. M.: The prognosis for malignant melanoma in the pregnant woman, *Cancer*, 4:324, 1951.
- ⁹⁰Palen, G. M.: Retroperitoneal lymphosarcoma complicating pregnancy, *J. Internat. Coll. Surgeons*, 13:376, 1950.
- ⁹¹Peckham, B., Greene, R. R., Chung, J. T., Bayley, M. A., and Benaron, B. H. B. W.: Epithelial abnormalities of the cervix during pregnancy, *Am. J. Obst. & Gynec.*, 67:21, 1954.
- ⁹²Peters, M. V.: Carcinoma of the breast associated with pregnancy, *Radiology*, 78:58, 1962.
- ⁹³Pommerenke, W. T.: Pregnancy following radical resection of the rectum for carcinoma, *Am. J. Obst. & Gynec.*, 31:522, 1936.
- ⁹⁴Putski, P. S., Scully, I. H., Kotz, J., Kaufman, M. S., and Torrey, W. E.: Carcinoma for colon producing acute intestinal obstruction during pregnancy, *Am. J. Surg.*, 77:749, 1949.
- ⁹⁵Rand, C. W. and Andler, M.: Tumors of the brain complicating pregnancy, *Arch. Neurol. & Psychiat.*, 63:1, 1950.
- ⁹⁶Reeves, D. L.: Tumors of the brain complicating pregnancy, *West. J. Surg.*, 60:211, 1952.
- ⁹⁷Reyes, G. R. and Perez, R. T.: Leukemia and pregnancy: observation of a case treated with Busulfan (Mylcran), *Blood*, 18:764, 1961.
- ⁹⁸Reynolds, R. P., Cantor, M. O., and Henderscn, H.: Carcinoma of the stomach in pregnancy, *Grace Hosp. Bull.*, 28:127, 1950.
- ⁹⁹Riva, H. L., Anderson, P. S. and O'Grady, J. W.: Pregnancy and Hodgkin's disease, *Am. J. Obst. & Gynec.*, 66:866, 1953.
- ¹⁰⁰Roddie, T. W.: Haemorrhage from primary carcinoma of the liver complicating pregnancy, *Brit. M. J.*, 1:31, 1957.
- ¹⁰¹Rubin, A. and Lewis, G. C.: Pregnancy and vaginal delivery following radical surgery for cancer of the vulva, *Am. J. Obst. & Gynec.*, 65:13747, 1953.
- ¹⁰²Russell, K. P., Rose, H., and Starr, P.: The effects of radioactive iodine on maternal and fetal thyroid function during pregnancy, *Surg., Gynec. & Obst.*, 104:560, 1957.
- ¹⁰³Sadugor, M. G., Palmer, J. P., and Reinhard, M. C.: Carcinoma of the cervix concomitant with pregnancy, *Am. J. Obst. & Gynec.*, 57:933, 1949.

- 104Schlemenson, M., Rubovits, F. E., and Abrams, B.: Cancer of the large intestine complicated by pregnancy, *J. Internat. Coll. Surg.*, 13:96, 1950.
- 105Schmitz, H. E., Isaacs, J. H., and Fetherston, W. C.: The value of routine cytologic smears in pregnancy, *Am. J. Obst. & Gynec.*, 79:910, 1960.
- 106Schmitz, H. E., Smith, C. J., and Isaacs, J. H.: Cancer of the cervix and cancer complicating pregnancy, *Trans. New England Obst. & Gynec. Soc.*, 11: 167, 1957.
- 107Schuyler, W. B. J.: Pregnancy complicated by carcinoma of the bowel arising in the site of a chronic granuloma, *Obst. & Gynec.*, 5:102, 1955.
- 108Sheehy, T. W.: An evaluation of the effect of pregnancy on chronic granulocytic leukemia, *Am. J. Obst. & Gynec.*, 75:788, 1958.
- 109Sinykin, M. B. and Kaplan, H.: Leukemia in pregnancy, *Am. J. Obst. & Gynec.*, 83:220, 1962.
- 110Smith, F. R.: The effect of pregnancy on malignant tumors, *Am. J. Obst. & Gynec.*, 34:616, 1937.
- 111Smith, G. N. and Niles, N. R.: Renal cell carcinoma in pregnancy, *Obst. & Gynec.*, 10:279, 1957.
- 112Smith, R. B. W., Sheehy, T. W., and Rothberg, H.: Hodgkin's disease and pregnancy, *Arch. Int. Med.*, 102:777, 1958.
- 113Sokal, J. E. and Lessmann, E. M.: Effects of cancer chemotherapeutic agents on the human fetus, *J.A.M.A.*, 172:1765, 1960.
- 114Southam, C. M., Diamond, H. D. and Carver, L. F.: Pregnancy during Hodgkin's disease, *Cancer*, 9:1141, 1956.
- 115Stander, R. W. and Lein, J. N.: Carcinoma of the cervix and pregnancy, *Am. J. Obst. & Gynec.*, 79:164, 1960.
- 116Steward, H. L. and Monto, R. W.: Hodgkin's disease and pregnancy, *Am. J. Obst. & Gynec.*, 63:570, 1952.
- 117Swartley, W. B., Newton, Z. B., Hartman, J. C., and Stayman, J. W.: Perforated carcinoma of the large intestine complicating pregnancy, *Ann. Surg.*, 125:251, 1947.
- 118Tuft, L.: Primary squamous cell carcinoma of the vagina complicating late pregnancy in a patient sixteen years old, *Arch. Path.*, 9:654, 1930.
- 119Waddington, H. K.: Renal cell carcinoma complicated by pregnancy, *Am. J. Obst. & Gynec.*, 72:441, 1956.
- 120Wall, J. A. and Lucci, J. A.: Adenocarcinoma of the corpus uteri and pelvic tuberculosis complicating pregnancy, *Obst. & Gynec.*, 2:629, 1953.
- 121Warren, R. P.: Carcinoma of the rectum and pregnancy, *Brit. J. Surg.*, 45: 61, 1957-8.
- 122Waters, E. G. and Fenimore, E. D.: Perforated carcinoma of the cecum in pregnancy, *Obst. & Gynec.*, 3:263, 1954.
- 123Weyland, R. D., MacCarty, C. S., and Wilson, R. B.: The effect of pregnancy on intracranial meningiomas occurring about the optic chiasm, *S. Clin. North America*, 31: No. 4, 1951.
- 124Whitcher, B. R.: Brain tumor complicating pregnancy, *J. M. Soc. New Jersey*, 53:112, 1956.
- 125White, L. G.: Bulsulfan in pregnancy, *J.A.M.A.*, 179:131, 1962.
- 126White, L. P., Linden, G., Breslow, L., and Harzfeld, L.: Studies on melanoma, *J.A.M.A.*, 177:51, 1961.
- 127White, T. T.: Carcinoma of the breast in the pregnant and the nursing patient, *Am. J. Obst. & Gynec.*, 69:1277, 1955.
- 128White, T. T.: Prognosis of breast cancer for pregnant and nursing women, (Analysis of 1,413 cases), *Surg., Gynec. & Obst.*, 100:661, 1955.
- 129White, T. T. and White, W. C.: Breast cancer and pregnancy report of 49 cases followed 5 years, *Ann. Surg.*, 144:384, 1956.
- 130Yahia, C., Hyman, G. A., and Phillips, L. L.: Acute leukemia and pregnancy, *Obst. & Gynec. Surv.*, 13:1, 1958.
- 131McGowan, L.: Cancer and pregnancy, *Obst. & Gynec. Surv.*, 19:285, 1964.