1						
2						
3						
4	Title:					
5						
6	DENITRIFICATION POTENTIAL OF DIFFERENT LANDUSE TYPES IN					
7	AN AGRICULTURAL WATERSHED, LOWER MISSISSIPPI VALLEY					
8						
9						
10	Paper Type: General Paper					
11						
12						
13	Running head:					
14	0					
15	Denitrification in Lower Mississippi					
16						
17						
18	Authors:					
19						
20	Sami Ullah ¹ * [†] and S.P. Faulkner ²					
21						
22						
23	1. Louisiana State University, Wetland Biogeochemistry Institute, Baton Rouge,					
24	Louisiana, USA.					
25						
26						
27	2. USGS National Wetlands Research Center, 700 Cajundome Blvd., Lafayette, LA					
28	70506, USA					
29						
30	*Author for Correspondence: Sami Ullah, Global Environmental and					
31	Climate Change Center, Department of Geography, McGill University, 610 Burnside					
32	Hall, 805 Sherbrooke St. W, Montreal, Quebec H3A 2K6, Canada					
33	Email: <u>sami.ullah@mcgill.ca</u> , Ph: +1-514-398-4957					
34	Fax: +1-514-398-7437					
35						
36	Key Word : Bottomland hardwoods; denitrification; forested wetlands; NO ₃ pollution					
5/	control; wetlands restoration.					
38						
<i>3</i> 9	Abbreviations: AMOC – Anaerobically mineralizable organic carbon; DP –					
40	Denitrification Potential; LMV- Lower Mississippi alluvial valley					

[†] Current address: Global Environmental and Climate Change Center, Department of Geography, McGill University, 610 Burnside Hall, 805 Sherbrooke St. W, Montreal, Quebec H3A 2K6, Canada

ABSTRACT

1 2

3 Expansion of agricultural land and excessive nitrogen (N) fertilizer use in the Mississippi 4 River watershed has resulted in a 3-fold increase in the nitrate load of the river since the 5 early 1950's. One way to reduce this nitrate load is to restore wetlands at suitable 6 locations between croplands and receiving waters to remove run-off nitrate through 7 denitrification. This research investigated denitrification potential (DP) of different land 8 uses and its controlling factors in an agricultural watershed in the lower Mississippi 9 valley (LMV) to help identify sites with high DP for reducing run-off nitrate. Soil 10 samples collected from seven land-use types of an agricultural watershed during spring, 11 summer, fall and winter were incubated in the laboratory for DP determination. Low-12 elevation clay soils in wetlands exhibited 6.3 and 2.5 times greater DP compared to high-13 elevation silt loam and low-elevation clay soils in croplands, respectively. DP of 14 vegetated-ditches was 1.3 and 4.2 times that of un-vegetated ditches and cultivated soils, 15 respectively. Soil carbon and nitrogen availability, bulk density, and soil moisture 16 significantly affected DP. These factors were significantly influenced in turn by 17 landscape position and land-use type of the watershed. It is evident from these results that 18 low-elevation, fine-textured soils under natural wetlands are the best locations for 19 mediating nitrate loss from agricultural watersheds in the LMV. Landscape position and 20 land-use types can be used as indices for the assessment/modeling of denitrification 21 potential and identification of sites for restoration for nitrate removal in agricultural 22 watersheds.

INTRODUCTION

1.

2	The primary source of increased nitrate in surface waters is nitrogen (N) fertilizer
3	applied to croplands (USEPA 1996). An increase in the nitrate concentration of water
4	bodies is correlated with increased agricultural activity in river watersheds (Smith et al.
5	1987; Galloway et al. 2003). Nitrogen fertilizer use in the US increased by 300% from
6	1961 to 1999 and current usage consumes 13% of the inorganic N fertilizer used globally
7	(Howarth et al. 2002). Thus, expansion of agricultural activities coupled with an
8	increased use of synthetic N fertilizer in the US has resulted in excessive accumulations
9	of reactive N in environments external to croplands (Galloway 2002; Howarth et al.
10	2002).
11	Extensive agricultural development and N fertilizer use over the past 200 years in
12	the Mississippi River basin has increased nitrate loading into the river and the northern
13	Gulf of Mexico (Turner and Rabalais 2003). Since the 1950's, N fertilizer use has
14	increased 20-fold in the basin (Battaglin and Goolsby 1996), which has contributed to a
15	3-fold increase in the nitrate load of the Mississippi River (Turner and Rabalais 1994;
16	Donner 2004). Agricultural run-off contributes about 74% of the current nitrate loading
17	carried by the Mississippi River (Rabalais et al. 2002) and the increased nitrate loading is
18	cited as one of the major causes of the extensive hypoxia in the northern Gulf of Mexico
19	(Rabalais et al. 2002). A 30% reduction of the N load delivered by the Mississippi River
20	has been recommended to reduce the hypoxia (EPA 2001; Mitsch et al. 2001).
21	The Lower Mississippi Alluvial Valley (LMV) has lost about 80% of its
22	bottomland hardwood forests to other land uses primarily agriculture (Allen et al. 2001).
23	The LMV was the largest floodplain ecosystem in the US covering about 23,300-km ²

1 area. Bottomland hardwood forests covered this floodplain and were flooded seasonally 2 as a result of over-bank flooding by the Mississippi River. Due to growth in agriculture, 3 the bottomland hardwoods were cleared, drained, ditched and cultivated for decades for 4 row crop cultivation. This practice not only led to the loss of the NO₃ sinks in the form of 5 greater denitrification rates of bottomland wetlands (Hunter and Faulkner 2001), but 6 enhanced its potential of loading additional nitrate into surface waters including the 7 Mississippi River through N fertilizer use, soil erosion (Mitsch et al. 2005; Rebich 2001; 8 ARS 2001), mineralization of organic nitrogen, and direct drainage of the cultivated 9 lands.

10 Measures and research recommended by Mitsch et al. (2001) and Mitsch and Day 11 (2006) for reducing the NO₃ loading of the Gulf of Mexico from the Mississippi River 12 basin include a) on-farm soil and N fertilizer management to enhance N use efficiency, b) 13 alternative cropping and management systems for reducing N loss from croplands, and c) 14 creation or restoration of wetlands and riparian ecosystems at suitable locations between 15 croplands and water bodies to remove run-off nitrate before its outfall into the river. Like 16 elsewhere in the basin, restoration of forested and riparian wetlands to reduce run-off 17 nitrate through plant uptake and denitrification (Lowrance et al. 1984, Comin et al. 1997) 18 in the LMV is recommended (Lindau et al. 1994). Moreover, re-connecting forested and 19 riparian wetlands with the rivers for over-bank flooding is another measure suggested for 20 nitrate removal from river water in the LMV (Mitsch and Day 2006; Lindau et al. 1994; 21 Lowrance et al. 1997). Denitrification is one of the major biological processes for nitrate 22 removal from soil and water. Soil organic carbon and NO₃ contents, moisture, 23 temperature and texture affect the rate and extent of denitrification (Galloway et al.

2003). The status of these physicochemical soil properties are the result of interactions of
 topography, soil hydrology and soil management at basin and sub-watershed scales in the
 landscape (Florinsky et al. 2004; Lowrance et al. 1997; Peterjohn and Correll 1984).
 Therefore, it is important to discern the effects of topographic and landuse attributes on
 denitrification potential of agricultural watersheds.

6 Agricultural watersheds in the LMV are not homogenous croplands, but are a 7 mosaic of land uses including well, moderately and poorly drained soils under row crop 8 cultivation, a network of drainage ditches and access roads, patches of bottomland 9 hardwood forests and depressional wetlands. Based on the current land use, hydrology, 10 and landscape position, these different land use types can either enhance or retard 11 denitrification. Maintaining environmentally sound crop production in the LMV and 12 reducing nitrate loading into aquatic ecosystems warrants investigation of landscape and 13 environmental factors regulating denitrification potential in agricultural watersheds. 14 Such research-based information is important for the assessment/modeling of 15 denitrification potential at watershed scale and identification of sites for wetland 16 restoration (White and Fennessy 2005) in the LMV. To our knowledge, there are no 17 scientific studies available on this topic in the LMV. Our objectives were to 1) determine 18 denitrification potential of different land use types of an agricultural watershed in the 19 LMV, and 2) identify some of the environmental and landscape/land-use management 20 factors regulating the denitrification potentials.

21

MATERIAL AND METHODS

22 2.1 Study Area

23 The study area is the 8.5 km² Beasley Lake watershed in Sunflower County,
24 Mississippi (Figure 1) in which about 0.25 km² area is covered by the Beasley Lake. The

2.

1	watershed is part of the Yazoo delta region of Northwestern Mississippi formed by the
2	alluvial deposits of the Mississippi River and its tributaries (Fisk 1951). Soils of the
3	watershed range from coarse-textured silty-loam and loam deposits to fine-textured clay
4	alluvium. Dominant soil series of the watershed are Sharkey clay (non-acidic
5	montmorrilinitic, Vertic Haplaquept), Dowling (Very-fine, smectitic, nonacid, thermic
6	Vertic Endoaquept), Alligator (Very-fine, smectitic, thermic Chromic Dystraquert),
7	Dundee (Fine-silty, mixed, active, thermic Typic Endoaqualf), Dubbs silt loam (Fine-
8	silty, mixed, active, thermic Typic Hapludalf), and Forestdale (Fine, smectitic, thermic
9	Typic Endoaqualf), (NRCS 1959).
10	The elevation gradient between the highest and lowest points in the watershed is 5.5
11	meters. Current land uses consist of high (Ag-high) and low (Ag-low) elevation
12	croplands, vegetated ditches (veg-ditches), un-vegetated ditches (unveg-ditches), natural
13	forested wetland and depressional wetlands. Forested wetlands are dominated by
14	bottomland hardwood tree species such as American elm (Ulmus americana), Water oak
15	(Quercus nigra), Pin oak (Quercus phellos), Green ash (Fraxinus pensylvanica) Red
16	maple (Acer ruburum), and Hackberry (Celtis leavigata). FW cover about 1.2 km ² area in
17	the watershed. Depressional wetlands ($\sim 0.1 \text{ km}^2$ area) are small depressions next to
18	Beasley Lake, which remain ponded during winter and spring and are dominated by
19	submerged and emergent wetland vegetation such as Potamogeton spp., Sagittaria spp.,
20	Scirpus spp. Typha spp., Nymphaea spp. Andropogon and Panicum species usually grow
21	on the drier banks of the depressions. These depressions are the remnants of the swale
22	and ridge topographic features of an ox-bow lake watershed where the swales developed
23	into depressional wetlands. Ag-high (~5.3 km ²) landuse covers mainly well-drained soils

1	while Ag-low (~ 1.5 km^2) covers poorly drained soils next to depressional wetlands and
2	forested wetlands. A low elevation natural ditch next to the forested wetland was
3	developed into a constructed wetland (~0.01 km ²) in spring of 2002 through excavation
4	and installation of a water control structure to increase the aerial extent of the flooded
5	soil. About 0.2 km ² high and low-elevation croplands of the watershed drained into the
6	constructed wetland. The heavy clay soil of the constructed wetland was similar to that
7	of the nearby natural wetlands. The Sharky and Dowling clay soil of the constructed
8	wetland supported Potamogton spp., Sagittaria spp. Typha spp. Panicum spp., and
9	Andropogon spp. The constructed wetland remained ponded in spring, fall and winter like
10	that of depressional wetlands.
11	The major management activity in the watershed is crop cultivation. Cotton and corn
12	are grown in the Ag.high croplands while soybean is grown in the Ag-low croplands with
13	conventional tillage system. Recreational land uses include fishing and hunting in the
14	Beasley Lake and in forested wetlands. Overhead irrigation is applied as needed for crop
15	cultivation. Irrigation run-off and rain water are drained through the ditches into Beasley
16	Lake. Due to cultivation and the shunting of agricultural runoff directly to the lake,
17	sedimentation in the lake has increased to a degree where it now threatens its ecology
18	(ARS 2001). Maintaining ditches under grass cover is one of the best management
19	practices (BMP) implemented in the watershed by USDA to help reduce sedimentation in
20	Beasley Lake (Rebich 2001). The dimension of the ditches ranges from 1 to 3 meters
21	wide and 1-2 meter deep in the high-elevation areas and upto 5 meters wide and 4 meters
22	deep in the low-elevation areas of the watershed. The banks of the veg.ditches are
23	stabilized by planting and maintaining switch grass (Panincum spp.)

1 Seven land use types of the watershed were selected for this research (Ag-high, Ag-2 low, veg-ditches, unveg-ditches, forested wetlands, depressional wetlands and the 3 constructed wetland). Eight sampling points were selected randomly in each land use 4 type. Four soil cores (0-10 cm deep; 3 cm dia.) were collected from each sampling point 5 of the seven land-use types of the watershed using a hand auger in March 2002, July 6 2002, October 2002 and January 2003. The four cores from each sampling point were 7 composited and were transferred to the laboratory on ice and refrigerated at 4 °C under 8 their original moisture content levels (field-moisture condition) for further analysis.

9

2.2 Denitrification Potential (DP)

10 The inherent capacity of a soil to reduce and denitrify nitrate to N₂ gas under an 11 unlimited supply of nitrate using organic carbon as an energy source under anaerobic 12 conditions is called denitrification enzyme assay (Beauchamp and Bergstrom 1993; 13 Groffman et al. 1999). This assay measures the amount of denitrification enzymes 14 available at the time of soil sampling. We modified this procedure by amending the soil 15 slurries only with nitrate as we were interested in the denitrification potential (DP) of the 16 different land use types under their existing soil carbon contents using the C_2H_2 block 17 technique (Hill and Cardaci 2004). It is well established that adding carbon to nitrate 18 amended slurries will increase denitrification rates (Hunter and Faulkner 2001; Groffman 19 and Crawford 2003), however there is no practical way of increasing soil carbon at the 20 landscape scale. Therefore, this approach is a more realistic evaluation of the existing 21 ability of the different landscape units to remove nitrate from surface or ground water. 22 For the purpose of this study DP is defined as the capacity of soil slurries to denitrify 23 nitrate under anoxic conditions at room temperature (22 degress Celcius). Field moist

1	soils were thoroughly homogenized by hand and brought to room temperature overnight
2	before incubation. The next morning, six sub samples (moist equivalent of 10 g dry soil)
3	of the homogenized soil from each of the eight soil samples from each of the seven land
4	use types were weighed into 6 replicate serum bottles (150 mL). Fifteen mL of 10 mg
5	$NO_3^{-}L^{-1}$ solution and 5 ml of de-ionized water were added to three of the six bottles to
6	deliver 15 μ g of NO ₃ ⁻ g ⁻¹ dry soil, while 20 mL of de-ionized water was added to the
7	remaining three bottles. The bottles were then capped airtight and purged with $O_2\mbox{-}free\ N_2$
8	gas for 20 minutes to induce anaerobic conditions. Ten percent of the serum bottle
9	headspace was replaced with cleaned C_2H_2 gas to block the bacterial conversion of N_2O
10	to N_2 gas. The bottles were then wrapped in Al foil and put on a reciprocating shaker for
11	continuous shaking at room temperature (22 to 25 °C). Headspace gas samples were
12	collected at 2, 4 and 6 hours with a syringe and stored in Beckton Dikinson
13	Vacutainers®. The gas samples were analyzed on a Tremetrics 9001GC having a
14	porapack Q column with ECD detector for N_2O concentration determination. The rate of
15	N_2O production was calculated in ug N- N_2O g ⁻¹ h ⁻¹ using the 3 gas sample readings
16	during the 6 hour incubation. Adjustments were made for soluble N_2O in the bottles using
17	a Bunsen absorption coefficient of 0.54 at 25 °C.

2.3 Anaerobically Mineralizable Organic Carbon (AMOC)

Denitrification depends directly on the amount of mineralizable organic C
available to the denitrifier population under anaerobic conditions (Singh et al. 1988; Gale
et al. 1992). Since denitrification is an anaerobic process, the amount of mineralizable
organic C available under anaerobic conditions would help explain any trend in the DP
among different land use types. Field moist soil (equivalents of 5 g oven-dried soil) from

1 each soil sample were weighed into 150 mL duplicate serum bottles. Twenty ml of 50 mg $NO_3^{-}L^{-1}$ solution was added into each bottle, which delivered 200 µg $NO_3^{-}g^{-1}$ soil. The 2 3 bottles were capped airtight and purged with oxygen-free N₂ gas for 20 minutes to induce 4 anaerobic conditions. After purging, the bottles were wrapped in aluminum foil and were 5 shaken for 15 minutes on a reciprocating shaker. After shaking, the bottles were stored at 6 room temperature (22-25 °C). The headspaces of the bottles were sampled with a syringe 7 at 1, 24, 48 and 72 hours of incubation and stored in 5-mL Beckton Dikinson 8 Vacutainers[®]. The gas samples were analyzed on a Tremetric 9001 GC fitted with a 9 methanizer and an FID detector for CO₂ concentration determination (Ullah et al. 2005). 10 The gas production over the length of incubation remained linear in all the landuse types. The amount of CO₂ produced was calculated as $\mu g C-CO_2 g^{-1} h^{-1}$. Corrections were made 11 12 for soluble CO_2 in the incubation bottles by using the Bunsen Adsorption coefficient of 0.752 at 25 °C. 13 14 2.4

Total soil carbon and nitrogen

15 Total soil carbon (C) and nitrogen (N) were determined using a Thermo Finnigan 16 CNS Analyzer. Soil samples were oven dried, pulverized and thoroughly homogenized. 17 A sub-sample of about 35 mg was weighed into a tin capsule for automated analysis to 18 determine concentrations of organic C and total N. These values and bulk density measurements were used to calculate Mt of C and N ha⁻¹. 19

20 2.5 Soil nitrate, bulk density, porosity, water-filled pore space and texture

21 Field-moist soil equivalents of 5 g oven-dried soil were weighed into 250 mL 22 duplicate bottles. Fifty mL of 2M KCL solution was added to each bottle. The bottles 23 were put on a reciprocating shaker for continuous shaking for 1 hour. After shaking, the

1 bottles were centrifuged at 3000 rpm for 5 min and were then filtered into 20 mL 2 scintillation vials through a No. 42 Whatman filter paper. The samples were stored in a 3 freezer until analysis for nitrate on a Lachat automated flow injection analyzer. Average values for each soil sample were determined and reported in mg N Kg⁻¹ oven-dried soil. 4 5 At each land use type eight intact soil cores (2.5 cm dia. x 10 cm long) were taken and 6 transferred to the lab for the determination of soil moisture and bulk density. Soil 7 porosity was determined using the equation of 1- (bulk density/particle density). Percent 8 water-filled pore space (WFPS) was determined for each landuse type for all season 9 according to Ullah et al. (2005). Soil particle size distribution was determined by the 10 filtration method according to Sheldrick and Wang (1993).

11

2.6

Statistical Analysis

12 Differences in DP among the landscape units within each season were analyzed 13 by a two-way analysis of variance using the General Linear Model in SAS (SAS Institute 14 1998). Landscape was treated as main effect, nitrate amendment was treated as a sub-plot 15 effect and season was treated as repeated measures variable in the ANOVA model. Post 16 ANOVA tests were conducted with Fisher's protected LSD at 5% significance level. 17 Linear regression of the yearly averaged DP on total soil C, N, and bulk density of the 7 18 land use types was done using SAS. Significant differences in physio-chemical properties 19 of soils of the 7 land use types were determined using one-way ANOVA. Pearson 20 correlation coefficients among DP, AMOC and soil moisture were calculated for each 21 season. The data were analyzed for normality and homogeneity of variance of the 22 residuals using the proc univariate procedure in SAS and Shapiro-wilk test of normality

23 of the residual at p > 0.05.

2

3. **RESULTS**

3 3.1 Landscape position and Landuse type effects on Denitrification Potential

4 The natural and constructed wetlands had significantly greater (p < 0.05) DP than 5 the Ag-high and Ag-low sites in all seasons (Figures 2). Forested wetlands showed the 6 highest DP while Ag-high showed the lowest during the four seasons when additional 7 NO_3 was added during the assay compared to their unamended soils (Figures 2 and 3). 8 The DP of forested wetlands was 4.5 to 11 times greater than the Ag-high during the 9 year. Similarly, the forested wetlands showed 2.4 to 4.3 times greater DP than the Ag-low 10 sites in all the seasons. Except in summer, Ag-high and Ag-low had similar DP values (p 11 > 0.05). On average, forested wetlands exhibited 3.0 and 2.7 times greater DP than those 12 of veg-ditches and unveg-ditches, respectively (Table 1).

Depressional and constructed wetlands showed similar DPs, except in summer when depressional wetlands had 1.5 times greater DP than the constructed wetland. On average both depressional and constructed wetlands had 3.7 to 7.4 and 1.2 to 3.7 times greater (p < 0.05) DP than that of the Ag-high and Ag-low respectively during the four seasons. Depressional and constructed wetlands also showed 1.2 and 1.6 times greater DP than veg.ditches and unveg-ditches, respectively, but were less than the DP of forested wetlands (Table 1).

When the DP of individual land use types were averaged for the four seasons (Table 1), the DP of veg-ditches was 4.1 times of the Ag-high (p <0.05) and 1.7 times of Ag-low, although statistically not significant (p > 0.05). Moreover, veg-ditches had 1.3 times greater DP than unveg-ditches. On the whole forested wetlands exhibited 1 significantly greater DP (p < 0.05) than the rest of the land use types. Similarly,

2 depressional and constructed wetlands had higher DP than unveg-ditches, Ag-high and

3 Ag-low sites (Table 1), but had lower DP than the forested wetlands. The DP of veg-

4 ditches was more variable and thus was not significantly different than Ag-low, unveg-

- 5 ditches, constructed and depressional wetlands (p > 0.05).
- 6

3.2 Environmental variables and Denitrification Potential

7 Among the environmental variables measured, nitrate amendment, season, 8 organic carbon availability, bulk density and %WFSP contents significantly influenced 9 DP of all the land use types. When amended with additional nitrate, forested and 10 depressional wetland soils showed 55% and 67% greater DP compared to the DP of soil 11 without nitrate additions (Figures 2 and 3). Constructed wetland and unveg-ditches 12 showed an increase in DP with NO_3 enrichment during fall and winter only while veg-13 ditches responded to nitrate addition in summer and fall. Ag-high and Ag-low soil did not 14 respond in terms of increased DP to nitrate amendment during the four seasons (Table 1), 15 which shows that their DPs were limited by factors other than nitrate, probably by the 16 availability of organic C and lack of anaerobosis. 17 Season significantly affected the DPs of all the land uses except Ag-high and 18 unveg-ditches (Table 1). When the DPs of all the land use types were averaged together

19 and tested for a seasonal effect, the winter DP was found the lowest than the rest of the

20 seasons. Lower winter soil temperatures are speculated to have limited the denitrifier

21 activity. Similar seasonal effect on denitrification potential of forested wetland soils in

the LMV was observed by Hunter and Faulkner (2001).

1	Significant differences in a number of soil properties among the different land use
2	types were observed (Table 2). Wetlands had 1.8 times greater total soil organic C than
3	the cultivated soils. Higher soil organic C in wetlands contributed to improved soil
4	structure resulting in lower bulk densities and high soil porosities in wetlands than the
5	Ag-high and Ag-low sites. Anaerobic incubation of soils from the 7 land use types
6	showed that the amount of anaerobically mineralizable organic carbon (AMOC) in
7	wetlands soil was 1.4 times those of the cultivated soils (Table 3). Similarly, veg-ditch
8	soil had a relatively lower soil bulk density, higher porosity (Table 2) and 1.3 time
9	greater AMOC values compared to those of the cultivated soils. Unveg-ditches had
10	similar soil bulk density, porosity and AMOC values to those of the cultivated soils. High
11	AMOC values observed in the wetland and veg-ditch soils supported greater DP
12	compared to those in the cultivated soils. AMOC showed a significant correlation with
13	DP of nitrate amended soils during the four seasons (p<0.05). AMOC values of the
14	spring, summer, fall and winter significantly correlated with DP (p <0.05) with r value of
15	0.51, 0.57, 0.75 and 0.81, respectively (Table 3). Simple linear regression identified
16	significant influence of total soil N, C and bulk density on DP of the nitrate amended
17	soils with r^2 values of 0.80 0.78, and 0.80 respectively. Soil moisture content also
18	correlated significantly with DP in each of the four seasons (Table 4). The Pearson's
19	correlation coefficients of spring, summer, fall and winter soil moisture with DP were
20	0.46, 0.44, 0.69 and 0.57 respectively at p <0.05.
21	4. DISCUSSION

Significant differences in the DPs of nitrate ameneded soils of different land use
types (Figures 2) are attributed to differences in the position and management of

1	different land use types in the watershed. Low-lying heavy clay soil (Sharkey, Dowling
2	and Alligator soil series) under forested and depressional wetlands showed 6.3-fold
3	greater DPs than the Ag-high silt-loam soils. Groffman and Tiedje (1989) reported
4	similar observation of significant influence of soil texture and drainage (surrogates for
5	landscape position) class in forest soils on denitrification potentials. Mohn et al. (2000)
6	also found low denitrification rates in drained mounds compared to wetland soils. Our
7	data shows that landscape position is a significant regulator of DP in soils as poorly-
8	drained, fine-textured soils in natural wetlands supported higher denitrifier activity than
9	the coarse textured Ag-high soils. It is worthy to note that Ag-low soil had an average 2.5
10	times higher DP than that of Ag-high soils, which implies that topographic position led to
11	the explicit differences in DP of these sites.
12	As expected, forested, depressional and constructed wetlands exhibited 3.0, 2.0
13	and 2.1 times greater DP than the pedogenically similar (Table 2) Ag-low soils,
14	respectively. Drainage and cultivation of the Ag-low soils over the years led to lower
15	available organic C substrate (Table 2) and moisture contents (Table 4), which resulted in
16	lower denitrifier activity compared to similar soils under wetlands in the watershed. In
17	another study in the same watershed, we observed significantly greater denitrification
18	rates in forested wetlands than the rates of an adjacent cultivated site under variable soil
19	moisture contents (Ullah et al. 2005), demonstrating that cultivation diminished the
20	capacity of these soils to denitrify nitrate at rates similar to forested wetlands. The natural
21	vegetation cover and the resulting soil litter production in forested and depressional
22	wetlands provide higher organic carbon substrate for supporting greater denitrifier
23	activity than similar textured Ag-low soils. This observation implies that pedogenically

similar soils at the same elevation in a watershed under row crops cultivation sustain
 lower denitrification potential than under wetlands.

The higher denitrifier activity in the veg-ditches is attributed to its maintenance as a grassed waterway, which resulted in maintaining high soil moisture regime and producing greater AMOC contents compared to croplands and unveg-ditches. This finding supports the current practices of maintaining veg-ditches as a BMP for erosion control and water quality improvement as recommended by the USDA (ARS 2001) in the region.

9 The DPs observed in spring, summer and fall of all the land use types were 10 significantly greater (p < 0.05) than their DPs of winter except Ag.high and unveg-ditch 11 soils (Table 2), respectively. This observation suggests that the lower average winter soil 12 temperatures (ranging from 6 to 9 °C) and higher % WFPS substantially reduced soil 13 denitrifier activity (Table 4). Lower winter temperatures and higher WFPS percentage 14 compared to the summer values suppressed microbial activity (Magg et al. 1997) and 15 reduced the supply of mineralizable organic C to denitrifiers (Table 3) (Mohn et al. 2000; 16 Mogge et al. 1998; Klein and Logtestijn 1996). Even though %WFPS of soils in winter 17 were relatively lower than the fall %WFPS, lower winter temperature led to significantly 18 lower AMOC production than the fall values of all the land use types (Table 3). This 19 indicates that lower soil temperature exerts significant controls over microbial activity. 20 Lower denitrifier activity may pose greater risk of nitrate loss from agricultural 21 watersheds in winter.

Unlike the Ag-high and Ag-low soils, nitrate availability was found limiting DPs
in forested and depressional wetlands, because nitrate additions to these soils led to

1 increases in DPs during the four seasons compared to DPs of Ag-high and Ag-low sites 2 (Figures 2 and 3). When averaged over the four seasons, forested and depressional 3 wetlands showed 3 and 4-fold increase in DP under additional nitrate compared to the 4 unamended soils. Higher denitrification rates under additional nitrate can happen, if the 5 process is not limited by the availability of organic C (Weier et al. 1993). No response of 6 Ag-high and Ag-low soils to nitrate amendment shows that DP in these land uses was 7 limited by the availability of organic C (Table 1). Constructed wetland responded with 8 increased DP to nitrate additions in fall and winter only. The increase in DP of the 9 constructed wetland in response to nitrate additions after 6-9 months of its construction 10 demonstrate that wetland restoration on abandoned marginal lands in low-elevation areas 11 of the watershed can enhance denitrifier activity within and/or after the first growing 12 season. Veg-ditches also responded with increased DPs to nitrate additions. These results 13 depicts that wetlands and veg-ditches can denitrify additional nitrate coming from 14 cultivated soils or other external sources. This finding is in agreement with the findings 15 of similar studies conducted regarding nitrate removal potential of riparian wetlands and 16 vegetated buffer-strips in agricultural watersheds (Lindau et al. 1994; Lowrance et al. 17 1995; Groffman and Crawford 2003; Groffman et al. 2002; Ingrid-Brettar et al. 2002; 18 Clement et al. 2002; DeLaune, et al. 2005). 19 As expected, higher total soil carbon, mineralizable organic C, wetter soil 20 conditions, greater soil porosity and fine-clay texture of natural and constructed wetlands

21 (Table 2) supported greater denitrifier activity than those observed in the Ag-high and the

22 pedogenically similar Ag-low and unveg-ditch soils. This result is in agreement with the

23 findings of Hill and Cardaci (2002), Davidsson and Stahl (2000), and Groffman and

1 Tiedje (1989) who reported significant control of denitrification potential by available 2 organic carbon, soil moisture contents and soil texture classes. At the biogeochemical 3 scale, soil moisture content, soil texture, available C and nitrate are the major regulators 4 of denitrifier activity in soils (Patrick and Reddy, 1976; Myrold et al. 1998; Weitz et al. 5 2001) and the same factors were found significantly affecting DPs in the Beasley 6 watershed. The status of these biogeochemical scale regulators of DP is a result of the 7 interaction of soil oxygen diffusion rate/dynamics, landscape position, plant community 8 structure, physical disruption (landuse), ecosystem type, microbial biomass, and organic 9 matter production characteristics of a given landscape unit (Myrold et al. 1998; Florinsky, 10 et al. 2004). Florinsky et al. (2004) reported significant control of denitrifier activity in 11 soils by landscape position of a site and its subsequent influence on soil moisture and 12 organic carbon availability. The authors concluded that higher denitrifying activity in 13 low-elevation areas was mostly affected by the redistribution and accumulation of soil 14 moisture and available organic carbon due to their additional gains from high-elevation 15 areas along the slope (Florinskey, et al. 2004). In the Beasley watershed, lower elevation 16 land uses under wetlands maintained higher soil moisture contents, greater soil organic 17 carbon pools, and lower bulk densities that resulted in higher DP in wetland soils than 18 either the high elevation or low elevation soils under cultivation. These findings show 19 that landscape position and landuse type exerts significant controls on denitrifying 20 activity in a watershed. Any effort to model and predict denitrification potential at watersheds scale such as the LMV should include topographic and land use aspects of a 21 22 watershed as key controllers of DP besides the biogeochemical scale variables in the 23 modeling process. Such an approach can help compute/predict relatively more realistic

DPs at watershed scale and identify sites for restoration in large river basins (White and
Fennessy, 2005). Application of digital terrain models for predicting microbial activities
of watersheds based on inputs of both biogeochemical and landscape scale variables into
the model have been found feasible in Canada (Florinsky et al. 2004).

5 The results of this research demonstrate that denitrification potential in soils is 6 regulated by more than one biogeochemical scale factor and is significantly influenced by 7 both the landscape position and land use types in a watershed. Marshal (1999), Groffman 8 et al. (1999), and Florinsky et al (2004) also reported that denitrification cannot be 9 predicted on the basis of one factor, rather a combination of factors are involved in the 10 regulation of the denitrification process. Therefore, it is suggested to consider landscape 11 scale variables (landscape position and land use type) in addition to biogeochemical scale 12 variables in assessing/modeling denitrification potential of soils and identification of sites 13 for wetland restoration for water quality improvement in the LMV. Our findings suggest 14 that fine-textured low-elevation sites in agricultural watersheds are the best candidates for 15 wetland restoration for nitrate removal in the LMV. Such sites can accumulate higher soil 16 organic carbon and retain higher soil moisture for sustaining persistent and enhanced 17 denitrification rates than coarse-textured Ag-high soils in agricultural watersheds.

18

ACKNOWLEDGEMENTS

The authors are thankful to Dr. Kewei Yu, Louisiana State University Wetland
Biogeochemistry Institute, Baton Rouge, Louisiana and Mr. Michael Baldwin, USGS
National Wetlands Research Center, Louisiana for their help during field sampling and
laboratory analysis. We also thank USDA National Sedimentation Laboratory, Oxford,

23 Mississippi for financially supporting this project.

1	REFERENCES
2 3 4 5	Allen, J. A., Keeland, B. D., Stauturf, J. A., Clewell, A. F., and Kennedy, H. E. Jr. 2001. A guide to bottomland hardwood restoration. Technical Report, USGS/BRD/ITR-2000- 0011, 132 p.
6 7 8 9	ARS. 2001. The Mississippi delta management systems evaluation areas project, 1995- 1999. Information Bulletin 337, Office of Agricultural Communication, Division of Agriculture, Forestry, and Veterinary Medicine, Mississippi State University.
10 11 12 13 14	Battaglin, W. A. and D. A. Goolsby. 1996. Spatial data on GIS format on agricultural chemical use, land use and cropping practices in the United States, U.S. Geological Survey, Water Research Institute, Rep. 94-4176. (available at www.water.usgs.gove/pubs/wri/wri944176).
15 16 17 18	Beauchamp, E.G. and D.W. Bergstrom. 1993. Denitrification, in M. R. Carter (Ed.) Soil Sampling and Methods of Analysis. Canadian Society of Soil Science, Lewis Publishers, Canada.
19 20 21	Bradley P. M., M. Fernadex, Jr. and F. H. Chapelle. 1992. Carbon limitation of denitrification rates in an anaerobic groundwater system. Environ. Sci.Tech. 26: 2377-2381.
23 24 25 26	Clement, J. C., G. Pinay, and P. Marmonier. 2002. Seasonal dynamics of denitrification along topohydrosequences in three different riparian wetlands. J. Environ. Qual. 31: 1025-1037.
20 27 28 29 30	Comin, A. F, J. A. Romero, V. Asorga, and C. Garcia. 1997. Nitrogen removal and cycling in restored wetlands used as filters of nutrients for agricultural run-off. Water Sci. Tech. 35: 255-261.
31 32 33	Davidsson, E.T. and Mattias Stahl. 2000. The influence of organic carbon on nitrogen transformations in five wetland soils. Soil.Sci.Soc.Am. J. 64:1129-1136.
34 35 36	DeLaune R.D., R.R. Boar, C.W. Lindau and B.A. Kleiss. 1996. Denitrification in bottomland hardwood wetland soils of the Cache River. Wetlands 16: 309-320.
37 38 39 40	DeLaune, R. D., A. Jugsujinda, J. L. West, C. B. Johnson, and M. Kongchum. 2005. A screening of the capacity of the Louisiana freshwater wetlands to process nitrate in diverted Mississippi River water. Ecological Engineering 25: 315-321.
41 42 43	Donner, S.D. 2004. Impact of changing land use practices on nitrate export by the Mississippi River. Global Biogeoch. Cycles: 18, GB1028, doi:10.1029/2003GB002093.

- 1 EPA. 2001. Action Plan for Reducing, Mitigating, and Controlling Hypoxia in the 2 Northern Gulf of Mexico. Washington DC: Office of Wetlands, Oceans, and Watersheds, 3 US EPA. 4 5 Fisk, N. H. 1951. Mississippi River geology relation to river regime. American Society of 6 Civil Engineers Transactions, paper No: 2511:667-682. 7 8 Florinsky, I. V., S. McMahon and D. L. Burton. 2004. Topographic control of soil 9 microbial activity: a case study of denitrifiers. Geoderma 119: 33-53. 10 11 Gale P.M., Reddy K.R. and D.A. Graetz. 1992. Mineralization of sediment organic 12 matter under anoxic conditions. J. Environ. Qual. 21:394-400. 13 14 Galloway, J. N. and E.B. Cowling. 2002. Reactive nitrogen and the world: Two hundred 15 years of change. Ambio 31: 64-71. 16 17 Galloway, J. N., J. D. Aber, J. W. Erisman, S. P. Seitzinger, R. W. Howarth, E. B. 18 Cowling and B. J. Cosby. 2003. The nitrogen cascade. BioScience 53: 341-356. 19 20 Groffman, P. M. and J. M. Tiedje. 1989. Denitrification in north temperate forests: spatial 21 and temporal patterns at the landscape and seasonal scales. Soil Biol.Biochem. 21: 613-22 620. 23 24 Groffman P.M., Holland E., Myrold D.D., Robertson G.P. and Zou X. 1999. 25 Denitrification. In Robertson G.P., Bledsoe C.S., Coleman D. C. and Sollins P. (eds), 26 Standard Soil Methods for Ecological Research. Oxford University Press, NY, pp. 272-27 288. 28 29 Groffman, P. M., N. J. Boulware, W. C. Zipperer, R. V. Pouyat, L. E. Band., and M. F. 30 Colosimo. 2002. Soil nitrogen cycle processes in urban riparian zones. Envorn. Sci. 31 Technol. 36: 4547-4552. 32 33 Groffman. P. M, and M. K. Crawford. 2003 Denitrification potential in urban riparian 34 zones. J. Environ. Qual. 32: 1144-1149. 35 36 Hill, R. A., and M. Cardaci. 2004. Denitrification and organic carbon availability in 37 riparian wetland soils and sub-surface sediments. Soil. Sci. Soc. Am. J. 68: 320-325. 38 39 Howarth, R. W., E. W. Boyer, W.J. Pabich and J. N. Galloway. 2002. Nitrogen use in the 40 United States from 1961-2000 and potential future trends. Ambio 31:88-96. 41
- 42 Hunter, R.G. and S.P. Faulkner. 2001. Denitrification potential in restored and natural
- 43 hardwood wetlands. Soil Sci. Soc.Am. J. 65:1865-1872. 44

1 2 3 4	Ingrid-Brettar, and M.G. Hofle. 2002. Close correlation between the nitrate elimination rate by denitrification and the organic matter content in hardwood forest soils of the upper Rhine floodplain (France). Wetlands 21: 214-224.
5 6 7 8	Jordan, E.T, D. E. Weller and D. L. Correl. 1998. Denitrification in surface soils of a riparian forest: Effects of water, nitrate and sucrose additions. Soil Biol.Biochem.30: 833-844.
9 10 11 12	Klein De C.A.M., and R.S.P. Van Logtestijn. 1996. Denitrification in grassland soils in the Netherlands in relation to irrigation, N-application rate, soil water content and soil temperature. Soil Biol.Bioch. 28:231-237.
13 14 15	Lindau, C. W., R. D. Delaune, and J. H. Pardue. 1994. Inorganic nitrogen processing and assimilation in forested wetland. Hydrobiologia 277: 171-178.
16 17 18 19 20 21	Locke, A. M., W.J. Staddon and R.M. Zablotowics. 2001. Nutrient distribution in reduced tillage and conventional tillage cotton soils. In Rebich, A. R., and S. Knight (editors). The Mississippi Delta Management Systems Evaluation Areas Project, 1995-1999. Information Bulletin 377, Office of Agricultural Communication, Mississippi State University.
22 22 23 24 25	Lowrance, R., R. Todd, J. Fail Jr., O. Hendrickson, Jr., R. Leonard, and L. Asmussen. 1984. Riparian forests as nutrient filters in agricultural watershed. BioScience 34:374-377.
26 27 28	Lowrance, R., G. Vellidis, and R. K. Hubbard. 1995. Denitrification in a restored riparian forest wetland. J. Environ. Qual. 24: 808-815.
20 29 30 31	Lowrance, R. et al. 1997. Water quality functions of riparian forest buffers in Chesapeake Bay watersheds. Environ. Manag. 21: 687-712.
32 33 34	Maag, M., M. Malinovsky, and S.M. Nielsen. 1997. Kinetics and temperature dependence of potential denitrification in riparian soils. J. Environ. Qual. 26:215-223.
35 36 37 38	Marshal, S.B., M.L. Cabrera, L.C. Braun, C.W. Wood, M.D. Mullen and E.A. Guertal. 1999. Denitrification from Fescue pastures in the Southeastern US fertilized with broiler litter. J. Environ. Qual. 28: 1978-1983.
39 40 41 42	Mitsch, W. J. and J. W. Day, Jr. 2006. Restoration of the Mississippi-Ohio-Missouri (MOM) River basins: Experience and needed research. Ecological Engineering 26: 55-69.
43 44	Mitsch, W. J., J. W. Day, Jr., L. Zhang, and R. R. Lane. 2005. Nitrate-nitrogen retention in the wetlands in the Mississippi River basin. Ecological Engineering 24: 267-278.

1 2 3	Mitsch, W.J, J.W. Day, Jr., J.W. Gilliam, P.M. Groffman, D.L. Hey. 2001. Reducing nitrogen loading to the Gulf of Mexico from the Mississippi River basin: Strategies to counter a persistent ecological problem. Bioscience 51:373-388.
5 6 7 8	Mogge, B., E. A. Kaiser, and J. C. Munch. 1998. Nitrous oxide emissions and denitrification N-losses from forest soils in the Bornhoved Lake region (Northern Germany). Soil Biol. Biochem. 30: 703-710.
9 10	Mohn J., A. Schürmann, F.Hagedorn, P.Schleppi and R. Bachofen. 2000. Increased rate of denitrification in nitrogen-treated forest soils. Forest Ecol.Manag.137:113-119.
11 12 13 14	Myrold, D. David.1998. Transformations of nitrogen. In Sylvia, D.M, et al (Eds.) Principles and Applications of Soil Microbiology, Prentice-Hall, Inc, New Jersey.
15 16 17	NRCS. 1959. Soil Survey of Sunflower County, Mississippi. Soil Conservation Service and Mississippi Agricultural Experiment Station.
17 18 19 20	Parkin, T. B., H. P. Kasper, A.J. Sexstone, and J. M. Tiedje. 1984. A gas-flow soil core method to measure field denitrification rates. Soil. Biol. Biochem. 16: 197-228.
20 21 22 23 24	Patrick, W.H. Jr. and K. R. Reddy. 1976. Nitrification-denitrification reactions in flooded soil and water bottoms: Dependence on oxygen supply and ammonium diffusion. J. Environ. Qual. 5: 469-472.
24 25 26 27	Perterjohn, W. T., and D. L. Corell.1984. Nutrient dynamics in an agricultural watershed: Observation on the role of riparian forests. Ecology 65: 1466-1475.
27 28 29 30	Rabalais, N. N., R. E. Turner, and D. Scavia. 2002. Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River. BioScience 52: 129-142.
30 31 32 33 34 35 26	Rebich, R. A. 2001. Quality of runoff in the Mississippi delta management systems evaluation areas project, published in The Mississippi Delta Management Systems Evaluation Areas Project, 1995-1999, Information Bulletin 337, Office of Agricultural Communication, Division of Agriculture, Forestry, and Veterinary Medicine, Mississippi State University.
30 37 38	SAS Institute. 1998. SAS User's Manual. SAS Inc. Carry, North Carolina, USA.
39 40 41	Sexstone, A.J., T.B. Parkin and J. M. Tiedje. 1985. Temporal response of soil denitrification rates to rainfall and irrigation. Soil Sci. Soc. Am. J. 49: 99-103.
42 43 44	Sheldrick B. H. and C. Wang. 1993. Particle size distribution. In Soil Sampling and Analysis, Edited by M.R. Carter, Canadian Society of Soil Science, Lewis Publishers.

- 1 Singh-Bijay, J.C. Ryden and D.C. Whitehead. 1988. Some relationships between
- 2 denitrification potential and fractions of organic carbon in air-dried and field moist soils.
- 3 Soil Biol.Biochem. 20: 737-741.
- 4
- 5 Skiba, U. M., L. J. Sheppard, J. MacDonald and D. Fowler. 1998. Some key
- 6 environmental variables controlling nitrous oxide emissions from agricultural and semi-
- 7 natural soils in the Scotland. Atmospheric Environment 32: 3311-3320.
- 9 Smith, R. A., R. B. Alexander, and M. G. Wolman. 1987. Water quality trends in the 10 nation's rivers. Science 235: 1608-1615.
- 11
- Turner, R. E., and N. N. Rabalais. 1994. Coastal eutrophication near the MississippiRiver delta. Nature 368: 619-621.
- 14
- Turner, R. E and N. N Rabalais. 2003. Linking landscape and water quality in the
 Mississippi River basin for 200 years. BioScience 53:563-572.
- 17
- Ullah, S., G.A. Breitenbeck and S. P. Faulkner. 2005. Denitrification and N₂O emissions
 from cultivated and forested alluvial clay soils. Biogeochemistry 73: 499-513.
- USEPA. 1996. Report to congress on the National Water Quality Inventory. US Gov.
 Print Office, Washington DC.
- 23

- Weier, K. L., J. W. Doran, J. F. Power and D. T. Walters. 1993. Denitrification and the
 dinitrogen/nitrous oxide ratio as affected by soil water, available carbon and nitrate. Soil
 Sci. Soc. Am. J. 57: 66-72.
- 27
- Weitz, A. M., E. Linder, S. Frolking, P.M. Crill, and M. Keller. 2001. N₂O emissions from humid tropical agricultural soils: effects of soil moisture, texture and nitrogen
- availability. Soil Biol.Biochem.33: 1077-1093.
- 31
- 32 White, D. and S. Fennessy. 2005. Modeling the suitability of wetland restoration
- 33 potential at watershed scale. Ecological Engineering 24: 359-377.
- 34

1 FIGURES LEGEND

2 Figure 1. Location map of the Beasley Lake watershed, Mississippi, USA.

3

4 Figure 2. Denitrification potential of different land use types amended with nitrate in

- 5 spring, summer, fall and winter with standard error of the means (same letters on top of
- 6 each bar show no significant differences in DPs among the land uses within each season 7 at p = 0.05).
- 8
- 9 Figure 3. Denitrification potential of different unamended land use types in spring,
- 10 summer, fall and winter with standard error of the means.



Figure 1. Location map of the Beasley Lake watershed, Mississippi, USA.

Soil parameters	Forested wetland	Dep.wetland	Constructed wetland	Veg-ditches	Unveg-	Ag-low	Ag-high
					ditches		
Bulk Density (g cm ⁻³)	0.90 (0.03) a	0.96 (0.01) a	1.20 (0.2) b	1.17 (0.03) b	1.24 (0.4) b	1.24 (0.9) bc	1.31 (0.4) c
Porosity cm ³ cm ⁻³	0.66 (0.98) a	0.64 (0.99)a	0.55 (0.91)b	0.56 (0.98) b	0.53 (0.98) b	0.53(0.96) bc	0.51(0.98) c
Clay (%)	51 (0.5) a	47 (1.2) a	45 (1.2) a	51 (2.5) a	45 (5.4) a	48 (2.1) a	23 (0.001) b
Silt (%)	47 (0.7) a	49 (3) a	45 (1.6) a	44 (3.5) a	33 (2.8) b	46 (1.3) a	65 (0.8) c
рН	5.4 (0.05) a	5.3 (0.03) a	5.5 (0.07) a	5.6 (0.07) a	5.6 (0.04) a	6.1 (0.05) b	6.3 (0.03) b
Total C 0-10 cm(Mt ha ⁻¹)	297 (30) a	167 (18) b	218 (9) b	189 (11) b	156 (44) bc	168 (13) bc	85 (26) c
Total N 0-10 cm (Mt ha ⁻¹)	26 (2) a	17 (1.4) b	23 (0.9) b	24 (2.2) b	13 (2.6) c	11 (1.0) c	5 (3.5) d
NO ₃ (mg kg soil ⁻¹)	5.7 (1.7) a	6.5 (2.5) a	6.1 (2.8) a	7.6 (3.2) a	7.4 (1.9) a	8.0 (1.6) a	7.7 (3.2) a

Table 2. Mean Physio-chemical properties of soils of the7 land use types in Beasley watershed. Means (S.E of the means) followed by the same lower-case letter indicate no significant difference (p > 0.05) among the different land use types.



Figure 2. Denitrification potential of different land use types amended with nitrate in spring, summer, fall and winter with standard error of the means (same letters on top of each bar show no significant differences in DPs among the land uses within each season at p = 0.05).



Figure 3. Denitrification potential of different land use types (no nitrate amended soils) in spring, summer, fall and winter with standard error of the means.

Table 1. Yearly average denitrification potential (DP) of the 7 landuse types with post-ANOVA comparisons between 7 land use types. Same lower-case letters following mean values shows no significant difference among the different land use types at p = 0.05. Effects of NO₃ amendment of soil slurries and season on denitrification potential of the 7 land use types are also shown using pooled variance two-sample t-test and ANOVA, respectively at p <0.05.

Land use types	DP of NO ₃ amended soils	NO ₃ addition	Seasonal
	$(ug N g^{-1} h^{-1})$	effects on DP	effects on DP
Forested wetland	1.18 (0.13) a	*	*
Depressional wetland	0.77 (0.03) b	*	*
Constructed wetland	0.82 (0.10)b	*	*
Veg-ditches	0.66 (0.17) bc	*	*
Unveg-ditches	0.50 (0.09) c	*	ns
Low-elevation Agric.	0.40 (0.08) cd	ns	*
High-elevation Agric.	0.16 (0.02) d	ns	ns

*: significant effect, ns: non-significant effect

Season	AMOC (μ g C-CO ₂ produced g ⁻¹ soil h ⁻¹)									
	Forested wetland	Dep. wetland	Constructed wetland	Veg- ditches	Unveg- ditches	Ag-low	Ag-high	Pearson's correlation with DP (n=57)		
Spring	0.78 (0.10)	0.63 (0.07)	0.82 (0.11)	0.77 (0.07)	0.14 (0.03)	0.46 (0.00)	0.35 (0.07)	0.51*		
Summer	1.26 (0.13)	1.29 (0.03)	1.21 (0.05)	0.91 (0.05)	0.85 (0.04)	0.88 (0.03)	0.89 (0.04)	0.57*		
Fall	1.03 (0.02)	1.04 (0.03)	0.99 (0.04)	0.98 (0.05)	0.89 (0.03)	0.91 (0.02)	0.66(0.01)	0.75*		
Winter	0.86 (0.04)	0.70 (0.04)	0.70 (0.04)	0.52 (0.03)	0.62 (0.03)	0.59 (0.06)	0.72 (0.02)	0.81*		

Table 3. Mean anaerobically mineralizable organic carbon content of the 7 land use types with standard error of the means and its Pearson's correlation coefficients with DP

* Significant difference at p< 0.05

	Water-filled pore space (%)						
Land use Types	Spring	Summer	Fall	Winter			
Forested wetland	62 (1.0)	48 (0.9)	59 (1.0)	53(1.0)			
Depressional wetland	63 (1.0)	28 (1.0)	52 (1.0)	44 (1.0)			
Constructed wetland	51(1.0)	29 (1.0)	55 (1.0)	48 (1.0)			
Veg-ditches	47 (1.0)	61 (1.0)	68 (.9)	50 (1.0)			
Unveg-ditches	49 (1.0)	60 (1.0)	58 (1.0)	51 (1.0)			
Low-elevation croplands	56 (1.0)	34 (1.0)	56 (1.0)	45 (1.0)			
High-elevation croplands	34 (1.0)	28 (1.0)	45 (1.0)	32 (1.0)			
Correlation of soil moisture with DP	0.46*	0.44*	0.69*	0.57*			
Soil Temp. ^o C	22	28	21	9			

Table 4. Seasonal water-filled pore space percentages with standard error of the mean and its correlation co-efficient with denitrification potential of the 7 land use types. Last row of the table shows seasonal average soil temperatures of Beasley watershed.

*Significant at p < 0.05, and n = 56