

1 **1. Introduction**

2 Ideas of interdisciplinarity have often found practical expression in research at the
3 interface of environmental science and policy. Indeed, ‘joined-up’ thinking is widely
4 considered a precondition for developing holistic and integrated responses to problems of
5 sustainability in society (Holdren, 2008); a process in which the natural and cultural
6 dimensions of environmental issues can be marshalled around a common sense of
7 intellectual purpose and endeavour. And yet, if it is one thing to assert the idea of
8 interdisciplinary research as central to the ambitions of sustainability science, it is quite
9 another to devise policy interventions that capture and reconcile often divergent
10 understandings of a given problem. After all, a common aspect of interdisciplinarity is
11 that it pulls together expertise in ways that make approaches to problem solving *not only*
12 multidimensional in outlook, but coherent in their conception of the world. This is a
13 daunting task not least because the issues upon which agendas for environmental science
14 and policy seek to act are often emergent and novel in form. Interdisciplinary research
15 means, in part, devising new theoretical constructs in which assessments of
16 environmental problems can be made.

17 One of the contexts in which this problem is keenly felt, and which forms the focus of
18 this paper, is in the development of holistic approaches to assess environmental risk.
19 Developing these assessments is arguably key to the political and regulatory culture of
20 contemporary environmental management, but as we shall see, the process of
21 conceptualising and operationalising them in an interdisciplinary fashion is fraught with
22 theoretical and methodological challenges. The purpose of this paper is to consider what
23 some of those challenges are by drawing on a recent research project examining the issue
24 of microbial pollution of surface waters.

1 In particular, the paper reflects critically upon efforts to develop an interdisciplinary
2 assessment of the factors that may affect the transfer of potential pathogens from
3 livestock farming systems to surface waters. This is a significant concern of the
4 contemporary environmental protection agenda (Kay et al., 2007). Manures and excreta
5 produced by grazing livestock are an important resource for farmers, but public exposure
6 to the pathogenic microorganisms contained within these can lead to gastro-intestinal
7 illness, for example infection caused by *E. coli* O157. Pathways to exposure may occur in
8 a variety of ways, such as by ingesting contaminated waters through downstream
9 recreational bathing (Graczyk et al., 2007) and through food consumption, for instance by
10 eating contaminated shellfish (Schets et al., 2007). Alongside these consequences for
11 human health, microbial pollution carries with it an economic burden. In the UK it has
12 been suggested that gastro-intestinal illness caused by *E. coli* O157 alone is estimated to
13 cost over £1 billion per annum in lost productivity and around £30 million annually in
14 healthcare (Jones, 1999). Moreover, the economic ‘knock-on’ effects of illness among
15 human populations are potentially very significant for rural areas where agricultural,
16 leisure and tourist economies are dependent on wider consumer confidence.

17 This paper explores the prospects for developing a holistic assessment of the factors
18 that underlie and create these microbial risks¹ as the basis for targeted mitigation at the
19 farm scale. It begins by introducing the wider policy and regulatory context in which a
20 concern to manage microbial risks is now situated, and goes on to explain how an
21 interdisciplinary approach might begin to capture the cultural² and natural processes that
22 come to influence them. In particular, the paper describes a procedure in which a
23 hypothetical construction of factors driving the transfer of microbial pollutants from land

¹ Microbial risk is understood in this paper as the product of likelihood of microbial contamination and magnitude of microbial impact on watercourses.

² In this paper we use the term ‘cultural’ as shorthand for the social and economic relations that define environmental risk, since in much environmental scholarship it is the term ‘culture’ that is dialectically related to the word ‘nature’, and with it, the concerns of natural science. See for example Whatmore (2002).

1 to surface waters was characterised and then subject to deeper rationalisation and
2 refinement through the technique of expert elicitation. As a result the paper addresses
3 some of the theoretical and methodological issues and conundrums that emerge from
4 attempting to conceptualise an operational risk system by bringing quite different
5 disciplinary positions to bear upon its construction, and with this, identifies some of the
6 issues arising for environmental researchers when seeking to turn specialist assessments
7 of risk into transdisciplinary - 'whole system' - knowledge. At the heart of the paper's
8 argument is the need to come to terms with contrasting assessments of how theoretical
9 representations of environmental risk work, and further, the need to address inherent
10 differences in how cultural and natural processes can be accounted for. As a result of this
11 analysis we move towards a conclusion that highlights the need for building critical
12 'unknowns' into assessments of 'known' risks. The paper suggests that sustainability
13 science, such as work conducted on the issue of pathogen transfer from livestock farming
14 practices, must recognise the partial nature of its assessments. How these uncertainties
15 are conveyed in the uptake and application of risk assessment tools and models is, we
16 suggest, just as important as seeking to render complex risks amenable to final prediction
17 by way of empirical knowledge.

18

19 **2. Research Context**

20 The insights of this paper were developed under the auspices of a research project
21 examining how farm management practices and circumstances can promote, limit and
22 prevent the risk of microbial watercourse contamination³. The study employed a Faecal

³ The research was funded under UK research councils 'Rural Economy and Land Use programme' (RELU), project code: RES-224-25-0086 (www.relu.ac.uk). Its focus was on the Taw Catchment in North Devon, UK, and has involved: an extensive survey of farmers across the Catchment eliciting attitudes and practices towards manure, land and livestock management; detailed microbial monitoring of water courses on ten of these farms, a review and mitigation methods; and a programme of public participation and

1 Indicator Organism (FIO) approach to researching and understanding pathogenic risks.
2 Present within all human and livestock faeces, FIOs are bacteria indicative of faecal
3 pollution and therefore potential pathogen presence. Central to the concerns of this
4 paper, the project has also sought to develop a FIO risk assessment tool for farms, one
5 designed to conceptualise the physical and cultural factors influencing the loss of faecal
6 bacteria as the basis for targeted policy inventions.

7 The water policy context to these efforts has been described elsewhere (see Kay et al.,
8 2008a). Designing risk assessment management plans are increasingly central to the
9 regulatory environment within which farm practices now take place. In the UK for
10 instance, state and quasi-state apparatus responsible for issues of environmental
11 protection have introduced a range of mandatory measures and guidance for helping
12 farmers to limit contamination of surface waters at the farm scale. These include the
13 Department of Environment, Food and Rural Affairs soil and manure management plans
14 (Defra, 2003; 2005a) and the Scottish Environmental Protection Agency's risk
15 assessment for manures and slurries (SEPA, 2004). These frameworks are underpinned
16 by more general responsibilities to 'codes of good agricultural and environmental
17 condition' and are central to the conditions under which farmers can participate in new
18 environmental stewardship programmes, such as Defra's Entry and Higher level
19 stewardship schemes (Defra, 2005b). The wider agricultural policy context driving these
20 developments in the UK and elsewhere has been well rehearsed (Wilson, 2007).
21 European farmers have been placed increasingly at the centre of a 'public goods' model
22 of the countryside where mandatory responsibilities for environmental protection at the
23 farm and field level exist alongside an increasing 'marketisation' of environmentally
24 friendly products (Buller and Morris, 2004).

debate in which emerging scientific evidence was explored in relation to wider understandings and assessments of environmental risk.

1 Within this policy context there is a need to develop sustainable food systems that, by
2 way of integrated approaches to land management, can effectively manage the wider, and
3 often diffuse, impacts of agriculture on society and economy. Managing for the human
4 health implications of ingesting pathogenic microorganisms either through recreational
5 bathing or the consumption of food is an important driver for fostering assessment
6 frameworks to assist both mandatory and voluntary action, not least given the exigencies
7 of emerging legislative frameworks such as the EU Water Framework Directive (Kaika,
8 2003) and the EU Bathing Water Directive (Georgiou and Bateman, 2005). Thus, one of
9 the purposes of our research has been to develop a conceptual framework within which
10 potential pathogenic risks can begin to be assessed. What this amounts to, in effect, is an
11 aspiration to construct a ‘whole system’ conception of microbial risk that can ultimately
12 inform the development of a prototypical risk assessment tool for practical use by policy
13 makers, environmental managers and farmers. A characterisation of this tool is reported
14 on elsewhere (Oliver et al., 2009). The purpose of this inquiry is to examine the issues
15 embedded in setting parameters around this tool’s underlying conception of risk and then
16 eliciting values for it.

17

18 **3. Characterising an Interdisciplinary Risk System**

19 As Chadwick *et al.* (2008) have argued specifically in the context of microbial
20 watercourse pollution, understandings of environmental risk need to be grounded in a
21 diversity of disciplinary perspectives. We are a research team comprising expertise in
22 microbiology, hydrochemistry, soil hydrology, manure management, agricultural
23 geography and rural sociology. We have worked on the basis that this expertise forms a
24 reasonably diverse ‘experience-base’ to help identify what the cultural and natural

1 parameters of a FIO risk system⁴ may, in principle, be. We say ‘experience base’ rather
2 than simply ‘evidence base’ because the process of identifying discrete factors or
3 variables that may constitute a set of whole system parameters means combining both
4 reliable knowledge into system variables (i.e. factors that we have confidence will have a
5 bearing on FIOs as a result of empirical study) but also working assumptions or
6 ‘hunches’, based on anecdotal evidence, about variables that may or may not have been
7 studied in the context of pathogens (i.e. factors that seem to be relevant to FIO behaviour
8 but of which we cannot be sure). In this we have found it useful to draw an analytical
9 distinction between *direct* and *indirect* drivers of FIO risk in defining the characteristics
10 of our risk system.

11 Direct drivers of risk concern variables in our hypothetical system that have an
12 immediate and tangible control over the potential occurrence of FIO transfers from land
13 to water. They are factors that operate at a practical and physical scale of expression, and
14 imply causal connections that are, in principle, observable and definable within farm
15 environments. Direct drivers presume, and to some extent may already encompass, an
16 emerging evidence base in order to validate their place within a risk system. In general
17 terms, the direct drivers we address in this study reflect the empirical work of natural
18 science research, and in our system is reflected in a four-fold categorisation. In particular
19 this categorisation draws inspiration from other environmental risk indexing work most
20 notably the empirically informed 1990 phosphorus index (see Lemunyon and Gilbert
21 1993). It encompasses issues of:

- 22 ▪ Source - incorporating those factors that determine the fundamental presence
23 and origin of FIOs within farm environments. These can include: livestock
24 grazing density (Aitken, 2003); faecal inputs and their associated FIO die-off

⁴ By ‘risk system’ we mean a definable and hypothetically bounded set of variables that characterise the drivers of an environmental risk, in this case, the transfer of FIOs from agricultural land to surface waters.

1 rates (Oliver et al., 2006); manure application methods (Hutchison et al., 2004);
2 and grazing duration (White et al., 2001).

- 3 ▪ Transfer - encompassing those factors encouraging FIOs to start their journey to
4 a watercourse including, for example: field slopes and shapes (Abu-Ashour and
5 Lee, 2000; Crowther et al., 2003), soil hydrological properties (Artz et al., 2005)
6 and cell interactions with soil particles (Muirhead et al., 2006).
- 7 ▪ Connectivity - encapsulating those variables that link source and transfer
8 mechanisms to a receiving water such as artificial field drainage (Oliver et al.,
9 2005), overland flow distance (Collins et al., 2005) and direct defecation of
10 excreta into streams (Davies-Colley et al., 2004).
- 11 ▪ Infrastructure - embodying human material interventions and constructions in
12 the landscape that underpin the presence and movement of FIOs in a given farm
13 environment (e.g. Edwards et al., 2008). These can include: manure storage
14 capacity and associated conditions (Smith et al., 2001) but also point source
15 inputs of FIOs associated with the hard standings and buildings used by
16 livestock (Kay et al., 2008b).

17 In contrast, indirect drivers of risk have a more tacit influence on FIO risks. They
18 encompass those factors that condition FIO risk more widely. For instance, while it may
19 be possible to infer that the source risk of 'grazing density' tangibly influences the
20 magnitude of FIO risks in a given farm landscape since variations here will govern the
21 number and rate of organisms excreted onto pasture, the issue of grazing density is, of
22 course, a land management choice shaped by a wider set of political, economic and
23 attitudinal influences. Indirect drivers are therefore designed to enhance the explanatory
24 potential of a risk system by acknowledging issues of both 'structure' (for instance, the
25 influence of policy measures and market incentives) and 'agency' (for example, the effect

1 of values) in processes of farmer decision making. This is a key, and longstanding,
2 analytical distinction within the study of social relations (Giddens, 1984) and central to
3 many theorisations of agricultural change (Wilson, 2007). Indeed, it is in the interplay of
4 these factors that rural studies have sought to make sense of where key determinants of
5 farmer action lie (see for instance, Wilson and Hart, 2000; Burton and Wilson, 2006). Yet
6 unlike issues of 'source', 'transfer', 'connectivity' and 'infrastructure', here it is more
7 awkward to identify a set of variables that can be packaged into a logical and discrete
8 sequence of influences on microbial watercourse pollution, or a set factors that can be
9 systematically linked to material processes at the farm level. Economic and political
10 drivers of FIO risk, for instance, originate in contexts far exceeding the farm gate, such as
11 political arenas that shape prevailing market and regulatory conditions, while farmer
12 attitudes and values are not only difficult to uncover, but tend to be multifaceted and
13 unstable (Burton and Wilson, 2006), and therefore defy neat typological prescription
14 (Fish et al., 2003). One of the vexed tasks facing the development of an interdisciplinary
15 risk system is therefore to translate a diffuse and nebulous set of issues into a functional
16 set of factors; ones amenable to the pragmatic demands of risk assessment. Accepting
17 that these structural and attitudinal influences are part of a spectrum of overlapping
18 influences, and that, unlike direct drivers, no evidence base exists for understanding
19 determinants of risk in a specifically FIO context, our research suggests that indirect
20 drivers of risk can be categorised in four key ways:

- 21 ▪ *Income and resources* - addressing the material and financial resources that
22 influence the uptake of measures to limit and prevent FIO risks occurring. Resource
23 constraints (such as access to labour) have been widely recognised to influence
24 farmer/enterprise decision making (e.g. Lobley and Potter 2004).

- 1 ▪ Knowledges and affiliations - encompassing the conditions under which the
2 intellectual capital to manage for FIO risks may be communicated, learnt and
3 ultimately deployed. This element attends to the wider social production and
4 exchange of information and knowledge amongst communities of farmers (such as
5 training and accreditation), a key determinant of enterprise behaviour (Tsouvalis et
6 al., 2000)
- 7 ▪ Non-agricultural activities - reflecting the influence of wider recreational
8 associations with landscape on management attitudes to water quality risks. This
9 component situates farm behaviour in the context of multifunctional land use (such
10 as recreational pursuits taking place on land); and therefore indexes a key
11 explanatory framework behind motivations to act in contemporary agricultural
12 systems (Wilson, 2007)
- 13 ▪ Farm structure and trajectory - incorporating variables that link capacities and
14 inclinations to manage for risk with prevailing household and enterprise
15 circumstances, (such as type of ownership and stage in lifecycle). This element
16 interprets decision making as a function of more tacit ‘path-dependencies’ (Wilson
17 2008), both economic and cultural in kind. Explicit recognition to the influence of
18 biography and economic transition is given credence in these aspects of the
19 characterisation.

20 A detailed overview of the variables that comprise these direct and indirect
21 categorisations is provided in Tables 1 and 2 respectively, and is accompanied by a
22 general rationale for their inclusion in our system.

23
24
25

1 **4. Creating an operational system through expert elicitation**

2 Designing a hypothetical risk system that draws on the distinction between direct and
3 indirect drivers is useful for it allows us to begin thinking systematically about what types
4 of evidence would be needed to operationalise the system in practice, and how
5 disciplinary responsibilities might begin to be assigned in such an undertaking. Perhaps
6 most significantly it provides a framework in which the explanatory potential of social
7 and natural sciences are understood to be co-dependent. By definition, the variables
8 which comprise any given system are open to critical scrutiny, but we suggest that, by
9 adopting a multidimensional, experience-based, approach to the task of system building,
10 a rich and plausible picture of factors driving FIO risk begins to emerge.

11 However, the process of constructing system parameters is not an end in itself, for it
12 tells us nothing about level of significance we might wish to attach to one particular
13 variable over another within the system. As such, if the first task for developing a FIO
14 risk tool is to establish a set of parameters within which microbial risks can be situated,
15 the second is to establish which elements are ‘more or less’ important to the system. To
16 do this we have followed a body of experimental methodological work that has sought to
17 develop heuristic assessments of problems through the use of expert elicitation
18 techniques, not least in the context of environmental science (e.g. Geneletti, 2005;
19 Nguyen and de Kok, 2007; Ye et al., 2008).

20 Expert elicitation is a means by which we can refine understandings of research
21 problems by exposing them to wider critical scrutiny. Like the issue under consideration
22 here, they are typically deployed in problem contexts where theoretical frameworks are
23 novel and empirical understanding is incomplete. Some applications of the technique,
24 such as through the use of the Delphi method or one of its many variants, are
25 characterised by a desire to reach consensus, building iteration, interaction and feedback

1 into its approach (Murry & Hammons, 1995; Chu and Hwang, 2008). Consequently, they
2 are designed to enable expert judgments to forecast and establish priorities in arenas of
3 science policy that would otherwise remain intractable. That is to say, they allow policy
4 programmes to be devised on a scientific basis in the absence of conventional forms of
5 scientific evidence. The argument here is that elicitation techniques allow policy makers
6 to craft judicious responses to emerging problems, not only because the process may
7 reveal strong areas of (otherwise unrecognised) agreement but also because they help
8 characterize and clarify regions of uncertainty (Rizak and Hruday, 2005). Yet, as von
9 Kraus et al. (2004 p.1515) note, “in situations where uncertainty is high, different experts
10 may have different interpretations of the data available, and the diversity of opinions may
11 be far larger than that which is suggested by consensus documents, such as risk
12 assessment reports”. Not surprisingly one of the issues facing those who use these
13 techniques is how to inform reasoned responses to policy problems without also
14 propagating the idea that scientific debates are somehow settled and resolved. As
15 Hoffman et al. (2008, p.697) have put it in the context of pathogenic risks: “Expert
16 elicitation ultimately is a complement, not a substitute, for primary research...[]... But it
17 can provide a practical aid to the better management of safety in complex systems where
18 perfect information is rarely available but better information might be”.

19 *4.1 Approach to elicitation*

20 In practical terms we initiated a process of expert consultation in which individuals with
21 different disciplinary expertise were asked to pass judgment on the relative importance of
22 possible controls on FIOs. The elicitation process followed Booker and McNamara
23 (2004, p.221) in understanding expert knowledge as that encompassing “what qualified
24 individuals know with respect to their technical practices, training and experience” and
25 made a distinction between those working in a natural science setting as having the expert

1 knowledge to gauge the relative importance of the direct bio-physical and management
2 variables, and those working in the context of social-science as able to pass judgment on
3 indirect cultural processes and structures.

4 Like the distinction between direct and indirect drivers made above, the purpose of
5 disentangling the consultation process into perspectives from the social and natural
6 sciences is designed to make an analytical and practical, rather than empirical and
7 substantive, distinction in how we can conceptualise and understand FIO risk. The
8 process of selecting participants in the elicitation process was designed to ensure that a
9 range of cognate science and social science areas of research would be represented. For
10 those working in a natural science context we solicited the views of those with expertise
11 and standing in academic microbiology, soil and environmental science, as well as
12 members of the policy community with professional investments and backgrounds in
13 microbial/waste management processes. For those working in a social science context we
14 solicited the views of those working in rural geography, agricultural studies, agricultural
15 economics, rural sociology and political science. Whilst those with natural science
16 expertise were required to have experience within the sphere of FIO research in order to
17 qualify for inclusion within the consortium (c.f. Cornelissen et al., 2003), this was not the
18 case in the social-science consultation, since as we have made clear above, no substantive
19 research areas exist to define an expert group in this way. Even so, all of the social
20 scientists included in the process were identified on the basis that their background
21 intersected with interests in rural environmental change, of which watercourse pollution
22 was a substantive, or at least, related concern.

23 At least one of the implications of this process is that different explanatory logics and
24 types of analytical and definitional precision have to be created for members of expert
25 consortiums. In the natural science consultation it was recognised, for instance, that

1 expert responses could be potentially variable depending on whether views were elicited
2 for *pathogenic* or *FIO* risks. In a technical sense, FIOs are one strand of a larger
3 pathogenic issue, a crucial difference to a respondent with precise expertise in this
4 research area (such as microbiology)⁵. In the social science context, this definitional
5 problem is inverted. We reasoned that FIOs would mean very little to those with training
6 in social science and care was taken to explain ‘FIOs’ as a framework for examining
7 potential *pathogenic* risks to watercourses.

8 The process was conducted through a standardised electronic scoring schedule, in
9 which experts were instructed to weight the relative influence of a given risk factor on a
10 continuous scale ranging from 0 (of no importance) through to 1 (of absolute
11 importance). If respondents felt unable to weight factors with confidence they were
12 instructed to leave potential entries blank. A qualitative dimension was also included in
13 that respondents were allowed to comment and elaborate on their choices should they
14 wish to do so. In total 28 experts participated in the elicitation process, representing a
15 response rate of approximately 80%.

16 It is of note at this juncture to recognise that this methodology is by no means the
17 only way in which the task of eliciting judgment can be approached, and a case has been
18 made for developing more interpretative and discursive approaches to risk and
19 uncertainty characterisation (EA, 2008). The approach we adopted in this study was
20 designed as a first approximation of expert knowledge, and was geared explicitly to the
21 producing responses that could be operationalised easily within a prototypical risk
22 assessment tool (Oliver et al., 2009). A deeper contextualisation of expert assessments of

⁵ One way of illustrating this point is with regard to our risk factor ‘animal type’. In an ‘FIO’ consultation, animal type might not be considered by experts a strong driver of risk, relative to other factors, because all livestock faeces contain considerable FIO content. But this is not necessarily the case should we have devised the consultation around ‘pathogens’ *per se*. Here animal type may be considered more important (for instance lambs are also large cryptosporidium reservoirs). The definitional formula is not merely a technicality: it could significantly influence measures arising from risk assessment.

1 risk through, for instance, depth and interviews would be a logical development on the
2 insight developments here (see for instance, Philips, 1999)

3 *4.2 Gauging the relative importance of system variables*

4 A general summary of statistics associated with the expert weighting exercise is shown in
5 Tables 3 and 4. As an initial layer of analysis, the project team calculated the arithmetic
6 mean of expert weightings for each variable as a way of assessing their relative
7 importance to the system as a whole. Assessing relative significance in this rudimentary
8 way begins the process of understanding where priorities for FIO risk assessment might
9 lie. Thus, following this approach the expert elicitation process suggests that ‘livestock
10 access to streams’ would be considered a key driver of risk on the basis of the natural
11 science consultation, for it achieved a mean expert weighting of 0.82 among experts, the
12 highest across all risk categories, and accommodates a strong negative skew in the
13 distribution of weightings received. Other salient factors included the transfer risk of
14 ‘runoff potential’ (0.73), the infrastructural risk of ‘drainage’ (0.72), and the source risks
15 of ‘stocking density’ and (0.72) ‘manure type’ (0.66). In the social science consultation,
16 key determinants of risk included whether a farmer had access to ‘technical grants for
17 waste management’ (0.83), the extent to which farming activities were currently
18 structured through ‘participation in agri-environmental schemes’ (0.69), and whether
19 practices were embedded in ‘organic farming’ regimes (0.68). With a mean weighting of
20 0.58 the ‘size of the holding’ also constituted a higher ranking determinant of risk in the
21 social science assessment.

22 By implication the consultation process also pointed to areas of our risk system where
23 the impact of variables was considered relatively weak. So, for instance, while the results
24 of the natural science consultation suggest that importance should be attached to
25 ‘livestock access to streams’, another source risk ‘livestock type’ achieved a mean

1 weighting of only 0.35, the lowest across *all* natural science risk categories. Likewise, the
2 importance experts attached to the transfer risk of ‘run-off potential’ can be contrasted
3 with issues of ‘erosion’ and ‘preferential flow’ potential, which returned mean weightings
4 of 0.38 and 0.56, respectively. In the social science consultation, the significance experts
5 assigned to ‘size of holding’ can be neatly contrasted with ‘tenure of holdings’, which
6 returned a mean expert weighting of only 0.30. Moreover, the social science consultation
7 placed little importance on lifecycle issues such as ‘succession’ and recreational factors
8 such as the ‘presence of public rights of ways on or bordering land’. Both these factors
9 achieved mean weightings of only 0.26. Of course, the range of expert weightings
10 received for each risk factor may be used in the future for uncertainty assessment
11 (Refsgaard et al., 2007) and fuzzy modelling approaches but that is beyond the scope of
12 this present paper.

13

14 *4.3 Areas of expert consensus and disagreement*

15 On the basis of this analysis our system begins to take shape as a framework in which
16 distinctions about FIO risk can begin to be drawn. However, making judgments in this
17 way tells us nothing about the extent to which experts agreed or disagreed about the
18 relative importance of risk factors. Risk factors may be ranked high or low, but variables
19 can have differing levels of consensus attached to them. This is an important underlying
20 issue, for as we have shown, in discriminating between the relative influence of given
21 factors, a risk assessment framework would also ideally aspire to focus on system
22 variables that have a high degree of consensus attached to them across disciplinary
23 boundaries.

24 The dimensions of this latter issue are expressed as box and whisker plots in
25 Figures 1 and 2. For the natural science consultation there were a number of high and low

1 ranking variables where experts displayed a high degree of consensus (i.e. a narrow
2 interquartile range). So, for instance, the highly ranking source factor of ‘stocking
3 density’ was accompanied by the lowest interquartile range and standard deviation
4 (σ) among experts (see Table 3), suggesting that collective confidence in the influence of
5 this variable was strong. Equally, there was close agreement among experts regarding
6 some factors which were otherwise considered to be weak in their influence. The
7 connectivity risk factor of ‘gateway location’, which returned a mean expert weighting of
8 0.39, would be a case in point here ($\sigma=0.21$). In the social science consultation there
9 were also generally strong levels of agreement surrounding high ranking factors such as
10 ‘technical grants’ ($\sigma=0.19$), but perhaps most notably, surrounding the influence of
11 responsibilities under ‘cross-compliance’⁶ ($\sigma=0.17$), a factor attaining a moderate expert
12 mean weighting of 0.49.

13 Yet these findings have to be squared with variables where expert opinion was largely
14 divided. In the natural science consultation there was considerable variability over the
15 importance we should assign to infrastructural factors. For example, while the mean
16 weighting for ‘farm tracks’ was ranked moderately at 0.40, some experts suggested its
17 weighting should be as high as 0.80, while others considered it of no importance at all in
18 driving risk, resulting in a standard deviation of 0.25 for this risk factor. How do we
19 explain such discrepancies? At one level it may be argued that contrasting visions reflect
20 an uncertain evidence base. For example, Edwards et al. (2008) note that a general
21 awareness exist regarding farmyards as sources and contributors of pollutants to
22 receiving waters but stress that few measurements have actually been made to quantify
23 contaminant flux from these areas. However, even where empirical research exists, wide
24 interquartile ranges can be returned. An interesting example of this was with regard to the

⁶ In the UK cross-compliance embodies a set of mandatory conditions regarding good agricultural conduct and underpins state support for agriculture in the form of a ‘single farm payment’.

1 source risk factor of ‘aged faecal material’. Consensus here was weak ($\sigma = 0.28$) despite
2 empirical research from the 1980s reporting on patterns of release of FIOs from ageing
3 faeces (Thelin and Gifford, 1983; Kress and Gifford, 1984).

4 Such discrepancies may well demonstrate the need to further consolidate
5 understanding of some drivers of microbial watercourse contamination not only through
6 future research agendas (Kay et al., 2007; Kay et al., 2008a) but in their effective
7 dissemination across different knowledge domains, but we suggest there may be a deeper
8 problematic at stake. At a more theoretical level this divergence may reflect tensions that
9 inherently exist between making ‘whole system’ assessments of risk and the disciplinary
10 nature of expertise partly informing the principle of expert judgements. In our risk
11 assessment, respondents were asked to comment on the *relative* importance of controls
12 across the risk system, but the basis of participation in the exercise was dictated, in part,
13 by their status as authority figures with ‘specialist’ types of expertise: that is, by their
14 knowledge of *particular* components of the risk system. And as the literature on expert
15 elicitation argues in this vein, when individuals respond as experts in this way they tend
16 to be overconfident in recognizing the influence of those factors that reflect their
17 professional remits and worldviews. Or as Granger Morgan et al. (2001, p.282) put it
18 “given their knowledge, ...[the]... subjective probability distributions ...[of experts] ...
19 tend to be too narrow”. In other words, precise notions of expertise tend to shape the
20 emphasis placed on particular aspects of risk assessment, leading to divergent accounts of
21 how a system works.

22 A telling example of this is in relation to two elements of the natural science
23 assessment: ‘preferential flow potential’ and ‘animal type’. Whereas respondents with a
24 background in microbiology weighted the former issue of lowest importance in the risk
25 system, those with soil science interest ranked this highest. Conversely, whereas a

1 number of microbiologists ranked ‘animal type’ as an issue of key importance, many
2 experts in soil science assigned it low significance. How far should we be surprised by
3 these divergent opinions? After all, preferential flow potential is a core concept of soil
4 science and will therefore be central to the professional remit those working in this area
5 of specialism. But this concern is outside of the formal scope of microbiology, where
6 animal-led explanations of risk are likely to prevail. The data provide further examples of
7 this issue. Responses to the risk assessment reveal that those natural science respondents
8 working directly in the arena of FIO policy intervention and development consistently
9 provide higher risk ratings to the importance of individual elements in the risk system
10 than academic scientists, perhaps reflecting how professional investments and awareness
11 in the given risk issue can tend to magnify *overall* understandings of risk. For example,
12 grouping the natural science expert panel into non-academic policy experts,
13 microbiologists, soil scientists and environmental scientists revealed that the policy-based
14 experts ranked 17 of the 22 risk factors with higher mean weightings, and on one of these
15 occasions (for the risk factor ‘farmyard drainage to watercourse’) the difference between
16 expert groups, derived using an analysis of variance test, was significant ($P < 0.05$). The
17 policy expert group weighted this risk factor as 0.98 (clearly illustrating near universal
18 agreement of the absolute importance of this risk factor), whereas the microbiologists,
19 soil scientists and environmental scientists groupings returned average weightings of
20 0.64, 0.60 and 0.65, respectively.

21 In other words this initial analysis of results suggests that disciplinary and
22 professional investments produce interesting distinctions in how FIO risks come to be
23 conceptualised, though in making this claim we feel it necessary to exercise some caution
24 here. Drawing conclusions in this way presumes that affiliations and expertise exist as a
25 series of tightly bounded, and mutually-exclusive, entities, when in practice, this may not

1 be the case. Disciplinary boundaries and identities are often quite fuzzy both in terms of
2 professional practices and research preoccupations; while the world of academic research
3 and policy driven science are often fundamentally entwined in terms of career pathways
4 and research priorities. Nonetheless, we suggest that these findings point to one way in
5 which divergent assessments of FIO risk might begin to be interpreted from an
6 interdisciplinary perspective.

7 Notwithstanding the need to understand why these divergent responses arise, one of
8 the key issues that such heterogeneity poses is how to arrive at overall assessments of risk
9 that would allow a future risk tool to be operationalised in practice. The use of mean
10 expert responses was our response to this approach, though it does not resolve the
11 underlying ontological problem here: how to create a common, as in collectively agreed
12 upon, vision of the risk system?

13 *4.4 The indeterminacy of risk*

14 To the theoretical conundrum of establishing a common vision we wish to add a further
15 and final dimension. This concerns the issue of indeterminacy. One of the tacit
16 assumptions underlying our analysis is that a given control on FIO risk has a consistent
17 relationship with how a risk is influenced. That is to say, experts may or may not agree
18 that one variable is more important than another in the system, but the assumption is that
19 the *direction* of risk is clear. In many cases this assumption is fair. Certainly it is true of
20 our direct risk factors. So for example it is clearly logical to suggest that the steeper a hill
21 slope the greater the potential for run-off, and therefore risk of FIO loss from land, just as
22 we can claim, with confidence, that the export of FIOs from land to water would be less
23 likely under extensive grazing regimes. However, in the case of our indirect drivers, this
24 principle does not necessarily follow. While it may be reasonable to infer that
25 ‘participation in an agri-environmental scheme’ or ‘receipt of technical grants’ would be

1 linked with a potential reduction in FIO risk, it seems much more problematic to make
2 such connections with a number of other indirect variables. So, for example, the variable
3 ‘length of time in farming’ is interesting in a social science context for it implies
4 capacities and inclinations that may influence risk in a variety of ways. Recent entrants to
5 farming may bring with them new models of working that are conducive to working
6 within the mandates of modern environmental protection, but so too may they lack the
7 very skill-sets that would come from longstanding involvement in a particular system of
8 livestock farming. In other words, the system variable ‘length of time in farming’
9 embodies in it a range of influences conducive to different types of outcome. This
10 indeterminacy seems to be true of ‘farm succession’, ‘size’ and ‘tenure’ of land farmed
11 and influence of ‘debt’. Contrasting narratives of FIO risk could be assigned to these
12 endemic characteristics and circumstance of a farm system.

13 Anticipating this response in the expert consultation, we examined this issue directly
14 with social scientists. Alongside the process of assigning *importance* to system elements,
15 we asked these experts to make judgments about how the *direction* of risk might unfold.
16 To do this, experts were presented with a series of propositions about the potential
17 influence of all indirect variables on FIO risk. Thus in the case of farm size, experts were
18 asked to consider the proposition ‘*The transfer of FIOs to water courses is less likely the*
19 *smaller the area farmed*’. Experts could then offer different responses to these
20 propositions to indicate the extent to which they agreed with the proposition or could
21 offer a meaningful response to it. A summary of these propositions and the framework
22 for expert responses is depicted in Table 5. The box and whisker plots, revealing extents
23 of agreement between respondents to this aspect of the consultation, are displayed in
24 Figure 4.

1 Not surprisingly the consultation identified some areas of social science agreement
2 where the directional influence of variables is clear. Collectively, the response of the
3 consultation suggests that ‘receipt of technical grants’ would be likely to lower FIO risks,
4 as would ‘responsibilities under cross compliance’, ‘participation in agri-environmental
5 schemes’ and ‘organic farming’, as well as ‘formal accreditation in land and livestock
6 management’. Yet as anticipated, in other instances, the consultation raised profound
7 disagreements over the directional influence of variables. Issues of ‘tenure’, ‘succession’
8 and most notably ‘farm size’ stand out here. While size of holding is a relatively high
9 ranking factor, there was little agreement about the direction of risk we might attach to a
10 particular operational farm scale. Many refused to draw neat conclusions. One suggested
11 that “those with a limited area may have more of a waste disposal problem”, but pointed
12 to alternative circumstances where size might imply farmers were “less intensive
13 smallholders” with lower risks.

14 The consultation suggests therefore, that there is no consistent way of assigning risks
15 to particular variants of some risk conditioning issues. We suggested in the section above
16 that a fundamental problem for those wishing to create holistic conceptions of risk is how
17 to ‘overcome’ differences in expert knowledge regarding how the system ‘works’. Here
18 the issue seems to be rather different. Such indeterminacies are problems produced by the
19 generic needs of risk assessment, not differences in expert opinion *per se*. It is tempting,
20 for instance, to ignore issues such as ‘size’, ‘tenure’ and ‘succession’ because they cannot
21 be standardized across a range of circumstances. But the message of this aspect of our
22 consultation is that risk assessment frameworks should not only be generic in design but
23 sensitive to context. In other words, size may well be an important generic variable, but
24 we can only draw conclusions about its *determinate* influence when studying precise sets
25 of circumstances.

1 *4.5 Quantifying potential errors associated with expert weightings*

2 The expert consultation exercise outlined in this study was designed to elicit indicative
3 responses from experts in relevant disciplines that could help make an on-farm risk tool
4 operational. Associated weightings derived through this process were eventually
5 embedded into a cross-disciplinary toolkit to assess the risk of faecal indicator loss from
6 grassland farm systems to surface waters (Oliver et al., 2009). While the statistical
7 analysis of the data presented in this study centres on traditional differences in
8 distributions of expert weightings for given risk factors there is a realisation that there is
9 likely to be a degree of interdependency between weighted risk factors when incorporated
10 into integrated risk management tools. It is when expert derived data is incorporated into
11 decision support tools (DST) that we can begin to undertake a more in-depth analysis of
12 variations in expert judgement. Whilst beyond the remit of this current study, it would be
13 wise to undertake a full sensitivity analysis of any DST that accommodates expert
14 judgement to determine the responsiveness of DST output to changes in individual risk
15 factor weightings. For example, if DSTs inform end-users of an index of relative risk of a
16 parcel of land contributing a pollutant to a watercourse then what would be the effect on
17 the final risk index of a change in any single risk factor weighting of, for example, 0.01
18 or 0.1? Furthermore, which risk factors are most sensitive to these changes?

19 Reporting on the range of original expert weightings embedded within management
20 tools enables researchers to assess the error placed on calculated risk indices when the
21 standard deviations (or confidence intervals) of the mean for each risk factor weighting
22 are taken into account. Given that different expert weightings (such as those derived in
23 this study) carry different ranges of opinion we can begin to assess which risk factors
24 contribute most to the errors intrinsic to expert weighted risk indexing approaches, thus
25 allowing a decision to be made as to whether a risk factor might be justifiably omitted.

1 Perhaps more importantly, the ranges of difference in expert weightings can highlight
2 risk factors to target for fundamental research to better align expert opinion.

3

4 **5. Concluding remarks**

5 In a paper speculating on future developments in the field of microbiology Curtis (2007)
6 argued recently for the creation of numerically reliable, and widely applicable,
7 explanatory frameworks for microbiological research; ones linked expressly to the needs
8 of wider political and social agendas and involving the development of a “consilient and
9 calibrated set of rules to describe and predict the behaviour of the microbial world as a
10 system” (*Ibid* p.1). In this paper we have sought to examine some of the complexities that
11 arise from seeking to foster such systemic and applied understandings of the microbial
12 world. If we were to appropriate Curtis’ vision for our particular purposes and take it to
13 its logical conclusion, then this would be a process in which researchers would be able to
14 calibrate real world livestock systems according to the intrinsic orderliness of potential
15 pathogen transfer within agricultural environments. In so doing, it may be possible to
16 start to make highly reliable judgments about the conditions under which microbial risks
17 are present and thereby controlled.

18 The paper has described an experimental process in which such a system might begin
19 to be imagined. Like many arenas of environmental sustainability there are few empirical
20 reference points in which to begin navigating a stable pathway through the essentially
21 complex and unbounded system of FIO risk. In this our research suggests that an
22 interdisciplinary approach can greatly enhance the plausibility of a hypothetical system
23 and can underpin the development of a new generation of study areas that are linking
24 ostensibly disparate research areas together. The idea of ‘diffusion pollution studies’
25 (Lane et al. 2006), of which an account of FIO risk would ultimately be a part, is a case in

1 point. Yet, so too has our research served to underline the substantive difficulties, both
2 theoretical and methodological, in arriving at a shared picture of the world. The idea of
3 building rule-governed, 'whole system', conceptions of environmental processes is itself
4 a distinct, and by no means unproblematic, theoretical aspiration. Assuming that we can
5 foster interdisciplinary teams to collaborate in this endeavour, the means by which we
6 then arrive at internally stable conceptions of risk is by no means accepted or understood.

7 As we have shown, one of the paradoxes of building interdisciplinary pictures is that
8 they necessarily emerge out of precise areas of scholarship, and this will mean that
9 conceptual and analytical differences over how a system works will tend to be magnified
10 as much as they will be diminished. One of the challenges facing those creating holistic
11 approaches to environmental risk assessment is therefore how to develop methodologies
12 by which such discrepancies can be overcome. In other senses, taking a whole system
13 approach begins to reveal complexities that question the way we structure our
14 representations of problems, and with this, the nature of our interventions. It may well be
15 that some drivers of risk are indeterminate, which fundamentally alters the needs of risk
16 assessment.

17 With these problems and challenges, we suggest, comes responsibility to design
18 procedures that can acknowledge, rather than bury, endemic system uncertainties. This
19 seems axiomatic given that theoretical representations of environmental problems reside
20 as much in a world of policy 'readiness' as they do in the long term agendas of scientific
21 research. Faced with the need to construct reasoned responses to emerging policy
22 questions, such as the management and mitigation of FIOs, we need to be circumspect
23 when translating highly tentative and contested constructions of environmental problems
24 into expert systems with 'ready-made' answers.

25

1 **Acknowledgements**

2 The authors would like to acknowledge the participation of the 28 anonymous experts for
3 their valuable contributions in the risk factor weighting exercise. This work was funded
4 as part of project RES-224-25-0086 by the Rural Economy and Land Use (RELU)
5 programme who receive financial support from the Biotechnology and Biological
6 Sciences Research Council (BBSRC), the Natural Environment Research Council
7 (NERC), the Economic and Social Research Council (ESRC), the Department for
8 Environment, Food and Rural Affairs (Defra) and the Scottish Office.

9

10 **References**

11 Abu-Ashour, J. and Lee, H. 2000. Transport of bacteria on sloping soil surfaces by
12 runoff. *Environmental Toxicology* 15, 149-153.

13

14 Aitken, M. N. 2003. Impact of agricultural practices and river catchment characteristics
15 on river and bathing water quality. *Water Science and Technology* 48, 217-224.

16

17 Artz, R. R. E., Townend, J., Brown, K., Towers, W. and Killham, K. 2005. Soil
18 macropores and compaction control the leaching potential of *Escherichia coli* O157:H7.
19 *Environmental Microbiology* 7, 241-248.

20

21 Booker, J.M., McNamara, L.A. 2004. Solving black box computation problems using
22 expert knowledge theory and methods. *Reliability Engineering and System Safety* 85,
23 331–340.

24

1 Burton, R.J.F and Wilson, G.A. (2006) Injecting social psychology theory into
2 conceptualisations of agricultural agency: Towards a post-productivist farmer self-
3 identity? . *Journal of Rural Studies*, 22, 95-
4
5 Buller, H., Morris, C., 2004. Growing goods: the market, the state and sustainable food
6 production. *Environment and Planning A* 36, 1065-1084.
7
8 Chadwick, D., Fish, R., Oliver, D. M, Heathwaite, L., Hodgson, C., Winter, M., 2008.
9 Management of livestock and their manure to reduce the risk of microbial transfers to
10 water – the case for an interdisciplinary approach. *Trends in Food Science and*
11 *Technology* 19, 240-247.
12
13 Chu, H.C., Hwang, G.J. 2008. A Delphi-based approach to developing expert systems
14 with the cooperation of multiple experts. *Expert Systems with Applications* 34, 2826-
15 2840.
16
17 Collins, R., Elliott, S. and Adams, R. 2005. Overland flow delivery of faecal bacteria to a
18 headwater pastoral stream. *Journal of Applied Microbiology* 99, 126-132.
19
20 Cornelissen, A.M.G., van den Berg J., Koops W.J., Kaymak, U., 2003. Elicitation of
21 expert knowledge for fuzzy evaluation of agricultural production systems. *Agriculture,*
22 *Ecosystems and Environment* 95, 1–18.
23
24 Crowther, J. Wyer, M. D., Bradford, M., Kay, D., Francis, C., A., and Knisel, W. G.
25 (2003). Modelling faecal indicator concentrations in large rural catchments using land use
26 and topographic data. *Journal of Applied Microbiology*. 94, 962-973.

1 Davies-Colley, R. J., Nagels, J. W., Smith, R. A., Young, R. G. and Phillips, C. J. 2004.
2 Water quality impact of a dairy cow herd crossing a stream. *New Zealand Journal of*
3 *Marine and Freshwater Research* 38, 569-576.

4

5 Defra 2003. Manure management plan: a step by step guide for farmers. Viewed on 18th
6 July 2008. Available online at:
7 www.defra.gov.uk/corporate/regulat/forms/agri_env/nvz/manureplan.pdf

8

9 Defra, 2005a. Producing a soil management plan for environmental stewardship. Viewed
10 on 18th July 2008. Available online at: [http://www.defra.gov.uk/erdp/pdfs/es/guidance/es-](http://www.defra.gov.uk/erdp/pdfs/es/guidance/es-soil-management-plan.pdf)
11 [soil-management-plan.pdf](http://www.defra.gov.uk/erdp/pdfs/es/guidance/es-soil-management-plan.pdf)

12

13 Defra, 2005b. Entry Level Stewardship Handbook. Viewed on 18th July 2008. Available
14 online at: <http://www.defra.gov.uk/erdp/pdfs/es/els-handbook.pdf>.

15

16 EA, 2008. Uncertainty Assessment of Phosphorus Risk to Surface Waters. Science
17 Report – SC050035.

18

19 Edwards, A. C., Kay, D., McDonald, A. T., Francis, C., Watkins, J., Wilkinson, J. R.,
20 Wyer, M. D., 2008. Farmyards, an overlooked source for highly contaminated runoff.
21 *Journal of Environmental. Management* 87, 51-59.

22

23 Fish, R. Seymour. S, and Watkins, C. (2003) Conserving English Landscapes: Land
24 managers and agri-environmental policy *Environment and Planning A*, 35, pp. 19-41

25

1 Geneletti, D., 2005. Formalising expert opinion through multi-attribute value functions:
2 An application in landscape ecology. *Journal of Environmental Management* 76, 255–
3 262.

4

5 Georgiou, S., Bateman., I., 2005. Revision of the EU Bathing Waters Directive:
6 economic costs and benefits. *Marine Pollution Bulletin* 50, 430-3438.

7

8 Giddens, A. (1984) *The Constitution of Society. Outline of the Theory of Structuration.*
9 Cambridge: Polity Press

10

11 Granger Morgan, M., Pitelka, L.F. , Shevliakova, E., 2001. Elicitation of expert
12 judgments of climate change impacts on forest ecosystems. *Climatic Change* 49: 279-307.

13

14 Graczyk, T. K., Sunderland, D., Tamang, L., Lucy, F. E. and Breysse, P. N., 2007. Bather
15 density and levels of *Cryptosporidium*, *Giardia*, and pathogenic microsporidian spores in
16 recreational bathing water. *Parasitology Research* 101, 1729-1731.

17

18 Hoffmann, S., Fischbeck, P., Krupnicka, A., McWilliams, M., 2008. Informing risk-
19 mitigation priorities using uncertainty measures derived from heterogeneous expert
20 panels: A demonstration using foodborne pathogens. *Reliability Engineering and System*
21 *Safety* 93, 687–698.

22

23 Holdren J. P., 2008. Science and technology for sustainable well-being. *Science* 319,
24 424-434.

25

1 Horlick-Jones. T., Sime J., 2004. Living on the border: knowledge, risk and
2 transdisciplinarity. *Futures* 36, 441–456.
3

4 Hutchison, M. L., Walters, L. D., Moore, A., Crookes, K. M., and Avery, S. M. 2004.
5 Effect of length of time before incorporation on survival of pathogenic bacteria present in
6 livestock wastes applied to agricultural soil. *Applied and Environmental Microbiology*
7 70, 5111-5118.
8

9 Kaika, M., 2003. The Water Framework Directive: A New Directive for a Changing
10 Social, Political and Economic European Framework” *European Planning Studies* 113,
11 299-316.
12

13 Kay, D., Crowther, J., Fewtrell, L., Francis, C.A., Hopkins, M., Kay, C., McDonald,
14 A.T., Stapleton, C.M., Watkins, J. Wilkinson, J., Wyer, M.D., 2008a. Quantification and
15 control of microbial pollution from agriculture: a new policy challenge? *Environmental*
16 *Science and Policy*. 11, 171-184.
17

18 Kay, D., Crowther J., Stapleton, C. M., Wyer, M. D., Fewtrell, L., Anthony, S., Bradford,
19 M., Edwards, A., Francis, C. A., Hopkins, M., Kay, C., McDonald, A. T., Watkins, J. and
20 Wilkinson, J., 2008b. Faecal indicator organism concentrations and catchment export
21 coefficients in the UK. *Water Res.* 42, 2649-2661.
22

23 Kay, D., Edwards, A. C., Ferrier, R. C., Francis, C., Kay, C., Rushby, L., Watkins, J.,
24 McDonald, A. T., Wyer, M., Crowther, J. , Wilkinson, J., 2007. Catchment microbial

1 dynamics: the emergence of a research agenda. Progress in Physical Geography. 31, 59-
2 76.

3

4 Kress, M. and Gifford, G. F.,1984. Fecal coliform release from cattle fecal deposits.
5 Water Resources Bulletin, 20, 61-66.

6

7 Lane, S.N., Brookes, C.J., Heathwaite, A.L. & Reaney, S.M., 2006. Surveillant Science:
8 Challenges for the Management of Rural Environments Emerging from the New
9 Generation Diffuse Pollution Models. Journal of Agricultural Economics. 57, 239-257.

10 Lemunyon, J. L., Gilbert, R. G., 1993. The concept and need for a phosphorus assessment
11 tool. J. Prod. Agric. 6, 483-486.

12

13 Lobley, M and Potter, C A (2004) Agricultural change and restructuring: recent evidence
14 from a survey of agricultural households in England. Journal of Rural Studies. 20, 499-
15 510

16

17 Muirhead, R. W., Collins, R. P., and Bremer, P. J. 2006. Interaction of *Escherichia coli*
18 and soil particles in runoff. Applied and Environmental Microbiology 72, 3406-3411.

19

20 Murry, J.W., Hammons, J.O., 1995. Delphi, a versatile methodology for conducting
21 qualitative research. Review of Higher Education 18, 423-436.

22

23 Nguyen, T.G., de Kok, J.L., 2007. Systematic testing of an integrated systems model for
24 coastal zone management using sensitivity and uncertainty analyses Environmental
25 Modelling & Software 22, 1572-1587.

1 Oliver, D. M., Heathwaite, A. L., Haygarth, P. M. and Clegg, C. D. 2005. Transfer of
2 *Escherichia coli* to water from drained and undrained grassland after grazing. Journal of
3 Environmental Quality, 34, 918-925.

4

5 Oliver, D. M., Haygarth, P. M., Clegg, C. D. and Heathwaite, A. L. 2006. Differential *E.*
6 *coli* die-off patterns associated with agricultural matrices. Environmental Science and
7 Technology, 40, 5710-5716.

8

9 Oliver, D. M., Fish, R. D., Hodgson, C. J., Heathwaite, A. L. Chadwick, D. R. and
10 Winter, M., 2009. A cross-disciplinary toolkit to assess the risk of faecal indicator loss
11 from grassland farm systems to surface waters. Agriculture, Ecosystems and
12 Environment, 129, 401-412.

13

14 Philips, L. D. 1999. Group elicitation of probability distributions: are many heads
15 better than one? Bayesian research conference; decision science and technology,
16 reflections on the contributions of ward edwards, Klumer Academic, Boston and London,
17 313-330.

18

19 Refsgaard, J. C., van der Sluijs, J. P., Hoberg, A. L., Vanrolleghem, P. A. 2007.
20 Uncertainty in the environmental modelling process – a framework and guidance.
21 Environmental Modelling and Software 22, 1543-1556.

22

23 Rizak, S., Hrudey, S.E., 2005. Interdisciplinary comparison of expert risk beliefs J.
24 Environmental Engineering Science 4, 173–185.

25

1 SEPA 2004. The 4 point plan for improved farm waste management. 2nd ed., Scottish
2 Environment Protection Agency, Scotland.

3

4 Schets, F. M., van den Berg, H. H. J. L., Engels, G. B., Lodder, W. J. and Husman, A. M.
5 D. R., 2007. *Cryptosporidium* and *Giardia* in commercial and non-commercial oysters
6 (*Crassostrea gigas*) and water from the Oosterschelde, the Netherlands. International
7 Journal of Food Microbiology. 113, 189-194.

8

9 Smith, K. A., Brewer, A. J., Dauven, A., and Wilson, D. W. (2001). A survey of the
10 production and use of animal manures in England and Wales. III. Cattle manures. Soil
11 Use Management. 17, 77-87.

12

13 Thelin, R., Gifford, G. F., 1983. Faecal coliform release patterns from faecal material of
14 cattle. Journal of Environmental Quality 12, 57-63.

15

16 Krayer von Krauss, M.P, Casman, E.A., Small, M.J. 2004. Elicitation of expert
17 judgments of uncertainty in the risk assessment of herbicide-tolerant oilseed crops. Risk
18 Analysis 24, 1515-1527.

19

20 Whatmore, S., 2002. Hybrid Geographies: Natures Cultures Spaces. Sage, London.

21 Wilson, G.A., 2007. Multifunctional Agriculture: A Transition Theory Perspective. CAB
22 International, Wallingford.

23

- 1 White, S. L., Sheffield, R. E., Washburn, S. P., King, L. D., and Green, J. T. (2001).
2 Spatial and time distribution of dairy cattle excreta in an intensive pasture system. Journal
3 of Environmental Quality 30, 2180-2187.
4
- 5 Wilson, G.A. and Hart, K. 2000. Financial imperative or conservation concern? EU
6 farmers' motivations for participation in voluntary agri-environmental schemes.
7 Environment and Planning A 32, 2161-2185.
8
- 9 Wilson, G.A. 2007: Multifunctional agriculture: a transition theory perspective. CAB
10 International Wallingford .
11
- 12 Ye, M., Pohlmann, K.F. Chapman, J. B. 2008. Expert elicitation of recharge model
13 probabilities for the Death Valley regional flow system. Journal of Hydrology 354, 102–
14 115.