

From THE DEPARTMENT OF CLINICAL NEUROSCIENCE
Karolinska Institutet, Stockholm, Sweden

ASSESSMENT OF DOPAMINE AND SEROTONIN RELEASE IN THE NON-HUMAN PRIMATE BRAIN USING PET

Sjoerd J. Finnema



**Karolinska
Institutet**

Stockholm 2011

Cover illustration:

MRI-coregistered PET summation images obtained after intravenous injection of the dopamine D₂/D₃ receptors agonist radioligand [¹¹C]MNPA into a cynomolgus monkey during baseline and post-amphetamine (1.0 mg/kg) conditions.

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by University Service US-AB, Stockholm, Sweden

© Sjoerd J. Finnema, 2011

ISBN 978-91-7457-214-8

ABSTRACT

The molecular imaging technique positron emission tomography (PET) allows for non-invasive examination of biochemical markers in the living brain. For over three decades PET studies have provided important insight into the relationship of monoaminergic neurotransmitter systems to brain functioning and psychiatric disorders. A more recent application of PET is the study of endogenous neurotransmitter release *in vivo*. Clinical relevance of such methods is found in studies demonstrating enhanced amphetamine-induced dopamine release in schizophrenia patients, whereas PET studies in non-human primates provide a translational model for evaluation of the pharmacological mechanisms before initiation of studies in man.

The first aim of this thesis was to develop improved methods for measurement of endogenous dopamine levels. In study I the potent D₂/D₃ receptors agonist (*R*)-(-)-2-methoxy-*N-n*-propyl-norapomorphine (MNPA) was radiolabeled with carbon-11 and found suitable for *in vivo* characterization of the high affinity state. In study II, amphetamine-induced displacement of [¹¹C]MNPA binding by dopamine was ~1.8 fold higher at four different doses than for the antagonist [¹¹C]raclopride and demonstrated that an agonist radioligand has improved sensitivity to endogenous neurotransmitter level. Study III aimed to further obtain *in vivo* support for the existence of two affinity states for the D₂/D₃ receptors. Receptor occupancy of the exogenous agonist apomorphine was determined with [¹¹C]MNPA and [¹¹C]raclopride. Binding of [¹¹C]MNPA and [¹¹C]raclopride was inhibited monophasic and approached full saturation. ID₅₀ and K_i values of apomorphine were indistinguishable when measured with the agonist or antagonist radioligand. Study III did not support the existence of two affinity states and a possible explanation could be that all D₂/D₃ receptors are in the high affinity state *in vivo*. In study IV, the new D₁/D₅ receptors partial agonist radioligand (*S*)-[¹¹C]*N*-methyl-NNC 01-0259 was found insensitive to dopamine levels, and receptor binding was inferior to previously developed antagonist radioligands. Moreover, a COMT formed radiometabolite was found to enter the brain but the formation could be prevented with the use of a COMT inhibitor. COMT inhibition provides a methodology enabling quantitative PET measurements with (*S*)-[¹¹C]*N*-methyl-NNC 01-0259.

The second aim of this thesis was to evaluate the sensitivity of the new 5-HT_{1B} receptor radioligand [¹¹C]AZ10419369 to alterations in endogenous serotonin concentration. Previous serotonergic PET radioligands have ambiguously shown sensitivity to serotonin level. In study V the effective serotonin releaser fenfluramine decreased the binding of [¹¹C]AZ10419369 in a dose-dependent manner. In study VI the effect of fenfluramine on [¹¹C]AZ10419369 binding was confirmed using an equilibrium approach with a bolus infusion protocol. The further developed methodology is suitable for exploring the sensitivity limit to serotonin levels as measured using [¹¹C]AZ10419369 and PET.

In conclusion, the present thesis demonstrates that the D₂/D₃ receptors agonist radioligand [¹¹C]MNPA is an improvement for measurement of dopamine release, when compared to previously used antagonist radioligands. Moreover, a novel methodology, using the 5-HT_{1B} receptor antagonist [¹¹C]AZ10419369 and PET, was developed for measurement of serotonin release in the living brain. These newly developed methodologies may help to further understand the treatment and pathophysiology of several major neurological and psychiatric disorders.

“Zing, vecht, huil, bid, lach, werk en bewonder”

[Sing, fight, cry, pray, laugh, work and admire]

Ramses Shaffy, 1933-2009

LIST OF PUBLICATIONS

This thesis is based on six studies performed at the Department of Clinical Neuroscience, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden. The studies are presented in the following papers and will be referred to in the text by their roman numerals:

- I. **Finnema SJ**, Seneca N, Farde L, Shchukin E, Sóvágó J, Gulyás B, Wikström HV, Innis RB, Neumeyer JL and Halldin C. (2005) A preliminary PET evaluation of the new dopamine D₂ receptor agonist [¹¹C]MNPA in cynomolgus monkey. *Nuclear Medicine and Biology* 32:353-360.
- II. Seneca N, **Finnema SJ**, Farde L, Gulyás B, Wikström HV, Halldin C* and Innis RB*. (2006) Effect of amphetamine on dopamine D₂ receptor binding in nonhuman primate brain: a comparison of the agonist radioligand [¹¹C]MNPA and antagonist [¹¹C]raclopride. *Synapse* 59:260-269.
- III. **Finnema SJ**, Halldin C, Bang-Andersen B, Gulyás B, Bundgaard C, Wikström HV and Farde L. (2009) Dopamine D_{2/3} receptor occupancy of apomorphine in the nonhuman primate brain – a comparative PET study with [¹¹C]raclopride and [¹¹C]MNPA. *Synapse* 63:378-389.
- IV. **Finnema SJ**, Bang-Andersen B, Jørgensen M, Gulyás B, Wikström HV, Farde L and Halldin C. Inhibition of catechol-*O*-methyltransferase enhances visualization of (*S*)-[¹¹C]*N*-methyl-NNC 01-0259 binding to dopamine D₁-like receptors in monkey brain. Manuscript.
- V. **Finnema SJ**, Varrone A, Hwang TJ, Gulyás B, Pierson ME, Halldin C* and Farde L*. (2010) Fenfluramine-induced serotonin release decreases [¹¹C]AZ10419369 binding to 5-HT_{1B}-receptors in the primate brain. *Synapse* 64:573-577.
- VI. **Finnema SJ**, Varrone A, Hwang TJ, Halldin C* and Farde L*. Confirmation of fenfluramine effect on 5-HT_{1B} receptor binding of [¹¹C]AZ10419369 using an equilibrium approach. Manuscript.

*Authors contributed equally to this work.

CONTENTS

1	Introduction	1
1.1	Major applications of PET	1
1.2	Principles of PET	1
1.3	Development of PET radioligands for CNS	2
1.4	Quantitative PET measurements	3
1.4.1	Quantification of radioligand receptor binding	3
1.4.2	Quantification of neurotransmitter release	4
1.5	G-protein coupled receptors	5
1.5.1	Families of G-protein coupled receptors	5
1.5.2	G-protein coupled receptor signalling	5
1.5.3	The high affinity binding state	6
1.5.4	GPCR models	7
1.5.5	Receptor internalization and β -arrestin	7
1.6	The dopamine system	8
1.6.1	Dopaminergic neurotransmission	8
1.6.2	Dopaminergic pathways	9
1.6.3	Dopamine receptor subtypes	9
1.6.4	PET studies on dopamine release	10
1.6.5	PET imaging with dopamine agonist radioligands	11
1.7	The serotonin system	13
1.7.1	Serotonergic neurotransmission	14
1.7.2	Serotonergic pathways	14
1.7.3	Serotonin receptor subtypes	14
1.7.4	PET studies on serotonin release	15
2	Aims	18
3	Materials and Methods	19
3.1	Radiochemistry	19
3.2	In vitro receptor assays	19
3.3	PET measurements in non-human primates	19
3.3.1	PET experimental procedures	19
3.3.2	PET acquisition in HR and HRRT	20
3.3.3	Determination of radiometabolites in plasma	20
3.3.4	Regions of interest	21
3.3.5	Quantitative PET data analysis	22
3.4	Experimental drugs	23
3.4.1	<i>D</i> -Amphetamine	23
3.4.2	(\pm)-Fenfluramine	24
3.4.3	(<i>R</i>)-Apomorphine	24
3.4.4	Tolcapone and entacapone	25
3.5	Statistical analysis	25
4	Results and comments	26
4.1	Study I: D_2/D_3 agonist PET radioligand development	26
4.2	Study II: Dopamine release measured with [^{11}C]MNPA	27
4.3	Study III: Apomorphine decreases D_2/D_3 radioligand binding	28
4.4	Study IV: D_1/D_5 agonist PET radioligand development	29

4.5	Study V: [¹¹ C]AZ10419369 is sensitive to serotonin release.....	31
4.6	Study VI: Confirmation of serotonin sensitivity of [¹¹ C]AZ10419369 ...	32
5	Methodological considerations	35
5.1	Resolution of the PET system	35
5.2	Reference tissue models	35
5.3	Sample size and test-retest reproducibility	36
5.4	Effects of anaesthesia	36
5.5	Ethical considerations.....	37
6	Summary of Findings	38
7	Future perspective and challenges	39
8	Acknowledgements	41
9	References.....	43

LIST OF ABBREVIATIONS

5-HT	Serotonin
5-HTP	5-hydroxy- <i>L</i> -tryptophan
AADC	Aromatic <i>L</i> -amino acid decarboxylase
AC	Adenylate cyclase
ADHD	Attention deficit hyperactivity disorder
ALDH	Aldehydedehydrogenase
AMPT	Alpha-methyl-para-tyrosine
ATP	Adenosine-5'-triphosphate
B_{max}	Receptor density
BBB	Blood-brain-barrier
<i>BP</i>	Binding potential
cAMP	Cyclic adenosine monophosphate
CNS	Central nervous system
COMT	Catechol- <i>O</i> -methyltransferase
DAT	Dopamine transporter
<i>ex vivo</i>	“Out of the living”, i.e. <i>in vivo</i> tissue studied outside an organism
FOV	Field of view
FWHM	Full width half maximum
G-protein	Guanine nucleotide binding protein
GDP	Guanosine diphosphate
GPCR	G-protein coupled receptor
GTP	Guanosine triphosphate
HPLC	High-performance liquid chromatography
HRRT	High resolution research tomograph
i.m.	Intramuscular
<i>in vitro</i>	“Within the glass”, i.e. in a test tube
<i>in vivo</i>	“Within the living”, i.e. in an organism
i.v.	Intravenous
K_D	Dissociation constant at equilibrium
K_i	Inhibition constant at equilibrium
<i>L</i> -DOPA	<i>L</i> -3,4-dihydroxyphenylalanine
LogD	Logarithm of partition coefficient of octanol/water at pH 7.4
MAO	Monoamine oxidase
MNPA	(<i>R</i>)-(-)-2-methoxy- <i>N</i> -n-propyl-norapomorphine
MRI	Magnetic resonance imaging
MRTM	Multilinear reference tissue model
PET	Positron emission tomography
PKA	Protein kinase A
ROI	Region of interest
SPECT	Single photon emission computerized tomography
SRTM	Simplified reference tissue model
SUV	Standard uptake value
TH	Tyrosine hydroxylase
TP	<i>L</i> -Tryptophan
TPH	Tryptophan hydroxylase
TYR	Tyrosine

1 INTRODUCTION

1.1 MAJOR APPLICATIONS OF PET

Molecular imaging is a rapidly expanding field and covers techniques that allow for the visualization of biochemical processes in living organisms. Prominent modalities are magnetic resonance imaging (MRI), optical imaging and nuclear imaging, with combination of techniques currently under development^{30,77}. Positron emission tomography (PET) is a nuclear imaging technique that allows for quantitative measurement of binding to specific proteins in tissue. The first PET studies of the human brain were performed in the early 1980s and focused on the study of neurotransmitter systems³⁴¹.

The dopamine and serotonin neurotransmitter systems have for several reasons attracted key interest in relation to the pathophysiology of several neurological and psychiatric disorders including, Parkinson's disease, schizophrenia, attention deficit hyperactivity disorder (ADHD) and depression. PET studies have provided the first quantification of receptors in the living brain and thereby allowed for studies of receptor density in relation to pathophysiology^{69,71,347}. A significant enhancement in the clinical utility of brain PET imaging is soon anticipated with recent development of biomarkers of pathophysiology. First examples are radioligands that bind β -amyloid aggregates, which are proteins thought to be heavily deposited in the brains of Alzheimer patients^{160,233}.

A second utility of PET imaging is to understand drug action and to facilitate drug development^{73,121,178}. Pioneering studies have explored dopamine receptor occupancy of antipsychotic drugs and examined the relationship between receptor occupancy, clinical efficacy and side-effects^{74,75,235,237}. Another type of approach undertaken makes use of the "microdosing concept" (typically $< 5\mu\text{g}$) in which a drug is radiolabeled to obtain a detailed description of its distribution in the body¹⁸. More recently, the microdosing approach is gaining increasing acceptance and interest from governmental approving agents and pharmaceutical industries.

Another application which uses PET and neuroreceptor radioligands is to study the competition between radioligand and endogenous neurotransmitter. Pioneering studies have evaluated amphetamine-induced changes in synaptic dopamine level in the primate brain^{57,74,139}, and the clinical relevance of such methodology has been demonstrated by increased dopamine release in patients with schizophrenia^{22,172}. Successful studies of endogenous neurotransmitter release have thus far been almost exclusively reported for dopamine, which is possibly related to the lack of radioligands sensitive to other endogenous neurotransmitters.

PET studies in non-human primates provide a translational model for evaluation of pharmacological mechanisms before initiation of studies in human subjects. Non-human primates are phylogenetically the closest relatives with humans: non-human primates share a large percentage of their DNA with humans and consequently yield physiological and neuroanatomical similarities. Similar complexities of adult human and monkey brains allow for translation from animal models to the human condition more readily than when using more phylogenetically distant animals^{11,230}.

1.2 PRINCIPLES OF PET

After intravenous (i.v.) administration of a substance labelled with a positron-emitting radionuclide (either radiotracer or radioligand), PET imaging allows for evaluation of regional distribution and quantification of radioactivity in the living body (*in vivo*). The radiolabeled substance contains a radionuclide, which at decay emits a

positron (β^+). The positron passes through surrounding tissue (one to a few millimetres) until it annihilates with an electron, resulting in the emission of two 511 keV γ -particles (photons), which travel approximately 180° apart (Figure 1).

The two photons comprise high energy and have therefore high probability to escape from the body. When both photons hit two γ -ray detectors of the PET system within a predefined time window, a “coincidence” event occurs. The coincidence event can be used to locate the positron-electron annihilation, and this location closely approximates of where the positron was emitted. A typical PET measurement consists of a large collection of coincidence events occurring after i.v. administration of a radiolabeled substance to animal or human. Quantitative images are then obtained after reconstruction and appropriate correction for absorption, scatter and random coincidences^{33,66}.




Figure 1. (Left) Summary of physics underlying the PET methodology, figure adopted from³³. (Right) HRRT PET system as used in routine practice at the Karolinska Institutet since 2008.

PET systems must be designed with high sensitivity and high spatial resolution for obtaining acquisition data with high accuracy. Spatial resolution can be defined as a measure of the smallest image area identifiable as a separate unit. The resolution of a PET system is typically expressed in terms of full width at half maximum (FWHM). A Gaussian function is used to represent a perfect point source, and the resolution is defined as the distance between the maximum signal to where the signal intensity is half of the maximum¹⁴. Specific brain dedicated PET systems are available with the high resolution research tomograph (HRRT) being the most recently developed system⁵⁰ (Figure 1). In this thesis two PET systems were used, the ECAT exact HR 47, which has a spatial resolution of 3.8 mm FWHM, and the HRRT, which has a resolution of 1.5 mm FWHM^{334,343}.

1.3 DEVELOPMENT OF PET RADIOLIGANDS FOR CNS

One of the key advantages of PET is that several available short-lived radionuclides are isotopes of elements very typically seen in biological materials, elements such as nitrogen, oxygen and carbon. Incorporation of the radionuclides into molecules does not affect the biological activity of the molecule and thereby provides a unique opportunity to study biological processes in living subjects. Short-lived radionuclides are typically produced onsite by a particle accelerator (cyclotron): the radionuclide used in the present thesis was carbon-11 ($t_{1/2} = 20.4$ min) (Table 1). The

short half-life of carbon-11 limits the possibility of transportation off site, but has the advantage to allow for multiple PET studies in the same subject on the same day.

Table 1. Main characteristics of frequently used radionuclides¹²¹.

	¹⁵ O	¹¹ C	¹⁸ F
Half-life (min)	2	20	110
Mode of decay (% β+)	100%	100%	97%
Maximal energy (MeV)	1.74	0.97	0.64
Penetration distance (mm)	8	4	2

The development of PET radioligands for imaging of new targets has often benefited from organic chemistry efforts in drug discovery. Whereas the properties of an optimal therapeutic drug and radioligand commonly differ, the drug programs provide large numbers of compounds, from structural different classes, of which some may eventually prove useful as drugs and others as radioligands. An example, relevant to this thesis, is the development of agonist radioligands selectively targeting D₁/D₅ or D₂/D₃ receptors. Dopamine agonists developed for the treatment of Parkinson's disease preferably have agonistic properties for both D₁ and D₂ receptor to achieve optimal treatment effect. A radioligand is in contrary preferably selective to one receptor subtype, thereby allowing evaluation of binding to the specific receptor subtype.

PET radioligands to be useful for the examination of targets in the central nervous system (CNS) must fulfil a range of criteria^{122,253}. In short, radioligands should display high selectivity and receptor affinity (K_D in nM range) to the binding target as well as sufficient lipophilicity to enable passage of the blood-brain-barrier (BBB). Too high lipophilicity may result in high non-specific binding in brain. Suitable radioligands for CNS are not substrates for efflux-transporters, such as P-glycoprotein (Pgp), as Pgp can drastically limit accumulation of radioligand in brain¹⁴⁹. In addition, radioligands should have appropriate metabolism avoiding formation of CNS-penetrating radiometabolites. For proper quantification of target binding, a suitable pharmacokinetic profile is of great importance. Optimal receptor binding kinetic properties for ¹¹C-labeled radioligands provide a peak in specific binding within 30-60 minutes after radioligand injection to provide reliable measurements. Finally, amenability for labelling with ¹¹C or ¹⁸F with sufficient high specific radioactivity should ensure straightforward radiosynthesis and the administration of minimal mass of radioligand to fulfil the demands of tracer conditions and safety requirements.

1.4 QUANTITATIVE PET MEASUREMENTS

1.4.1 Quantification of radioligand receptor binding

In pharmacology, the binding reaction between substance (drug or radioligand) and receptor is conventionally described according to equation 1, in which equilibrium exists between available receptors (R), concentration of free substance (F) and concentration of receptor bound substance (B).



At equilibrium conditions, the relationship between receptor binding and the concentration of substance can be described by the Michaelis-Menten equation, which

includes the receptor density (B_{\max}) and the equilibrium dissociation constant (K_D) (equation 2).

$$B = \frac{B_{\max}F}{K_D + F} \quad (2)$$

In applied studies this hyperbolic function can, for instance, be used to describe the relationship between receptor occupancy of drug, maximum receptor occupancy (Occ_{\max}), drug concentration (C_D) and inhibition constant (K_i) (equation 3).

$$\text{Receptor occupancy} = \frac{Occ_{\max}C_D}{K_i + C_D} \quad (3)$$

Neurochemical techniques using radioligands administered at minimal mass, such as PET, allow for redefinition of equation 2. During tracer conditions, often defined as less than 5% receptor occupancy, the concentration of free substance (F) is considered much smaller than the dissociation constant (K_D) resulting in the expression:

$$\frac{B}{F} = \frac{B_{\max}}{K_D} \quad (4)$$

In 1984 Mintun and co-workers used established concepts for *in vitro* radioligand binding to introduce the term binding potential (BP) for quantification of radioligand receptor binding with PET²¹². The BP was defined as the ratio of B_{\max} to K_D . At tracer dose conditions, the BP definition of Mintun equals the ratio of B/F at equilibrium. Recently, the nomenclature for *in vivo* imaging of radioligands with reversible binding was further established and the definition of BP was specified in relation to the reference concentration used to define an affinity constant¹⁴⁰. Shortly, affinity can be expressed by comparison to three distinct reference concentrations, the free plasma concentration of radioligand (BP_F), the total plasma concentration of radioligand (BP_P) or the non-displaceable concentration of radioligand (BP_{ND}). The main outcome parameter in this thesis was BP_{ND} , which represents the product of receptor density (B_{\max}), apparent affinity (K_D^{-1}) and the free fraction of radioligand in the non-displaceable tissue compartment (f_{ND})¹⁴⁰.

1.4.2 Quantification of neurotransmitter release

The application of PET to study neurotransmitter release was first proposed in 1984⁸⁴. Investigations conducted since have resulted in a large number of studies confirming that modification of neurotransmitter concentration can be measured using this technology¹⁷⁵. During PET measurements, the radioligand competes with the neurotransmitter for binding to receptors. When including the free neurotransmitter concentration (F_{NT}) and the neurotransmitter dissociation constant (K_{NT}) in equation 4 it can be rewritten as:

$$BP_{ND \text{ Baseline}} = f_{ND} \frac{B_{\max}}{K_D \left(1 + \frac{F_{NT}}{K_{NT}}\right)} \quad (5)$$

A change in free neurotransmitter concentration (ΔF_{NT}), induced by a pharmacological or physiological intervention, results in a change in BP_{ND} due to the altered competition between radioligand and neurotransmitter to the receptor. In the competition model, it is assumed that the B_{\max} , K_D , F_{NT} and K_{NT} are not modified when

compared to the baseline measurement. The BP_{ND} in the challenge study can accordingly be described as:

$$BP_{ND \text{ Challenge}} = f_{ND} \frac{B_{max}}{K_D(1 + \frac{F_{NT} + \Delta F_{NT}}{K_{NT}})} \quad (6)$$

Comparison of the BP_{ND} obtained during baseline and challenge conditions provides initial evaluation of the sensitivity of a radioligand to neurotransmitter release. By combining equation 5 and 6 the relative change in BP_{ND} (Δ) observed in the challenge condition can be expressed as:

$$\Delta = \frac{BP_{ND \text{ Baseline}} - BP_{ND \text{ Challenge}}}{BP_{ND \text{ Baseline}}} = \frac{\Delta F_{NT}}{K_{NT} + F_{NT} + \Delta F_{NT}} \quad (7)$$

From equation 7 it becomes evident that a significant change in BP_{ND} (Δ) can be expected when the change in neurotransmitter concentration (ΔF_{NT}) is much larger than the sum of the affinity of the neurotransmitter (K_{NT}) and the baseline neurotransmitter concentration (F_{NT}). In addition, it can be concluded that the observed Δ is independent of radioligand characteristics and solely depends on the affinity of the neurotransmitter to the receptor. Development of radioligands suitable for study of neurotransmitter release should therefore preferably target receptors for which the neurotransmitter has high affinity²⁴⁸.

It has previously been argued that “low-affinity” radioligands could be more suitable tools for measurement of neurotransmitter release²⁹⁰. As seen from equation 7, the affinity of the radioligand to the target receptor (K_D) does however not influence the change in BP_{ND} (Δ)⁷². A prerequisite of the validity of equation 7 is that receptor binding is obtained during equilibrium and at tracer dose conditions. Some PET experiments are, on the other hand, performed under rapid changes in neurotransmitter concentration and, under these conditions, it has been suggested that the K_D , or more specifically the K_{off} , is an important parameter allowing for rapid adjustment to the changes in neurotransmitter concentration⁶⁴. Also, the rate constant k_2 has been suggested to be an important radioligand characteristic when the radioligand is to be used for measurement of neurotransmitter release during dynamic conditions^{64,187,216}. No systematic study has thus far been conducted to support these theories with experimental results.

1.5 G-PROTEIN COUPLED RECEPTORS

1.5.1 Families of G-protein coupled receptors

G-protein coupled receptors (GPCRs) are the largest family of membrane proteins and mediate the majority of cellular responses to hormones and neurotransmitters. All GPCRs contain seven membrane-spanning segments, which are separated by alternating intra- and extracellular loop regions. In vertebrates, GPCRs are commonly divided into five families: rhodopsin, secretin, glutamate, adhesion and Frizzled/Taste²⁷⁶.

1.5.2 G-protein coupled receptor signalling

It has been estimated that more than half of available drugs on the market target GPCRs⁸¹. Despite intensive academic and industrial research efforts, the structural basis of GPCR functioning is still not fully understood. Guanine nucleotide binding proteins (G-proteins) are heterotrimeric and exist out of three subunits, α , β and γ , with

the α -subunit containing a guanine nucleotide binding site. In the ternary models, agonist binding is thought to modulate the proportion of receptors that are in an active conformation to those that are inactive and not signalling. An agonist induced conformational change results in activation of the associated heterotrimeric G-protein, involving exchange of bound guanosine diphosphate (GDP) for guanosine triphosphate (GTP) by the α -subunit and thereby causing dissociation of the heterotrimeric complex. The dissociated α - and $\beta\gamma$ -subunits separately promote cellular signalling by second messenger systems, such as adenylate cyclase (AC). Signal transduction is terminated when bound GTP is hydrolyzed to GDP, and the heterotrimeric complex is reunited and coupled back to the receptor^{241,242,275} (Figure 2).




Figure 2. Schematic representation of GPCR activation and deactivation²⁶⁶.

1.5.3 The high affinity binding state

Three decades ago, *in vitro* binding studies on tissue homogenate indicated that GPCRs exist in two affinity states for agonist binding. Saturation studies using an antagonist radioligand showed biphasic displacement after addition of increasing concentrations of an agonist. The high affinity state was thought of as coupled to the G-protein, since the addition of guanine nucleotide altered the biphasic displacement curve into a monophasic curve with similar affinity as the low affinity state. Conversely, antagonist binding was shown as being insensitive to GTP addition and as binding with only one affinity^{52,300,342}. Moreover, Scatchard analyses demonstrated different receptor densities (B_{max}) when obtained with agonist and antagonist D_2/D_3 receptors radioligands¹⁷. The proportion of receptors in the respective states may have implications for function and disease. For instance, the high affinity state of D_2 receptors has been suggested to represent the functional state of the receptor⁹², and it has been proposed that proportions may vary in such CNS-disorders as schizophrenia^{294,296}.

1.5.4 GPCR models

Several models have been used to describe the agonist activation of GPCRs (Figure 3). The ternary model was proposed in 1980 and described the interaction between agonist (A), receptor (R) and G-protein (G)⁵¹. In this model, agonists promote and stabilize the ternary complex (ARG). The model accounts for the heterogeneity of agonist binding and for the high affinity receptor state being sensitive to guanine nucleotides.

In 1993, an extension of the ternary model was proposed. The extension was based on observations that GPCRs have basal activity in the absence of an agonist, and that mutant GPCRs can enhance the agonistic-independent activity^{34,181,281}. In the extended ternary complex, an equilibrium is described between the inactive receptor state (R) and the active receptor state (R*). Agonistic efficacy was thought to be reflected in the ability to modify the equilibrium between R and R*, with R* having potential to bind the G-protein. The extended ternary complex can account for different classes of drug activity, including full agonists, partial agonists, neutral antagonists and inverse agonists.

Over the last two decades it has become clear that not all GPCR properties can be explained by the extended ternary complex^{155,156}. The existence of multiple conformational states has gained support and in this model the inactive receptor state R is, after agonist binding, proposed to gradually conform into the active R* state by intermediate states R' and R''⁹³. Several functional and biophysical studies now support that most GPCRs sample multiple conformations¹⁶¹. The existence and functional relevance of multiple conformational states has been mainly demonstrated *in vitro* and awaits further *in vivo* support.




Figure 3. Three models describing GPCR activation by agonist (A), receptor (R) and G-protein (G). (A) Classical ternary complex. (B) Extended ternary complex. (C) Multistate model^{51,93,181}.

1.5.5 Receptor internalization and β -arrestin

After agonist-induced G-protein dissociation, the receptor can undergo desensitization by phosphorylation and β -arrestin binding to the receptor. This initial desensitization is followed by receptor internalization in which β -arrestin functions as a scaffolding protein. Receptor internalization is a process in which cells internalize cell surface located receptors into plasma membrane vesicles, a process considered to maintain cellular homeostasis. After internalization, the receptor may be recycled or degraded^{78,79}. Recent work in cell lines suggests that β -arrestin not only regulates desensitization, but is a multifunctional adaptor protein also involved in the signalling cascade. Receptor ligands have been shown as unbalanced in stimulating the G-protein and β -arrestin signalling pathways, with some receptor systems activating only one pathway; this mechanism is referred to as “biased agonism”^{192,261,338}. These new insights in GPCR signalling pathways should be taken into account when studying neurotransmitter release with PET.

1.6 THE DOPAMINE SYSTEM

Arvid Carlsson discovered in 1958 that dopamine is of great importance for the functioning of the healthy brain²⁷. Dopamine was characterized as a catecholamine with neurotransmitter function in the CNS. The dopamine system is one of the most widely studied neurotransmitter systems, and has been examined in an extensive number of studies with molecular imaging techniques⁴³. Dopamine is thought to play an important role in physiological functions such as cognition, movement, reward, emotional expression, prolactin secretion and cardiovascular function. The dopamine system is degenerated in Parkinson's disease and Huntington's disease and dopamine dysfunction has been postulated to play a role in ADHD and schizophrenia.

1.6.1 Dopaminergic neurotransmission

Dopamine does not pass the BBB and is synthesised in brain from the essential amino acid tyrosine (TYR). TYR is converted by tyrosine hydroxylase (TH) into *L*-3,4-dihydroxyphenylalanine (*L*-DOPA) and subsequent decarboxylation by the *L*-aromatic acid decarboxylase-enzyme (AADC) results in dopamine. After synthesis dopamine is stored in terminal vesicles and upon neurological firing dopamine is released into the synaptic cleft. Activation by dopamine is primarily terminated by reuptake by the dopamine transporter (DAT). Degradation of dopamine occurs by three enzymes, monoamine oxidase (MAO), aldehydedehydrogenase (ALDH) and catechol-*O*-methyltransferase (COMT)²¹⁵. MAO and ALDH are mainly membrane bound enzymes and are predominately located on the outer layer of mitochondria in neurons and glial cells²⁰. COMT is present in periphery and CNS and located in the cytoplasm of neurons and glial cells in brain¹⁹⁶ (Figure 4).




Figure 4. (Left) Schematic representation of a dopaminergic neuron with cell body (top) and the synaptic terminal region, including synaptically located dopamine receptors (bottom). (Right) Dopaminergic pathways in the human brain, modified from¹³³.

1.6.2 Dopaminergic pathways

The dopaminergic neurons in the midbrain (substantia nigra and ventral tegmental area) and the hypothalamus give origin to the four main dopaminergic pathways (Figure 4). The nigrostriatal pathway projects from the substantia nigra to the dorsal striatum, is important for the control of movement and is the system primarily involved in movement disorders such as Parkinson's disease. The mesocortical and mesolimbic pathways originate from the ventral tegmental area and project, respectively, to the neocortex and to limbic structures, such as the nucleus accumbens, amygdala and hippocampus. Both these pathways have been suggested to have a role in memory, reward and desire, as well as addiction and emotion, and have been proposed to be affected in the pathophysiology of schizophrenia. The fourth pathway is the tuberoinfundibular pathway between the hypothalamus and the pituitary gland. The tuberoinfundibular pathway plays a role in hormonal regulation, maternal behaviour and sensory processes³³⁰. In recent years, the existence of a novel dopaminergic system targeting the thalamus has been described for macaques and humans using immunolabelling techniques^{90,91,282}.

1.6.3 Dopamine receptor subtypes

The effects of dopamine are mediated through five receptor subtypes, divided into two families, the D₁-like receptors (D₁ and D₅) and the D₂-like receptors (D₂, D₃ and D₄), based on pharmacological and structural properties^{152,309,330}. Two splice variants of the D₂ receptor exist, D_{2-short} and D_{2-long} which differ by an insertion of 29 amino acids¹⁰³. The D_{2-short} is mainly located presynaptically and proposed to function as autoreceptor, while the D_{2-long} is mainly located postsynaptically³²⁹.

Activation of the D₁-like receptor class results in the stimulation of G α_s or G α_{olf} , which induces activation of AC. AC catalyzes the conversion of adenosine-5'-triphosphate (ATP) to cyclic adenosine monophosphate (cAMP) and consequently activates the disinhibition of protein kinase A (PKA). PKA in turn causes phosphorylation of several downstream targets, including CREB and DARPP-32. *Post-mortem* autoradiography studies have shown that the D₁ receptor is the most abundant dopamine receptor in the human brain, with high densities in striatum and moderate density in substantia nigra and neocortex^{113,114}. Currently, there are no pharmacological tools available that differentiate between the D₁ and D₅ receptor. Comparison of mRNA levels in monkey brain shows high levels for the D₁ receptor in striatum with lower levels in cortex and amygdala, whereas the D₅ receptor expression is high in cortex^{35,184}. Studies using subtype-specific antibodies have mapped D₅ receptors in several brain regions, including striatum, hippocampus and substantia nigra in rat and human brain¹⁵⁹.

Activation of the D₂-like receptor class results in stimulation of G α_i or G α_o , and inhibition of cAMP production⁴⁹. Consequently, downstream effects of cAMP, such as PKA and phosphorylation of DARPP-32, are thereby inhibited. While D₁ and D₂ receptors have opposite effects at the molecular level, they often have a synergistic action when more complex outputs are considered. *Post-mortem* autoradiography studies have shown that the D₂ receptor density is high in striatum^{113,114} with low levels in neocortex and thalamus^{115,158}. The D₃ receptor has been shown mainly located in the ventral striatum, including the nucleus accumbens, although most studies have been performed with D₂/D₃-receptors unselective radioligands making selective study of the less abundant D₃ receptor cumbersome¹¹⁶. In rodents, the D₄ receptor has been shown to be located in limbic and motor areas using autoradiography⁵³. A recent PET study indicates, however, high D₄ receptor binding in retina but low binding in the monkey brain¹⁶⁷.

1.6.4 PET studies on dopamine release

1.6.4.1 D_2/D_3 receptors antagonist radioligands

PET radioligands developed for imaging of the D_2/D_3 receptors in brain have primarily been antagonist radioligands, such as [^{11}C]raclopride^{61,68} and [^{11}C]FLB 457^{76,118}. A large number of studies have shown that dopamine release induced by a pharmacological or cognitive/behavioural intervention can be studied using D_2/D_3 receptors radioligands and PET or single photon emission computerized tomography (SPECT)^{59,175}. Typically, amphetamine has been used to enhance and reserpine or alpha-methyl-para-tyrosine (AMPT) to decrease dopamine levels, thereby respectively decreasing or increasing D_2/D_3 receptors binding of radioligand in primates¹⁷⁵. Two early observations in human subjects greatly stimulated the use of this methodology for further applications. First an enhanced amphetamine effect on [^{123}I]IBZM and [^{11}C]raclopride binding was observed in patients with schizophrenia^{22,172}. Secondly, videogame playing was shown to cause a significant decrease in [^{11}C]raclopride binding^{59,163}. It has more recently been shown that dopamine release can be detected in extra-striatal regions in some studies using [^{18}F]fallypride^{40,218,219,269} or [^{11}C]FLB 457^{2,3,31,83,228}, but not when using [^{11}C]fallypride²²⁸.

1.6.4.2 D_1/D_5 receptors antagonist radioligands

Previous studies using D_1/D_5 receptors antagonist radioligands, such as (*R*)-[^{11}C]SCH 23390 and (+)-[^{11}C]NNC 112 did not show changes in *BP* after modification of dopamine levels^{4,36}. These observations were not anticipated as electrical induced dopamine release did decrease [^3H]SCH 23390 binding *in vitro*⁹⁵. Several reasons have been proposed to explain this discrepancy.

Firstly, dopamine affinity to the D_1 receptor has been reported to be of the same order as for the D_2 receptor (30 vs. 6 nM, $K_{i\text{ high}}$)²⁹⁵, but more recent studies have shown lower D_1 receptor affinity (897 vs. 64 nM, $K_{i\text{ high}}$)¹¹³. The large variation between results obtained from different assays indicates the dependence to assay conditions. Secondly, the D_1/D_5 receptors have been found to be located predominantly extra-synaptically^{128,183}, which is possibly a mismatch when amphetamine increases dopamine mainly in the synapse. This possible mismatch does not, however, explain the lack of effect on radioligand binding after the long-lasting effect of reserpine. Thirdly, a rather small percentage of D_1 receptors are in the high affinity state, *in vitro* measured as 20-40%^{195,201,271}. An agonist radioligand selectively binding the high affinity state may therefore provide a more sensitive approach to further understand the reported lack of dopamine sensitivity of D_1/D_5 receptors antagonist radioligands.

1.6.4.3 Ceiling effect

The interpretation of the neurotransmitter release studies along the competition model was supported by the relationship between changes in dopamine release measured with microdialysis and observed changes in *BP* with PET^{22,62,173,319,320}. Enhanced dopamine level, however, only modified *BP* values to a small extent, as well as a 44% increase in dopamine level, causing 1% change in *BP*. This modest displacement suggested a maximum effect smaller than that observed for antipsychotics, which could completely inhibit radioligand binding. This effect became referred to as “ceiling effect”. The ceiling effect was proposed to be related to the two affinity states of the D_2/D_3 receptors and the displaceable part of the antagonist radioligand binding was described as binding to the high affinity state. Agonist radioligands, specifically targeting the high affinity state were therefore expected to be more effectively competing with endogenous dopamine than antagonist radioligands,

thereby providing more sensitive tools for measurement of dopamine release (Figure 5)¹⁷⁵.




Figure 5. The proposed pharmacological model for agonist and antagonist radioligand binding to D_2/D_3 receptors, modified from¹⁷⁵.

1.6.5 PET imaging with dopamine agonist radioligands

PET imaging with agonist radioligands was expected to confirm the existence of the high affinity state *in vivo*. In addition, the agonist radioligands were suggested as superior tools for measurement of changes in endogenous dopamine level.

1.6.5.1 Agonist radioligands for the D_1/D_5 receptors

The development of PET radioligands targeting the D_1/D_5 receptors has thus far focused on the 1-phenyl-3-benzazepines scaffold^{8,231}, as other potent D_1/D_5 receptors agonists, such as dihydrexidine¹⁸⁸ and dinapsoline^{94,302}, cannot be easily radiolabeled. DaSilva and co-workers compared (+)-[¹¹C]SKF 75670 and (±)-[¹¹C]SKF 82957⁴⁴ and found (±)-[¹¹C]SKF 82957 to be more suitable⁴⁵. Several studies using [¹¹C]SKF 82957 have been reported, including dopamine challenge studies in rat and baboon^{46,109,174,283,314} as well as pioneering PET studies in humans⁴⁷. However, in 2003, it was shown that a lipophilic radiometabolite accumulated in rat brain after i.v. injection of (+)-[¹¹C]SKF 82957⁴⁸. The formation of this lipophilic radiometabolite most likely confound the quantification of D_1/D_5 receptors binding in brain, and therefore drastically limited the further application of (+)-[¹¹C]SKF 82957 in PET studies⁴⁸.

1.6.5.2 Agonist radioligands for the D_2/D_3 receptors

The development of an agonist radioligand for D_2/D_3 receptors has been pursued for about two decades and includes the radiolabeling of compounds originating from numerous scaffolds, such as the ergolines, aporphines, 2-aminotetralins, benzoquinolines and naphthoxazines. In Appendix I a detailed review is provided on all D_2/D_3 receptors agonist PET radioligands reported in the literature. This section only shortly discusses the three D_2/D_3 receptors agonist radioligands that have been evaluated in human subjects, [¹¹C]PHNO, [¹¹C]NPA and [¹¹C]MNPA (Figure 6).




Figure 6. The three most widely used D_2/D_3 receptors agonist PET radioligands. From left to right: [¹¹C]PHNO, [¹¹C]NPA and [¹¹C]MNPA.

[¹¹C]PHNO binding has been studied in rat⁸⁵, cat¹⁰⁰, non-human primate,²²⁵ and human subjects^{101,105,344}. [¹¹C]PHNO has high D₃ receptor affinity and several studies have confirmed that the [¹¹C]PHNO signal in brain mainly represents D₃ binding^{107,260,289,325}. [¹¹C]PHNO provides a very good binding signal in striatum ($BP_{ND} \sim 3$). One disadvantage of [¹¹C]PHNO, however, is the radiolabeling by ¹¹C-propionylation,³⁴⁶ which typically results in a mass administered of >1 µg. This mass can be considered as a non-tracer dose and may sometimes induce adverse events, such as transient nausea (mass dose of >0.03 µg/kg)²¹³. [¹¹C]PHNO has already been used in clinical studies evaluating D₃ receptor function in neurologic and psychiatric disorders^{19,106}. An optimization of labelling strategy is ongoing and may further advance the use of [¹¹C]PHNO^{60,89}.

Another potent class of D₂/D₃ receptors agonists is the aporphines scaffold. [¹¹C]NPA¹³⁶ was the first aporphine extensively studied in non-human primates^{137,223,224} and has now been evaluated in humans^{177,227,229}. [¹¹C]MNPA is the 2-methoxy derivative of NPA and has been reported to have higher D₂ receptor binding affinity (K_i , 0.17 vs. 0.80 nM)⁸⁷. PET studies have, however, reported comparable striatal BP_{ND} values (0.8-1.0) for [¹¹C]MNPA and [¹¹C]NPA in monkey^{297,315} and human subjects^{162,246}, as possibly partly explained by higher non-displaceable binding due to the higher lipophilicity of [¹¹C]MNPA (3.69 vs. 3.37 cLogD). A clear advantage of [¹¹C]MNPA is the radiolabeling with ¹¹C-methylation, when compared to ¹¹C-propionylation. ¹¹C-Methylation is more routinely used, more automated, more GMP-compliant and achieved with higher specific radioactivity. The agonist radioligand of choice in this thesis has therefore been [¹¹C]MNPA.

1.6.5.3 PET imaging of the high affinity agonist binding site

During the time period of this thesis work several approaches have been undertaken to demonstrate that the D₂/D₃ receptors exist *in vivo* in two affinity states. Scatchard analyses have been performed directly comparing the B_{max} for agonist and antagonist radioligands. Two studies reported no major differences in B_{max} using two-point Scatchard plots in cat and baboon^{100,224}. A different approach in rodents made use of dopaminergic supersensitivity models, in which altered fractions of receptors in the high affinity state were reported using *in vitro* binding techniques. Among four different models, no differences were observed between agonist and antagonist radioligand binding measured either *ex vivo*²⁰³ or *in vivo*³⁰⁴. A third approach taken was to use saturation studies to examine different doses of exogenous dopamine receptor agonists that displace antagonist and agonist radioligand binding. Receptor occupancy of exogenous agonists was found monophasic when measured with [¹¹C]/[³H]raclopride with indistinguishable K_i values as obtained with agonist radioligands^{164,202,250}. Despite this variety of studies attempting different approaches, the existence of the high and low affinity state *in vivo* has not been conclusively disputed or confirmed.

1.6.5.4 Agonist radioligands as improved tools for study of dopamine release

A second rationale for the development of agonist radioligands is to obtain tools that are more sensitive to changes in endogenous dopamine level than the already available D₂/D₃ receptors antagonist radioligands. PET studies using dopamine level altering drugs, such as amphetamine, have consistently reported a more pronounced effect on agonist radioligand binding than on antagonist radioligand binding^{41,42,100,223}. Interestingly, the amphetamine studies showed a similar, amphetamine-dose-consistent ratio of improved sensitivity for agonist radioligands, when compared to antagonist radioligands. Based on the consistent ratio observed, it was suggested that a constant

fraction of receptors was in the high affinity state (60-80%)²²³. The findings stand in contrast to earlier mentioned studies directly investigating the existence of two affinity state *in vivo*.

Importantly, a small number of studies have now shown that anaesthesia and stress modify agonist radioligand binding to D₂/D₃ receptors^{204,240,322}. Direct comparison in humans is required to fully understand the confounding effects of anaesthesia. The first comparative study in man used [¹¹C]NPA and [¹¹C]raclopride, and indicates that the agonist radioligand [¹¹C]NPA was more sensitive to amphetamine-induced changes in dopamine concentration²²⁹. Amphetamine has already been shown to displace [¹¹C]PHNO binding in man³⁴⁵. Although a full comparative study with [¹¹C]PHNO and [¹¹C]raclopride in human subjects has thus far only been reported in abstract form, the preliminary results confirm improved sensitivity to dopamine concentration for the agonist [¹¹C]PHNO²⁹⁹. Confirmation of the enhanced sensitivity of agonist radioligands to endogenous dopamine level in man will encourage the development and future application of agonist radioligands.

1.6.5.5 PET studies supporting the internalization model

Agonist induced internalization of the D₁ and D₂ receptor has been consistently demonstrated *in vitro*^{16,58,318}. Three *in vivo* observations suggest a contribution of internalization and raise concern about the validity of the competition model for interpretation of dopamine challenges studies^{99,175}. First, a temporal discrepancy has been observed between the amphetamine-induced dopamine pulse and the prolonged effect on agonist and antagonist radioligand binding^{26,100,131,173,226}. Second, amphetamine does not decrease receptor binding of all D₂/D₃ receptors radioligands (e.g., butyrophenones)^{125,244}. Third, amphetamine has been shown to decrease [¹¹C]raclopride B_{max} values *in vivo* in cat⁹⁸ and *ex vivo* in rat³¹¹.

To account for these observations, part of the change in radioligand binding may be related to receptor internalization¹⁷⁵. The internalization model was initially proposed by Chugani and co-workers for paradoxical observations using [³H]spiperone³⁷. The model proposes that internalization of receptors includes a relocalisation of receptors into an intracellular compartment resulting in reduced accessibility of radioligands which is reflected in a decrease in B_{max} . Conversely, an elegant *in vitro* study recently conducted observed a decrease in receptor affinity for radioligands after receptor internalization¹¹¹. Whereas the exact contributing factors remain unclear, the internalization model can help to explain the three previously mentioned inconsistencies of the competition model observed in PET studies. Recently, the contribution of internalization in dopamine challenge studies was confirmed using PET. Four hours after amphetamine administration, a decrease in D₂/D₃ receptors radioligand binding was still observed in wild-type mice, but not in arrestin-3 knockout mice, which are animals that lack the capacity to internalize D₂/D₃ receptors³⁰³.

1.7 THE SEROTONIN SYSTEM

Serotonin (5-hydroxytryptamine, 5-HT) was first identified in serum as a vasoconstrictive substance. In 1953, serotonin was found to be present in the mammalian brain³²⁴, and its function in brain has since then been widely studied and an extensive number of studies with molecular imaging techniques have been performed³³⁵. Serotonin is thought to play an important role in normal physiological processes such as appetite, pain perception, sleep and thermoregulation, and serotonin dysfunction has been implicated in addiction, anxiety, depression, migraine, obsessive compulsive disorders and schizophrenia.

1.7.1 Serotonergic neurotransmission

Serotonin does not pass the BBB and is therefore synthesised in serotonergic neurons in brain. The biosynthesis starts from *L*-tryptophan (TP): the initial step is conversion to 5-hydroxy-*L*-tryptophan (5-HTP) by tryptophan hydroxylase (TPH). 5-HTP is then consecutively converted by the enzyme AADC to 5-HT. Both *L*-tryptophan and 5-HTP can pass the BBB, and are integral to our dietary intake. After synthesis, serotonin is stored in terminal vesicles, and upon neurological firing released into the synaptic cleft. Activation by serotonin is terminated by reuptake by the serotonin transporter (SERT), and degradation occurs by MAO and ALDH (Figure 7).




Figure 7. (Left) Schematic representation of a serotonergic neuron with cell body in the raphe nuclei (top) and the synaptic terminal region, including synaptically located 5-HT receptors (bottom). (Right) Serotonergic pathways in the human brain, modified from¹³³.

1.7.2 Serotonergic pathways

Serotonergic innervations in brain arise from cell bodies concentrated in the raphe nuclei. Two distinct subdivisions of raphe nuclei are recognized: the rostral nuclei (located in midbrain and rostral pons) and the caudal nuclei (located primarily in the medulla oblongata). Most brain serotonergic innervations originate from the rostral raphe nuclei, which includes the dorsal raphe nucleus and the median raphe nucleus. From the rostral raphe nuclei, axons ascend to the cerebral cortex, the limbic regions and to the basal ganglia. Serotonergic nuclei in the caudal raphe nuclei give rise to descending axons, some of which terminate in the medulla, while others descend to the spinal cord (Figure 7)^{11,317}.

1.7.3 Serotonin receptor subtypes

The serotonin system is one of the oldest phylogenetic neurotransmitter/hormone systems, and includes a diverse group of receptors. Thus far, fourteen mammalian receptor subtypes have been characterized based on distinct structural and pharmacological properties. The receptor subtypes are assigned to seven families, 5-HT₁₋₇ and all serotonin receptors are G-protein coupled, except for the 5-HT₃

receptor^{15,132}. The receptors can be further categorized into four groups according to their main second messenger system: the 5-HT₁ receptors coupled to G_{α_i}/G_{α_o} proteins (5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E}, 5-HT_{1F}), the 5-HT₂ receptors coupled to G_{α_q} proteins (5-HT_{2A}, 5-HT_{2B}, 5-HT_{2C}), the 5-HT₄, 5-HT₆, and 5-HT₇ receptors coupled to G_{α_s} proteins, and the 5-HT₅ receptors (5-HT_{5A} and 5-HT_{5B}) for which the coupling is still uncertain²⁶⁴.

1.7.4 PET studies on serotonin release

Whereas PET imaging of dopamine release has been successful, this approach has not been successfully extended to the serotonin system²⁴⁸. PET has long been recognized as a promising methodology for measuring serotonin release, thereby providing further understanding of the pathophysiology and treatment of such common CNS-diseases as mood, anxiety, sleep and food disorders. Several suitable radioligands have been developed for PET imaging of the serotonin system, specifically for the 5-HT_{1A}, 5-HT_{1B}, 5-HT_{2A} and 5-HT₄ receptor and SERT. To date, most of these radioligands have been tested for sensitivity to serotonin level, but with limited success.

1.7.4.1 5-HT_{1A} receptor radioligands

An extensive number of radioligands have been developed for the 5-HT_{1A} receptor, with the antagonist [*carbonyl*-¹¹C]WAY-100635 being the most commonly used^{165,245,252}. Several studies have attempted to evaluate the effect of pharmacological-induced changes in serotonin release on [¹¹C]WAY-100635 binding, but have shown inconsistent effects in rodents^{129,135,193,270} and no effect in man²⁵⁹. The development of new radioligands having potential for enhanced sensitivity to serotonin has resulted in [¹⁸F]FPWAY, [¹⁸F]FCWAY and [¹⁸F]MPPF, of which [¹⁸F]MPPF is perhaps the most promising with regards to serotonin sensitivity^{102,142,298}.

Although initial studies with [¹⁸F]MPPF indicated serotonin susceptibility in rat and cat^{12,13,265,268,327,351-353}, the effect has not been consistently confirmed in non-human primates³²⁸ or humans^{56,256,301,326}. The contradictory findings are suggested to partly originate from differences in regional response to serotonin release. 5-HT_{1A} autoreceptors in the raphe nuclei were shown to undergo internalization, while postsynaptic receptors in the hippocampus and cortex do not internalize^{12,268}. A report on fluoxetine effect on serotonin level being restricted to raphe nuclei in man is in line with these regional differences in the serotonin response³⁰¹. The raphe nuclei is, however, a small region, and determination of *BP* has proven challenging and has thus far not resulted in optimal reproducibility^{147,189}.

1.7.4.2 5-HT_{2A} and 5-HT₄ receptor radioligands

Radioligands selectively targeting 5-HT_{2A} or 5-HT₄ receptors have also been tested for sensitivity to modified serotonin release: radioligands targeting 5-HT_{2A} receptors, for example, include the non-selective radioligands [¹⁸F]setoperone, [³H]NMSP and the selective radioligands [¹⁸F]altanserin and [¹¹C]MDL-100907. However, PET studies have shown no evident susceptibility of these radioligands to modified serotonin level in rat^{130,270} and human^{169,199,207,254,306,349}. Currently, only [¹¹C]SB207145 is available for imaging of 5-HT₄ receptors using PET¹⁹⁷, but citalopram did not modify [¹¹C]SB207145 binding in control subjects¹⁹⁸. It can therefore be concluded that none of the serotonin receptor targeting radioligands has unambiguously been shown to be sensitive to serotonin level.

1.7.4.3 SERT radioligands

Another approach for measurement of serotonin release is the use of radioligands targeting SERT. This approach is more challenging as most serotonin challenge studies utilize drugs acting on SERT. However, several attempts have been reported using a non-selective MAO inhibitor, 5-HTP, or TP depletion. After elevation of serotonin level, consistent decreases in [¹¹C]DASB binding has been shown in rat^{190,191}, cat⁹⁷ and non-human primate^{191,348}. Milak and colleagues, however, reported that tryptophan depletion decreased *BP* values, possibly related to serotonin induced SERT internalization²⁰⁸. In two studies in control subjects no effect of TP depletion on [¹¹C]DASB binding has been shown^{257,313}. In summary, although elevation of serotonin levels did effect [¹¹C]DASB binding in animals, this approach has not been found suitable for humans.

1.7.4.4 State of the art for PET imaging of serotonin release

It can be concluded that after two decades of serotonergic radioligand development, no radioligand has been found suitable for study of modified serotonin release in clinical studies in man. Potential reasons for this lack of success were recently reviewed²⁴⁸. Considering the basal serotonin concentration and receptor affinity, imaging of serotonin release may be feasible, but more cumbersome than for dopamine. Comparison of the fraction of receptors in high affinity state has indicted lower fractions for the 5-HT_{1A} (20-40%) and 5-HT_{2A} receptor (40-60%) when compared to D₂ dopamine receptors (70%), possibly explaining the observed low sensitivity to serotonin for radioligands targeting these receptors. Moreover, the internalized receptor pool may vary between receptors, thereby limiting endogenous serotonin competition with the radioligand binding. This may be a particular problem for the 5-HT_{2A} receptor (80-90% internalized).

1.7.4.5 PET imaging of 5-HT_{1B} receptors

Radioligands targeting the 5-HT₇ or 5-HT_{1B} receptor have been suggested as more promising because endogenous serotonin has relative high affinity to these receptor subtypes. Initially, it was claimed that the 5-HT_{1B} receptor only existed in rodents²⁴⁹ but was later demonstrated to be the homologous species of the human 5-HT_{1Dβ} receptor^{6,123,132}. Several physiological functions have been shown to be mediated through 5-HT_{1B} receptors, and animal studies have shown a role in aggression^{243,262,287}, feeding^{157,179}, learning⁷ and locomotion²⁶². The 5-HT_{1B} receptors are also implicated in the pathophysiology and potential treatment of several neuropsychiatric disorders, including anxiety disorders¹⁸⁵, depression²⁷⁹, migraine³³⁷, anorexia¹⁵⁷ and substance abuse^{39,273,274}.

The 5-HT_{1B} receptor acts as autoreceptor^{65,104,200,307} and heteroreceptor⁸⁰ when located at serotonergic or non-serotonergic neurons, respectively. On serotonergic neurons the 5-HT_{1B} receptor is not located on cell bodies, like the 5-HT_{1A} autoreceptors, but on terminals^{21,267,285,286}. Endogenous serotonin has relative high 5-HT_{1B} receptor affinity (~1 nM)²¹¹ and a large proportion of receptors has been shown to be G-protein coupled and thus in the high affinity state^{10,25,108,211,239}. Radioligands targeting the 5-HT_{1B} receptor may thus have potential for measurement of serotonin release. However, until recently, the role of 5-HT_{1B} receptors could not be studied in the living human brain because no suitable PET radioligand was available.

During the time of this thesis work, two novel radioligands suitable for study of the 5-HT_{1B} receptor were reported, [¹¹C]AZ10419369²⁵¹ and [¹¹C]P943⁸⁶. [¹¹C]AZ10419369 is a selective antagonist radioligand for the 5-HT_{1B} receptor subtype

(K_D is 0.8 nM) as confirmed by autoradiography and PET studies in monkey and human^{194,251,333}. [¹¹C]AZ10419369 binding is high in 5-HT_{1B} receptor rich regions, including the globus pallidus, the substantia nigra and occipital cortex. The receptor binding of [¹¹C]AZ10419369 has been shown to be reversible, making this radioligand suitable for quantitative PET measurements of 5-HT_{1B} receptors and drug occupancy *in vivo*^{251,333}.

[¹¹C]P943 is a selective antagonist radioligand for the 5-HT_{1B} receptor and the brain distribution of [¹¹C]P943 has been shown to be very similar to [¹¹C]AZ10419369 in primates^{86,221}. Though, no direct comparison of [¹¹C]P943 and [¹¹C]AZ10419369 has thus far been reported, the reported BP_{ND} values of [¹¹C]AZ10419369 are approximately 30-40% higher³³³. Initial clinical studies using [¹¹C]P943 indicate the clinical relevance of 5-HT_{1B} receptor imaging in alcohol-dependence¹³⁴ and depression²²⁰. The 5-HT_{1B} receptor radioligand of choice in this thesis is [¹¹C]AZ10419369.

2 AIMS

The overall aim of the present thesis was to evaluate drug-induced changes in dopamine and serotonin release in the non-human primate brain using PET.

The specific aims of the program were as follows:

1. To develop improved methods for measurement of endogenous dopamine level, with the following sub-aims:
 - a. To radiolabel and to perform an *in vivo* PET evaluation of the new D₂/D₃ receptors agonist radioligand [¹¹C]MNPA.
 - b. To compare the sensitivity of [¹¹C]MNPA and [¹¹C]raclopride to stimulant-induced dopamine release.
 - c. To apply [¹¹C]MNPA to further obtain *in vivo* support for the existence of two affinity states for the D₂/D₃ receptors.
 - d. To evaluate the sensitivity of the newly developed D₁/D₅ receptors partial agonist (*S*)-[¹¹C]*N*-methyl-NNC 01-0259 to alterations in endogenous dopamine concentration.
2. To evaluate the sensitivity of the new 5-HT_{1B}-receptor radioligand [¹¹C]AZ10419369 to alterations in endogenous serotonin concentration.

3 MATERIALS AND METHODS

The present chapter provides a description of the general methods used during this thesis work. For complete details of the experimental work, the reader is referred to the full papers and manuscripts as indicated elsewhere in this book.

3.1 RADIOCHEMISTRY

All irradiations were performed on a GEMS PETtrace cyclotron (GE, Uppsala, Sweden) equipped with a [¹¹C]methane ([¹¹C]CH₄) target filled with nitrogen gas containing 10% hydrogen. [¹¹C]Methyl iodide ([¹¹C]CH₃I) was prepared according to previously reported methods^{9,170,284}. In short, [¹¹C]CH₄ was collected in a Porapak Q trap cooled with liquid nitrogen and subsequently released into a recirculation system. The [¹¹C]CH₄ was mixed with vapours from iodine crystals at 60°C and then reacted at 720°. After the reaction, [¹¹C]CH₃I was collected in a Porapak Q trap at room temperature and the unreacted [¹¹C]CH₄ was recirculated for three minutes. [¹¹C]Methyl triflate ([¹¹C]CH₃OTf) was prepared by sweeping [¹¹C]CH₃I vapour through a heated glass column containing silver-triflate-impregnated graphitized carbon, as previously described²²².

The produced [¹¹C]CH₃I ([¹¹C]MNPA and (*S*)-[¹¹C]*N*-methyl-NNC 01-0259) or [¹¹C]CH₃OTf ([¹¹C]raclopride and [¹¹C]AZ10419369) was trapped into a reaction vessel containing the corresponding mixture of precursor, solvent and base. After possible heating, the reaction mixture was diluted with mobile phase and injected into the semi-preparative HPLC system for purification. The fraction from the semi-preparative HPLC that contained the product was evaporated to dryness under reduced pressure and the residue dissolved into 8 mL sterile physiological phosphate buffer solution (pH = 7.4). The solution was finally filtered through a Millipore Millex®GV filter unit (0.22 µm).

3.2 IN VITRO RECEPTOR ASSAYS

All *in vitro* receptor affinity and efficacy assays were performed at H. Lundbeck A/S, Valby, Denmark, except for the 5-HT_{2A} receptor assay (Cerep, Paris, France). Data obtained in the assays at H. Lundbeck A/S were measured in a minimum of two full concentration-response curves using 10 concentrations of drugs (covering 4 decades). The results are provided as K_i values (nM) derived from computer fitted IC₅₀ values converted to K_i values using the Cheng-Prusoff equation ($K_i = IC_{50}/(1+(L/K_D))$).

3.3 PET MEASUREMENTS IN NON-HUMAN PRIMATES

3.3.1 PET experimental procedures

Cynomolgus monkeys (*Macaca fascicularis*) are housed in the Astrid Fagraeus Laboratory of the Swedish Institute for Infectious Disease Control (SMI), Solna, Sweden. All studies were approved by the Animal Ethics Committee of the Swedish Animal Welfare Agency (Dnrs: 245/04, 147/05, 16/06, 260/07, 145/08 and 399/08) and were performed according to “Guidelines for planning, conducting and documenting experimental research” (Dnr 4820/06-600) of the Karolinska Institutet as well as the “Guide for the Care and Use of Laboratory Animals”³⁸.

On all experimental days, anaesthesia was initiated by intramuscular (i.m.) injection of ketamine hydrochloride (~10 mg/kg, Ketaminol®, Intervet AB). Following transportation to the PET facility, the anaesthesia was maintained in the research setting by either a mixture of ketamine and xylazine or by sevoflurane. For study I-IV, anaesthesia was maintained for the duration of the experiment with i.m. injections of a

mixture of ketamine hydrochloride (3.75 mg/kg/h, Ketalar®, Pfizer) and xylazine hydrochloride (1.5 mg/kg/h, Rompun® Vet., Bayer). In study V and VI, the anaesthesia was maintained for the duration of the experiment by a mixture of sevoflurane (1.5-8%, Abbott Scandinavia AB), oxygen (~30%) and medical air after endotracheal intubation.

During all experimental days, the head was immobilized with a fixation device¹⁵¹. Body temperature was maintained by Bair Hugger Model 505 (Arizant Healthcare Inc., Eden Prairie, MN, USA) and continuously monitored by an oral or rectal thermometer. Cardiac and respiratory rates were initially monitored manually at least every 20 minutes during the duration of the experiment (study I-IV). For study V and VI, the ECG, heart rate, respiratory rate and blood gasses were continuously monitored throughout the experiment and blood pressure was monitored at least every 15 minutes.

3.3.2 PET acquisition in HR and HRRT

In study I-V, each PET measurement used a sterile physiological phosphate buffer solution (pH = 7.4) containing radioligand that was injected as a bolus (4 mL) into a surreal vein during 5 seconds with simultaneous start of the PET acquisition. As described in study VI, a bolus plus constant infusion approach was applied and included in addition to the bolus injection an infusion of radioligand with a speed of ~10 mL/hour at different K_{bol} values.

In study I-IV, radioactivity in brain was measured continuously with the Siemens ECAT EXACT HR system (Siemens, Knoxville, TN, USA). All acquisitions were acquired in 3D-mode³⁴³. A three-ring detector block architecture gives a 15-cm wide field of view. The transversal resolution in the reconstructed image is about 3.8 mm FWHM and an axial resolution of 3.125 mm. The attenuation correction of the data was obtained with three rotating ⁶⁸Ge rod sources. Raw PET data were then reconstructed using standard filtered back projection consisting of the following reconstruction parameters: 2-mm Hanning filter, scatter correction, a zoom factor of 2.17, and a 128 3 128 matrix size³⁴³. Emission data were collected continuously for 93 min, according to a pre-programmed series of 20 frames starting immediately after i.v. injection of radioligand.

In study V and VI, PET measurements were conducted using the High Resolution Research Tomograph (HRRT) (Siemens Molecular Imaging, Knoxville, TN, USA). List-mode data were reconstructed using the ordinary Poisson-3D-ordered subset expectation maximization (OP-3D-OSEM) algorithm, with 10 iterations and 16 subsets including modelling of the point spread function (PSF). The corresponding in-plane resolution with OP-3D-OSEM PSF was 1.5 mm FWHM in the centre of the field of view (FOV) and 2.4 mm at 10-cm off-centre directions³³⁴. Attenuation correction data was acquired before every PET measurement, with a six minutes transmission measurement using a single ¹³⁷Cs source. List mode data were acquired continuously for 125 or 155 minutes starting at the injection of [¹¹C]AZ10419369 and PET images were then reconstructed with a series of frames.

3.3.3 Determination of radiometabolites in plasma

The evaluation of metabolism of radioligands measured in plasma was performed with slight modification of HPLC methods which were previously described¹¹⁹. In short, venous blood samples (1-2 mL) were obtained from the monkey at several time points after injection of radioligand. After centrifugation at 2000 g for 2 min, plasma was obtained (0.5 mL) and mixed with acetonitrile (0.7 mL). The mixture was centrifuged at 2000 g for 2 minutes and the supernatant (1 mL) was injected to a HPLC system. The radioactivity in blood and plasma were measured in a sodium iodide (NaI)

well counter. The unchanged radioligand fraction was calculated by the integration of the corresponding radioactivity peak and its area was expressed as a percentage of the sum of the areas of all radioactive peaks.

3.3.4 Regions of interest

In study I-III, a preliminary set of anatomical regions of interest (ROIs) for striatum and cerebellum was manually defined on summation images, representing mean radioactivity measured between 9 and 93 min during baseline conditions. Preliminary parametric images of BP_{ND}^{140} and relative blood flow (R_1) were generated from the original reconstructed PET data by use of the two-parameter multilinear reference tissue model (MRTM2)¹³⁸. The final set of ROIs was manually delineated on the fused preliminary R_1 and BP_{ND} images, according to an atlas of a cryosected cynomolgus monkey head in situ¹⁵¹. The final set of ROIs was verified by visual inspection of the delineation on the summation images and applied to all PET studies performed on the same day. ROI volumes of all regions were kept similar in size between experimental days.

In study IV-VI, magnetic resonance images were available of the individual monkeys. Brain magnetic resonance imaging was performed in a 1.5-T GE Signa system (General Electric, Milwaukee, WI, USA). A T1 weighted image was obtained for co-registration with PET and delineation of anatomic brain regions. The T1 sequence was a 3D spoiled gradient recalled (SPGR) protocol with the following settings: repetition time (TR) 21 ms, flip angle 35°; FOV 12.8; matrix 256x256x128; 128x1.0 mm slices; 2 NEX. The sequence was optimized for trade-off between a minimum of scanning time and a maximum of spatial resolution and contrast between gray and white matter.




Figure 8. ROIs delineated on an individual monkey MRI orientated in the horizontal projection. Presented ROIs: dorsal lateral prefrontal cortex (red), caudate nucleus (green), putamen (blue), thalamus (yellow) and occipital cortex (pink).

Before delineation of ROIs, the orientation of the brain was spatially normalized by having the high-resolution T1-weighted magnetic resonance images reoriented according to the line defined by the anterior and posterior commissures being parallel to the horizontal plane and the interhemispheric plane being parallel to the sagittal plane. The standardized T1-weighted MR images were then resliced and used as an individual anatomical template for each monkey. ROIs were defined manually on the reoriented MR images (Figure 8). Mean PET images representing mean radioactivity between 0-57 minutes of the baseline PET measurement were co-registered to the magnetic resonance images using PMOD (PMOD technologies Ltd, Zurich, Switzerland). The generated transformation matrices were then applied to the dynamic emission data sets of all PET measurements obtained on the same day.

3.3.5 Quantitative PET data analysis

All quantitative analyses were based on the assumption that all radioactivity in brain represents unchanged radioligand²¹². Time activity curves were obtained by calculation of regional radioactivity for each frame, corrected for decay and plotted versus time. For quantification of receptor binding, a mathematical model is often required. Typically, this model describes the relationship between input to brain and the brain response to the input. The input can be directly measured using arterial blood sampling or indirectly estimated using a reference region approach. A well accepted mathematical model is the compartmental model (Figure 9). The compartmental model assumes homogenous pools of radioligand concentration, which are assumed to exchange radioligand according to a prescribed set of equations. The values of the parameters of the model can then be estimated by fitting of the data. Several approaches can be used for fitting of the data, including kinetic, equilibrium and graphical analysis. In this thesis work, no arterial blood data was obtained from the monkeys. The kinetic models used for quantification of PET data are therefore reference tissue models.




Figure 9. Multicompartment model. C refers to concentration of radioligand in the defined compartment. C_P = plasma, C_{FT} = free in tissue, C_{NS} = non-specifically bound, C_S = specifically bound, C_{ND} = non-displaceable. K_1 - k_6 refer to kinetic rate constants.

3.3.5.1 The simplified reference tissue model

In study IV the BP_{ND} values were estimated with the Simplified Reference Tissue Model (SRTM), using the cerebellum as reference region. The SRTM model contains only three parameters (R_1 , k_2 and BP_{ND}) in which R_1 is introduced to control for difference in K_1 and K'_1 . The kinetic model is based on two main assumptions. First, the volume of distribution of the non-displaceable bound radioligand is the same in both target and reference region ($C_{ND} = C'_{ND}$). Second, the free concentration (C_{FT}) and non-specific concentration (C_{NS}) equilibrate quickly and can therefore be fitted satisfactorily to a single tissue compartment model. The final expression includes the BP_{ND} and is solved in a convolution manner using least square fitting of the data¹⁶⁸.

3.3.5.2 The multilinear reference tissue model

In study II and III, BP_{ND} values were estimated with MRTM2. This approach is a variation of the graphical method of Logan¹⁸⁶ which derives the ratio of radioligand distribution volumes¹³⁸. The method makes use of linear least squares estimation

algorithms, which is a less computer consuming approach than using non-linear square fitting (such as SRTM) and MRTM is therefore well suited for parametric imaging. Preliminary application of MRTM was used to estimate the cerebellum clearance rate (k'_2). The k'_2 value was then fixed for all voxels (MRTM2) and BP_{ND} and R_1 maps were generated¹³⁸. Final BP_{ND} values were obtained by ROI analysis of the BP_{ND} maps.

3.3.5.3 The transient and late equilibrium approach


In study I and V, the transient and late equilibrium approach were applied, respectively. The occurrence of the transient equilibrium is defined as when the derivative of the specific binding equals zero and at this specific time point the ratio C_S/C_{ND} equals BP_{ND} ⁷². To improve reliability of quantification, a time interval method has been used in which the time interval includes a short period before and after the occurrence of the transient equilibrium²³⁶. Moving the time interval to the late part of the time activity curves^{24,272,339} has been referred to as the late time method¹⁴¹.

3.3.5.4 The continuous infusion equilibrium approach

In study VI a continuous infusion approach was applied. When using this method the BP_{ND} is defined by the ratio of C_S and C_{ND} during the time interval of equilibrium^{28,176}. In study VI $C_{CER}(t)$ was used as estimate for $C_{ND}(t)$. Calculations were performed for a time period at which $C_{CER}(t)$ and $C_S(t)$ were constant.

3.4 EXPERIMENTAL DRUGS


3.4.1 D-Amphetamine



Dextroamphetamine (*D*-amphetamine) and substituted amphetamines, such as methamphetamine (METH) and methylenedioxyamphetamine (MDMA, ecstasy), are widely abused psychostimulant drugs. Amphetamines are substrates for the transporters of the monoamines dopamine, norepinephrine and serotonin. Administration of amphetamines promotes the release of monoamines by several pathways, mainly along two primary mechanisms: reversal of monoamine transporter function and modification of monoamine redistribution between synaptic vesicles and cytosol³¹⁰.


Microdialysis studies in rodents have shown that *D*-amphetamine mainly increases dopamine levels¹⁴⁸. *D*-amphetamine is still widely prescribed for ADHD¹²⁷ and narcolepsy²³² and has negligible affinity to the D_2 receptor¹³⁹ and could therefore be safely used as a test drug in PET studies evaluating dopamine release with D_2/D_3 receptors radioligands in non-human primates^{22,29,36,57,96,125,139,173} and human subjects^{22,73,171,172}. Microdialysis studies in non-human primates^{22,214,288} have shown a dose-dependent relation between amphetamine and increase in dopamine levels, ~500% at 0.27 mg/kg, ~1300% at 0.6 mg/kg, ~1600% at 1.0 mg/kg and ~1800% at 1.5 mg/kg¹⁷³. In study II, *D*-amphetamine was administered intravenously approximately 20 minutes prior to radioligand injection at four doses (0.1, 0.2, 0.5 and 1.0 mg/kg). *D*-amphetamine was obtained from Apoteket, Sweden.

3.4.2 (±)-Fenfluramine




(±)-Fenfluramine is generally considered a substrate for the SERT and may increase extracellular serotonin levels by SERT inhibition and by promotion of a process of carrier-mediated exchange^{88,278}. Microdialysis studies in unanaesthetized monkeys have shown that an i.v. dose of 5 and 10 mg/kg (±)-fenfluramine increases serotonin levels by 20- and 35-fold, respectively³²⁸. Fenfluramine has previously been on the market for treatment of obesity but has been withdrawn due to increased risks of valvular heart disease, possibly related to 5-HT_{2B} receptor affinity²⁷⁷. PET studies evaluating the sensitivity to serotonin levels of 5-HT_{1A} receptor radioligands have previously made use of fenfluramine in rodents and non-human primates^{135,193,351}. (±)-Fenfluramine has been reported to have lower affinity to the 5-HT_{1B} receptor than to the 5-HT_{1A} receptor (16.3 vs. 3.7 μM, IC₅₀)²⁰⁵. In study V and VI (±)-fenfluramine was i.v. administered to the monkeys at doses of 1.0 and 5.0 mg/kg. (±)-Fenfluramine was provided by H. Lundbeck A/S.

3.4.3 (R)-Apomorphine



(R)-Apomorphine is an exogenous dopamine agonist and has *in vitro* high affinity for the D₄ receptor, moderate affinities for D₂, D₃, and D₅ and similar or lower affinity for the D₁ receptor²¹⁰. It has been shown that (R)-apomorphine is a full agonist for both splice variants D_{2-short} and D_{2-long}^{112,206}, and competition studies have demonstrated two binding sites with 34–58 times selectivity to the high affinity state. This selectivity is similar to that reported for dopamine *in vitro* (65–79)^{52,300}. (R)-Apomorphine is currently indicated for symptomatic treatment of recurring episodes of hypomotility (“off” episodes) in patients with advanced Parkinson’s disease⁵⁵. Importantly, by contrast to dopamine, (R)-apomorphine readily passes the BBB. (R)-Apomorphine was, therefore suitable as a test drug for *in vivo* studies and was administered in study III at i.v. doses of 0.01, 0.05, 0.15, 0.5, 1.0 and 3.0 mg/kg. (R)-Apomorphine was obtained from Apoteket, Sweden.

3.4.4 Tolcapone and entacapone



Tolcapone and entacapone are COMT inhibitors¹⁴⁶ and are currently used in conjunction with dopamine agents in the treatment of Parkinson's disease¹⁸⁰. COMT inhibitors are used to decrease metabolism of *L*-DOPA and thereby to improve the efficacy of *L*-DOPA treatment^{144,145,280}. Tolcapone and entacapone inhibit equally rat liver COMT but tolcapone has better brain penetration than entacapone⁸². The effect of COMT inhibitors *in vivo* has been widely studied using 6-[¹⁸F]fluoro-*L*-DOPA^{110,124,182,258} and these PET studies have here been used for dose-selection for study IV. Tolcapone was i.v. administered at 1, 5, 10 and 30 mg/kg and entacapone at 1 and 10 mg/kg. Tolcapone and entacapone were provided by H. Lundbeck A/S.

3.5 STATISTICAL ANALYSIS

In study II, IV and VI, analyses of variance (ANOVA) or repeated measures analyses of variance (RM ANOVA) were performed to test for group differences in BP_{ND} values. In study VI, subsequent paired *t*-tests were performed to test individual regions for fenfluramine effect. The minimum level of significance was designated as $P < 0.05$.

4 RESULTS AND COMMENTS

4.1 STUDY I: D₂/D₃ AGONIST PET RADIOLIGAND DEVELOPMENT

In this study, an agonist radioligand was prepared to evaluate the high affinity state of the D₂/D₃ receptors in the living brain using PET. (*R*)-(-)-2-methoxy-*N-n*-propyl-norapomorphine (MNPA) is a potent D₂/D₃ receptors agonist ($K_i = 0.17$ nM) with high selectivity over the D₁ receptor ($K_i = 1.8$ μ M, 10500 fold)⁸⁷. [¹¹C]MNPA was prepared by direct methylation using [¹¹C]CH₃I and (*R*)-(-)-2-hydroxy-*N-n*-propyl-norapomorphine as precursor. Derivatization of [¹¹C]MNPA and comparison to reference standards on HPLC confirmed a selective labelling of the 2-hydroxy-position. An improved radiosynthesis was more recently developed in which [¹¹C]MNPA is produced in a two-step synthesis starting from (*R*)-(-)-2-hydroxy-10,11-acetonide-*N-n*-propyl-norapomorphine³⁰⁸.

PET measurements after i.v. injection of [¹¹C]MNPA in cynomolgus monkey showed high uptake in D₂/D₃ receptors rich regions, such as putamen and caudate nucleus, and striatum to cerebellum ratios reached a maximum value of 2.2 (Figure 10 and 11).




Figure 10. Colour-coded PET images showing the distribution of radioactivity in the monkey brain after i.v. injection of about 57 MBq [¹¹C]MNPA (summation image of 9-93 min).




Figure 11. (A) Time course for regional radioactivity (nCi/cc) in the brain of a cynomolgus monkey after i.v. injection of [¹¹C]MNPA and corresponding binding ratios (B). (C) Time course for specific binding (nCi/cc) in the brain of a cynomolgus monkey after i.v. injection of [¹¹C]MNPA in baseline and pretreatment conditions. In the pretreatment measurement, raclopride (1 mg/kg) was administered i.v. ten minutes before injection of [¹¹C]MNPA.

Administration of raclopride reduced the specific binding in striatum (Figure 11C) and striatum to cerebellum ratios decreased to 1.3 confirming specific binding of [¹¹C]MNPA to D₂/D₃ receptors. The fraction of total radioactivity in monkey plasma representing unchanged [¹¹C]MNPA was 20% at 45 minutes after injection of radioligand and no radiometabolites were observed more lipophilic than [¹¹C]MNPA, as measured with gradient HPLC. This initial characterization demonstrated that [¹¹C]MNPA has potential as an agonist radioligand for examination of D₂/D₃ receptors in human subjects.

4.2 STUDY II: DOPAMINE RELEASE MEASURED WITH [¹¹C]MNPA

Experimental studies using pharmacological challenges and PET allow for an *in vivo* assessment of synaptic neurotransmitter levels in a non-invasive manner. Amphetamine is a strong dopamine releaser and has been consistently shown to decrease D₂/D₃ receptors binding of some antagonist radioligands in animals and humans. The purpose of this study was to compare the effect of amphetamine-enhanced dopamine level on D₂/D₃ receptors binding of the agonist radioligand [¹¹C]MNPA and the antagonist radioligand [¹¹C]raclopride. Four cynomolgus monkeys were examined with both radioligands before and after i.v. administration of a single dose of amphetamine. Finally, the results obtained were used in an attempt to estimate the proportion of D₂/D₃ receptors in the high and low affinity state.

During baseline conditions, i.v. injection of [¹¹C]MNPA and [¹¹C]raclopride resulted in high accumulation of radioactivity in striatum while concentrations of radioactivity in cerebellum were lower. Comparison of striatal BP_{ND} values showed significantly higher values for [¹¹C]raclopride (5.76 ± 0.95 , $n = 8$) when compared to [¹¹C]MNPA (1.31 ± 0.21 , $n = 8$) (RM ANOVA, $P = 0.00$). Comparison of the BP_{ND} values between the four monkeys showed no significant inter-subject differences for [¹¹C]MNPA and [¹¹C]raclopride (ANOVA, $P > 0.10$). Administration of amphetamine caused a dose dependent reduction in [¹¹C]MNPA striatal BP_{ND} values of 4% at 0.1 mg/kg, 23% at 0.2 mg/kg, 25% at 0.5 mg/kg, and 46% at 1.0 mg/kg. Reductions in [¹¹C]raclopride striatal BP_{ND} values were less than for [¹¹C]MNPA, 2% at 0.1 mg/kg, 16% at 0.2 mg/kg, 15% at 0.5 mg/kg, and 23% at 1.0 mg/kg. The amphetamine effect was found significantly greater for [¹¹C]MNPA than compared to [¹¹C]raclopride (ANOVA, $P = 0.024$) (Figure 12).




Figure 12. BP_{ND} maps of [¹¹C]raclopride and [¹¹C]MNPA estimated by MRTM2 at baseline and post-amphetamine conditions.

The observed improvement in sensitivity to endogenous dopamine level for [¹¹C]MNPA was consistent at all evaluated amphetamine doses, illustrated by the ratio of change in striatal BP_{ND} of [¹¹C]MNPA and [¹¹C]raclopride being 1.85 at 0.1 mg/kg, 1.51 at 0.2 mg/kg, 1.70 at 0.5 mg/kg, and 2.03 at 1.0 mg/kg, with an average of 1.77. Previously, Narendran et al. have proposed that this ratio of changes in BP_{ND} of both tracers could be used to calculate the fraction of receptors in the high affinity state according to equation 8²²³.

$$\frac{\Delta BP_{\text{Rac}}}{\Delta BP_{\text{MNPA}}} = \frac{R_{\text{high avail}}}{B_{\text{max avail}}} \quad (8)$$

Assuming that 10% of the D₂/D₃ receptors are occupied by endogenous dopamine, it was calculated for the current study that 61% of the D₂/D₃ receptors were configured in the high affinity state. Furthermore, by assuming that escalating doses of amphetamine would not further decrease radioligand binding, it was calculated that only 23% of the total pool of D₂/D₃ receptors was in the high affinity state and susceptible to dopamine, so called “synaptic” located receptor sites.

4.3 STUDY III: APOMORPHINE DECREASES D₂/D₃ RADIOLIGAND BINDING

Binding studies *in vitro* have indicated that the D₂ receptor may exist in two affinity states for agonists^{52,300}. *In vitro* studies have further shown that agonists induce measurable D₂ receptor occupancy at clinically relevant concentrations, but only when measured at the high affinity state^{293,295}. Recently developed PET-radioligands, such as [¹¹C]MNPA, have now made it possible to directly study agonist binding *in vivo*. The aim of this study was to obtain further *in vivo* support for the existence of two affinity states for the D₂/D₃ receptors, by comparison of the inhibition by apomorphine of agonist and antagonist radioligand binding *in vivo*. A total of 36 PET measurements were performed with [¹¹C]raclopride or [¹¹C]MNPA in two cynomolgus monkeys. On each study day, a baseline measurement was followed by two consecutive pretreatment studies with rising doses of apomorphine (0.01, 0.05, 0.15, 0.5, 1.0, and 3.0 mg/kg). BP_{ND} values were calculated for the striatum with cerebellum as reference region.




Figure 13. BP_{ND} maps of [¹¹C]raclopride and [¹¹C]MNPA estimated by MRTM2 at baseline and post-apomorphine conditions.




Figure 14: (A) Inhibition of striatal BP_{ND} of [¹¹C]raclopride and [¹¹C]MNPA by pretreatment with apomorphine in two monkeys (A and B). The lines represent an unconstrained four-parameter logistic fit. (B) D₂/D₃ receptors occupancy of apomorphine as a function of plasma concentration 30 min after radioligand injection (ng/mL). The lines represent a one-site binding fit constrained to B_{max} of 100%.

The binding of the antagonist [^{11}C]raclopride as well as the agonist [^{11}C]MNPA was inhibited to a high degree and approached full saturation (Figure 13 and 14). Multifarious analysis methods indicated that the inhibition by apomorphine was monophasic, and the Hill slope coefficients were close to unity. Brain homogenate binding studies *in vitro* have previously demonstrated that apomorphine binds with 30 to 60-fold selectivity to the high affinity state over the low affinity state^{52,300}.

In this study, we obtained non-distinguishable ID_{50} and K_i values of apomorphine for the D_2/D_3 receptors when measured with the antagonist or the agonist radioligand (0.26 and 0.50 mg/kg and 29 and 31 ng/mL, respectively). This PET study provides no support for the existence of two affinity states of the D_2/D_3 receptors. A possible explanation is that almost all D_2/D_3 receptors are in the high affinity state *in vivo*. The results of this study are not directly in line with the outcome of study III. The differences observed could possibly be related to dissimilarity in cellular concentration of agonist or by differences in agonist-induced receptor internalization.

4.4 STUDY IV: D_1/D_5 AGONIST PET RADIOLIGAND DEVELOPMENT

A radioligand with D_1 receptor agonistic properties may provide new understanding of the reasons for lack of sensitivity to dopamine shown for D_1/D_5 receptors antagonist radioligands, such as (*R*)-[^{11}C]SCH 23390^{54,70,117,263} and (+)-[^{11}C]NNC 112¹²⁰. The utility of the previous reported partial D_1/D_5 receptors agonist radioligand (+)-[^{11}C]SKF 82957^{44,45} has been limited due to a brain-penetrant radiometabolite, being formed by COMT⁴⁸. In this study, we radiosynthesised and performed an extensive *in vitro* and *in vivo* evaluation of the new D_1/D_5 receptors partial agonist radioligand (*S*)-[^{11}C]N-methyl-NNC 01-0259 ((*S*)-[^{11}C]1) (Figure 15).

(*S*)-1 has high affinity to the D_1 receptor ($K_i = 4.9$ nM), which is in a similar range as for the antagonists (*R*)-SCH 23390 and (+)-NNC 112 ($K_i = 2.1$ and 1.5 nM, respectively). A functional D_1 receptor assay confirmed high potency of (*S*)-1, but indicated only partial agonistic activity ($\text{EC}_{50} = 1.9$ nM, 35% of dopamine).




Figure 15. Two compounds synthesised in the current study, the partial D_1/D_5 receptors agonist [^{11}C]N-methyl-NNC 01-0259 ((*S*)-[^{11}C]1) and the 7-methoxy analogue of (*S*)-[^{11}C]N-methyl-NNC 01-0259, (*S*)-[^{11}C]2.

The regional distribution of radioactivity after injection of (*S*)-[^{11}C]2 was in accordance with the known distribution of the D_1/D_5 receptors, with high concentration of radioactivity in striatum, moderate in neocortex and lowest in cerebellum. The peak in specific binding in the striatum was observed at ~30 minutes and specific binding ratios reached a maximum value of about 2.0 approximately sixty minutes after radioligand injection (Figure 16).




Figure 16. (A) Regional brain radioactivity (%SUV) after i.v. administration of (*S*)-[¹¹C]1 in cynomolgus monkey. (B) Specific binding in striatum and neocortex. (C) Specific binding ratios in striatum and neocortex. All values represent the mean and standard deviation (presented to above) of nine measurements.

Administration of the dopamine releaser *D*-amphetamine did not have a significant effect on receptor binding of (*S*)-[¹¹C]1. The specificity and selectivity of (*S*)-[¹¹C]1 receptor binding was tested with pretreatment studies using the D₁/D₅ receptors antagonist (*R*)-SCH 23390 and the selective 5-HT_{2A} receptor antagonist MDL-100907. (*R*)-SCH 23390 decreased BP_{ND} in the striatum with 97% to 0.06, while the BP_{ND} value in neocortex only decreased with 77% to 0.12. The mean decrease in BP_{ND} value after MDL-100907 was 19% in striatum and 30% in neocortex.

During the PET measurements in monkeys radiometabolism of (*S*)-[¹¹C]1 was measured in plasma and demonstrated the formation of a radiometabolite, which was more lipophilic than (*S*)-[¹¹C]1. The lipophilic radiometabolite was identified by HPLC as (*S*)-7-methoxy-[¹¹C]*N*-methyl-NNC 01-0259 ((*S*)-[¹¹C]2) (Figure 15). An *in vitro* binding assay demonstrated that (*S*)-2 has affinity for D₁ receptors ($K_i = 52$ nM). A PET study after injection of (*S*)-[¹¹C]2 in monkey confirmed that (*S*)-[¹¹C]2 passes the BBB and concentrates in D₁/D₅ receptors rich brain regions (BP_{ND} in striatum is 0.33). The second part of this project therefore aimed for avoidance of the formation of the confounding lipophilic radiometabolite (*S*)-[¹¹C]2 by the use of COMT inhibitors tolcapone and entacapone. COMT inhibition indeed reduced the formation of (*S*)-[¹¹C]2 measured in plasma and tolcapone administration significantly increased the regional BP_{ND} values, measured after injection of (*S*)-[¹¹C]1 in monkey (unpaired *t*-test, $p < 0.05$) (Figure 17). These results are consistent with a recent study showing that systemic COMT inhibition enabled (+)-[¹¹C]SKF 82957 receptor studies in rodents²⁴⁷.




Figure 17. Striatal BP_{ND} values measured after i.v. injection of (*S*)-[¹¹C]*N*-methyl-NNC 01-0259 in monkeys pretreated with COMT inhibitor entacapone or tolcapone. *Indicates significant difference compared to BP_{ND} of baseline ($p < 0.05$, unpaired *t*-test).

From this study, two main conclusions can be made. Firstly, (*S*)-[¹¹C]1 receptor binding is insensitive to altered dopamine level and (*S*)-[¹¹C]1 is an inferior radioligand for imaging of D₁/D₅ receptors when compared to previous reported antagonist radioligands. Secondly, quantification is complicated by the *in vivo* formation of (*S*)-

[¹¹C]2 but quantitative measurements with PET and (*S*)-[¹¹C]1 are feasible when performed in combination with a COMT inhibitor. As a catechol group is most often critical for D₁ receptor agonism, the COMT inhibition approach may indeed be needed as a more general approach when using a D₁/D₅ receptors agonist radioligand and, importantly, the approach seems suitable for extension to humans.

4.5 STUDY V: [¹¹C]AZ10419369 IS SENSITIVE TO SEROTONIN RELEASE

The need for PET-radioligands that are sensitive to changes in endogenous serotonin levels in brain is recognized in experimental and clinical psychiatric research. Of available serotonergic radioligands, the 5-HT_{1A} receptor antagonist [¹⁸F]MPPF has been the most promising with regard to changes in endogenous serotonin levels in the rat and cat brain^{12,351,352}, but the findings have not been consistently confirmed in monkey and human subjects^{56,256,301,326,328}. Examination of serotonin levels *in vivo* by PET has therefore possibly been precluded by the lack of suitable radioligands. [¹¹C]AZ10419369 is a novel PET radioligand highly selective for the 5-HT_{1B} receptor and recently developed by our group in collaboration with AstraZeneca^{194,251,333}. Of the fourteen 5-HT-receptor subtypes, the 5-HT_{1B} receptor is of particular interest for radioligand development since this primarily presynaptic autoreceptor regulates the release of serotonin^{65,104,200} and thus should be sensitive to serotonin levels.

In this PET study the sensitivity of [¹¹C]AZ10419369 to altered endogenous serotonin levels was examined in cynomolgus monkeys. Serotonin levels were enhanced with the test compound fenfluramine which has been shown to increase basic serotonin levels by 20-fold (5 mg/kg) in monkey³²⁸. The effect of fenfluramine on [¹¹C]AZ10419369 receptor binding was studied in three monkeys using a displacement paradigm. Fenfluramine (1 and 5 mg/kg) was administered in the displacement study during the period of 15 and 20 minutes after radioligand injection.




Figure 18. PET images of mean radioactivity 45–123 min after i.v. injection of [¹¹C]AZ10419369 in a cynomolgus monkey at baseline (top left) and after displacement with 5 mg/kg (±)-fenfluramine (top right). In the bottom corresponding MR images overlaid with PET images.

After administration of fenfluramine there was an evident decrease in [¹¹C]AZ10419369 binding. There was no evident effect of fenfluramine on the radioactivity concentration in the reference region cerebellum, whereas there was a rapid decrease in specific [¹¹C]AZ10419369 binding in all other regions examined. The mean specific binding ratio (SBR) was calculated by use of the area under the curve for time interval 45-123 min (Figure 18 and 19).




Figure 19. (A) Mean radioactivity in the occipital cortex (closed symbols) and cerebellum (open symbols) at baseline and displacement with fenfluramine (1.0 and 5.0 mg/kg). (B) Specific [^{11}C]AZ10419369 binding in the occipital cortex. (C) Specific binding ratios in the occipital cortex. Fenfluramine was administered between 15 and 20 min after radioligand injection and the time interval is illustrated by two vertical lines.

The fenfluramine effect appeared dose-dependent because the decrease was more pronounced after a dose of 5 mg/kg than after 1 mg/kg. Fenfluramine-induced serotonin release decreased the SBR in a dose-dependent fashion with a regional average of 27% after 1 mg/kg and 50% after 5 mg/kg (Figure 20).




Figure 20. (A) Regional mean specific binding ratios (SBR) during baseline and after displacement with fenfluramine (1 and 5 mg/kg). (B) Relative change in SBR after fenfluramine. Bars represent mean \pm SD.

This preliminary study supports that [^{11}C]AZ10419369 is sensitive to endogenous serotonin levels *in vivo* and may serve as a tool to examine the pathophysiology and treatment of major psychiatric disorders.

4.6 STUDY VI: CONFIRMATION OF SEROTONIN SENSITIVITY OF [^{11}C]AZ10419369

The displacement paradigm is attractive for quantification of neurotransmitter release because it allows for direct observation of changes in neuronal activity during pharmacological or physiological stimulation, which are reflected in modified radioligand binding. A methodology considered suitable for quantification of neurotransmitter release, when using a displacement paradigm, is the equilibrium approach, which utilizes the administration of radioactivity by a bolus and constant infusion (BI-protocol)²⁸. The aim of the present study was to develop an updated methodology for measurement of drug-induced serotonin release using [^{11}C]AZ10419369 and PET in non-human primates and to apply this setup to confirm

our previous findings of fenfluramine-induced decreases in [¹¹C]AZ10419369 receptor binding.

A total of 24 PET measurements were conducted, including six preparative PET measurements to assess suitable K_{bol} values. During baseline conditions the C_S/C_{ND} ratios became stable after approximately 50 minutes in the occipital cortex as well as in all other brain regions. K_{bol} values of 180-240 minutes allowed for rapid achievement of steady state and BP_{ND} was calculated between 50-80 minutes (equilibrium before fenfluramine) and between 117-153 minutes (equilibrium after fenfluramine) (Figure 21).




Figure 21. Time activity courses of radioactivity in cerebellum (left) and occipital cortex (middle) after i.v. administration of [¹¹C]AZ10419369 using a BI-protocol in monkey during baseline and displacement conditions with fenfluramine (1.0 and 5.0 mg/kg, FEN). (Right) Corresponding time activity courses for the C_S/C_{ND} values for the occipital cortex are shown on the right. All values represent the mean and standard deviation (presented to above) of three or six measurements. In the displacement studies fenfluramine was administered during the time period indicated by the vertical lines (80-85 min).

During baseline measurements ($n=3 \times 2$) [¹¹C]AZ10419369 BP_{ND} values were stable over time and did not significantly differ during time frames 51'-81' vs. 117'-153' (RM ANOVA, $P = 0.2842$). [¹¹C]AZ10419369 BP_{ND} values ($n=3 \times 2$) of baseline and displacement measurements obtained during 51'-81' showed no significant difference (RM ANOVA, $P = 0.9929$), thus demonstrating good test-retest variability (Figure 22).




Figure 22. BP_{ND} values obtained after i.v. administration of [¹¹C]AZ10419369 using a BI-protocol in monkey during baseline and displacement conditions with fenfluramine (1.0 mg/kg (left) and 5.0 mg/kg (right), FEN). All values represent the mean and standard deviation of three subjects. In the displacement studies fenfluramine was administered during the time period of 80-85 min after the start of BI-protocol. The effect of fenfluramine is therefore only represented in the BP_{ND} values corresponding to FEN (EM 117'-153') values. *Indicates significant difference compared to BP_{ND} of method FEN (EM 51'-81') ($p < 0.05$, two-tailed paired t -test).

Fenfluramine caused a dose-dependent decrease in [^{11}C]AZ10419369 binding in all evaluated regions except for in the cerebellum. In the displacement paradigm fenfluramine significantly reduced [^{11}C]AZ10419369 BP_{ND} (FEN EM 117'-153') when compared to control BP_{ND} (FEN EM 51'-81'), with no significant ROI effect, at both 1.0 mg/kg and 5.0 mg/kg, respectively (RM ANOVA, $P < 0.0001$). Among the different regions, the effect of fenfluramine was significant in the occipital cortex (paired t -test, $p < 0.05$) when [^{11}C]AZ10419369 BP_{ND} (FEN EM 117'-153') were compared to control BP_{ND} (FEN EM 51'-81') for both fenfluramine doses.

To further improve counting statistics, pretreatment studies with fenfluramine were performed, which enabled the use of the majority of the PET measurement for quantification of BP_{ND} values. In the pretreatment paradigm, fenfluramine (5.0 mg/kg) significantly reduced [^{11}C]AZ10419369 BP_{ND} (FEN EM 51'-123') when compared to baseline BP_{ND} (Bsln EM 51'-123') with a significant ROI effect (RM ANOVA, $P < 0.0001$). A significant effect of 5.0 mg/kg fenfluramine on [^{11}C]AZ10419369 BP_{ND} (FEN EM 51'-123') was found in the globus pallidus, occipital cortex, midbrain, thalamus and hippocampus (paired t -test, $p < 0.05$) when compared to baseline BP_{ND} (Bsln EM 51'-123') (Figure 23).




Figure 23. (Left) Time activity courses of C_S/C_{ND} values for the occipital cortex after i.v. administration of [^{11}C]AZ10419369 using a BI-protocol in monkey during baseline and pretreatment conditions with fenfluramine (5.0 mg/kg, FEN). All values represent the mean and standard deviation (presented to both sides) of three subject. (Right) Effect of fenfluramine on BP_{ND} values obtained for eight different brain regions of three subjects. *Indicates significant difference compared to baseline measurements ($p < 0.05$, two-tailed paired t -test).

This study confirms that the new 5-HT_{1B}-receptor radioligand [^{11}C]AZ10419369 is sensitive to fenfluramine-induced changes in endogenous serotonin levels *in vivo*. The effect of fenfluramine on [^{11}C]AZ10419369 BP_{ND} was dose-dependent in the displacement paradigm, but more reliably estimated with the pretreatment paradigm. After pretreatment administration of fenfluramine (5.0 mg/kg), the mean BP_{ND} of the occipital cortex decreased by 39%, from 1.43 ± 0.04 to 0.87 ± 0.08 .

The further developed methodology is suitable for exploring the sensitivity limit to serotonin release as measured using [^{11}C]AZ10419369 and PET. The developed methodology can accordingly be applied to further examine the effect of drugs on brain endogenous serotonin level, and to study serotonin-related brain functions and psychiatric disorders in man.

5 METHODOLOGICAL CONSIDERATIONS

5.1 RESOLUTION OF THE PET SYSTEM

When considering the small volume of the cynomolgus monkey brain (~65 cc), partial volume and spillover effects must be taken into account, in particular in small brain regions.

In this thesis, baseline and drug-challenge PET studies were performed on the same day. Due to the use of a head fixation device the monkey head was maintained in the same position during the day. This approach makes it possible to use the same ROIs for sequentially performed PET measurements. Importantly, potential differences in partial volume and spill over effects were therefore minimized between PET measurements and should therefore not influence the calculated effects significantly. Over the course of the thesis work, two PET systems were used. The ECAT EXACT HR has a in plane resolution of 3.8 mm FWHM³⁴³ and the HRRT system has an in plane resolution of 2.3 mm FWHM³³⁴. The resolution of the HRRT has been further enhanced to 1.5 mm FWHM by software advancements which include the point spread function (PSF) of the PET system³³⁴. The use of a combination of the HRRT and PSF reconstruction provides imaging with improved quantification and reduced partial volume effects^{312,334}. Future studies will benefit from this improvement in methodology.

5.2 REFERENCE TISSUE MODELS

PET measurements in cynomolgus monkeys were performed without arterial cannulation. The quantification of radioligand receptor binding has therefore been limited to the use of reference tissue models. Studies performed with [¹¹C]MNPA, [¹¹C]raclopride, (*S*)-[¹¹C]*N*-methyl NNC 01-0259 and [¹¹C]AZ10419369 have all used cerebellum as a reference region. This approach is based on two main assumptions: that cerebellum contains negligible specific binding and that the non-displaceable component is comparable in cerebellum and other brain regions.

The cerebellum has been shown to contain a negligible density of D₁/D₅ and D₂/D₃ receptors in *post-mortem* autoradiography¹¹⁴⁻¹¹⁶. The use of the cerebellum as a reference region for [¹¹C]raclopride has previously been extensively discussed^{63,72}. The binding of [¹¹C]MNPA in cerebellum was evaluated in study I and III. The absence of specific binding in the cerebellum was confirmed by the lack of change in [¹¹C]MNPA binding in cerebellum after injection of raclopride (study I) and apomorphine (study III). Recent PET studies in non-human primates²⁹⁷ and humans²⁴⁶ further confirmed with kinetic modelling that [¹¹C]MNPA has no specific binding in the cerebellum and that cerebellum can therefore be used as a suitable reference region.

Previous studies using the D₁/D₅ receptors antagonist (*R*)-[¹¹C]SCH 23390 have shown that the cerebellum functions as a suitable reference region^{32,70}. The lack of specific binding in cerebellum after injection of (*S*)-[¹¹C]*N*-methyl-NNC 01-0259 was confirmed in study IV, in which injection of (*R*)-SCH 23390 did not decrease radioligand binding in cerebellum. In addition, administration of the inactive enantiomer (*R*)-[¹¹C]*N*-methyl NNC 01-0259 resulted in radioactivity levels in striatum and neocortex similar to observed as in the cerebellum after injection of (*S*)-[¹¹C]*N*-methyl-NNC 01-0259.

[¹¹C]AZ10419369 has recently been developed²⁵¹ and the complete validation of quantitative methods is currently emerging. Autoradiography studies have previously indicated a negligible 5-HT_{1B} receptor density in cerebellum^{331,332}. The use of the cerebellum as a reference region has been confirmed in studies demonstrating that 5-

HT_{1B} receptor antagonists do not effect binding in cerebellum, when measured with PET in monkey²⁵¹ and with autoradiography in guinea pig¹⁹⁴. Full kinetic modelling in humans has demonstrated that [¹¹C]AZ10419369 binding in cerebellum can be described by a one tissue compartment model³³³, further confirming no specific binding in cerebellum. In study V and VI, fenfluramine did not cause changes in [¹¹C]AZ10419369 binding in cerebellum thereby further confirming the use of cerebellum as a reference region.

5.3 SAMPLE SIZE AND TEST-RETEST REPRODUCIBILITY

A limitation of PET studies is that they typically do not include a large sample size. In this thesis, studies were performed in two or three monkeys per investigated condition.

A small sample size can be accepted when reproducibility between PET measurements is high. Typically, an experimental day included a baseline measurement followed by one or two drug-challenge measurements. To obtain good reproducibility, a head fixation device was used and the monkey head was thereby located in the same position during sequential PET measurements during the day. For PET data analysis the same ROIs could also be applied for both measurements. In a small test-retest study, four cynomolgus monkeys were examined two times after administration of [¹¹C]MNPA two hours apart. The individual test-retest difference in striatal BP_{ND} values ranged from -6.4 to +4.2%, with a mean difference of $-1.1 \pm 4.6\%$ (unpublished results). The high reproducibility is in line with previous reported reproducibility of 3-7% for measurements with (+)-[¹¹C]NNC 112 in monkey³⁶ and with [¹¹C]raclopride²³⁴ and (*R*)-[¹¹C]SCH 23390 in man³². This observation in a small number of animals supports that the measurement of [¹¹C]MNPA binding is reliable.

During the time period of this thesis, MR images of the individual monkeys were obtained. Individual MR image-based ROI templates were generated and used for data analysis in study IV-VI. Co-registration of the baseline PET measurements to the MR images allowed for determination of the transformation matrix for all PET measurements performed on the same day. This method allows for the use of the same ROIs for PET measurements performed on separate days. The reproducibility of [¹¹C]AZ10419369 binding during baseline conditions was examined for six experimental days in one monkey using a BI-protocol. The individual occipital cortex BP_{ND} value differences to the mean value ranged from -3.1% to +4.6%, with a mean difference of $0.0 \pm 3.2\%$ (unpublished results). These results indicate that measurement of [¹¹C]AZ10419369 binding is reliably estimated between experimental days.

5.4 EFFECTS OF ANEASTHESIA

Study I-IV used radioligands targeting dopamine receptors and were performed using i.m. injections of a mixture of ketamine and xylazine. Ketamine is an antagonist for the *N*-methyl-*D*-aspartate (NMDA) receptor and has been reported to bind to the D_{2-long} receptor ($K_i = 55$ nM) *in vitro*²⁹². Ketamine has been reported to exhibit partial or full agonistic properties¹⁵⁰, although this has not been confirmed by others^{143,238}. In addition to direct effects on the receptor, ketamine has been found to increase extracellular dopamine concentrations in the prefrontal cortex in conscious rat, but this increase was blunted or not observed in the striatum of conscious rat and monkey^{5,320,336}.

Further support of the lack of ketamine effect on D₂/D₃ receptors antagonist radioligand binding can be found in PET and SPECT studies. Although initial studies indicated a small reduction in [¹¹C]raclopride binding after ketamine administration^{23,305,320,340} more recent studies have shown no effect on [¹²³I]IBZM or

[¹¹C]raclopride binding^{1,126,153,154}. Taken together there are several but inconsistent observations suggesting that ketamine may influence antagonist radioligand binding to the D₂/D₃ receptors.

Ketamine/xylazine anaesthesia has, however, more recently been shown to increase D₂/D₃ receptors binding of the agonist radioligand [¹¹C]MNPA in monkeys²⁴⁰. In addition, the enhanced sensitivity of [¹¹C]MNPA binding to dopamine release was not observed in unanaesthetized monkeys. In a following study, the authors demonstrated that D₂/D₃ receptors radioligand binding in unanaesthetized animals is related to stress, but differs in direction when measured with an agonist or antagonist radioligand³²³. Taken together it can be concluded that the effect of ketamine and stress on agonist radioligands is complex and warrants further investigation.

The use of another anaesthetic could have been considered for the dopamine studies but, at clinical concentrations, several anaesthetics (e.g., isoflurane, halothane, ketamine and ethanol) have already been shown to inhibit the high affinity states of the D₂ receptor, as well as other GPCRs *in vitro* and *ex vivo*²⁹¹. Moreover, isoflurane was shown to facilitate the actions of various dopamine related pharmacological stimulants, such as amphetamine, cocaine, and nicotine *in vivo*^{202,204,319,321}. Further studies in man are warranted to fully address the possible differential effects of anaesthesia on agonist and antagonist radioligand binding.

Study V and VI evaluated serotonin release in sevoflurane anesthetized monkeys. Microdialysis studies have demonstrated that serotonin level is lower during sleep than during awake conditions²⁵⁵. Comparison of serotonin level during slow-wave sleep and isoflurane anaesthesia showed a similar reduction in serotonin level, to 21-44% of awake condition, when measured with microdialysis in rats²¹⁷. Serotonin levels are therefore likely decreased in anaesthetized animals, independent of the type of anaesthesia. Importantly, in study VI no significant differences were observed between [¹¹C]AZ10419369 *B*P_{ND} values at baseline and during displacement measurements obtained during 51-81 minutes, indicating no differences in anaesthesia effects during the experimental day.

5.5 ETHICAL CONSIDERATIONS

The use of non-human primates in medical research is under constant debate. These unique experiments should be conducted with the highest possible quality and have therefore inherently been subject of improvements during the time of this work. The monkeys were always continuously monitored during and between PET measurements. Introduction of tracheal intubation and gas anaesthesia techniques has further enhanced animal safety and welfare and allows for detailed adjustments in anaesthesia. Temperature, ECG, heart rate, respiratory rate and blood gasses are now being continuously monitored during experimental days and provide valuable information. Out of 150 monkey PET measurements in the current work, no complications were related to experimental procedures.

6 SUMMARY OF FINDINGS

The present thesis focused on drug-induced changes in endogenous neurotransmitter release measured with PET in the living non-human primate brain.

The successful development of the agonist radioligand [^{11}C]MNPA allowed for *in vivo* evaluation of the high affinity state of D₂/D₃ receptors with PET. [^{11}C]MNPA was found more sensitive than the antagonist radioligand [^{11}C]raclopride to amphetamine-induced changes in dopamine level. The improved sensitivity was ~1.8 fold and consistent across four doses. It was estimated that about 60% of the total pool of D₂/D₃ receptors was in the high affinity state.

The existence of two affinity states for D₂/D₃ receptors *in vivo* was further studied using the exogenous agonist apomorphine. D₂/D₃ receptors occupancy was found indistinguishable when measured with [^{11}C]MNPA and [^{11}C]raclopride. This study provided therefore no support for the existence of two affinity states for D₂/D₃ receptors and it was speculated that all receptors are in the high affinity state at *in vivo* conditions. The differences observed between the amphetamine and apomorphine study could possibly be related to dissimilarity in cellular concentration of agonist or by differences in agonist-induced receptor internalization.

To further understand the study of dopamine release with PET, the D₁/D₅ receptors partial agonist (*S*)-[^{11}C]N-methyl-NNC 01-0259 was developed. Evaluation in monkey indicated that the radioligand is insensitive to dopamine level and inferior to previous reported antagonist radioligands in respect to binding signal. COMT inhibition was developed as a method to allow for proper quantification of receptor binding.

Studies on serotonin release made use of the newly developed 5-HT_{1B} receptor antagonist radioligand [^{11}C]AZ10419369. It was demonstrated that the effective serotonin releasing agent fenfluramine caused a dose-dependent reduction in [^{11}C]AZ10419369 receptor binding. The effect of fenfluramine on [^{11}C]AZ10419369 receptor binding was further confirmed in a study with an equilibrium approach using a bolus infusion protocol. If confirmed in human subjects, the developed methodology can be used to study the treatment and pathophysiology of major psychiatric disorders.

7 FUTURE PERSPECTIVE AND CHALLENGES

PET assessment of neurotransmitter release

The study of neurotransmitter release with PET is rapidly gaining interest and benefits from recent improvements in PET methodology and available radioligands. Although the studies have been mainly limited to dopamine, an extension to other neurotransmitter systems is expected in the near future. Understanding of the molecular underpinning of neurotransmitter-induced changes in radioligand binding has proven challenging, and including biological and chemical aspects. The combination of microPET imaging with molecular-modified animals provides new pathways to evaluate underlying mechanisms. Finally, for the methodology to provide clinical utility, improvements in sensitivity are required to allow for investigation of dynamic neurotransmitter concentrations at more physiological relevant levels.

Based on this thesis some specific challenges can be discussed:

Two affinity states in vivo

The existence of two affinity states for the D₂/D₃ receptors has not been demonstrated *in vivo* consistently and additional studies are required. Recent development of radioligands targeting the 5-HT_{1A}, 5-HT_{2A} and opioid κ-receptors make it possible to extent comparison of agonist and antagonist radioligand binding to other neurotransmitter systems^{67,166,209,316,350}.

D₂/D₃ agonist radioligands as improved tools for measurement of dopamine release

PET studies in anaesthetized animals have consistently reported a more pronounced effect of dopamine concentration on D₂/D₃ agonist radioligand binding than on D₂/D₃ antagonist radioligand binding. Future studies are required to understand the mechanisms of enhanced dopamine level sensitivity of agonist radioligands. Finally, studies in human subjects are needed to confirm that agonist radioligands have increased sensitivity to dopamine levels and to elucidate the potential contribution of stress.

Improvements for agonist radioligands targeting D₂/D₃ receptors

Existing D₂/D₃ receptors agonist PET radioligands are not optimal yet. For further improvement, several aspects can be considered including: receptor selectivity, receptor affinity, receptor kinetics, non-specific binding, specific radioactivity and ease of radiolabelling.

Development of a full D₁ receptor agonist radioligand

As a catechol group is most often critical for D₁ receptor agonism, the developed COMT inhibition approach provides a more general method for full quantitative measurement with D₁/D₅ receptors agonist radioligands. A full D₁ agonist radioligand is not yet available and is required to further understand the lack of sensitivity of antagonist and partial agonist radioligands to changes in endogenous dopamine level. The so far radiolabeled D₁/D₅ receptors agonist and antagonist PET radioligands almost all originate from the 1-phenyl-3-benzazepines scaffold. A radiolabeled agonist originating from another scaffold may be of interest to rule out possible scaffold-related insensitivity to dopamine level.

Measurement of serotonin release using [¹¹C]AZ10419369 and PET

The results of study V and VI demonstrate that [¹¹C]AZ10419369 binding is sensitive to major changes in endogenous serotonin level. Challenges causing smaller changes in serotonin concentration are required to explore the sensitivity of the methodology. Finally, the findings need to be confirmed in control human subjects to explore the possibilities of prospective clinical studies.

Improvements for radioligands targeting 5HT_{1B} receptors

The development of a 5-HT_{1B} receptor agonist radioligand would be of interest, as it may have enhanced sensitivity to serotonin concentration. In addition, radioligands with higher affinity could allow for study of serotonin concentration in 5-HT_{1B} receptor low regions, such as the hippocampus.

8 ACKNOWLEDGEMENTS

Hereby, I express my sincere gratitude and appreciation to everyone who has made this thesis possible. I am especially grateful to:

Professor Christer Halldin, my main supervisor, for guiding my way into the world of radiochemistry and PET, and for your endless optimism, enthusiasm and open-mindedness to exploring new scientific challenges.

Professor Lars Farde, my co-supervisor, for the inspiration offered by sharing your tremendous analytical and critical scientific skills, as well as your impressive knowledge on neuropsychopharmacology.

Professor Håkan Wikström, my co-supervisor, for guiding my first steps into neuroscience and for paving the way to Sweden and the Karolinska Institutet. Your tremendous scientific drive and deep understanding of dopaminergic neurotransmission will always inspire me.

Dr. Benny Bang-Andersen, my co-supervisor, for fruitful scientific discussions and an unlimited supply of goodwill and chemical resources. I look forward to see some additional shared projects come through!

Associate Professor Bálazs Gulyás and Gudrun Nylén, for many shared hours spent together with our non-human primate friends and for our many discussions on life and science. Your individual and combined dedication has yielded an artful methodological approach to this work, my sincerest gratitude to you both for teaching me how to perform monkey PET studies. Thanks also to the SMI staff for excellent animal house keeping and to all our primate cousins for participation.

All current and previous colleagues at the radiochemistry group at the Karolinska Institutet: Dr. Jan Andersson, Dr. Magnus Schou and Carsten Steiger for friendship, discussions on life and pleasant conference attendance; my first officemate, Phong Truong, for lasting friendship even as you travel the globe; Dr. Vladimir Stepanov, for your scientific brightness and shared interest in receptor kinetics; Drs. Evgeny Shchukin and Anu Airaksinen, for introducing me into the world of radiochemistry and for help during the early days of this thesis; Ithamar Brinkman and Hans Buitter for MNPA chemistry and significant contributions to the “Dutch colonization” of KI. For excellent technical assistance in this thesis work, I thank especially Dr. Zhisheng Jia, Guennadi Jogolev, Arsalan Amir, Kenneth Dahl, Dr. Jonas Bergström and Dr. Fabienne Gourand. All other members of the radiochemistry group which have made life at work easygoing: Mahabuba Jahan, Sangram Nag, Dr. Peter Johnström, Anna Sumic, Linda Bergman, Dr. Ryuji Nakao, Siv Eriksson, Dr. Raisa Krasikova, Mirkka Sarparanta, Dr. Sean Donohue and Jari Tarkiainen.

All current and previous colleagues at the PET imaging group at the Karolinska Institutet: Dr. Nicholas Seneca, for teaching me the first steps of PET data analyses, for your impressive high-speed-data generation and manuscript preparation, and valuable

friendship. To the growing list of roommates, I thank my Japanese friends Drs. Akihiro Takano, Shigeyuki Yamamoto, Yoko Ikoma, Tomoyuki Saijo, Tetsya Ichimiya, and Swedish “kamrater” Drs. Anton Forsberg and Johan Lundberg for interesting discussions, and for creating a nice work environment. To Dr. Tzung-Jeng Hwang, for your contribution to the serotonin studies, may we spend more time together in the future. To Dr. Andrea Varrone, for discussions on kinetic modeling as well as your excellent contribution to the bolus infusion studies. To Dr. Judit Sóvágó, for early help with monkey studies. To Dr. Zsolt Cselényi and Dr. Katarina Varnäs, for interesting discussions on PET data analyses, and friendship. To all other members of the group for unique brainpower: Professor Anna-Lena Nordström, Dr. Per Kalsson, Dr. Jacqueline Borg, Dr. Simon Cervenka, Dr. Bengt Andree, Dr. Mirjam Talvik, Dr. Mikael Tiger, Dr. Hristina Jovanovic, Dr. Aurelija Jucaite, Dr. Patrik Hansson, Dr. Hans Olsson, Dr. Nina Erixon-Lindroth, Dr. Magdalena Nord, Martin Schain, Miklos Toth and Pavitra Kannan.

Current and past staff members of the PET group, who have made a difference during these years: Karin Zahir, Ulla-Kajsa Pehrson, Pia Schönbeck, Sofia Sjödin, Johan Molin, Kjerstin Lind and Jan Everhov. Thanks to Nils Sjöholm, Julio Gabriel, Urban Hansson and Dr. Stefan Pauli for providing always functioning PET systems and generating high-quality, and sometimes even spinning, images.

Collaborators at H. Lundbeck A/S (Valby), Dr. Morten Jørgensen, Dr. Christoffer Bundgaard and Dr. Lise Tøttrup Brennum, for a great collaboration on dopamine agonists; at AstraZeneca (Wilmington/Södertälje), Dr. Edward Pierson, Dr. Svante Nyberg and Dr. Charlotte Ahlgren, for fruitful collaboration; at NIMH (Bethesda), Dr. Robert Innis, Dr. Victor Pike, and former KI-NIMH PhD program students, Dr. Mette Skinbjerg and Dr. Garth Terry for fruitful collaboration, in particular on [¹¹C]MNPA; at NIRS (Chiba), Dr. Tetsya Suhura, Dr. Kazutoshi Suzuki, Dr. Hiroshi Ito and Dr. Makoto Higuchi, for providing me the opportunity to perform PET studies on awake animals and Dr. Masaki Tokunaga, for helping me around in Japan during the research training month, and for sharing your great scientific skills.

Members of the radiochemistry group of the Karolinska pharmacy, Professor Sharon Stone-Elander, Dr. Jan-Olov Thorell, Dr. Anna Fredriksson, Erik Samen and Emma Jussing for smooth co-operation in the radiochemistry labs.

Friends from home, especially Ronald and Katarina, Theo and Afke, Sytse and Elena, Djoek and Marlies, Gerard, Bas, and Manne. Although the distance has been there over the years it has never felt like it. Thanks for all support and travels in both directions.

De famylje thús yn Fryslân, fan Beppes oan't Finette, tige tank foar alle stipe trôch de jierren hinne. It is fijn dat der altyd plak foar ús is, wert wij ek mar wei moatte komme.

Finally, though not least, I especially thank my family, Pauline and Noah, for making Sweden our home, and for your eternal support, inspiration and love. Dearest Noah, by outwitting science, you have put things into the right perspective.

9 REFERENCES

1. Aalto S., J. Hirvonen, J. Kajander, H. Scheinin, K. Nagren, H. Vilkmann, L. Gustafsson, E. Syvalahti and J. Hietala. **2002**. Ketamine does not decrease striatal dopamine D2 receptor binding in man. *Psychopharmacology (Berl)* 164(4):401-406.
2. Aalto S., A. Bruck, M. Laine, K. Nagren and J.O. Rinne. **2005**. Frontal and temporal dopamine release during working memory and attention tasks in healthy humans: a positron emission tomography study using the high-affinity dopamine D2 receptor ligand [11C]FLB 457. *J Neurosci* 25(10):2471-2477.
3. Aalto S., J. Hirvonen, V. Kaasinen, N. Hagelberg, J. Kajander, K. Nagren, T. Seppala, J.O. Rinne, H. Scheinin and J. Hietala. **2009**. The effects of d-amphetamine on extrastriatal dopamine D2/D3 receptors: a randomized, double-blind, placebo-controlled PET study with [11C]FLB 457 in healthy subjects. *Eur J Nucl Med Mol Imaging* 36(3):475-483.
4. Abi-Dargham A., N. Simpson, L. Kegeles, R. Parsey, D.R. Hwang, S. Anjilvel, Y. Zea-Ponce, I. Lombardo, R. Van Heertum, J.J. Mann, C. Foged, C. Halldin and M. Laruelle. **1999**. PET studies of binding competition between endogenous dopamine and the D1 radiotracer [11C]NNC 756. *Synapse* 32(2):93-109.
5. Adams B.W., C.W. Bradberry and B. Moghaddam. **2002**. NMDA antagonist effects on striatal dopamine release: microdialysis studies in awake monkeys. *Synapse* 43(1):12-18.
6. Adham N., P. Romanienko, P. Hartig, R.L. Weinshank and T. Branchek. **1992**. The rat 5-hydroxytryptamine1B receptor is the species homologue of the human 5-hydroxytryptamine1D beta receptor. *Mol Pharmacol* 41(1):1-7.
7. Ahlander-Luttgen M., N. Madjid, P.A. Schott, J. Sandin and S.O. Ogren. **2003**. Analysis of the role of the 5-HT1B receptor in spatial and aversive learning in the rat. *Neuropsychopharmacology* 28(9):1642-1655.
8. Andersen P.H. and J.A. Jansen. **1990**. Dopamine receptor agonists: selectivity and dopamine D1 receptor efficacy. *Eur J Pharmacol* 188(6):335-347.
9. Andersson J., P. Truong and C. Halldin. **2009**. In-target produced [C-11]methane: Increased specific radioactivity. *Applied Radiation and Isotopes* 67(1):106-110.
10. Audinot V., S. Lochon, A. Newman-Tancredi, G. Lavielle and M.J. Millan. **1997**. Binding profile of the novel 5-HT1B/1D receptor antagonist, [3H]GR 125,743, in guinea-pig brain: a comparison with [3H]5-carboxamidotryptamine. *Eur J Pharmacol* 327(2-3):247-256.
11. Azmitia E.C. and P.J. Gannon. **1986**. The primate serotonergic system: a review of human and animal studies and a report on *Macaca fascicularis*. *Adv Neurol* 43:407-468.
12. Aznavour N., L. Rbah, M. Riad, A. Reilhac, N. Costes, L. Descarries and L. Zimmer. **2006**. A PET imaging study of 5-HT(1A) receptors in cat brain after acute and chronic fluoxetine treatment. *Neuroimage* 33(3):834-842.
13. Aznavour N. and L. Zimmer. **2007**. [18F]MPPF as a tool for the in vivo imaging of 5-HT1A receptors in animal and human brain. *Neuropharmacology* 52(3):695-707.
14. Baily D.L., D.W. Townsend and P.E. Valk. **2005**. Positron Emission Tomography. Maisey MN, editor. London: Springer-Verlag.
15. Barnes N.M. and T. Sharp. **1999**. A review of central 5-HT receptors and their function. *Neuropharmacology* 38(8):1083-1152.
16. Barton A.C., L.E. Black and D.R. Sibley. **1991**. Agonist-induced desensitization of D2 dopamine receptors in human Y-79 retinoblastoma cells. *Mol Pharmacol* 39(5):650-658.
17. Battaglia G. and M. Titeler. **1982**. [3H]N-propylapomorphine and [3H]spiperone binding in brain indicate two states of the D2-dopamine receptor. *Eur J Pharmacol* 81(3):493-498.

18. Bergstrom M., A. Grahnen and B. Langstrom. **2003**. Positron emission tomography microdosing: a new concept with application in tracer and early clinical drug development. *Eur J Clin Pharmacol* 59(5-6):357-366.
19. Boileau I., M. Guttman, P. Rusjan, J.R. Adams, S. Houle, J. Tong, O. Hornykiewicz, Y. Furukawa, A.A. Wilson, S. Kapur and S.J. Kish. **2009**. Decreased binding of the D3 dopamine receptor-preferring ligand [11C]-(+)-PHNO in drug-naïve Parkinson's disease. *Brain* 132(Pt 5):1366-1375.
20. Bortolato M., K. Chen and J.C. Shih. **2008**. Monoamine oxidase inactivation: from pathophysiology to therapeutics. *Adv Drug Deliv Rev* 60(13-14):1527-1533.
21. Boulenguez P., R. Pinard and L. Segu. **1996**. Subcellular localization of 5-HT1B binding sites in the stratum griseum superficiale of the rat superior colliculus: An electron microscopic quantitative autoradiographic study. *Synapse* 24(3):203-212.
22. Breier A., T.P. Su, R. Saunders, R.E. Carson, B.S. Kolachana, A. de Bartolomeis, D.R. Weinberger, N. Weisenfeld, A.K. Malhotra, W.C. Eckelman and D. Pickar. **1997**. Schizophrenia is associated with elevated amphetamine-induced synaptic dopamine concentrations: evidence from a novel positron emission tomography method. *Proc Natl Acad Sci U S A* 94(6):2569-2574.
23. Breier A., C.M. Adler, N. Weisenfeld, T.P. Su, I. Elman, L. Picken, A.K. Malhotra and D. Pickar. **1998**. Effects of NMDA antagonism on striatal dopamine release in healthy subjects: application of a novel PET approach. *Synapse* 29(2):142-147.
24. Brooks D.J., V. Ibanez, G.V. Sawle, E.D. Playford, N. Quinn, C.J. Mathias, A.J. Lees, C.D. Marsden, R. Bannister and R.S. Frackowiak. **1992**. Striatal D2 receptor status in patients with Parkinson's disease, striatonigral degeneration, and progressive supranuclear palsy, measured with 11C-raclopride and positron emission tomography. *Ann Neurol* 31(2):184-192.
25. Bruinvels A.T., J.M. Palacios and D. Hoyer. **1993**. Autoradiographic characterisation and localisation of 5-HT1D compared to 5-HT1B binding sites in rat brain. *Naunyn Schmiedebergs Arch Pharmacol* 347(6):569-582.
26. Cardenas L., S. Houle, S. Kapur and U.E. Busto. **2004**. Oral D-amphetamine causes prolonged displacement of [11C]raclopride as measured by PET. *Synapse* 51(1):27-31.
27. Carlsson A. **1959**. The occurrence, distribution and physiological role of catecholamines in the nervous system. *Pharmacol Rev* 11:490-493.
28. Carson R.E., M.A. Channing, R.G. Blasberg, B.B. Dunn, R.M. Cohen, K.C. Rice and P. Herscovitch. **1993**. Comparison of bolus and infusion methods for receptor quantitation: application to [18F]cyclofoxy and positron emission tomography. *J Cereb Blood Flow Metab* 13(1):24-42.
29. Carson R.E., A. Breier, A. de Bartolomeis, R.C. Saunders, T.P. Su, B. Schmall, M.G. Der, D. Pickar and W.C. Eckelman. **1997**. Quantification of amphetamine-induced changes in [11C]raclopride binding with continuous infusion. *J Cereb Blood Flow Metab* 17(4):437-447.
30. Cassidy P.J. and G.K. Radda. **2005**. Molecular imaging perspectives. *J R Soc Interface* 2(3):133-144.
31. Cervenka S., C. Halldin and L. Farde. **2008**. Age-related diurnal effect on D2 receptor binding: a preliminary PET study. *Int J Neuropsychopharmacol* 11(5):671-678.
32. Chan G.L., J.E. Holden, A.J. Stoessl, D.J. Doudet, Y. Wang, T. Dobko, K.S. Morrison, J.M. Huser, C. English, B. Legg, M. Schulzer, D.B. Calne and T.J. Ruth. **1998**. Reproducibility of the distribution of carbon-11-SCH 23390, a dopamine D1 receptor tracer, in normal subjects. *J Nucl Med* 39(5):792-797.
33. Cherry S.R. **2001**. Fundamentals of positron emission tomography and applications in preclinical drug development. *J Clin Pharmacol* 41(5):482-491.
34. Chidiac P., T.E. Hebert, M. Valiquette, M. Dennis and M. Bouvier. **1994**. Inverse agonist activity of beta-adrenergic antagonists. *Mol Pharmacol* 45(3):490-499.

35. Choi W.S., C.A. Machida and O.K. Ronnekleiv. **1995**. Distribution of dopamine D1, D2, and D5 receptor mRNAs in the monkey brain: ribonuclease protection assay analysis. *Brain Res Mol Brain Res* 31(1-2):86-94.
36. Chou Y.H., P. Karlsson, C. Halldin, H. Olsson and L. Farde. **1999**. A PET study of D(1)-like dopamine receptor ligand binding during altered endogenous dopamine levels in the primate brain. *Psychopharmacology (Berl)* 146(2):220-227.
37. Chugani D.C., R.F. Ackermann and M.E. Phelps. **1988**. In vivo [3H]spiperone binding: evidence for accumulation in corpus striatum by agonist-mediated receptor internalization. *J Cereb Blood Flow Metab* 8(3):291-303.
38. Clark J.D., R.L. Baldwin, K.A. Bayne, M.J. Brown, G.F. Gebhart, J.C. Gonder, J.K. Gwathmey, M.E. Keeling, D.F. Kohn, J.W. Robb, O.A. Smith, J.-A.D. Steggerda and J.L. VandeBergh. **1996**. Guide for the care and use of laboratory animals. Washington D.C.: The National Academies Press.
39. Crabbe J.C., T.J. Phillips, D.J. Feller, R. Hen, C.D. Wenger, C.N. Lessov and G.L. Schafer. **1996**. Elevated alcohol consumption in null mutant mice lacking 5-HT1B serotonin receptors. *Nat Genet* 14(1):98-101.
40. Cropley V.L., R.B. Innis, P.J. Nathan, A.K. Brown, J.L. Sangare, A. Lerner, Y.H. Ryu, K.E. Sprague, V.W. Pike and M. Fujita. **2008**. Small effect of dopamine release and no effect of dopamine depletion on [18F]fallypride binding in healthy humans. *Synapse* 62(6):399-408.
41. Cumming P., D.F. Wong, R.F. Dannals, N. Gillings, J. Hilton, U. Scheffel and A. Gjedde. **2002**. The competition between endogenous dopamine and radioligands for specific binding to dopamine receptors. *Ann N Y Acad Sci* 965:440-450.
42. Cumming P., D.F. Wong, N. Gillings, J. Hilton, U. Scheffel and A. Gjedde. **2002**. Specific binding of [(11)C]raclopride and N-[(3)H]propyl-norapomorphine to dopamine receptors in living mouse striatum: occupancy by endogenous dopamine and guanosine triphosphate-free G protein. *J Cereb Blood Flow Metab* 22(5):596-604.
43. Cumming P. **2009**. Imaging dopamine. New York: Cambridge University Press. 340 p.
44. DaSilva J.N., A.A. Wilson, J.N. Nobrega, D. Jiwa and S. Houle. **1996**. Synthesis and autoradiographic localization of the dopamine D-1 agonists [11C]SKF 75670 and [11C]SKF 82957 as potential PET radioligands. *Appl Radiat Isot* 47(3):279-284.
45. DaSilva J.N., A.A. Wilson, C.M. Valente, D. Hussey, D. Wilson and S. Houle. **1996**. In vivo binding of [11C]SKF 75670 and [11C]SKF 82957 in rat brain: two dopamine D-1 receptor agonist ligands. *Life Sci* 58(19):1661-1670.
46. DaSilva J.N., E. Greenwald, R.A. Schwartz, A.A. Wilson and S. Houle. **1998**. Chronic reserpine differentially alters *in vivo* binding of D₁ agonists R/S- and R-[¹¹C]SKF 82957 as compared to [¹¹C]SCH 23390 in rat brain. *Soc Neurosci Abstr* 24:22.
47. DaSilva J.N., R.A. Schwartz, D. Hussey, K. Cheung, A.A. Wilson and S. Houle. **1999**. Human PET imaging with the dopamine D1 agonist R-[11C]-SKF 82957. *J Cereb Blood Flow Metab* 19:S332??
48. DaSilva J.N., H. Cheung, A.A. Wilson and S. Houle. **2003**. Metabolism of dopamine D₁ agonist R-(+)-[¹¹C]SKF 82957 produces a radiolabeled metabolite in rat brain. *J Label Compd Radiopharm* 46:S368.
49. De Camilli P., D. Macconi and A. Spada. **1979**. Dopamine inhibits adenylate cyclase in human prolactin-secreting pituitary adenomas. *Nature* 278(5701):252-254.
50. de Jong H.W., F.H. van Velden, R.W. Kloet, F.L. Buijs, R. Boellaard and A.A. Lammertsma. **2007**. Performance evaluation of the ECAT HRRT: an LSO-LYSO double layer high resolution, high sensitivity scanner. *Phys Med Biol* 52(5):1505-1526.
51. De Lean A., J.M. Stadel and R.J. Lefkowitz. **1980**. A ternary complex model explains the agonist-specific binding properties of the adenylate cyclase-coupled beta-adrenergic receptor. *J Biol Chem* 255(15):7108-7117.

52. De Lean A., B.F. Kilpatrick and M.G. Caron. **1982**. Dopamine receptor of the porcine anterior pituitary gland. Evidence for two affinity states discriminated by both agonists and antagonists. *Mol Pharmacol* 22(2):290-297.
53. Defagot M.C. and M.C. Antonelli. **1997**. Autoradiographic localization of the putative D4 dopamine receptor in rat brain. *Neurochem Res* 22(4):401-407.
54. DeJesus O.T., G.J. Van Moffaert and A.M. Friedman. **1987**. Synthesis of [¹¹C]SCH 23390 for dopamine D1 receptor studies. *Int J Rad Appl Instrum A* 38(5):345-348.
55. Deleu D., Y. Hanssens and M.G. Northway. **2004**. Subcutaneous apomorphine : an evidence-based review of its use in Parkinson's disease. *Drugs Aging* 21(11):687-709.
56. Derry C., C. Benjamin, P. Bladin, D. le Bars, H. Tochon-Danguy, S.F. Berkovic, L. Zimmer, N. Costes, R. Mulligan and D. Reutens. **2006**. Increased serotonin receptor availability in human sleep: evidence from an [¹⁸F]MPPF PET study in narcolepsy. *Neuroimage* 30(2):341-348.
57. Dewey S.L., G.S. Smith, J. Logan, J.D. Brodie, J.S. Fowler and A.P. Wolf. **1993**. Striatal binding of the PET ligand 11C-raclopride is altered by drugs that modify synaptic dopamine levels. *Synapse* 13(4):350-356.
58. Dumartin B., I. Caille, F. Gonon and B. Bloch. **1998**. Internalization of D1 dopamine receptor in striatal neurons in vivo as evidence of activation by dopamine agonists. *J Neurosci* 18(5):1650-1661.
59. Egerton A., M.A. Mehta, A.J. Montgomery, J.M. Lappin, O.D. Howes, S.J. Reeves, V.J. Cunningham and P.M. Grasby. **2009**. The dopaminergic basis of human behaviors: A review of molecular imaging studies. *Neurosci Biobehav Rev* 33(7):1109-1132.
60. Egerton A., E. Hirani, R. Ahmad, D.R. Turton, D. Brickute, L. Rosso, O.D. Howes, S.K. Luthra and P.M. Grasby. **2010**. Further evaluation of the carbon11-labeled D(2/3) agonist PET radiotracer PHNO: reproducibility in tracer characteristics and characterization of extrastriatal binding. *Synapse* 64(4):301-312.
61. Ehrin E., L. Farde, T. de Paulis, L. Eriksson, T. Greitz, P. Johnstrom, J.E. Litton, J.L. Nilsson, G. Sedvall, S. Stone-Elander and et al. **1985**. Preparation of 11C-labelled Raclopride, a new potent dopamine receptor antagonist: preliminary PET studies of cerebral dopamine receptors in the monkey. *Int J Appl Radiat Isot* 36(4):269-273.
62. Endres C.J., B.S. Kolachana, R.C. Saunders, T. Su, D. Weinberger, A. Breier, W.C. Eckelman and R.E. Carson. **1997**. Kinetic modeling of [¹¹C]raclopride: combined PET-microdialysis studies. *J Cereb Blood Flow Metab* 17(9):932-942.
63. Endres C.J., S. Swaminathan, O.T. DeJesus, M. Sievert, A.E. Ruoho, D. Murali, S.G. Rommelfanger and J.E. Holden. **1997**. Affinities of dopamine analogs for monoamine granular and plasma membrane transporters: implications for PET dopamine studies. *Life Sci* 60(26):2399-2406.
64. Endres C.J. and R.E. Carson. **1998**. Assessment of dynamic neurotransmitter changes with bolus or infusion delivery of neuroreceptor ligands. *J Cereb Blood Flow Metab* 18(11):1196-1210.
65. Engel G., M. Gothert, D. Hoyer, E. Schlicker and K. Hillenbrand. **1986**. Identity of inhibitory presynaptic 5-hydroxytryptamine (5-HT) autoreceptors in the rat brain cortex with 5-HT1B binding sites. *Naunyn Schmiedebergs Arch Pharmacol* 332(1):1-7.
66. Eriksson L., M. Dahlbom and L. Widen. **1990**. Positron emission tomography-- a new technique for studies of the central nervous system. *J Microsc* 157(Pt 3):305-333.
67. Ettrup A., M. Palner, N. Gillings, M.A. Santini, M. Hansen, B.R. Kornum, L.K. Rasmussen, K. Nagren, J. Madsen, M. Begtrup and G.M. Knudsen. **2010**. Radiosynthesis and Evaluation of 11C-CIMBI-5 as a 5-HT2A Receptor Agonist Radioligand for PET. *J Nucl Med* 51(11):1763-1770.
68. Farde L., E. Ehrin, L. Eriksson, T. Greitz, H. Hall, C.G. Hedstrom, J.E. Litton and G. Sedvall. **1985**. Substituted benzamides as ligands for visualization of

- dopamine receptor binding in the human brain by positron emission tomography. *Proc Natl Acad Sci U S A* 82(11):3863-3867.
69. Farde L., H. Hall, E. Ehrin and G. Sedvall. **1986**. Quantitative analysis of D2 dopamine receptor binding in the living human brain by PET. *Science* 231(4735):258-261.
 70. Farde L., C. Halldin, S. Stone-Elander and G. Sedvall. **1987**. PET analysis of human dopamine receptor subtypes using 11C-SCH 23390 and 11C-raclopride. *Psychopharmacology (Berl)* 92(3):278-284.
 71. Farde L., F.A. Wiesel, H. Hall, C. Halldin, S. Stone-Elander and G. Sedvall. **1987**. No D2 receptor increase in PET study of schizophrenia. *Arch Gen Psychiatry* 44(7):671-672.
 72. Farde L., L. Eriksson, G. Blomquist and C. Halldin. **1989**. Kinetic analysis of central [11C]raclopride binding to D2-dopamine receptors studied by PET--a comparison to the equilibrium analysis. *J Cereb Blood Flow Metab* 9(5):696-708.
 73. Farde L. and H. Hall. **1992**. Positron emission tomography--examination of chemical transmission in the living human brain. Development of radioligands. *Arzneimittelforschung* 42(2A):260-264.
 74. Farde L., A.L. Nordstrom, F.A. Wiesel, S. Pauli, C. Halldin and G. Sedvall. **1992**. Positron emission tomographic analysis of central D1 and D2 dopamine receptor occupancy in patients treated with classical neuroleptics and clozapine. Relation to extrapyramidal side effects. *Arch Gen Psychiatry* 49(7):538-544.
 75. Farde L. and A.L. Nordstrom. **1993**. PET examination of central D2 dopamine receptor occupancy in relation to extrapyramidal syndromes in patients being treated with neuroleptic drugs. *Psychopharmacol Ser* 10:94-100.
 76. Farde L., T. Suhara, S. Nyberg, P. Karlsson, Y. Nakashima, J. Hietala and C. Halldin. **1997**. A PET-study of [11C]FLB 457 binding to extrastriatal D2-dopamine receptors in healthy subjects and antipsychotic drug-treated patients. *Psychopharmacology (Berl)* 133(4):396-404.
 77. Fatemi-Ardekani A., N. Samavati, J. Tang and M.V. Kamath. **2009**. Advances in multimodality imaging through a hybrid PET/MRI system. *Crit Rev Biomed Eng* 37(6):495-515.
 78. Ferguson S.S., W.E. Downey, 3rd, A.M. Colapietro, L.S. Barak, L. Menard and M.G. Caron. **1996**. Role of beta-arrestin in mediating agonist-promoted G protein-coupled receptor internalization. *Science* 271(5247):363-366.
 79. Ferguson S.S. **2001**. Evolving concepts in G protein-coupled receptor endocytosis: the role in receptor desensitization and signaling. *Pharmacol Rev* 53(1):1-24.
 80. Fink K.B. and M. Gothert. **2007**. 5-HT receptor regulation of neurotransmitter release. *Pharmacol Rev* 59(4):360-417.
 81. Flower D.R. **1999**. Modelling G-protein-coupled receptors for drug design. *Biochim Biophys Acta* 1422(3):207-234.
 82. Forsberg M., M. Lehtonen, M. Heikkinen, J. Savolainen, T. Jarvinen and P.T. Mannisto. **2003**. Pharmacokinetics and pharmacodynamics of entacapone and tolcapone after acute and repeated administration: a comparative study in the rat. *J Pharmacol Exp Ther* 304(2):498-506.
 83. Frankle W.G., N.S. Mason, E.A. Rabiner, K. Ridler, M.A. May, D. Asmonga, C.M. Chen, S. Kendro, T.B. Cooper, C.A. Mathis and R. Narendran. **2010**. No effect of dopamine depletion on the binding of the high-affinity D 2/3 radiotracer [11C]FLB 457 in the human cortex. *Synapse* 64(12):879-885.
 84. Friedman A.M., O.T. DeJesus, J. Revenaugh and R.J. Dinerstein. **1984**. Measurements in vivo of parameters of the dopamine system. *Ann Neurol* 15 Suppl:S66-76.
 85. Galineau L., A.A. Wilson, A. Garcia, S. Houle, S. Kapur and N. Ginovart. **2006**. In vivo characterization of the pharmacokinetics and pharmacological properties of [11C]-(+)-PHNO in rats using an intracerebral beta-sensitive system. *Synapse* 60(2):172-183.
 86. Gallezot J.D., N. Nabulsi, A. Neumeister, B. Planeta-Wilson, W.A. Williams, T. Singhal, S. Kim, R.P. Maguire, T. McCarthy, J.J. Frost, Y. Huang, Y.S. Ding

- and R.E. Carson. **2010**. Kinetic modeling of the serotonin 5-HT_{1B} receptor radioligand [(11C)P943] in humans. *J Cereb Blood Flow Metab* 30(1):196-210.
87. Gao Y.G., R.J. Baldessarini, N.S. Kula and J.L. Neumeyer. **1990**. Synthesis and dopamine receptor affinities of enantiomers of 2-substituted apomorphines and their N-n-propyl analogues. *J Med Chem* 33(6):1800-1805.
 88. Garattini S., T. Mennini, C. Bendotti, R. Invernizzi and R. Samanin. **1986**. Neurochemical mechanism of action of drugs which modify feeding via the serotonergic system. *Appetite* 7 Suppl:15-38.
 89. Garcia-Arguello S.F., E. Arstad, D. Brickute, S.K. Luthra, D.R. Turton, M. Glaser, R. Fortt and E.G. Robins. **2010**. Automated synthesis of [11C]-(+)-PHNO from [11C]methyl iodide. *Neuroimage* 52:S218-S218.
 90. Garcia-Cabezas M.A., B. Rico, M.A. Sanchez-Gonzalez and C. Cavada. **2007**. Distribution of the dopamine innervation in the macaque and human thalamus. *Neuroimage* 34(3):965-984.
 91. Garcia-Cabezas M.A., P. Martinez-Sanchez, M.A. Sanchez-Gonzalez, M. Garzon and C. Cavada. **2009**. Dopamine innervation in the thalamus: monkey versus rat. *Cereb Cortex* 19(2):424-434.
 92. George S.R., M. Watanabe, T. Di Paolo, P. Falardeau, F. Labrie and P. Seeman. **1985**. The functional state of the dopamine receptor in the anterior pituitary is in the high affinity form. *Endocrinology* 117(2):690-697.
 93. Gether U. and B.K. Kobilka. **1998**. G protein-coupled receptors. II. Mechanism of agonist activation. *J Biol Chem* 273(29):17979-17982.
 94. Ghosh D., S.E. Snyder, V.J. Watts, R.B. Mailman and D.E. Nichols. **1996**. 9-Dihydroxy-2,3,7,11b-tetrahydro-1H-naph[1,2,3-de]isoquinoline: a potent full dopamine D₁ agonist containing a rigid-beta-phenyldopamine pharmacophore. *J Med Chem* 39(2):549-555.
 95. Gifford A.N., S.J. Gatley and C.R. Ashby, Jr. **1996**. Endogenously released dopamine inhibits the binding of dopaminergic PET and SPECT ligands in superfused rat striatal slices. *Synapse* 22(3):232-238.
 96. Ginovart N., L. Farde, C. Halldin and C.G. Swahn. **1999**. Changes in striatal D₂-receptor density following chronic treatment with amphetamine as assessed with PET in nonhuman primates. *Synapse* 31(2):154-162.
 97. Ginovart N., A.A. Wilson, J.H. Meyer, D. Hussey and S. Houle. **2003**. [11C]-DASB, a tool for in vivo measurement of SSRI-induced occupancy of the serotonin transporter: PET characterization and evaluation in cats. *Synapse* 47(2):123-133.
 98. Ginovart N., A.A. Wilson, S. Houle and S. Kapur. **2004**. Amphetamine pretreatment induces a change in both D₂-Receptor density and apparent affinity: a [11C]raclopride positron emission tomography study in cats. *Biol Psychiatry* 55(12):1188-1194.
 99. Ginovart N. **2005**. Imaging the dopamine system with in vivo [11C]raclopride displacement studies: understanding the true mechanism. *Mol Imaging Biol* 7(1):45-52.
 100. Ginovart N., L. Galineau, M. Willeit, R. Mizrahi, P.M. Bloomfield, P. Seeman, S. Houle, S. Kapur and A.A. Wilson. **2006**. Binding characteristics and sensitivity to endogenous dopamine of [11C]-(+)-PHNO, a new agonist radiotracer for imaging the high-affinity state of D₂ receptors in vivo using positron emission tomography. *J Neurochem* 97(4):1089-1103.
 101. Ginovart N., M. Willeit, P. Rusjan, A. Graff, P.M. Bloomfield, S. Houle, S. Kapur and A.A. Wilson. **2007**. Positron emission tomography quantification of [11C]-(+)-PHNO binding in the human brain. *J Cereb Blood Flow Metab* 27(4):857-871.
 102. Giovacchini G., L. Lang, Y. Ma, P. Herscovitch, W.C. Eckelman and R.E. Carson. **2005**. Differential effects of paroxetine on raphe and cortical 5-HT_{1A} binding: a PET study in monkeys. *Neuroimage* 28(1):238-248.
 103. Giros B., P. Sokoloff, M.P. Martres, J.F. Riou, L.J. Emorine and J.C. Schwartz. **1989**. Alternative splicing directs the expression of two D₂ dopamine receptor isoforms. *Nature* 342(6252):923-926.
 104. Gothert M., E. Schlicker, K. Fink and K. Classen. **1987**. Effects of RU 24969 on serotonin release in rat brain cortex: further support for the identity of

- serotonin autoreceptors with 5-HT_{1B} sites. *Arch Int Pharmacodyn Ther* 288(1):31-42.
105. Graff-Guerrero A., M. Willeit, N. Ginovart, D. Mamo, R. Mizrahi, P. Rusjan, I. Vitcu, P. Seeman, A.A. Wilson and S. Kapur. **2008**. Brain region binding of the D_{2/3} agonist [¹¹C]-(+)-PHNO and the D_{2/3} antagonist [¹¹C]raclopride in healthy humans. *Hum Brain Mapp* 29(4):400-410.
 106. Graff-Guerrero A., R. Mizrahi, O. Agid, H. Marcon, P. Barsoum, P. Rusjan, A.A. Wilson, R. Zipursky and S. Kapur. **2009**. The dopamine D₂ receptors in high-affinity state and D₃ receptors in schizophrenia: a clinical [¹¹C]-(+)-PHNO PET study. *Neuropsychopharmacology* 34(4):1078-1086.
 107. Graff-Guerrero A., L. Redden, W. Abi-Saab, D.A. Katz, S. Houle, P. Barsoum, A. Bhathena, R. Palaparthi, M.D. Saltarelli and S. Kapur. **2010**. Blockade of [¹¹C](+)-PHNO binding in human subjects by the dopamine D₃ receptor antagonist ABT-925. *Int J Neuropsychopharmacol* 13(3):273-287.
 108. Granas C., J. Nordquist, N. Mohell and D. Larhammar. **2001**. Site-directed mutagenesis of the 5-HT_{1B} receptor increases the affinity of 5-HT for the agonist low-affinity conformation and reduces the intrinsic activity of 5-HT. *Eur J Pharmacol* 421(2):69-76.
 109. Greenwald E., J.N. DaSilva, C. Valente, A.A. Wilson and S. Houle. **1998**. In vivo evaluation of R/S-[C-11]SKF 82957 in rats as a D-1 agonist tracer for PET: effect of endogenous dopamine and chronic D-1 and D-2 antagonists. *J Nucl Med* 39(5):236P.
 110. Gunther I., M. Psylla, G.N. Reddy, A. Antonini, P. Vontobel, H.W. Reist, A. Zollinger, R.J. Nickles, H.F. Beer, P.A. Schubiger and et al. **1995**. Positron emission tomography in drug evaluation: influence of three different catechol-O-methyltransferase inhibitors on metabolism of [NCA] 6-[¹⁸F]fluoro-L-dopa in rhesus monkey. *Nucl Med Biol* 22(7):921-927.
 111. Guo N., W. Guo, M. Kralikova, M. Jiang, I. Schieren, R. Narendran, M. Slifstein, A. Abi-Dargham, M. Laruelle, J.A. Javitch and S. Rayport. **2010**. Impact of D₂ receptor internalization on binding affinity of neuroimaging radiotracers. *Neuropsychopharmacology* 35(3):806-817.
 112. Hall D.A. and P.G. Strange. **1999**. Comparison of the ability of dopamine receptor agonists to inhibit forskolin-stimulated adenosine 3'5'-cyclic monophosphate (cAMP) accumulation via D_{2L} (long isoform) and D₃ receptors expressed in Chinese hamster ovary (CHO) cells. *Biochem Pharmacol* 58(2):285-289.
 113. Hall H., L. Farde and G. Sedvall. **1988**. Human dopamine receptor subtypes--in vitro binding analysis using 3H-SCH 23390 and 3H-raclopride. *J Neural Transm* 73(1):7-21.
 114. Hall H., G. Sedvall, O. Magnusson, J. Kopp, C. Halldin and L. Farde. **1994**. Distribution of D₁- and D₂-dopamine receptors, and dopamine and its metabolites in the human brain. *Neuropsychopharmacology* 11(4):245-256.
 115. Hall H., L. Farde, C. Halldin, Y.L. Hurd, S. Pauli and G. Sedvall. **1996**. Autoradiographic localization of extrastriatal D₂-dopamine receptors in the human brain using [¹²⁵I]epidepride. *Synapse* 23(2):115-123.
 116. Hall H., C. Halldin, D. Dijkstra, H. Wikstrom, L.D. Wise, T.A. Pugsley, P. Sokoloff, S. Pauli, L. Farde and G. Sedvall. **1996**. Autoradiographic localisation of D₃-dopamine receptors in the human brain using the selective D₃-dopamine receptor agonist (+)-[³H]PD 128907. *Psychopharmacology (Berl)* 128(3):240-247.
 117. Halldin C., S. Stone-Elander, L. Farde, E. Ehrin, K.J. Fasth, B. Langstrom and G. Sedvall. **1986**. Preparation of ¹¹C-labelled SCH 23390 for the in vivo study of dopamine D-1 receptors using positron emission tomography. *Int J Rad Appl Instrum A* 37(10):1039-1043.
 118. Halldin C., L. Farde, T. Hogberg, N. Mohell, H. Hall, T. Suhara, P. Karlsson, Y. Nakashima and C.G. Swahn. **1995**. Carbon-11-FLB 457: a radioligand for extrastriatal D₂ dopamine receptors. *J Nucl Med* 36(7):1275-1281.
 119. Halldin C., C.-G. Swahn, L. Farde and G. Sedvall. **1995**. Radioligand disposition and metabolism - Key information in early drug development. In:

- Comar D, editor. PET for Drug Development and Evaluation. Dordrecht: Kluwer Academic Publishers. p. 55-65.
120. Halldin C., C. Foged, Y.H. Chou, P. Karlsson, C.G. Swahn, J. Sandell, G. Sedvall and L. Farde. **1998**. Carbon-11-NNC 112: a radioligand for PET examination of striatal and neocortical D1-dopamine receptors. *J Nucl Med* 39(12):2061-2068.
 121. Halldin C., B. Gulyas and L. Farde. **2001**. PET studies with carbon-11 radioligands in neuropsychopharmacological drug development. *Curr Pharm Des* 7(18):1907-1929.
 122. Halldin C., B. Gulyas, O. Langer and L. Farde. **2001**. Brain radioligands--state of the art and new trends. *Q J Nucl Med* 45(2):139-152.
 123. Hartig P.R., D. Hoyer, P.P. Humphrey and G.R. Martin. **1996**. Alignment of receptor nomenclature with the human genome: classification of 5-HT_{1B} and 5-HT_{1D} receptor subtypes. *Trends Pharmacol Sci* 17(3):103-105.
 124. Hartvig P., K.J. Lindner, J. Tedroff, P. Bjurling, K. Hornfelt and B. Langstrom. **1992**. Regional brain kinetics of 6-fluoro-(beta-¹¹C)-L-dopa and (beta-¹¹C)-L-dopa following COMT inhibition. A study in vivo using positron emission tomography. *J Neural Transm Gen Sect* 87(1):15-22.
 125. Hartvig P., R. Torstenson, J. Tedroff, Y. Watanabe, K.J. Fasth, P. Bjurling and B. Langstrom. **1997**. Amphetamine effects on dopamine release and synthesis rate studied in the Rhesus monkey brain by positron emission tomography. *J Neural Transm* 104(4-5):329-339.
 126. Hassoun W., M. Le Cavorsin, N. Ginovart, L. Zimmer, V. Gualda, F. Bonnefoi and V. Leviel. **2003**. PET study of the [¹¹C]raclopride binding in the striatum of the awake cat: effects of anaesthetics and role of cerebral blood flow. *Eur J Nucl Med Mol Imaging* 30(1):141-148.
 127. Heal D.J., S.C. Cheetham and S.L. Smith. **2009**. The neuropharmacology of ADHD drugs in vivo: insights on efficacy and safety. *Neuropharmacology* 57(7-8):608-618.
 128. Hersch S.M., B.J. Ciliax, C.A. Gutekunst, H.D. Rees, C.J. Heilman, K.K. Yung, J.P. Bolam, E. Ince, H. Yi and A.I. Levey. **1995**. Electron microscopic analysis of D1 and D2 dopamine receptor proteins in the dorsal striatum and their synaptic relationships with motor corticostriatal afferents. *J Neurosci* 15(7 Pt 2):5222-5237.
 129. Hirani E., J. Opacka-Juffry, R. Gunn, I. Khan, T. Sharp and S. Hume. **2000**. Pindolol occupancy of 5-HT_{1A} receptors measured in vivo using small animal positron emission tomography with carbon-11 labeled WAY 100635. *Synapse* 36(4):330-341.
 130. Hirani E., T. Sharp, M. Sprakes, P. Grasby and S. Hume. **2003**. Fenfluramine evokes 5-HT_{2A} receptor-mediated responses but does not displace [¹¹C]MDL 100907: small animal PET and gene expression studies. *Synapse* 50(3):251-260.
 131. Houston G.C., S.P. Hume, E. Hirani, J.L. Goggi and P.M. Grasby. **2004**. Temporal characterisation of amphetamine-induced dopamine release assessed with [¹¹C]raclopride in anaesthetised rodents. *Synapse* 51(3):206-212.
 132. Hoyer D. and G. Martin. **1997**. 5-HT receptor classification and nomenclature: towards a harmonization with the human genome. *Neuropharmacology* 36(4-5):419-428.
 133. <http://www.cnsforum.com/educationalresources/imagebank/>.
 134. Hu J., S. Henry, J.D. Gallezot, J. Ropchan, J.F. Neumaier, M.N. Potenza, R. Sinha, J.H. Krystal, Y. Huang, Y.S. Ding, R.E. Carson and A. Neumeister. **2010**. Serotonin 1B receptor imaging in alcohol dependence. *Biol Psychiatry* 67(9):800-803.
 135. Hume S., E. Hirani, J. Opacka-Juffry, R. Myers, C. Townsend, V. Pike and P. Grasby. **2001**. Effect of 5-HT on binding of [¹¹C] WAY 100635 to 5-HT_{1A} receptors in rat brain, assessed using in vivo microdialysis and PET after fenfluramine. *Synapse* 41(2):150-159.
 136. Hwang D.R., L.S. Kegeles and M. Laruelle. **2000**. (-)-N-[¹¹C]propyl-norapomorphine: a positron-labeled dopamine agonist for PET imaging of D₂ receptors. *Nucl Med Biol* 27(6):533-539.

137. Hwang D.R., R. Narendran and M. Laruelle. **2005**. Positron-labeled dopamine agonists for probing the high affinity states of dopamine subtype 2 receptors. *Bioconjug Chem* 16(1):27-31.
138. Ichise M., J.S. Liow, J.Q. Lu, A. Takano, K. Model, H. Toyama, T. Suhara, K. Suzuki, R.B. Innis and R.E. Carson. **2003**. Linearized reference tissue parametric imaging methods: application to [¹¹C]DASB positron emission tomography studies of the serotonin transporter in human brain. *J Cereb Blood Flow Metab* 23(9):1096-1112.
139. Innis R.B., R.T. Malison, M. al-Tikriti, P.B. Hoffer, E.H. Sybirska, J.P. Seibyl, S.S. Zoghbi, R.M. Baldwin, M. Laruelle, E.O. Smith and et al. **1992**. Amphetamine-stimulated dopamine release competes in vivo for [¹²³I]IBZM binding to the D2 receptor in nonhuman primates. *Synapse* 10(3):177-184.
140. Innis R.B., V.J. Cunningham, J. Delforge, M. Fujita, A. Gjedde, R.N. Gunn, J. Holden, S. Houle, S.C. Huang, M. Ichise, H. Iida, H. Ito, Y. Kimura, R.A. Koeppe, G.M. Knudsen, J. Knuuti, A.A. Lammertsma, M. Laruelle, J. Logan, R.P. Maguire, M.A. Mintun, E.D. Morris, R. Parsey, J.C. Price, M. Slifstein, V. Sossi, T. Suhara, J.R. Votaw, D.F. Wong and R.E. Carson. **2007**. Consensus nomenclature for in vivo imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab* 27(9):1533-1539.
141. Ito H., J. Hietala, G. Blomqvist, C. Halldin and L. Farde. **1998**. Comparison of the transient equilibrium and continuous infusion method for quantitative PET analysis of [¹¹C]raclopride binding. *J Cereb Blood Flow Metab* 18(9):941-950.
142. Jagoda E.M., L. Lang, J. Tokugawa, A. Simmons, Y. Ma, C. Contoreggi, D. Kiesewetter and W.C. Eckelman. **2006**. Development of 5-HT_{1A} receptor radioligands to determine receptor density and changes in endogenous 5-HT. *Synapse* 59(6):330-341.
143. Jordan S., R. Chen, R. Fernalld, J. Johnson, K. Regardie, J. Kambayashi, Y. Tadori, H. Kitagawa and T. Kikuchi. **2006**. In vitro biochemical evidence that the psychotomimetics phencyclidine, ketamine and dizocilpine (MK-801) are inactive at cloned human and rat dopamine D₂ receptors. *Eur J Pharmacol* 540(1-3):53-56.
144. Jorga K., B. Fotteler, G. Sedek, T. Nielsen and J. Aitken. **1998**. The effect of tolcapone on levodopa pharmacokinetics is independent of levodopa/carbidopa formulation. *J Neurol* 245(4):223-230.
145. Jorga K.M., B. Fotteler, P. Heizmann and G. Zurcher. **1998**. Pharmacokinetics and pharmacodynamics after oral and intravenous administration of tolcapone, a novel adjunct to Parkinson's disease therapy. *Eur J Clin Pharmacol* 54(5):443-447.
146. Kaakkola S., A. Gordin and P.T. Mannisto. **1994**. General properties and clinical possibilities of new selective inhibitors of catechol O-methyltransferase. *Gen Pharmacol* 25(5):813-824.
147. Kalbitzer J., C. Svarer, V.G. Frokjaer, D. Erritzoe, W.F. Baare, J. Madsen, S.G. Hasselbalch and G.M. Knudsen. **2009**. A probabilistic approach to delineating functional brain regions. *J Nucl Med Technol* 37(2):91-95.
148. Kankaanpaa A., E. Meririnne, P. Lillsunde and T. Seppala. **1998**. The acute effects of amphetamine derivatives on extracellular serotonin and dopamine levels in rat nucleus accumbens. *Pharmacol Biochem Behav* 59(4):1003-1009.
149. Kannan P., C. John, S.S. Zoghbi, C. Halldin, M.M. Gottesman, R.B. Innis and M.D. Hall. **2009**. Imaging the function of P-glycoprotein with radiotracers: pharmacokinetics and in vivo applications. *Clin Pharmacol Ther* 86(4):368-377.
150. Kapur S. and P. Seeman. **2002**. NMDA receptor antagonists ketamine and PCP have direct effects on the dopamine D₂ and serotonin 5-HT₂ receptors-implications for models of schizophrenia. *Mol Psychiatry* 7(8):837-844.
151. Karlsson P., L. Farde, C. Halldin, C.G. Swahn, G. Sedvall, C. Foged, K.T. Hansen and B. Skrumager. **1993**. PET examination of [¹¹C]NNC 687 and [¹¹C]NNC 756 as new radioligands for the D₁-dopamine receptor. *Psychopharmacology (Berl)* 113(2):149-156.
152. Keibarian J.W. and D.B. Calne. **1979**. Multiple receptors for dopamine. *Nature* 277(5692):93-96.

153. Kegeles L.S., A. Abi-Dargham, Y. Zea-Ponce, J. Rodenhiser-Hill, J.J. Mann, R.L. Van Heertum, T.B. Cooper, A. Carlsson and M. Laruelle. **2000**. Modulation of amphetamine-induced striatal dopamine release by ketamine in humans: implications for schizophrenia. *Biol Psychiatry* 48(7):627-640.
154. Kegeles L.S., D. Martinez, L.D. Kochan, D.R. Hwang, Y. Huang, O. Mawlawi, R.F. Suckow, R.L. Van Heertum and M. Laruelle. **2002**. NMDA antagonist effects on striatal dopamine release: positron emission tomography studies in humans. *Synapse* 43(1):19-29.
155. Kenakin T. **1995**. Agonist-receptor efficacy. II. Agonist trafficking of receptor signals. *Trends Pharmacol Sci* 16(7):232-238.
156. Kenakin T. **1997**. Agonist-specific receptor conformations. *Trends Pharmacol Sci* 18(11):416-417.
157. Kennett G.A., C.T. Dourish and G. Curzon. **1987**. 5-HT_{1B} agonists induce anorexia at a postsynaptic site. *Eur J Pharmacol* 141(3):429-435.
158. Kessler R.M., W.O. Whetsell, M.S. Ansari, J.R. Votaw, T. de Paulis, J.A. Clanton, D.E. Schmidt, N.S. Mason and R.G. Manning. **1993**. Identification of extrastriatal dopamine D₂ receptors in post mortem human brain with [¹²⁵I]epidepride. *Brain Res* 609(1-2):237-243.
159. Khan Z.U., A. Gutierrez, R. Martin, A. Penafiel, A. Rivera and A. de la Calle. **2000**. Dopamine D₅ receptors of rat and human brain. *Neuroscience* 100(4):689-699.
160. Klunk W.E., H. Engler, A. Nordberg, Y. Wang, G. Blomqvist, D.P. Holt, M. Bergstrom, I. Savitcheva, G.F. Huang, S. Estrada, B. Ausen, M.L. Debnath, J. Barletta, J.C. Price, J. Sandell, B.J. Lopresti, A. Wall, P. Koivisto, G. Antoni, C.A. Mathis and B. Langstrom. **2004**. Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol* 55(3):306-319.
161. Kobilka B.K. and X. Deupi. **2007**. Conformational complexity of G-protein-coupled receptors. *Trends Pharmacol Sci* 28(8):397-406.
162. Kodaka F., H. Ito, H. Takano, H. Takahashi, R. Arakawa, M. Miyoshi, M. Okumura, T. Otsuka, K. Nakayama, C. Halldin, L. Farde and T. Suhara. **2010**. Effect of risperidone on high-affinity state of dopamine D₂ receptors: a PET study with agonist ligand [¹¹C](R)-2-CH₃O-N-n-propylnorapomorphine. *Int J Neuropsychopharmacol*:1-7.
163. Koepp M.J., R.N. Gunn, A.D. Lawrence, V.J. Cunningham, A. Dagher, T. Jones, D.J. Brooks, C.J. Bench and P.M. Grasby. **1998**. Evidence for striatal dopamine release during a video game. *Nature* 393(6682):266-268.
164. Kortekaas R., R.P. Maguire, T.I. Cremers, D. Dijkstra, A. van Waarde and K.L. Leenders. **2004**. In vivo binding behavior of dopamine receptor agonist (+)-PD 128907 and implications for the "ceiling effect" in endogenous competition studies with [(11)C]raclopride—a positron emission tomography study in Macaca mulatta. *J Cereb Blood Flow Metab* 24(5):531-535.
165. Kumar J.S. and J.J. Mann. **2007**. PET tracers for 5-HT_{1A} receptors and uses thereof. *Drug Discov Today* 12(17-18):748-756.
166. Kumar J.S., J. Prabhakaran, V.J. Majo, M.S. Milak, S.C. Hsiung, H. Tamir, N.R. Simpson, R.L. Van Heertum, J.J. Mann and R.V. Parsey. **2007**. Synthesis and in vivo evaluation of a novel 5-HT_{1A} receptor agonist radioligand [O-methyl-¹¹C]2-(4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl)-4-methyl-1,2,4-triazine-3,5(2H,4H)dione in nonhuman primates. *Eur J Nucl Med Mol Imaging* 34(7):1050-1060.
167. Lacivita E., P. De Giorgio, I.T. Lee, S.I. Rodeheaver, B.A. Weiss, C. Fracasso, S. Caccia, F. Berardi, R. Perrone, M.R. Zhang, J. Maeda, M. Higuchi, T. Suhara, J.A. Schetz and M. Leopoldo. **2010**. Design, synthesis, radiolabeling, and in vivo evaluation of carbon-11 labeled N-[2-[4-(3-cyanopyridin-2-yl)piperazin-1-yl]ethyl]-3-methoxybenzamide, a potential positron emission tomography tracer for the dopamine D₄ receptors. *J Med Chem* 53(20):7344-7355.
168. Lammertsma A.A. and S.P. Hume. **1996**. Simplified reference tissue model for PET receptor studies. *Neuroimage* 4(3 Pt 1):153-158.
169. Larisch R., A. Klimke, K. Hamacher, U. Henning, S. Estalji, T. Hohlfeld, H. Vosberg, M. Tosch, W. Gaebel, H.H. Coenen and H.W. Muller-Gartner. **2003**.

- Influence of synaptic serotonin level on [18F]altanserin binding to 5HT₂ receptors in man. *Behav Brain Res* 139(1-2):21-29.
170. Larsen P., J. Ulin, K. Dahlstrom and M. Jensen. **1997**. Synthesis of [¹¹C]Iodomethane by Iodination of [¹¹C]Methane. *Applied Radiation and Isotopes* 48(2):153-157.
 171. Laruelle M., A. Abi-Dargham, C.H. van Dyck, W. Rosenblatt, Y. Zea-Ponce, S.S. Zoghbi, R.M. Baldwin, D.S. Charney, P.B. Hoffer, H.F. Kung and et al. **1995**. SPECT imaging of striatal dopamine release after amphetamine challenge. *J Nucl Med* 36(7):1182-1190.
 172. Laruelle M., A. Abi-Dargham, C.H. van Dyck, R. Gil, C.D. D'Souza, J. Erdos, E. McCance, W. Rosenblatt, C. Fingado, S.S. Zoghbi, R.M. Baldwin, J.P. Seibyl, J.H. Krystal, D.S. Charney and R.B. Innis. **1996**. Single photon emission computerized tomography imaging of amphetamine-induced dopamine release in drug-free schizophrenic subjects. *Proc Natl Acad Sci U S A* 93(17):9235-9240.
 173. Laruelle M., R.N. Iyer, M.S. al-Tikriti, Y. Zea-Ponce, R. Malison, S.S. Zoghbi, R.M. Baldwin, H.F. Kung, D.S. Charney, P.B. Hoffer, R.B. Innis and C.W. Bradberry. **1997**. Microdialysis and SPECT measurements of amphetamine-induced dopamine release in nonhuman primates. *Synapse* 25(1):1-14.
 174. Laruelle M., A. Abi-Dargham, N. Simpson, L. Kegeles, R. Parsey, D.R. Hwang, Y. Zea-Ponce, I. Lombardo, R. Weiss, R. Van Heertum and J.J. Mann. **1998**. PET studies of binding competition between endogenous dopamine and D1 antagonists and agonists. *Soc Neurosci Abstr* 24:22.
 175. Laruelle M. **2000**. Imaging synaptic neurotransmission with in vivo binding competition techniques: a critical review. *J Cereb Blood Flow Metab* 20(3):423-451.
 176. Lassen N.A. **1992**. Neuroreceptor quantitation in vivo by the steady-state principle using constant infusion or bolus injection of radioactive tracers. *J Cereb Blood Flow Metab* 12(5):709-716.
 177. Laymon C.M., N.S. Mason, W.G. Frankle, J.P. Carney, B.J. Lopresti, M.Y. Litschge, C.A. Mathis, J.M. Mountz and R. Narendran. **2009**. Human biodistribution and dosimetry of the D2/3 agonist 11C-N-propylnorapomorphine (11C-NPA) determined from PET. *J Nucl Med* 50(5):814-817.
 178. Lee C.M. and L. Farde. **2006**. Using positron emission tomography to facilitate CNS drug development. *Trends Pharmacol Sci* 27(6):310-316.
 179. Lee M.D. and K.J. Simansky. **1997**. CP-94, 253: a selective serotonin1B (5-HT_{1B}) agonist that promotes satiety. *Psychopharmacology (Berl)* 131(3):264-270.
 180. Lees A.J. **2008**. Evidence-based efficacy comparison of tolcapone and entacapone as adjunctive therapy in Parkinson's disease. *CNS Neurosci Ther* 14(1):83-93.
 181. Lefkowitz R.J., S. Cotecchia, P. Samama and T. Costa. **1993**. Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins. *Trends Pharmacol Sci* 14(8):303-307.
 182. Leger G., A. Gjedde, H. Kuwabara, M. Guttman and P. Cumming. **1998**. Effect of catechol-O-methyltransferase inhibition on brain uptake of [18F]fluorodopa: implications for compartmental modelling and clinical usefulness. *Synapse* 30(4):351-361.
 183. Levey A.I., S.M. Hersch, D.B. Rye, R.K. Sunahara, H.B. Niznik, C.A. Kitt, D.L. Price, R. Maggio, M.R. Brann and B.J. Ciliax. **1993**. Localization of D1 and D2 dopamine receptors in brain with subtype-specific antibodies. *Proc Natl Acad Sci U S A* 90(19):8861-8865.
 184. Lidow M.S., F. Wang, Y. Cao and P.S. Goldman-Rakic. **1998**. Layer V neurons bear the majority of mRNAs encoding the five distinct dopamine receptor subtypes in the primate prefrontal cortex. *Synapse* 28(1):10-20.
 185. Lin D. and L.H. Parsons. **2002**. Anxiogenic-like effect of serotonin(1B) receptor stimulation in the rat elevated plus-maze. *Pharmacol Biochem Behav* 71(4):581-587.

186. Logan J., J.S. Fowler, N.D. Volkow, A.P. Wolf, S.L. Dewey, D.J. Schlyer, R.R. MacGregor, R. Hitzemann, B. Bendriem, S.J. Gatley and et al. **1990**. Graphical analysis of reversible radioligand binding from time-activity measurements applied to [N-11C-methyl]-(-)-cocaine PET studies in human subjects. *J Cereb Blood Flow Metab* 10(5):740-747.
187. Logan J., S.L. Dewey, A.P. Wolf, J.S. Fowler, J.D. Brodie, B. Angrist, N.D. Volkow and S.J. Gatley. **1991**. Effects of endogenous dopamine on measures of [18F]N-methylspiperidol binding in the basal ganglia: comparison of simulations and experimental results from PET studies in baboons. *Synapse* 9(3):195-207.
188. Lovenberg T.W., W.K. Brewster, D.M. Mottola, R.C. Lee, R.M. Riggs, D.E. Nichols, M.H. Lewis and R.B. Mailman. **1989**. Dihydroxydopamine, a novel selective high potency full dopamine D-1 receptor agonist. *Eur J Pharmacol* 166(1):111-113.
189. Lundberg J., C. Halldin and L. Farde. **2006**. Measurement of serotonin transporter binding with PET and [11C]MADAM: a test-retest reproducibility study. *Synapse* 60(3):256-263.
190. Lundquist P., H. Wilking, A.U. Hoglund, J. Sandell, M. Bergstrom, P. Hartvig and B. Langstrom. **2005**. Potential of [11C]DASB for measuring endogenous serotonin with PET: binding studies. *Nucl Med Biol* 32(2):129-136.
191. Lundquist P., M. Roman, S. Syvanen, P. Hartvig, G. Blomquist, M. Hammarlund-Udenaes and B. Langstrom. **2007**. Effect on [11C]DASB binding after tranylcypramine-induced increase in serotonin concentration: positron emission tomography studies in monkeys and rats. *Synapse* 61(6):440-449.
192. Luttrell L.M. and D. Gesty-Palmer. **2010**. Beyond desensitization: physiological relevance of arrestin-dependent signaling. *Pharmacol Rev* 62(2):305-330.
193. Maeda J., T. Suhara, M. Ogawa, T. Okauchi, K. Kawabe, M.R. Zhang, J. Semba and K. Suzuki. **2001**. In vivo binding properties of [carbonyl-11C]WAY-100635: effect of endogenous serotonin. *Synapse* 40(2):122-129.
194. Maier D.L., C. Sobotka-Briner, M. Ding, M.E. Powell, Q. Jiang, G. Hill, J.R. Heys, C.S. Elmore, M.E. Pierson and L. Mrzljak. **2009**. [N-methyl-3H3]AZ10419369 binding to the 5-HT1B receptor: in vitro characterization and in vivo receptor occupancy. *J Pharmacol Exp Ther* 330(1):342-351.
195. Mamelak M., S. Chiu and R.K. Mishra. **1993**. High- and low-affinity states of dopamine D1 receptors in schizophrenia. *Eur J Pharmacol* 233(1):175-176.
196. Mannisto P.T. and S. Kaakkola. **1999**. Catechol-O-methyltransferase (COMT): biochemistry, molecular biology, pharmacology, and clinical efficacy of the new selective COMT inhibitors. *Pharmacol Rev* 51(4):593-628.
197. Marner L., N. Gillings, R.A. Comley, W.F. Baare, E.A. Rabiner, A.A. Wilson, S. Houle, S.G. Hasselbalch, C. Svarer, R.N. Gunn, M. Laruelle and G.M. Knudsen. **2009**. Kinetic modeling of 11C-SB207145 binding to 5-HT4 receptors in the human brain in vivo. *J Nucl Med* 50(6):900-908.
198. Marner L., N. Gillings, K. Madsen, D. Erritzoe, W.F. Baare, C. Svarer, S.G. Hasselbalch and G.M. Knudsen. **2010**. Brain imaging of serotonin 4 receptors in humans with [11C]SB207145-PET. *Neuroimage* 50(3):855-861.
199. Matusch A., R. Hurlmann, E. Rota Kops, O.H. Winz, D. Elmenhorst, H. Herzog, K. Zilles and A. Bauer. **2007**. Acute S-ketamine application does not alter cerebral [18F]altanserin binding: a pilot PET study in humans. *J Neural Transm* 114(11):1433-1442.
200. Maura G., E. Roccatagliata and M. Raiteri. **1986**. Serotonin autoreceptor in rat hippocampus: pharmacological characterization as a subtype of the 5-HT1 receptor. *Naunyn Schmiedeberg's Arch Pharmacol* 334(4):323-326.
201. McCauley P.G., K.M. O'Boyle and J.L. Waddington. **1995**. Dopamine-induced reduction in the density of guanine nucleotide-sensitive D1 receptors in human postmortem brain in the absence of apparent D1: D2 interactions. *Neuropharmacology* 34(7):777-783.
202. McCormick P.N., S. Kapur, P. Seeman and A.A. Wilson. **2008**. Dopamine D2 receptor radiotracers [(11C)(+)-PHNO and [(3)H]raclopride are indistinguishably inhibited by D2 agonists and antagonists ex vivo. *Nucl Med Biol* 35(1):11-17.

203. McCormick P.N., S. Kapur, G. Reckless and A.A. Wilson. **2009**. Ex vivo [11C]-(+)-PHNO binding is unchanged in animal models displaying increased high-affinity states of the D2 receptor in vitro. *Synapse* 63(11):998-1009.
204. McCormick P.N., N. Ginovart and A.A. Wilson. **2010**. Isoflurane Anaesthesia Differentially Affects the Amphetamine Sensitivity of Agonist and Antagonist D2/D3 Positron Emission Tomography Radiotracers: Implications for In Vivo Imaging of Dopamine Release. *Mol Imaging Biol*.
205. Mennini T., A. Bizzi, S. Caccia, A. Codegoni, C. Fracasso, E. Frittoli, G. Guiso, I.M. Padura, C. Taddei, A. Uslenghi and et al. **1991**. Comparative studies on the anorectic activity of d-fenfluramine in mice, rats, and guinea pigs. *Naunyn Schmiedebergs Arch Pharmacol* 343(5):483-490.
206. Mercier D., P. Falardeau and D. Levesque. **2001**. Autoreceptor preference of dopamine D2 receptor agonists correlates with preferential coupling to cyclic AMP. *Neuroreport* 12(7):1473-1479.
207. Meyer J.H., R. Cho, S. Kennedy and S. Kapur. **1999**. The effects of single dose nefazodone and paroxetine upon 5-HT_{2A} binding potential in humans using [18F]-setoperone PET. *Psychopharmacology (Berl)* 144(3):279-281.
208. Milak M.S., R.T. Ogden, D.N. Vinocur, R.L. Van Heertum, T.B. Cooper, J.J. Mann and R.V. Parsey. **2005**. Effects of tryptophan depletion on the binding of [11C]-DASB to the serotonin transporter in baboons: response to acute serotonin deficiency. *Biol Psychiatry* 57(1):102-106.
209. Milak M.S., A.J. Severance, R.T. Ogden, J. Prabhakaran, J.S. Kumar, V.J. Majo, J.J. Mann and R.V. Parsey. **2008**. Modeling considerations for 11C-CUMI-101, an agonist radiotracer for imaging serotonin 1A receptor in vivo with PET. *J Nucl Med* 49(4):587-596.
210. Millan M.J., L. Maiofiss, D. Cussac, V. Audinot, J.A. Boutin and A. Newman-Tancredi. **2002**. Differential actions of antiparkinson agents at multiple classes of monoaminergic receptor. I. A multivariate analysis of the binding profiles of 14 drugs at 21 native and cloned human receptor subtypes. *J Pharmacol Exp Ther* 303(2):791-804.
211. Millan M.J., A. Newman-Tancredi, S. Lochon, M. Touzard, S. Aubry and V. Audinot. **2002**. Specific labelling of serotonin 5-HT_{1B} receptors in rat frontal cortex with the novel, phenylpiperazine derivative, [3H]GR125,743. A pharmacological characterization. *Pharmacol Biochem Behav* 71(4):589-598.
212. Mintun M.A., M.E. Raichle, M.R. Kilbourn, G.F. Wooten and M.J. Welch. **1984**. A quantitative model for the in vivo assessment of drug binding sites with positron emission tomography. *Ann Neurol* 15(3):217-227.
213. Mizrahi R., S. Houle, I. Vitcu, A. Ng and A.A. Wilson. **2010**. Side effects profile in humans of (11)C-(+)-PHNO, a dopamine D_{2/3} agonist ligand for PET. *J Nucl Med* 51(3):496-497.
214. Moghaddam B., C.W. Berridge, P.S. Goldman-Rakic, B.S. Bunney and R.H. Roth. **1993**. In vivo assessment of basal and drug-induced dopamine release in cortical and subcortical regions of the anesthetized primate. *Synapse* 13(3):215-222.
215. Molinoff P.B. and J. Axelrod. **1971**. Biochemistry of catecholamines. *Annu Rev Biochem* 40:465-500.
216. Morris E.D. and K.K. Yoder. **2007**. Positron emission tomography displacement sensitivity: predicting binding potential change for positron emission tomography tracers based on their kinetic characteristics. *J Cereb Blood Flow Metab* 27(3):606-617.
217. Mukaida K., T. Shichino, S. Koyanagi, S. Himukashi and K. Fukuda. **2007**. Activity of the serotonergic system during isoflurane anesthesia. *Anesth Analg* 104(4):836-839.
218. Mukherjee J., Z.Y. Yang, R. Lew, T. Brown, S. Kronmal, M.D. Cooper and L.S. Seiden. **1997**. Evaluation of d-amphetamine effects on the binding of dopamine D-2 receptor radioligand, 18F-fallypride in nonhuman primates using positron emission tomography. *Synapse* 27(1):1-13.
219. Mukherjee J., B.T. Christian, T.K. Narayanan, B. Shi and D. Collins. **2005**. Measurement of d-amphetamine-induced effects on the binding of dopamine D-

- 2/D-3 receptor radioligand, 18F-fallypride in extrastriatal brain regions in non-human primates using PET. *Brain Res* 1032(1-2):77-84.
220. Murrough J.W., S. Henry, J. Hu, J.D. Gallezot, B. Planeta-Wilson, J.F. Neumaier and A. Neumeister. **2010**. Reduced ventral striatal/ventral pallidal serotonin(1B) receptor binding potential in major depressive disorder. *Psychopharmacology (Berl)*.
221. Nabulsi N., Y. Huang, D. Weinzimmer, J. Ropchan, J.J. Frost, T. McCarthy, R.E. Carson and Y.S. Ding. **2010**. High-resolution imaging of brain 5-HT 1B receptors in the rhesus monkey using [11C]P943. *Nucl Med Biol* 37(2):205-214.
222. Nagren K., L. Muller, C. Halldin, C.G. Swahn and P. Lehtikoinen. **1995**. Improved synthesis of some commonly used PET radioligands by the use of [11C]methyl triflate. *Nucl Med Biol* 22(2):235-239.
223. Narendran R., D.R. Hwang, M. Slifstein, P.S. Talbot, D. Erritzoe, Y. Huang, T.B. Cooper, D. Martinez, L.S. Kegeles, A. Abi-Dargham and M. Laruelle. **2004**. In vivo vulnerability to competition by endogenous dopamine: comparison of the D2 receptor agonist radiotracer (-)-N-[11C]propyl-norapomorphine ([11C]NPA) with the D2 receptor antagonist radiotracer [11C]-raclopride. *Synapse* 52(3):188-208.
224. Narendran R., D.R. Hwang, M. Slifstein, Y. Hwang, Y. Huang, J. Ekelund, O. Guillin, E. Scher, D. Martinez and M. Laruelle. **2005**. Measurement of the proportion of D2 receptors configured in state of high affinity for agonists in vivo: a positron emission tomography study using [11C]N-propyl-norapomorphine and [11C]raclopride in baboons. *J Pharmacol Exp Ther* 315(1):80-90.
225. Narendran R., M. Slifstein, O. Guillin, Y. Hwang, D.R. Hwang, E. Scher, S. Reeder, E. Rabiner and M. Laruelle. **2006**. Dopamine (D2/3) receptor agonist positron emission tomography radiotracer [11C]-(+)-PHNO is a D3 receptor preferring agonist in vivo. *Synapse* 60(7):485-495.
226. Narendran R., M. Slifstein, D.R. Hwang, Y. Hwang, E. Scher, S. Reeder, D. Martinez and M. Laruelle. **2007**. Amphetamine-induced dopamine release: duration of action as assessed with the D2/3 receptor agonist radiotracer (-)-N-[(11)C]propyl-norapomorphine ([11C]NPA) in an anesthetized nonhuman primate. *Synapse* 61(2):106-109.
227. Narendran R., W.G. Frankle, N.S. Mason, C.M. Laymon, B.J. Lopresti, J.C. Price, S. Kendro, S. Vora, M. Litschge, J.M. Mountz and C.A. Mathis. **2009**. Positron emission tomography imaging of D(2/3) agonist binding in healthy human subjects with the radiotracer [(11)C]-N-propyl-norapomorphine: preliminary evaluation and reproducibility studies. *Synapse* 63(7):574-584.
228. Narendran R., W.G. Frankle, N.S. Mason, E.A. Rabiner, R.N. Gunn, G.E. Searle, S. Vora, M. Litschge, S. Kendro, T.B. Cooper, C.A. Mathis and M. Laruelle. **2009**. Positron emission tomography imaging of amphetamine-induced dopamine release in the human cortex: a comparative evaluation of the high affinity dopamine D2/3 radiotracers [11C]FLB 457 and [11C]fallypride. *Synapse* 63(6):447-461.
229. Narendran R., N.S. Mason, C.M. Laymon, B.J. Lopresti, N.D. Velasquez, M.A. May, S. Kendro, D. Martinez, C.A. Mathis and W.G. Frankle. **2010**. A comparative evaluation of the dopamine D(2/3) agonist radiotracer [11C](-)-N-propyl-norapomorphine and antagonist [11C]raclopride to measure amphetamine-induced dopamine release in the human striatum. *J Pharmacol Exp Ther* 333(2):533-539.
230. Nelson E.E. and J.T. Winslow. **2009**. Non-human primates: model animals for developmental psychopathology. *Neuropsychopharmacology* 34(1):90-105.
231. Neumeyer J.L., N.S. Kula, J. Bergman and R.J. Baldessarini. **2003**. Receptor affinities of dopamine D1 receptor-selective novel phenylbenzazepines. *Eur J Pharmacol* 474(2-3):137-140.
232. Nishino S. and M. Okuro. **2010**. Emerging treatments for narcolepsy and its related disorders. *Expert Opin Emerg Drugs* 15(1):139-158.
233. Nordberg A., J.O. Rinne, A. Kadir and B. Langstrom. **2010**. The use of PET in Alzheimer disease. *Nat Rev Neurol* 6(2):78-87.

234. Nordstrom A.L., L. Farde, S. Pauli, J.E. Litton and C. Halldin. **1992**. PET analysis of central [¹¹C]raclopride binding in healthy young adults and schizophrenic patients - reliability and age effects. *Psychopharmacology (Berl)* 7:157-165.
235. Nordstrom A.L., L. Farde, F.A. Wiesel, K. Forslund, S. Pauli, C. Halldin and G. Uppfeldt. **1993**. Central D2-dopamine receptor occupancy in relation to antipsychotic drug effects: a double-blind PET study of schizophrenic patients. *Biol Psychiatry* 33(4):227-235.
236. Nyberg S., L. Farde, L. Eriksson, C. Halldin and B. Eriksson. **1993**. 5-HT₂ and D₂ dopamine receptor occupancy in the living human brain. A PET study with risperidone. *Psychopharmacology (Berl)* 110(3):265-272.
237. Nyberg S., B. Eriksson, G. Oxenstierna, C. Halldin and L. Farde. **1999**. Suggested minimal effective dose of risperidone based on PET-measured D₂ and 5-HT_{2A} receptor occupancy in schizophrenic patients. *Am J Psychiatry* 156(6):869-875.
238. Odagaki Y. and R. Toyoshima. **2006**. Dopamine D₂ receptor-mediated G protein activation assessed by agonist-stimulated [35S]guanosine 5'-O-(gamma-thiotriphosphate) binding in rat striatal membranes. *Prog Neuropsychopharmacol Biol Psychiatry* 30(7):1304-1312.
239. Offord S.J., G.A. Ordway and A. Frazer. **1988**. Application of [125I]iodocyanopindolol to measure 5-hydroxytryptamine_{1B} receptors in the brain of the rat. *J Pharmacol Exp Ther* 244(1):144-153.
240. Ohba H., N. Harada, S. Nishiyama, T. Kakiuchi and H. Tsukada. **2009**. Ketamine/xylazine anesthesia alters [11C]MNPA binding to dopamine D₂ receptors and response to methamphetamine challenge in monkey brain. *Synapse* 63(6):534-537.
241. Oldham W.M. and H.E. Hamm. **2007**. How do receptors activate G proteins? *Adv Protein Chem* 74:67-93.
242. Oldham W.M. and H.E. Hamm. **2008**. Heterotrimeric G protein activation by G-protein-coupled receptors. *Nat Rev Mol Cell Biol* 9(1):60-71.
243. Olivier B., J. Mos, R. van Oorschot and R. Hen. **1995**. Serotonin receptors and animal models of aggressive behavior. *Pharmacopsychiatry* 28 Suppl 2:80-90.
244. Onoe H., O. Inoue, K. Suzuki, H. Tsukada, T. Itoh, N. Mataga and Y. Watanabe. **1994**. Ketamine increases the striatal N-[11C]methylspiperone binding in vivo: positron emission tomography study using conscious rhesus monkey. *Brain Res* 663(2):191-198.
245. Osman S., C. Lundkvist, V.W. Pike, C. Halldin, J.A. McCarron, C.G. Swahn, L. Farde, N. Ginovart, S.K. Luthra, R.N. Gunn, C.J. Bench, P.A. Sargent and P.M. Grasby. **1998**. Characterisation of the appearance of radioactive metabolites in monkey and human plasma from the 5-HT_{1A} receptor radioligand, [carbonyl-C-11]WAY-100635 - Explanation of high signal contrast in PET and an aid to biomathematical modelling. *Nuclear Medicine and Biology* 25(3):215-223.
246. Otsuka T., H. Ito, C. Halldin, H. Takahashi, H. Takano, R. Arakawa, M. Okumura, F. Kodaka, M. Miyoshi, M. Sekine, C. Seki, R. Nakao, K. Suzuki, S. Finnema, Y. Hirayasu, T. Suhara and L. Farde. **2009**. Quantitative PET analysis of the dopamine D₂ receptor agonist radioligand 11C-(R)-2-CH₃O-N-n-propylnorapomorphine in the human brain. *J Nucl Med* 50(5):703-710.
247. Palner M., P. McCormick, J. Parkes, G.M. Knudsen and A.A. Wilson. **2010**. Systemic catechol-O-methyl transferase inhibition enables the D(1) agonist radiotracer R-[(11)C]SKF 82957. *Nucl Med Biol* 37(7):837-843.
248. Paterson L.M., R.J. Tyacke, D.J. Nutt and G.M. Knudsen. **2010**. Measuring endogenous 5-HT release by emission tomography: promises and pitfalls. *J Cereb Blood Flow Metab* 30(10):1682-1706.
249. Pedigo N.W., H.I. Yamamura and D.L. Nelson. **1981**. Discrimination of multiple [3H]5-hydroxytryptamine binding sites by the neuroleptic spiperone in rat brain. *J Neurochem* 36(1):220-226.
250. Peng T., J. Zysk, P. Dorff, C.S. Elmore, P. Strom, J. Malmquist, M. Ding, D. Tuke, J. Werkheiser, D. Widzowski, L. Mrzljak and D. Maier. **2010**. D₂ receptor occupancy in conscious rat brain is not significantly distinguished with

- [(3)H]-MNPA, [(3)H]-(+)-PHNO, and [(3)H]-raclopride. *Synapse* 64(8):624-633.
251. Pierson M.E., J. Andersson, S. Nyberg, D.J. McCarthy, S.J. Finnema, K. Varnas, A. Takano, P. Karlsson, B. Gulyas, A.M. Medd, C.M. Lee, M.E. Powell, J.R. Heys, W. Potts, N. Seneca, L. Mrzljak, L. Farde and C. Halldin. **2008**. [11C]AZ10419369: a selective 5-HT1B receptor radioligand suitable for positron emission tomography (PET). Characterization in the primate brain. *Neuroimage* 41(3):1075-1085.
 252. Pike V.W., J.A. McCarron, A.A. Lammertsma, S. Osman, S.P. Hume, P.A. Sargent, C.J. Bench, I.A. Cliffe, A. Fletcher and P.M. Grasby. **1996**. Exquisite delineation of 5-HT1A receptors in human brain with PET and [carbonyl-11 C]WAY-100635. *Eur J Pharmacol* 301(1-3):R5-7.
 253. Pike V.W. **2009**. PET radiotracers: crossing the blood-brain barrier and surviving metabolism. *Trends Pharmacol Sci* 30(8):431-440.
 254. Pinborg L.H., K.H. Adams, S. Yndgaard, S.G. Hasselbalch, S. Holm, H. Kristiansen, O.B. Paulson and G.M. Knudsen. **2004**. [18F]altanserin binding to human 5HT2A receptors is unaltered after citalopram and pindolol challenge. *J Cereb Blood Flow Metab* 24(9):1037-1045.
 255. Portas C.M., B. Bjorvatn and R. Ursin. **2000**. Serotonin and the sleep/wake cycle: special emphasis on microdialysis studies. *Prog Neurobiol* 60(1):13-35.
 256. Praschak-Rieder N., D. Hussey, A.A. Wilson, A. Carella, M. Lee, E. Dunn, M. Willeit, R.M. Bagby, S. Houle and J.H. Meyer. **2004**. Tryptophan depletion and serotonin loss in selective serotonin reuptake inhibitor-treated depression: an [(18)F] MPPF positron emission tomography study. *Biol Psychiatry* 56(8):587-591.
 257. Praschak-Rieder N., A.A. Wilson, D. Hussey, A. Carella, C. Wei, N. Ginovart, M.J. Schwarz, J. Zach, S. Houle and J.H. Meyer. **2005**. Effects of tryptophan depletion on the serotonin transporter in healthy humans. *Biol Psychiatry* 58(10):825-830.
 258. Psylla M., I. Gunther, A. Antonini, P. Vontobel, H.W. Reist, A. Zollinger and K.L. Leenders. **1997**. Cerebral 6-[18F]fluoro-L-DOPA uptake in rhesus monkey: pharmacological influence of aromatic amino acid decarboxylase (AAAD) and catechol-O-methyltransferase (COMT) inhibition. *Brain Res* 767(1):45-54.
 259. Rabiner E.A., C. Messa, P.A. Sargent, K. Husted-Kjaer, A. Montgomery, A.D. Lawrence, C.J. Bench, R.N. Gunn, P. Cowen and P.M. Grasby. **2002**. A database of [(11)C]WAY-100635 binding to 5-HT(1A) receptors in normal male volunteers: normative data and relationship to methodological, demographic, physiological, and behavioral variables. *Neuroimage* 15(3):620-632.
 260. Rabiner E.A., M. Slifstein, J. Nobrega, C. Plisson, M. Huiban, R. Raymond, M. Diwan, A.A. Wilson, P. McCormick, G. Gentile, R.N. Gunn and M.A. Laruelle. **2009**. In vivo quantification of regional dopamine-D3 receptor binding potential of (+)-PHNO: Studies in non-human primates and transgenic mice. *Synapse* 63(9):782-793.
 261. Rajagopal S., K. Rajagopal and R.J. Lefkowitz. **2010**. Teaching old receptors new tricks: biasing seven-transmembrane receptors. *Nat Rev Drug Discov* 9(5):373-386.
 262. Ramboz S., F. Saudou, D.A. Amara, C. Belzung, L. Segu, R. Misslin, M.C. Buhot and R. Hen. **1996**. 5-HT1B receptor knock out--behavioral consequences. *Behav Brain Res* 73(1-2):305-312.
 263. Ravert H.T., A.A. Wilson, R.F. Dannals, D.F. Wong and H.N. Wagner, Jr. **1987**. Radiosynthesis of a selective dopamine D-1 receptor antagonist: R(+)-7-chloro-8-hydroxy-3-[11C]methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine ([11C]SCH 23390). *Int J Rad Appl Instrum A* 38(4):305-306.
 264. Raymond J.R., Y.V. Mukhin, A. Gelasco, J. Turner, G. Collinsworth, T.W. Gettys, J.S. Grewal and M.N. Garnovskaya. **2001**. Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacol Ther* 92(2-3):179-212.

265. Rbah L., V. Leviel and L. Zimmer. **2003**. Displacement of the PET ligand 18F-MPPF by the electrically evoked serotonin release in the rat hippocampus. *Synapse* 49(4):239-245.
266. Repapetilo. 2010. Cartoon depicting the GPCR-G-protein activation/deactivation cycle. In: cycle.jpg G, editor. Adobe Illustrator CS3.
267. Riad M., S. Garcia, K.C. Watkins, N. Jodoin, E. Doucet, X. Langlois, S. el Mestikawy, M. Hamon and L. Descarries. **2000**. Somatodendritic localization of 5-HT1A and preterminal axonal localization of 5-HT1B serotonin receptors in adult rat brain. *J Comp Neurol* 417(2):181-194.
268. Riad M., L. Zimmer, L. Rbah, K.C. Watkins, M. Hamon and L. Descarries. **2004**. Acute treatment with the antidepressant fluoxetine internalizes 5-HT1A autoreceptors and reduces the in vivo binding of the PET radioligand [18F]MPPF in the nucleus raphe dorsalis of rat. *J Neurosci* 24(23):5420-5426.
269. Riccardi P., R. Li, M.S. Ansari, D. Zald, S. Park, B. Dawant, S. Anderson, M. Doop, N. Woodward, E. Schoenberg, D. Schmidt, R. Baldwin and R. Kessler. **2006**. Amphetamine-induced displacement of [18F] fallypride in striatum and extrastriatal regions in humans. *Neuropsychopharmacology* 31(5):1016-1026.
270. Rice O.V., S.J. Gatley, J. Shen, C.L. Huemmer, R. Rogoz, O.T. DeJesus, N.D. Volkow and A.N. Gifford. **2001**. Effects of endogenous neurotransmitters on the in vivo binding of dopamine and 5-HT radiotracers in mice. *Neuropsychopharmacology* 25(5):679-689.
271. Richfield E.K., J.B. Penney and A.B. Young. **1989**. Anatomical and affinity state comparisons between dopamine D1 and D2 receptors in the rat central nervous system. *Neuroscience* 30(3):767-777.
272. Rinne J.O., A. Laihininen, U.K. Rinne, K. Nagren, J. Bergman and U. Ruotsalainen. **1993**. PET study on striatal dopamine D2 receptor changes during the progression of early Parkinson's disease. *Mov Disord* 8(2):134-138.
273. Risinger F.O., N.M. Bormann and R.A. Oakes. **1996**. Reduced sensitivity to ethanol reward, but not ethanol aversion, in mice lacking 5-HT1B receptors. *Alcohol Clin Exp Res* 20(8):1401-1405.
274. Rocha B.A., K. Searce-Levie, J.J. Lucas, N. Hiroi, N. Castanon, J.C. Crabbe, E.J. Nestler and R. Hen. **1998**. Increased vulnerability to cocaine in mice lacking the serotonin-1B receptor. *Nature* 393(6681):175-178.
275. Rodbell M. **1997**. The complex regulation of receptor-coupled G-proteins. *Adv Enzyme Regul* 37:427-435.
276. Rosenbaum D.M., S.G. Rasmussen and B.K. Kobilka. **2009**. The structure and function of G-protein-coupled receptors. *Nature* 459(7245):356-363.
277. Rothman R.B. and M.H. Baumann. **2009**. Serotonergic drugs and valvular heart disease. *Expert Opin Drug Saf* 8(3):317-329.
278. Rudnick G. and J. Clark. **1993**. From synapse to vesicle: the reuptake and storage of biogenic amine neurotransmitters. *Biochim Biophys Acta* 1144(3):249-263.
279. Ruf B.M. and Z. Bhagwagar. **2009**. The 5-HT1B receptor: a novel target for the pathophysiology of depression. *Curr Drug Targets* 10(11):1118-1138.
280. Ruottinen H.M. and U.K. Rinne. **1996**. A double-blind pharmacokinetic and clinical dose-response study of entacapone as an adjuvant to levodopa therapy in advanced Parkinson's disease. *Clin Neuropharmacol* 19(4):283-296.
281. Samama P., S. Cotecchia, T. Costa and R.J. Lefkowitz. **1993**. A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model. *J Biol Chem* 268(7):4625-4636.
282. Sanchez-Gonzalez M.A., M.A. Garcia-Cabezas, B. Rico and C. Cavada. **2005**. The primate thalamus is a key target for brain dopamine. *J Neurosci* 25(26):6076-6083.
283. Sanci V., S. Houle and J.N. DaSilva. **2002**. No change in dopamine D1 receptor in vivo binding in rats after sub-chronic haloperidol treatment. *Can J Physiol Pharmacol* 80(1):36-41.
284. Sandell J., O. Langer, P. Larsen, F. Dolle, F. Vaufrey, C. Demphel, C. Crouzel and C. Halldin. **2000**. Improved specific radioactivity of the PET radioligand [11C]FLB457 by use of the GE medical systems PETtrace MeI MicroLab. *J Labelled Compd Radiopharm* 43:331-338.

285. Sari Y., K. Lefevre, M. Bancila, M. Quignon, M.C. Miquel, X. Langlois, M. Hamon and D. Verge. **1997**. Light and electron microscopic immunocytochemical visualization of 5-HT_{1B} receptors in the rat brain. *Brain Res* 760(1-2):281-286.
286. Sari Y., M.C. Miquel, M.J. Brisorgueil, G. Ruiz, E. Doucet, M. Hamon and D. Verge. **1999**. Cellular and subcellular localization of 5-hydroxytryptamine_{1B} receptors in the rat central nervous system: immunocytochemical, autoradiographic and lesion studies. *Neuroscience* 88(3):899-915.
287. Saudou F., D.A. Amara, A. Dierich, M. LeMeur, S. Ramboz, L. Segu, M.C. Buhot and R. Hen. **1994**. Enhanced aggressive behavior in mice lacking 5-HT_{1B} receptor. *Science* 265(5180):1875-1878.
288. Saunders R.C., B.S. Kolachana and D.R. Weinberger. **1994**. Local pharmacological manipulation of extracellular dopamine levels in the dorsolateral prefrontal cortex and caudate nucleus in the rhesus monkey: an in vivo microdialysis study. *Exp Brain Res* 98(1):44-52.
289. Searle G., J.D. Beaver, R.A. Comley, M. Bani, A. Tziortzi, M. Slifstein, M. Mugnaini, C. Griffante, A.A. Wilson, E. Merlo-Pich, S. Houle, R. Gunn, E.A. Rabiner and M. Laruelle. **2010**. Imaging dopamine D₃ receptors in the human brain with positron emission tomography, [¹¹C]PHNO, and a selective D₃ receptor antagonist. *Biol Psychiatry* 68(4):392-399.
290. Seeman P., H.C. Guan and H.B. Niznik. **1989**. Endogenous dopamine lowers the dopamine D₂ receptor density as measured by [³H]raclopride: implications for positron emission tomography of the human brain. *Synapse* 3(1):96-97.
291. Seeman P. and S. Kapur. **2003**. Anesthetics inhibit high-affinity states of dopamine D₂ and other G-linked receptors. *Synapse* 50(1):35-40.
292. Seeman P., F. Ko and T. Tallero. **2005**. Dopamine receptor contribution to the action of PCP, LSD and ketamine psychotomimetics. *Mol Psychiatry* 10(9):877-883.
293. Seeman P., F. Ko, M. Willeit, P. McCormick and N. Ginovart. **2005**. Antiparkinson concentrations of pramipexole and PHNO occupy dopamine D₂(high) and D₃(high) receptors. *Synapse* 58(2):122-128.
294. Seeman P., J. Schwarz, J.F. Chen, H. Szechtman, M. Perreault, G.S. McKnight, J.C. Roder, R. Quirion, P. Boksa, L.K. Srivastava, K. Yanai, D. Weinschenker and T. Sumiyoshi. **2006**. Psychosis pathways converge via D₂high dopamine receptors. *Synapse* 60(4):319-346.
295. Seeman P. **2007**. Antiparkinson therapeutic potencies correlate with their affinities at dopamine D₂(High) receptors. *Synapse* 61(12):1013-1018.
296. Seeman P. **2010**. All Roads to Schizophrenia Lead to Dopamine Supersensitivity and Elevated Dopamine D₂ Receptors. *CNS Neurosci Ther* 10.1111/j.1755-5949.2010.00162.x.
297. Seneca N., M. Skinbjerg, S.S. Zoghbi, J.S. Liow, R.L. Gladding, J. Hong, P. Kannan, E. Tuan, D.R. Sibley, C. Halldin, V.W. Pike and R.B. Innis. **2008**. Kinetic brain analysis and whole-body imaging in monkey of [¹¹C]MNPA: a dopamine agonist radioligand. *Synapse* 62(9):700-709.
298. Shiue C.Y., G.G. Shiue, P.D. Mozley, M.P. Kung, Z.P. Zhuang, H.J. Kim and H.F. Kung. **1997**. P-[¹⁸F]-MPPF: a potential radioligand for PET studies of 5-HT_{1A} receptors in humans. *Synapse* 25(2):147-154.
299. Shotbolt P., A. Tziortzi, S. Miller, G. Searle, J. Van der Aart, S. Abanades, C. Plisson, M. Huiban, G. Searle, J. Beaver, R. Gunn, M. Laruelle and E.A. Rabiner. **2010**. Within-subject comparison of the sensitivity of [¹¹C]-(+)-PHNO and [¹¹C]raclopride to amphetamine induced changes in endogenous dopamine in healthy human volunteers. *Neuroimage* 52:S42-S43.
300. Sibley D.R., A. De Lean and I. Creese. **1982**. Anterior pituitary dopamine receptors. Demonstration of interconvertible high and low affinity states of the D-2 dopamine receptor. *J Biol Chem* 257(11):6351-6361.
301. Sibon I., C. Benkelfat, P. Gravel, N. Aznavour, N. Costes, S. Mzengeza, L. Booij, G. Baker, J.P. Soucy, L. Zimmer and L. Descarries. **2008**. Decreased [¹⁸F]MPPF binding potential in the dorsal raphe nucleus after a single oral dose of fluoxetine: a positron-emission tomography study in healthy volunteers. *Biol Psychiatry* 63(12):1135-1140.

302. Sit S.Y., K. Xie, S. Jacutin-Porte, M.T. Taber, A.G. Gulwadi, C.D. Korpinen, K.D. Burris, T.F. Molski, E. Ryan, C. Xu, H. Wong, J. Zhu, S. Krishnananthan, Q. Gao, T. Verdoorn and G. Johnson. **2002**. (+)-Dinapsoline: an efficient synthesis and pharmacological profile of a novel dopamine agonist. *J Med Chem* 45(17):3660-3668.
303. Skinbjerg M., J.S. Liow, N. Seneca, J. Hong, S. Lu, A. Thorsell, M. Heilig, V.W. Pike, C. Halldin, D.R. Sibley and R.B. Innis. **2010**. D2 dopamine receptor internalization prolongs the decrease of radioligand binding after amphetamine: a PET study in a receptor internalization-deficient mouse model. *Neuroimage* 50(4):1402-1407.
304. Skinbjerg M., N. Seneca, J.S. Liow, J. Hong, D. Weinschenker, V.W. Pike, C. Halldin, D.R. Sibley and R.B. Innis. **2010**. Dopamine beta-hydroxylase-deficient mice have normal densities of D(2) dopamine receptors in the high-affinity state based on in vivo PET imaging and in vitro radioligand binding. *Synapse* 64(9):699-703.
305. Smith G.S., R. Schloesser, J.D. Brodie, S.L. Dewey, J. Logan, S.A. Vitkun, P. Simkowitz, A. Hurley, T. Cooper, N.D. Volkow and R. Cancro. **1998**. Glutamate modulation of dopamine measured in vivo with positron emission tomography (PET) and ¹¹C-raclopride in normal human subjects. *Neuropsychopharmacology* 18(1):18-25.
306. Staley J.K., C.H. Van Dyck, P.Z. Tan, M. Al Tikriti, Q. Ramsby, H. Klump, C. Ng, P. Garg, R. Soufer, R.M. Baldwin and R.B. Innis. **2001**. Comparison of [(18F)]altanserin and [(18F)]deuteroaltanserin for PET imaging of serotonin(2A) receptors in baboon brain: pharmacological studies. *Nucl Med Biol* 28(3):271-279.
307. Starke K., M. Gothert and H. Kilbinger. **1989**. Modulation of neurotransmitter release by presynaptic autoreceptors. *Physiol Rev* 69(3):864-989.
308. Steiger C., S.J. Finnema, L. Raus, M. Schou, R. Nakao, K. Suzuki, V.W. Pike, H.V. Wikstrom and C. Halldin. **2009**. A two-step one-pot radiosynthesis of the potent dopamine D2/D3 agonist PET radioligand [¹¹C]MNPA. *J Label Compd Radiopharm* 52:158-165.
309. Strange P.G. **1993**. New insights into dopamine receptors in the central nervous system. *Neurochem Int* 22(3):223-236.
310. Sulzer D., M.S. Sonders, N.W. Poulsen and A. Galli. **2005**. Mechanisms of neurotransmitter release by amphetamines: a review. *Prog Neurobiol* 75(6):406-433.
311. Sun W., N. Ginovart, F. Ko, P. Seeman and S. Kapur. **2003**. In vivo evidence for dopamine-mediated internalization of D2-receptors after amphetamine: differential findings with [³H]raclopride versus [³H]spiperone. *Mol Pharmacol* 63(2):456-462.
312. Sureau F.C., A.J. Reader, C. Comtat, C. Leroy, M.J. Ribeiro, I. Buvat and R. Trebossen. **2008**. Impact of image-space resolution modeling for studies with the high-resolution research tomograph. *J Nucl Med* 49(6):1000-1008.
313. Talbot P.S., W.G. Frankle, D.R. Hwang, Y. Huang, R.F. Suckow, M. Slifstein, A. Abi-Dargham and M. Laruelle. **2005**. Effects of reduced endogenous 5-HT on the in vivo binding of the serotonin transporter radioligand ¹¹C-DASB in healthy humans. *Synapse* 55(3):164-175.
314. Tang C., D.M. Tomkins, V. Sanci, S. Houle and J.N. DaSilva. **2000**. Chronic ethanol increases binding of dopamine D₁ agonist R-[¹¹C]SKF 82957 in vivo in rat brain. *Soc Neurosci Abstr* 26:786.
315. Tokunaga M., N. Seneca, R.M. Shin, J. Maeda, S. Obayashi, T. Okauchi, Y. Nagai, M.R. Zhang, R. Nakao, H. Ito, R.B. Innis, C. Halldin, K. Suzuki, M. Higuchi and T. Suhara. **2009**. Neuroimaging and physiological evidence for involvement of glutamatergic transmission in regulation of the striatal dopaminergic system. *J Neurosci* 29(6):1887-1896.
316. Tomasi G., S.F. Lin, M. Zheng, B. Planeta-Wilson, S. Henry, N. Goebel, J. Tauscher, H. Huang, A. Neumeister and R. Carson. **2010**. Modeling analysis of the new kappa opioid receptor antagonist tracer [¹¹C]LY2795050 in humans. *Neuroimage* 52:S39-S39.

317. Tork I. **1990**. Anatomy of the serotonergic system. *Ann N Y Acad Sci* 600:9-34; discussion 34-35.
318. Trogadis J.E., G.Y. Ng, B.F. O'Dowd, S.R. George and J.K. Stevens. **1995**. Dopamine D1 receptor distribution in Sf9 cells imaged by confocal microscopy: a quantitative evaluation. *J Histochem Cytochem* 43(5):497-506.
319. Tsukada H., S. Nishiyama, T. Kakiuchi, H. Ohba, K. Sato, N. Harada and S. Nakanishi. **1999**. Isoflurane anesthesia enhances the inhibitory effects of cocaine and GBR12909 on dopamine transporter: PET studies in combination with microdialysis in the monkey brain. *Brain Res* 849(1-2):85-96.
320. Tsukada H., N. Harada, S. Nishiyama, H. Ohba, K. Sato, D. Fukumoto and T. Kakiuchi. **2000**. Ketamine decreased striatal [(11)C]raclopride binding with no alterations in static dopamine concentrations in the striatal extracellular fluid in the monkey brain: multiparametric PET studies combined with microdialysis analysis. *Synapse* 37(2):95-103.
321. Tsukada H., K. Miyasato, T. Kakiuchi, S. Nishiyama, N. Harada and E.F. Domino. **2002**. Comparative effects of methamphetamine and nicotine on the striatal [(11)C]raclopride binding in unanesthetized monkeys. *Synapse* 45(4):207-212.
322. Tsukada H., H. Ohba, S. Nishiyama and T. Kakiuchi. **2010**. Differential effects of stress on [(11)C]raclopride and [(11)C]MNPA binding to striatal D(2)/D(3) dopamine receptors: A PET study in conscious monkeys. *Synapse* 65(1):84-89.
323. Tsukada H., H. Ohba, S. Nishiyama and T. Kakiuchi. **2011**. Differential effects of stress on [(1)(1)C]raclopride and [(1)(1)C]MNPA binding to striatal D/D dopamine receptors: a PET study in conscious monkeys. *Synapse* 65(1):84-89.
324. Twarog B.M. and I.H. Page. **1953**. Serotonin content of some mammalian tissues and urine and a method for its determination. *Am J Physiol* 175(1):157-161.
325. Tziortzi A.C., G.E. Searle, S. Tzimopoulou, C. Salinas, J.D. Beaver, M. Jenkinson, M. Laruelle, E.A. Rabiner and R.N. Gunn. **2010**. Imaging dopamine receptors in humans with [11C]-(+)-PHNO: dissection of D3 signal and anatomy. *Neuroimage* 54(1):264-277.
326. Udo de Haes J.I., F.J. Bosker, A. Van Waarde, J. Pruijm, A.T. Willemsen, W. Vaalburg and J.A. Den Boer. **2002**. 5-HT(1A) receptor imaging in the human brain: effect of tryptophan depletion and infusion on [(18)F]MPPF binding. *Synapse* 46(2):108-115.
327. Udo de Haes J.I., T.I. Cremers, F.J. Bosker, F. Postema, T.D. Tiemersma-Wegman and J.A. den Boer. **2005**. Effect of increased serotonin levels on [18F]MPPF binding in rat brain: fenfluramine vs the combination of citalopram and ketanserin. *Neuropsychopharmacology* 30(9):1624-1631.
328. Udo de Haes J.I., N. Harada, P.H. Elsinga, R.P. Maguire and H. Tsukada. **2006**. Effect of fenfluramine-induced increases in serotonin release on [18F]MPPF binding: a continuous infusion PET study in conscious monkeys. *Synapse* 59(1):18-26.
329. Usiello A., J.H. Baik, F. Rouge-Pont, R. Picetti, A. Dierich, M. LeMeur, P.V. Piazza and E. Borrelli. **2000**. Distinct functions of the two isoforms of dopamine D2 receptors. *Nature* 408(6809):199-203.
330. Vallone D., R. Picetti and E. Borrelli. **2000**. Structure and function of dopamine receptors. *Neurosci Biobehav Rev* 24(1):125-132.
331. Varnas K., H. Hall, P. Bonaventure and G. Sedvall. **2001**. Autoradiographic mapping of 5-HT(1B) and 5-HT(1D) receptors in the post mortem human brain using [(3)H]GR 125743. *Brain Res* 915(1):47-57.
332. Varnas K., C. Halldin and H. Hall. **2004**. Autoradiographic distribution of serotonin transporters and receptor subtypes in human brain. *Hum Brain Mapp* 22(3):246-260.
333. Varnas K., S. Nyberg, C. Halldin, A. Varrone, A. Takano, P. Karlsson, J. Andersson, D. McCarthy, M. Smith, M.E. Pierson, J. Soderstrom and L. Farde. **2010**. Quantitative analysis of [(11)C]AZ10419369 binding to 5-HT(1B) receptors in human brain. *J Cereb Blood Flow Metab*.
334. Varrone A., N. Sjöholm, L. Eriksson, B. Gulyas, C. Halldin and L. Farde. **2009**. Advancement in PET quantification using 3D-OP-OSEM point spread function

- reconstruction with the HRRT. *Eur J Nucl Med Mol Imaging* 36(10):1639-1650.
335. Veltman D.J., H.G. Ruhe and J. Booij. **2010**. Investigating serotonergic function using positron emission tomography: overview and recent findings. *Curr Pharm Des* 16(18):1979-1989.
336. Verma A. and B. Moghaddam. **1996**. NMDA receptor antagonists impair prefrontal cortex function as assessed via spatial delayed alternation performance in rats: modulation by dopamine. *J Neurosci* 16(1):373-379.
337. Villalon C.M., D. Centurion, L.F. Valdivia, P. de Vries and P.R. Saxena. **2003**. Migraine: pathophysiology, pharmacology, treatment and future trends. *Curr Vasc Pharmacol* 1(1):71-84.
338. Violin J.D. and R.J. Lefkowitz. **2007**. Beta-arrestin-biased ligands at seven-transmembrane receptors. *Trends Pharmacol Sci* 28(8):416-422.
339. Volkow N.D., J.S. Fowler, G.J. Wang, S.L. Dewey, D. Schlyer, R. MacGregor, J. Logan, D. Alexoff, C. Shea, R. Hitzemann and et al. **1993**. Reproducibility of repeated measures of carbon-11-raclopride binding in the human brain. *J Nucl Med* 34(4):609-613.
340. Vollenweider F.X., P. Vontobel, I. Oye, D. Hell and K.L. Leenders. **2000**. Effects of (S)-ketamine on striatal dopamine: a [¹¹C]raclopride PET study of a model psychosis in humans. *J Psychiatr Res* 34(1):35-43.
341. Wagner H.N., Jr., H.D. Burns, R.F. Dannals, D.F. Wong, B. Langstrom, T. Duelfer, J.J. Frost, H.T. Ravert, J.M. Links, S.B. Rosenbloom, S.E. Lukas, A.V. Kramer and M.J. Kuhar. **1983**. Imaging dopamine receptors in the human brain by positron tomography. *Science* 221(4617):1264-1266.
342. Wei J.W. and P.V. Sulakhe. **1979**. Agonist-antagonist interactions with rat atrial muscarinic cholinergic receptor sites: differential regulation by guanine nucleotides. *Eur J Pharmacol* 58(1):91-92.
343. Wienhard K., M. Dahlbom, L. Eriksson, C. Michel, T. Bruckbauer, U. Pietrzyk and W.D. Heiss. **1994**. The ECAT EXACT HR: performance of a new high resolution positron scanner. *J Comput Assist Tomogr* 18(1):110-118.
344. Willeit M., N. Ginovart, S. Kapur, S. Houle, D. Hussey, P. Seeman and A.A. Wilson. **2006**. High-affinity states of human brain dopamine D2/3 receptors imaged by the agonist [¹¹C]-(+)-PHNO. *Biol Psychiatry* 59(5):389-394.
345. Willeit M., N. Ginovart, A. Graff, P. Rusjan, I. Vitcu, S. Houle, P. Seeman, A.A. Wilson and S. Kapur. **2008**. First human evidence of d-amphetamine induced displacement of a D2/3 agonist radioligand: A [¹¹C]-(+)-PHNO positron emission tomography study. *Neuropsychopharmacology* 33(2):279-289.
346. Wilson A.A., P. McCormick, S. Kapur, M. Willeit, A. Garcia, D. Hussey, S. Houle, P. Seeman and N. Ginovart. **2005**. Radiosynthesis and evaluation of [¹¹C]-(+)-4-propyl-3,4,4a,5,6,10b-hexahydro-2H-naphtho[1,2-b][1,4]oxazin-9-ol as a potential radiotracer for in vivo imaging of the dopamine D2 high-affinity state with positron emission tomography. *J Med Chem* 48(12):4153-4160.
347. Wong D.F., H.N. Wagner, Jr., L.E. Tune, R.F. Dannals, G.D. Pearlson, J.M. Links, C.A. Tamminga, E.P. Broussolle, H.T. Ravert, A.A. Wilson, J.K. Toung, J. Malat, J.A. Williams, L.A. O'Tuama, S.H. Snyder, M.J. Kuhar and A. Gjedde. **1986**. Positron emission tomography reveals elevated D2 dopamine receptors in drug-naïve schizophrenics. *Science* 234(4783):1558-1563.
348. Yamamoto S., H. Onoe, H. Tsukada and Y. Watanabe. **2007**. Effects of increased endogenous serotonin on the in vivo binding of [¹¹C]DASB to serotonin transporters in conscious monkey brain. *Synapse* 61(9):724-731.
349. Yatham L.N., P.F. Liddle, I.S. Shiah, R.W. Lam, M.J. Adam, A.P. Zis and T.J. Ruth. **2001**. Effects of rapid tryptophan depletion on brain 5-HT(2) receptors: a PET study. *Br J Psychiatry* 178:448-453.
350. Zheng M.Q., N. Nabulsi, G. Tomasi, C. Mitch, S. Quimby, V. Barth, K. Rash, J. Masters, A. Navarro, E. Seest, R.E. Carson and Y.Y. Huang. **2010**. Synthesis and evaluation of [¹¹C]LY2795050, an antagonist PET imaging tracer for the kappa opioid receptors. *Neuroimage* 52:S120-S120.

351. Zimmer L., G. Mauger, D. Le Bars, G. Bonmarchand, A. Luxen and J.F. Pujol. **2002**. Effect of endogenous serotonin on the binding of the 5-HT1A PET ligand 18F-MPPF in the rat hippocampus: kinetic beta measurements combined with microdialysis. *J Neurochem* 80(2):278-286.
352. Zimmer L., L. Rbah, F. Giacomelli, D. Le Bars and B. Renaud. **2003**. A reduced extracellular serotonin level increases the 5-HT1A PET ligand 18F-MPPF binding in the rat hippocampus. *J Nucl Med* 44(9):1495-1501.
353. Zimmer L., M. Riad, L. Rbah, A. Belkacem-Kahlouli, D. Le Bars, B. Renaud and L. Descarries. **2004**. Toward brain imaging of serotonin 5-HT1A autoreceptor internalization. *Neuroimage* 22(3):1421-1426.